


Journal of
Pakistan Association of Dermatologists

JPAD

www.jp.ad.com.pk

SPECIAL FOCUS;

Tinea corporis caused by *microsporum canis* in children with HIV



A Publication of

PAKISTAN ASSOCIATION OF DERMATOLOGISTS

About the Journal

Editor-in-Chief

Prof. Ijaz Hussain

Editor

Prof. Zahida Rani

Associate Editors

Prof. Faria Asad

Prof. Shehla Shaukat

Assistant Editors

Dr. Ghazala Butt
Dr. Bushra Bashir
Dr. Sadia Jabeen
Dr. Rutaba Kiran

Dr. Mahwish Zahoor
Dr. Umara Siddique
Dr. Wajieha Saeed

Dr. Amina Afzal
Dr. Zareen Saqib
Dr. Nabeela Shehzadi

JPAD, the official journal of Pakistan Association of Dermatologists is published quarterly, four issues per volume and one volume per year (ISSN 1560-9014, print; ISSN 2413-7650, online). The journal is recognized by Pakistan Medical Commission, Higher Education Commission of Pakistan and is indexed in College of Physicians and Surgeons Pakistan MEDLIP; Ulrich's International Periodical Directory, USA; ExtraMED, London; EMBASE/Excerpta Medica, PakMediNet, The Netherlands; and Index Medicus, WHO Alexandria, Egypt.

Vol. 33 No. 1 (2023): January-March

Published: 2023-02-05

Editorial

Emergency of skin diseases during flood

Mamoona Ghafoor

1-2

PDF

Original Articles

Evaluation of methotrexate and cyclosporine for severe psoriasis: A retrospective analysis from tertiary hospital in Indonesia

Menul Ayu Umborowati, Made Putri Hendaria, Sylvia Anggraeni, Irmadita Citrashanty, Damayanti, Afif Nurul Hidayati, Anang Endaryanto, Ingrid S Surono, Cita Rosita Sigit Prakoeswa

3-8

PDF

The picture of cutaneous tuberculosis today: New diverse clinical manifestations in the absence of the old lupus vulgaris in a series of 72 patients

Khalifa E. Sharquie, Raed I. Jabbar

9-17

PDF

The The risk of occupational contact dermatitis in COVID-19 isolation ward setting

Damayanti, Sylvia Anggraeni, Menul Ayu Umborowati, Andre Yuindartanto, Sawitri, Astindari, Cita Rosita Sigit Prakoeswa

18-23

PDF

Wistar Rat as Photoaging Mouse Model

Damayanti, Cita Rosita Sigit Prakoeswa, Djoko Agus Purwanto, Anang Endaryanto, Muhammad Yulianto Listiawan, Yohanes Widodo Wirohadjoyo, Soetjipto, Siswandono, Budi Utomo

24-29

PDF

Characteristics of patch test patients in Yogyakarta, Indonesia, 2011-2020 Study Period

Agnes Rosarina Prita Sari, Sri Awalia Febriana, Fajar Waskito, Hardyanto Soebono, Niken Indrastuti
30-36

[PDF](#)

Clinico-epidemiological study of vitiligo in a tertiary care hospital of Eastern India

Jayanta Kumar Barua, Kaushiki Hazra, Prabal Samanta, Param Satsangi, Tirtharaj Saha, Professor Saswati Halder
37-42

[PDF](#)

Vitamin D deficiency in patients with pemphigus vulgaris: A cross sectional study

Amara Amin, Naima Luqman, Sumiya Ashraf, Uzma Almas
43-48

[PDF](#)

Clinico-Etiological Description of Annular Skin Eruptions

Basman Medhat Fadheel, Noor J. Rashid
49-58

[PDF](#)

Evaluation of quality of life in patients with Mycosis fungoides in Isfahan in 2017-2019

Farahnaz Fatemi Naeini, Fatemeh Mohaghegh, Farshad Riahi, Fatemeh Mokhtari, Neda Adibi
59-64

[PDF](#)

The spectrum of alopecia areata: Combination treatment with oral azathioprine and I/M injection triamcinolone acetonide as a bridging strategy

Attiya Tareen, Irfan Ullah, Muzna Gillani, Ammara Safdar, Habiba Yaqub Khan, Hadia Yaqub Khan
65-71

[PDF](#)

Triple therapy of Kaposi's sarcoma using oral propranolol, oral zinc sulfate and oral acyclovir as new initiative study

Khalifa E. Sharquie, Raed I. Jabbar
72-77

[PDF](#)

Comparison of Mean Serum Prolactin Levels between Healthy and Psoriatic Patients

Sabahat Shehzadi, Ghazala Butt, Wajieha Saeed, Huma Saleem, Anam Mehmood, Ijaz Hussain
78-83

PDF

Dermatology quality of life and psychological morbidity in patients with chronic dermatophytosis

Anushree LS, Anupama YG, Praveena C, Dadapeer HJ

84-90

PDF

Prevalence of elevated testosterone levels among female patients of acne vulgaris

Naila Tabassum, Hira Tariq, Saelah Batool, Zahra Arooba, Azam Bukhari, Shahbaz Aman

91-95

PDF

Alcoholism is still a cause of pellagra in sunny countries

Fatema Aljaralla, Khalifa E. Sharquie, Salwa F. Taha

96-100

PDF

Relationship between severity of psoriasis vulgaris based on Psoriasis Area And Severity Index (PASI) scores and depression

Elok Nurfaiqoh, Annisa Marsha Evanti, Pratiwi Prasetya Primisawitri, Muhamad Eko Irawanto

101-107

PDF

Bacteria causing community acquired superficial skin infections in tertiary care hospitals and their antibiotics susceptibility patterns

Aiman Waheed, Irfanullah Afridi, Attiya Tareen, Mehran Khan, Dawood Khan

108-115

PDF

Adverse cutaneous reactions post Covid-19 vaccination- Case series from a tertiary care centre in Eastern India

Nisha Agrawal, Kaushiki Hajra, Uddalak Chakraborty, Saswati Halder

116-122

PDF

Skin appendageal tumors: A three-year dermatopathologic institutional study in Martin, Slovakia

Vladimír Bartoš

123-131

PDF

Comparison of efficacy of oral ivermectin and topical permethrin 5% in the treatment of scabies

Maimona Ilyas, Atif Shahzad, Saadiya Siddiqui; Mohsina Awais, Maryam Qayyum; Nadia Ali Azafar

132-136

[PDF](#)

Immunoregulatory factors (IRF3 and IRF7) serum levels and its relation to disease activity in Vitiligo patients

Maha Anani, Omayma Hassan, Fadia Attia , Moustafa Eyada , Ghada Mohammed

137-141

[PDF](#)

Clinico-sociodemographic Profile of Leprosy Neuropathy in Medan, Indonesia

Dina Arwina Dalimunthe, Duma Wenty Irene Sinambela, Syahril Rahmat Lubis

142-148

[PDF](#)

Therapeutic Efficacy and Safety of Fractional Carbon dioxide Laser versus Q-Switched Nd:YAG 1064 nm Laser in the treatment of Melasma- A Comparative Interventional Study

Kiran Irfan Ali, Aisha Ghias, Wajieha Saeed, Zaima Sohail Chaudhry, Kainat Zaffar Laghari, Zaib, Shahbaz Aman

149-156

[PDF](#)

Association between stress and acne among patients visiting THQ Gujar Khan, Rawalpindi

Bazala Ali, Saman Waqar, Choudhry Tahir Ali, Mehwish Ali

157-163

[PDF](#)

Awareness regarding sun exposure and use of sunscreen amongst general population in a local hospital setting

Syeda Muzna Gillani, Attiya Tareen, Mehreen Baig; Nimrah Bukhari; Rabia Mehreen, Anum Zafar

165-171

[PDF](#)

Analysis of clinico-etiological pattern of adverse cutaneous drug reactions

Warda Orf Rida Naeem, Hira Tariq, Saelah Batool, Shahbaz Aman

172-177

[PDF](#)

A The role of body mass index and age on pelvic floor muscle strength

Nita Damayanti; Devi Artami Susetiati, Adissa Tiara Yulinvia, Satiti Retno Pudjiati
178-183

PDF

Understanding Vitiligo as a Psychosocial Dilemma

Salman Haseeb, Hira Tariq, Saelah Batool, Natasha Rehman, Sumara Rashid, Faria Asad, Shahbaz Aman, Atif Majeed
184-189

PDF

Comparison of amitriptyline and carbamazepine effectiveness in the treatment of postherpetic neuralgia

Mehran Khan, Naimat Ullah, Irfan Ullah, Dawood Khan, Hurriya Khan
190-195

PDF

Serum vitamin levels in Psoriasis and Acne: Is there a relationship with disease? A Comparative cross sectional study in local population

Attiya Tareen , Sohaib Asghar, Hadia Yaqub Khan, Habiba Yaqub Khan, Ammara Safdar
196-203

PDF

A clinco-epidemiological study and assessment of Life Quality Index among patients with lichen planus at a tertiary care centre in South India

Cinthana Palanichamy Sathianesan, Seethalakshmi Ganga vellaisamy, Kannan Gopalan
204-214

PDF

RISK FACTORS OF ERYTHEMA NODOSUM LEPROSUM IN DERMATOVENEREOLOGY OUTPATIENTS

Nur Rachmat Muliando; Fatimah Fitriani, Rieska Widyaswari, Sesia Pradestine
215-219

PDF

Review Articles

Epidemiology of Occupational Contact Dermatitis (OCD) on health workers in Covid-19

Sawitri, Astindari, Andre Yuindartanto, Cita Rosita Sigit Prakoeswa, Damayanti, Sylvia Anggraeni, Menul Ayu Umborowati, Iskandar Zulkarnain, M. Yulianto Listiawan, Afif Nurul Hidayati

220-234

PDF

Vitamin D and autoimmune dermatoses: A review

Paulus Mario Christopher, Muljani Enggalhardjo

235-243

[PDF](#)

Issues related to topical immunotherapy for the treatment of warts: A review of the literature

Ghada Farouk Mohammed, Mohammed Saleh Al-Dhubaibi, Saleh Salem Bahaj, Ahmed Ibrahim AbdElneam

244-250

[PDF](#)

The role of diet in chronic urticaria

Siti Efrida Fiqnasyani, Suci Widhiati, Harijono Kariosentono, Muhammad Eko Irawanto, Triasari Oktavriana

251-264

[PDF](#)

Post-burn pruritus: A brief review

Sajad Ahmad Salati

265-274

[PDF](#)

The role of epigenetics in immunosuppression in patients with infectious diseases as a fascinating applied observation

Inas K. Sharquie, Khalifa E. Sharquie

275-282

[PDF](#)

Case Reports

Gynecomastia caused by efavirenz: Two cases with different severity

Sri Mariyani, Yulia Eka Irmawati, Satiti Retno Pudjiati, Kristiana Etnawati

283-288

[PDF](#)

Intrahepatic cholestasis of pregnancy in a chronic kidney disease patient: A case report

Sri Mariyani, Yulia Eka Irmawati, Kristiana Etnawati, Agnes Sri Siswati

289-293

[PDF](#)

Diffuse Large B-Cell Lymphoma with secondary cutaneous involvement

Praneet P. Awake, Anand Bhosale , Rajan Bindu , Sayali Deshmukh , Pallavi Chaudhari , Vivek Dugad
294-297

PDF

Challenge in diagnosis and management of periorificial dermatitis: A case report

Danny Surya, Sri Linuwih Menaldi, Sandra Widaty
298-301

PDF

Canaliform Median Raphe cyst: A rare variant of a rare diagnosis

Vedika Barman, Manjunath P
302-304

PDF

Coinfection of gonorrhea and chlamydia trachomatis: A case report

Anita Indah Sari, Idrianti Idrus, Safruddin Amin, Muji Iswanty, Anni Adriani, Muhlis
305-309

PDF

Successful treatment of three cases of pediatric alopecia areata using a combination of corticosteroids, minoxidil, immunomodulator and light-emitting diode therapy

Eyleny Meisyah Fitri, Lili Legiawati, Irma Bernadette S. Sitohang, Lis Surachmiati Suseno
310-316

PDF

Severe ichthyosis vulgaris with vitamin D insufficiency and multiple flexion contractures

Bobby Febrianto, Suci Widhiati, Endra Yustin Ellistasari, Indah Julianto, Annang Giri Moelyo, Adhitya Indra Pradhana
317-321

PDF

Multiple trichoepithelioma: A case report

Sebastian Hendry Wibowo, Sthefanie Mariska Gaghana, Pieter Levinus Suling, Shienty Gaspersz, Thigita Aga Pandaleke
322-325

PDF

Steroid-induced rosacea in systemic lupus erythematosus

Windy Keumala Budianti, Nevi Yasnova, Arlha Aporia Debinta, Shafira Anindya
326-330

PDF

A fatal case of Lucio phenomenon in untreated Lucio leprosy

Irwan Saputra Batubara, Melani Marissa, Sri Linuwih Menaldi, Yufanti Sujudi
331-334

PDF

Mid-face Toddler Excoriation Syndrome (MiTES): A new conundrum in the field?

an enigma in sensory neuropathy

Abhineetha Hosthota, Swapna Bondade, Bhavana V, Sanjay Thejesvi
335-338

PDF

Actinic granuloma over photo exposed and covered parts

Bhavya Mani K, Haritha Thiruveedhula, Haritha Samanthula
339-341

PDF

Combination of light-emitting diode with minoxidil 2%, topical corticosteroid and oral immunomodulator induced hair regrowth in a pediatric alopecia areata

Dina Evyana, Lili Legiawati, Githa Rahmayunita, Irma Bernadette S. Sitohang, Lis Surachmiati, Dina Kusumawardhani
342-344

PDF

A neglected pyoderma gangrenosum following COVID-19: A case report

Thamir Kubaisi
345-349

PDF

Erythroderma caused by anti tuberculoid drug in pulmonary tuberculosis and HIV infected patient: A case report

Nugrohoaji Dharmawan; Adniana Nareswari , Fatimah Fitriani
350-353

PDF

Congenital presentation of linear morphea in a 4 year old boy: A rare case report with literature review

Nava Kumar, Kannan Gopalan, Narmadha Arumugam, Seethalakshmi Ganga Vellaisamy, Anbu Lenin Kulandaivel
354-357

PDF

Wistar Rat as Photoaging Mouse Model

Damayanti^{1,2}, Cita Rosita Sigit Prakoeswa², Djoko Agus Purwanto³, Anang Endaryanto⁴, Muhammad Yulianto Listiawan², Yohanes Widodo Wirohadidjoyo⁵, Soetjipto⁶, Siswandono³, Budi Utomo⁷

¹ Doctoral Study Program of Medical Science, Faculty of Medicine, Universitas Airlangga, Indonesia.

² Department of Dermatology and Venereology, Faculty of Medicine, Universitas Airlangga, Indonesia.

³ Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Universitas Airlangga, Indonesia.

⁴ Department of Pediatrics, Faculty of Medicine, Universitas Airlangga, Indonesia.

⁵ Department of Dermatology and Venereology, Faculty of Medicine, Universitas Gajah Mada, Indonesia.

⁶ Department of Biochemistry, Faculty of Medicine, Universitas Airlangga, Indonesia.

⁷ Department of Community Medicine, Faculty of Medicine, Universitas Airlangga, Indonesia.

Abstract

Background The increasing of human life expectancy may cause aging problems, including photoaging. Photoaging raises new problems in community, because its influence in human self-esteem and quality of life. Studies about photoaging have been extensively performed to enhance the basic mechanism of photoaging. Mouse skin is usually used as cutaneous aging model, because it is genetically similar to human skin.

Methods This study was true experimental study, post test only-control group design that was aimed to evaluate the role of ultraviolet (UV) B irradiation to produce photoaging mouse model. Twenty-four Wistar rats were divided randomly allocation into 2 groups, namely non-UV irradiated group and UV irradiated group. Ultraviolet B irradiation were performed for 5 weeks until reaching the total dose 3100 mJ/cm². Dermal collagen density and transepidermal water loss (TEWL) level were evaluated after the intervention.

Results Dermal collagen density in non-UV irradiated group and UV irradiated group were 38.21±2.67% and 27.73±0.77%. TEWL level in non-UV irradiated group and UV irradiated group were 7.55±0.89 g/m²/h and 17.53±3.94 g/m²/h. There was significant difference between the groups in dermal collagen density and TEWL level (p<0.05).

Conclusion Wistar rat in UV irradiated group that were given total dose 3100 mJ/cm² of UV B for 5 weeks has been proven to become photoaging mouse model.

Key words

Photoaging, mouse model, wistar rat, UV B, life expectancy.

Introduction

Photoaging is caused by long term effect of ultraviolet (UV) irradiation. Photoaging and chronological skin aging speed up the occurrence of skin aging with dry skin, lack of elasticity, lack of firmness, and lack of skin smoothness as the clinical manifestation. Photoaging is 80-90% of all skin aging, that can

be prevented. The increasing of human life

Address for correspondence

Prof. Cita Rosita Sigit Prakoeswa, MD, PhD.
Department of Dermatology and Venereology,
Faculty of Medicine Universitas Airlangga/ Dr. Soetomo
General Academic Hospital, Surabaya, Indonesia.
Prof Dr. Moestopo Number 47, Surabaya,
East Java, Indonesia.
Phone: +62-811328199
Email: cita-rosita@fk.unair.ac.id

expectancy may cause aging problems, including photoaging. Photoaging raises new problems in community, because its influence in human self-esteem and quality of life.¹⁻³

Studies about photoaging have been extensively performed to enhance the basic mechanism of photoaging and to develop its prevention. Several studies were done in *in vivo* study using mouse model. Mouse model in photoaging studies is used as cutaneous aging model, because mouse skin is genetically similar to human skin.⁴

Hwang *et al.* showed that collagen synthesis velocity in the skin of 24 months-old mouse decreased 10 times, comparing with 1 month-old mouse. Meanwhile, collagen degradation velocity in mouse model increased by the age. Collagen degradation velocity was 6.4% in 1 month-old mouse and 56% in 24 months-old mouse.⁴

Photon energy from ultraviolet B was 1000 times higher than ultraviolet A. Ultraviolet B is the main cause of DNA destruction and inflammation (activation of prostaglandin (PG) via activation of cyclooxygenase-2 (COX-2)). Ultraviolet B induce the production of reactive oxygen species (ROS), that activate keratinocyte to express transcription factor, namely Nuclear Factor kappa B (NF-κB) and Activator Protein-1 (AP-1). NF-κB and AP-1 induce the activation of matrix-metalloproteinase (MMP), that cause collagen degradation cascade.^{1,2,5-7}

Movement of water from stratum corneum to the atmosphere is called transepidermal water loss (TEWL). The definition of TEWL is water content per skin surface area unit and per duration of time which loss from the skin to the atmosphere through diffusion and evaporation, with normal value 0-20 g/m²/h. Skin barrier regulates water evaporation from the skin. The

disruption of skin barrier may cause disruption of these mechanisms. ROS production caused by ultraviolet irradiation may cause the destruction of lipid in stratum corneum, corneocyte proteins, and skin barrier. These process induce inflammatory response by increasing the production of proinflammatory cytokines, which cause photoaging manifestation, showed by the elevation of TEWL level.⁸⁻¹⁰

Methods

This study was true experimental study, post test only-control group design, aimed to prove the role of ultraviolet B irradiation to produce photoaging mouse model. Ultraviolet-irradiated male Wistar rats were used in this study as photoaging animal model. Male Wistar rats (*Rattus norvegicus*) were used to avoid the effect of estrogen level reduction in photoaging. Ten to twelve weeks-old Wistar rats were used in this study to avoid effect of intrinsic skin aging before ultraviolet B irradiation, in order to get photoaging animal model from the only effect of ultraviolet B irradiation.¹¹

Twenty-four Wistar rats were divided randomly allocation into 2 groups, namely non-UV irradiated group and UV irradiated group. All subjects were given the same intake and environment. The fur on dorsal area of the rat was shaved in 3x3 cm² size. No intervention was performed in non-UV irradiated group. Dermal collagen density and transepidermal water loss (TEWL) level were evaluated after intervention.

Ultraviolet irradiation

In UV irradiated group, Wistar rats were performed ultraviolet B irradiation using Ultraviolet B Broadband TL lamps 20W. Ultraviolet B irradiation were performed for 5 weeks until reaching the total dose 3100 mJ/cm². The dose of ultraviolet B were increased



Figure 1 Ten to twelve weeks-old Wistar rats.

Table 1 Dermal collagen density differences between UV irradiated and non-UV irradiated Wistar rat group.

Groups	Dermal collagen density (%)	p
Non-UV irradiated group	38.21±2.67	0.0000
UV irradiated group	27.73±0.77	

gradually from 100 to 400 mJ/cm². On the first week, ultraviolet B irradiation were performed 3 times (100 mJ/cm²/time). On the second and third week, ultraviolet B irradiation were performed 3 times per week (200 mJ/cm²/time). On the fourth and fifth week, ultraviolet B irradiation were performed 2 times per week (400 mJ/cm²/time).^{12,13}

Histopathology examination

The skin on the back part of the male Wistar rats were obtained from sacrifice animal under anesthesia at the end of the study. The skin specimens were fixed in formalin buffer 10% and cryosections were prepared. Masson Trichrome staining is staining technique using acid fuchsin and methyl blue in order to evaluate the dermal collagen density. Dermal collagen will be blue stained. All of these evaluation used

light microscopy Nikon H600L completed with digital camera DS Fi2 300 megapixel and image analyzer software Nikon Image System. Dermal collagen density was evaluated by calibrated Image J software. The evaluation was performed in 200x magnification.¹⁴

TEWL level evaluation

Evaluation of TEWL level in this study was performed with Cutometer MPA 580 and Tewameter[®] TM 300 probe. TEWL level evaluation was performed in 20-22°C room temperature and 40-60% room humidity. The evaluation was performed 3 times in 30 seconds duration in each evaluation.¹⁵⁻¹⁷

Results

The mean of dermal collagen density in non-UV irradiated group and UV irradiated group were 38.21±2.67% and 27.73±0.77%. There was significant difference between the groups (p<0.05) (Table 1). Figure 2 showed histopathology result in Masson Trichrome staining.




Figure 2 Histopathology result in Masson Trichrome staining showing dermal collagen density (200x magnification). A. In non-UV irradiated group. B. In UV irradiated group.

Table 2 TEWL level differences between UV irradiated and non-UV irradiated Wistar rat group.

Groups	TEWL level (g/m ² /h)	p
Non-UV irradiated group	7.55±0.89	0.0324
UV irradiated group	17.53±3.94	

There was high density of dermal collagen in non-UV irradiated group, meanwhile the dermal collagen density in UV irradiated group was low.

The mean of TEWL level in non-UV irradiated group and UV irradiated group were 7.55±0.89 g/m²/h and 17.53±3.94 g/m²/h. There was significant difference between the groups (p<0.05) (Table 2).

Discussion

Photoaging is one of new problems that arise in community due to the increase in life expectancy. Photoaging is the process of skin aging caused by external factors, mainly due to exposure to ultraviolet radiation. Unlike intrinsic aging, photoaging has better respond to interventions and preventive measures. Typical clinical features of photoaging which are mostly caused by ultraviolet radiation include wrinkle, weakness, roughness, increased fragility, and telangiectasia. In addition, the skin will appear darker and have a mottled pigmentation.^{1,2,4}

Sunlight is consist of infrared light, visible light, and ultraviolet (UV) light, with most UV rays are blocked by the earth's atmosphere. UV irradiation that reaches the earth's surface, consists of UVA (320 to 400 nm) in more than 95% and UVB (280 to 320 nm) in about 5%. UVA and UVB contributes to skin aging. UVB, which spreads to the epidermis and upper dermis, is major source of direct DNA damage, inflammation, and immunosuppression. On the other hand, UVA penetrate the skin to the lower

dermis, and considered a greater contributor to skin aging than UVB, due to the greater penetration depth and higher percentage of surface sunlight.^{2,5}

Many studies on photoaging have been carried out to learn about the basic mechanisms of photoaging and to develop agents for the prevention and treatment of photoaging, which often starting with *in vivo* studies, using mouse models.^{1,2,4}

Collagen is the majority structural protein in dermal layer. Two main collagen regulators are transforming growth factor β (TGF β) and AP-1. TGF β increase the synthesis of procollagen and collagen; while AP-1 inhibit synthesis of procollagen and collagen. Ultraviolet induce ROS production, that activate AP-1 and inhibit TGF β . These mechanisms result in decreasing of dermal collagen count.¹⁸⁻²¹

This study showed decreasing of dermal collagen count in group receiving 3100 mJ/cm² ultraviolet B irradiation for 5 weeks, significantly (p<0.05). Study of Kim *et al.* (2015) showed decreasing of dermal collagen density, elevation of JNK and elevation of MMP-9 in hairless mice after receiving ultraviolet irradiation significantly (p<0.05).¹² Ultraviolet B irradiation decrease dermal collagen count via MMP activation, especially MMP-1, that was showed by study of Dong *et al.* (2008).²²

Ultraviolet can increase mitochondrial DNA mutations which will decrease mitochondrial function and the formation of reactive oxygen species (ROS). Ultraviolet can cause damage to the basement membrane at the dermal-epidermal junction. In addition, photoaging can increase matrix metalloproteinase (MMP).^{2,5}

Structural changes in the aging skin layer,

including collagen as the main component of the dermis, increase rigidity and impaired wound healing process. Reduction of collagen type I and III in intrinsic skin aging will be accelerated by the photoaging process. Collagen fragmentation also plays a role in the downregulation of collagen synthesis in intrinsic skin aging and photoaging. Collagen degradation by MMP, increases with the increasing age. MMP-1, MMP-3, and MMP-9 were elevated by the effect of acute UV irradiation, which mainly acts on the epidermis layer. Meanwhile, MMP-1, MMP-2, MMP-3, MMP-9, MMP-11, MMP-17, and MMP-27, which produced by dermal fibroblasts, were elevated in chronically photodamaged skin. This will cause a decrease in the amount of collagen due to increased collagen degradation and decreased collagen production, because fragmented collagen interferes with fibroblast function.^{2,5,21}

Evaluation of TEWL level in this study was performed with Cutometer MPA 580 and Tewameter[®] TM 300 probe. The elevation of TEWL level in photoaging was proved in this study, showed by higher TEWL level in UV irradiated group significantly. Ultraviolet B irradiation induced TEWL level in Wistar rats, which elevated the TEWL level 1.5 times higher than in non-UV irradiated group. This result was similar to studies from Hung *et al.* (2015) and Choi *et al.* (2019), which showed higher TEWL level in ultraviolet B-irradiated hairless mice group ($p < 0.001$).^{15,23}

Skin is the first body protection from environment, protects the body from mechanical destruction, pathogen penetration and ultraviolet irradiation. Skin also play role in homeostatic mechanism in controlling TEWL through stratum corneum. TEWL is water content per skin surface area unit and per duration of time, which loss from the skin to the atmosphere,

through diffusion and evaporation. Water evaporation from the skin is regulated by the skin barrier. The disruption of skin barrier function may cause disruption of water retention and increasing the water evaporation. TEWL level describe the skin barrier function, that was used as the parameter in evaluating photoaging process in this study.^{8,9,24}

Ultraviolet irradiation results in disruption of skin barrier function, caused by ROS production. Disruption of skin barrier function may induce inflammatory response through elevation of proinflammatory cytokines and decreasing of water binding capacity in ultraviolet-exposed area. These mechanisms manifest as dry skin in photoaging, that could be evaluated from elevation of TEWL level.^{10,23}

Conclusion

The total dose 3100 mJ/cm² of ultraviolet B for 5 weeks was able to decrease the dermal collagen density and increase the TEWL level significantly, as the parameters in photoaging in this study, and has been proven to be photoaging mouse model.

Acknowledgement

We gratefully thank Directorate of Research and Community Service - Directorate General of Research and Development-Ministry of Research, Technology and Higher Education (Direktorat Riset dan Pengabdian Masyarakat-Direktorat Jenderal Riset dan Pengembangan-Kementerian Riset, Teknologi dan Pendidikan Tinggi/ Kemenristekdikti) Indonesia for supporting this research.

References

1. Puizina IN. Skin aging. *Acta Dermatovenereol Alp Pannonica Adriat.* 2008;17(21):47-54.

2. Yaar M, Gilchrest BA. Aging of the skin. Fitzpatrick's in General Medicine 8th ed., New York: The McGraw Hill Companies; 2012. p 1213-26.
3. Nabila YA, Damayanti, Handayani H, Setyaningrum T. The Effect of Lifestyle on Skin Aging. *Berkala Ilmu Kesehatan Kulit dan Kelamin*. 2021;**33(2)**:110-5.
4. Hwang KA, Yi BR, Choi KC. Molecular mechanism and in vivo mouse models of skin aging associated with dermal matrix alterations. *Lab Anim Res*. 2011;**27(1)**:1-8.
5. Gonzaga ER. Role of UV light in photodamage, skin aging, and skin cancer. *Am J Clin Dermatol*. 2009;**10(1)**:19-24.
6. Lephart ED. Skin aging and oxidative stress: equol's anti-aging effects via biochemical and molecular mechanism. *Ageing Res Rev*. 2016;**31**:36-54.
7. Ahmad Z, Damayanti. Skin Aging: Pathophysiology and Clinical Manifestation. *Berkala Ilmu Kesehatan Kulit dan Kelamin*. 2018;**30(3)**:208-15.
8. Spencer TS. Transepidermal water loss. Methods for cutaneous investigation, New York: Marcell Dekker; 1990. p 191-217.
9. Schaefer H, Redelmeir TE, Nohynek GJ, Lademann J. Pharmacokinetics and topical applications of drugs. Fitzpatrick's in General Medicine 8th ed., New York: The McGraw Hill Companies; 2012. p 2652-8.
10. Agustyniak AM, Rotsztein H, Bartnicka E, Budzisz. Effects of reactive oxygen species on skin photoaging. *Przegl Dermatol*. 2016;**103**:233-9.
11. Chang KCN, Wang Y, Oh IG, Jenkins S, Freedman LP, Thompson CC *et al*. Estrogen receptor β is a novel therapeutic target for photoaging. *Mol Pharmacol*. 2010;**77(5)**:744-50.
12. Kim YJ, Kim HN, Shin MS, Choi BY. Thread embedding acupuncture inhibits ultraviolet B irradiation-induced skin photoaging in hairless mice. *Evid Based Complement Alternat Med*. 2015;1-9. doi: 10.1115/2015/539172.
13. Kwon S, Jung SH. Topical administration of manuka oil prevents UV-B irradiation-induced cutaneous photoaging in mice. *Evid Based Complement Alternat Med*. 2013;**2013**:1-10. doi: 10.1155/2013/930857.
14. Chen Y, Yu Q, Bao Xu C. A convenient method for quantifying collagen fibers in atherosclerotic lesions by ImageJ software. *Int J Clin Exp Med*. 2017;**10(10)**:14904-10.
15. Choi SI, Jung TD, Cho BY, Choi SH, Sim WS, Han X, *et al*. Anti-photoaging effect of fermented agricultural by-products on ultraviolet B-irradiated hairless mouse skin. *Int J Mol Med*. 2019;**44**:559-68.
16. Plessis J, Stefaniak A, Eloff F, John S, Agner T, Chou TC, *et al*. International guidelines for the in vivo assessment of skin properties in non-clinical settings: part 2. transepidermal water loss and skin hydration. *Skin Res Technol*. 2013;**19(3)**:265-78.
17. Gardien KLM, Baas DC, Vet HCW, Middelkoop E. Transepidermal water loss measured with the Tewameter TM300 in burn scar. *Burns*. 2016;**30**:1-8.
18. Rittie L, Fisher GJ. UV-light-induced signal cascades and skin aging. *Ageing Res Rev*. 2002;**1**:705-20.
19. Chiu HW, Chen CH, Chen YJ, Hsu YH. Far infrared suppresses in skin photoaging in ultraviolet B-exposed fibroblast and hairless mice. *PLoS One*. 2017;**12(3)**:1-15.
20. Poon F, Kang S, Chien AL. Mechanism and treatments of photoaging. *Photodermatol Photoimmunol Photomed*. 2014;**31**:65-74.
21. Pittayapruek P, Meehasan J, Prapapan O, Komine M, Ohtsuki M. Role of matrix metalloproteinases in photoaging and photocarcinogenesis. *Int J Mol Sci*. 2016;**17**:868(1-20).
22. Dong KK, Damaghi N, Picart SD, Markova NG, Obayashi K, Okano Y, *et al*. UV-induced DNA damage initiates release of MMP-1 in human skin. *Exp Dermatol*. 2008;**17**:1037-44.
23. Hung CF, Chen WY, Aljuffali IA, Lin YK, Shih HC, Fang JY. Skin aging modulates percutaneous drug absorption: the impact of ultraviolet irradiation and ovariectomy. *Age*. 2015;**37(21)**:1-14.
24. Lee KO, Kim SN, Kim YC. Anti-wrinkle effect of water extracts of teas in hairless mouse. *Toxicol Res*. 2014;**30(4)**:283-9.