

T1_22_Completeness_of_Prescription_Contributes

by Djazuly Chalidyanto

Submission date: 07-Aug-2023 09:43AM (UTC+0800)


Submission ID: 2142324501

File name: T1_22_Completeness_of_Prescription_Contributes_to.2.pdf (623.56K)

Word count: 3096

Character count: 17226

Completeness of Prescription Contributes to Prescribing Errors in Hospitals: Is this the Doctor's Negligence?

Shely Oktavia Puspita Ningrum¹, Mardhina Ratna Prabasari¹, Djazuly Chalidyanto¹ 

¹Department of Health Administration and Policy, Public Health Faculty, Universitas Airlangga Surabaya, Indonesia

ABSTRACT

Introduction: Incident reporting is the primary key to implement patient safety. One indicator of patient safety is to reduce prescription errors. Errors in writing prescriptions can cause the treatment process to be disrupted and even lead to malpractice and ethical violations. This study aimed to know the effect of individual and organization factors on prescribing errors in the Outpatient Unit of Nganjuk Public Hospital. **Methods:** This research was a cross-sectional observational study in the Outpatient Clinic of Nganjuk State General Hospital. This study used a questionnaire and prescriptions with a total sample of 362 prescriptions from 24 doctors, with exclusion criteria: doctors with an expired license at the time of the study and doctors with <2 years of service. Prescriptions were also used as a sample to assess the incidence of prescribing errors. This research used descriptive analysis with cross-tabulations. **Results:** A doctor's knowledge about writing complete prescriptions is the most related factor to prescribing errors ($n = 0.159$) in individual factors. Likewise, the doctor's perception variable related to prescribing policy has the most substantial relationship compared to other variables on organizational factors ($n = 0.235$). **Conclusions:** Several factors influence prescribing errors, such as the physician's skill and knowledge. Doctors with high subjective and objective workloads, doctors' perceptions regarding the incident reporting system, prescribing policies, and medication management also affect prescribing errors. Training and reduction of doctors' workload could be the solution to reduce prescribing errors.

KEYWORDS: Patient safety, prescribing error, prescribing knowledge

Submitted: 16-Aug-2022
Revised: 19-Oct-2022
Accepted: 29-Oct-2022
Published: 09-Dec-2022

INTRODUCTION

Incident reporting is essential to service quality and prevention.^{1,2} Medication errors occur in four stages: prescribing, transcribing, dispensing, and administration errors.³⁻⁵ Prescribing errors can occur when medication is selected incorrectly, defining is ambiguous, medication names are abbreviated, and patient information is incomplete. Errors in prescription writing can lead to prescription reading errors or transcribing errors.⁶⁻⁸ Based on internal data about the incident report from 2016 to 2019 at Nganjuk Hospital, the type of incident that occurred the most was medication error, with an average percentage of 31.11% and constantly increasing yearly.^{9,10} The incidence of prescribing errors is increasing, so it is necessary to research the factors that increase the incidence of prescribing errors.

Reporting prescribing errors can identify trends and reduce the risk of the reoccurrence of incidents; however, underreporting is common.¹¹⁻¹³ Factors affecting drug administration accuracy are organizational, unit management, and individual factors.^{14,15} The subjective workload is the main latent factor for prescribing errors because it is related to doctors' physical and psychological burdens, so errors tend to occur when writing prescription drugs.^{16,17} Workload correlates with the occurrence of medication errors. The prevention of medication errors can also be done by increasing the awareness of officers regarding the importance of writing a complete prescription.^{18,19}

Address for correspondence: Mrs. Shely Oktavia Puspita Ningrum, Kolibri 17 Street, Malang, East Java Province, Indonesia.
E-mail: shely.oktavia.puspita-2021@fkm.unair.ac.id

Access this article online	
Quick Response Code: 	Website: https://journals.lww.com/bhsj
	DOI: 10.4103/bhsj.bhsj_27_22

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Ningrum SO, Prabasari MR, Chalidyanto D. Completeness of prescription contributes to prescribing errors in hospitals: Is this the doctor's negligence? *Biomol Health Sci J* 2022;5:77-80.

This study aimed to know the effect of individual and organizational factors on prescribing errors in the Outpatient Unit of Nganjuk Public Hospital. Determining these factors is essential to prevent prescribing errors in hospital services.

METHODS

This study was an observational cross-sectional research design. The population in this study was all doctors who provide services in the outpatient clinic with a total sample of 362 prescriptions from 24 doctors, with exclusion criteria: doctors with an expired license at the time of the study and doctors with <2 years of service. The dependent variable of this study was incidents of prescribing errors, namely, errors in writing doses, drug names, drinking rules, patient names, and illegible prescriptions. The independent variable of this study was individual factors and organizational factors. Individual factors, namely, doctor's knowledge, skills, objective workload, and awareness of the need to write a complete prescription. At the same time, organizational factors include incident reporting systems, prescribing policies, and medication management. The scoring for each variable is categorized as high if it is more than equal to the mean value and categorized as low if it is less than the mean. Data analysis was done using descriptive analysis with cross-tabulations. Variables are considered correlated if they have a difference of >20% between categories of independent variable while assessing the strength of the relationship that can be seen from the value of the contingency coefficient between variables. If the contingency coefficient approaches the value of 1, the relationship between variables gets stronger and vice versa. This study has received ethical approval from the Ethics Committee of Nganjuk General Hospital with number 893/02/411.801/2021.

RESULTS

Table 1 shows that the respondent characteristics were mainly female (54.2%). The leading age group

Table 1: Respondent's characteristics

Variable	Description	Frequency (%)
Gender	Male	11 (45.8)
	Female	13 (54.2)
Age	30-50	18 (75.0)
	>50	6 (25.0)
Qualification	General practitioner	2 (8.3)
	Specialist	22 (91.7)
Years of service (years)	≤5	9 (37.5)
	>5	15 (62.5)
Further training	Received	14 (58.3)
	Did not receive	10 (41.7)

was 30–50 years (75%). Most of the respondents are specialists (91.7%). Most respondents had worked for more than 5 years (62.5%). Most respondents received further prescription training after medical school (58.3%).

Table 2 shows that the doctor's knowledge of prescribing error has the highest correlation with the incidence of prescribing error (contingency coefficient = 0.159) compared to other variables on individual factors. The doctor's perception variable related to prescribing policy has the highest correlation with the incidence of prescribing error (contingency coefficient = 0.235) compared to other organizational factors.

DISCUSSION

The knowledge of prescribing error has the highest correlation with the incidence of prescribing error in individual factors. Doctors with the knowledge of prescribing errors and a low objective workload tend to have a high incidence of prescribing errors. Increasing knowledge is a strategic step to achieve zero accidents through awareness.²⁰ The incidence of prescribing errors was closely related to writing prescriptions that did not follow the standard operating procedure. Individuals with insufficient knowledge of prescribing errors tend to make prescribing errors.^{16,21} The main factor causing prescribing errors in hospitals is the insufficient knowledge of doctors regarding prescribing errors.²²⁻²⁴ Individual factors regarding knowledge related to high prescribing errors have been shown to reduce the incidence of prescribing errors.²⁵⁻²⁷

The prescribing policy has the highest level of correlation with the incidence of prescribing error on organizational factors. An excellent reliable reporting system will reduce the occurrence of prescribing errors. Several studies show that hospitals with good reporting systems will have zero accidents in the prescribing error phase. A good reporting system is essential in reducing medication errors.²⁸⁻³¹ In prescribing policies, the aspects assessed are related to the standard operating procedures applied and the factors that hinder the implementation of policies in hospitals.³² Prescribing policy is one of the organizational factors that influence the motivation of doctors in writing prescriptions.³³ The policy's success in motivating doctors refers to the mechanism, human resources, and the relationship between prescribers and policy actors.³⁴ An excellent prescribing policy can reduce prescribing errors.

The results from organizational factors also show that low prescribing policies and medication management tend to have high prescribing errors; this is in line with the previous study, which showed that implementing good policies and medication management positively affects the performance.³⁵

Table 2: Crosstab and coefficient contingency

	Prescribing error incident		Coefficient contingency
	Low, n (%)	High, n (%)	
Individual factors			
Doctor's knowledge			0.159
Low	3 (12.5)	10 (41.7)	
High	7 (29.2)	4 (16.7)	
Doctor's skill			0.070
Low	4 (16.7)	10 (41.7)	
High	6 (25.0)	4 (16.7)	
Doctor's objective workload			
Low	0	0	
High	10 (41.7)	14 (58.3)	
Doctor's subjective workload			
Low	10 (41.7)	14 (58.3)	
High	0	0	
Awareness of the need to write a complete prescription			0.094
Low	5 (20.8)	12 (50.0)	
High	5 (20.8)	2 (8.3)	
Organizational factors			
Incident reporting systems			0.156
Low	7 (29.2)	12 (50.0)	
High	3 (12.5)	12 (50.0)	
Prescribing policies			0.235
Low	4 (16.7)	16 (66.7)	
High	6 (25.0)	4 (16.7)	
Medication management			0.022
Low	4 (16.7)	9 (37.5)	
High	6 (25.0)	5 (20.8)	

In addition, prescribing errors from doctors are fatal errors.³⁶ To avoid these mistakes, formulate a good prescription writing policy that is agreed on by the hospital's internals.³⁷ To increase the capacity to write prescriptions, doctors must be trained. The role of other health workers is also needed to reduce prescribing errors by doctors. Health workers such as nurses, midwives, and pharmacists can help cross-check.^{38,39} Solid teamwork and reminding each other can reduce errors in prescription writing.^{40,41}

In addition, it is necessary to formulate policies to develop electronic prescriptions to reduce errors. Implementing prescription writing with an electronic system is an alternative to reduce errors in prescription writing.⁴² Electronic prescription writing systems are proven to reduce prescription errors.⁴³ Hospitals with a high incidence of prescribing errors should consider using an electronic writing system to reduce prescribing errors due to typographical errors or misinterpretation.

CONCLUSIONS

Several factors influence prescribing errors, such as the physician's skill and knowledge. Doctors with high subjective and objective workloads tend to make prescribing

errors. Doctors' perceptions regarding the incident reporting system, prescribing policies, and medication management also affect prescribing errors. The prescribing error must be approached systematically to minimize error. Training and reduction of doctors' workload could be the solution to reduce prescribing errors.

Limitation

This study only looked at individual and organizational factors. Many other factors, such as workload, must be researched to complete this research.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Zheng S, Tucker AL, Ren ZJ, Heineke J, McLaughlin A, Podell AL. The impact of internal service quality on preventable adverse events in hospitals. *Prod Oper Manag* 2018;27:2201-12.
- Hasanah D, Aira D, Widiastuti F. Hubungan beban kerja perawat dengan penerapan keselamatan pasien di ruang rawat inap rumah sakit umum daerah raden mattaheer. *J Manaj Terap dan Keuang* 2018;7:68-78.
- Riaz MK, Riaz M, Latif A. Review – Medication errors and

- strategies for their prevention. *Pak J Pharm Sci* 2017;30:921-8.
4. Goedecke T, Ord K, Newbould V, Brosch S, Arlett P. Medication errors: New EU good practice guide on risk minimisation and error prevention. *Drug Saf* 2016;39:491-500.
 5. Kementerian Kesehatan Republik Indonesia. Peraturan Menteri Kesehatan Republik Indonesia Nomor 72 tahun 2016 Tentang Standar Pelayanan Kefarmasian di Rumah Sakit. Jakarta: Kementerian Kesehatan RI; 2017.
 6. Dyasanoor S, Urooge A. Insight into quality of prescription writing – An institutional study. *J Clin Diagn Res* 2016;10:C61-4.
 7. Batta A, Singh B. Rational approach to prescription writing: A preview. *Neurol India* 2018;66:928-33.
 8. Sheikh D, Mateti UV, Kabekkodu S, Sanal T. Assessment of medication errors and adherence to WHO prescription writing guidelines in a tertiary care hospital. *Futur J Pharm Sci* 2017;3:60-4.
 9. Regency Regional Financial and Asset Management Agency of Nganjuk Regency. The Government Agencies Performance Accountability Report of Nganjuk Regency at 2019. 2019;62.
 10. Patient safety committee. Nganjuk Hospital patient safety incident report in 2019. Nganjuk-Indonesia; 2019.
 11. Burns N, Alkaisy Z, Sharp E. Doctors attitudes towards medication errors at 2002 & 2015. *Int J Health Care Qual Assur* 2018;31:451-63.
 12. Holmström AR. Learning from Medication Errors in Healthcare: How to Make Medication Error Reporting Systems Work? Vol. 119. Helsinki: Health Policy; 2015.
 13. Bullen K, Hall N, Sherwood J, Wake N, Donovan G. Prescribing error reporting in primary care: A narrative synthesis systematic review. *Integr Healthc J* 2020;2:e000026.
 14. Lin JC, Lee TT, Mills ME. Evaluation of a barcode medication administration information system. *Comput Inform Nurs* 2018;36:596-602.
 15. Bayang AT, Pasinringi S. Faktor penyebab Medication Error di RSUD Anwar Makkatutu Kabupaten Bantaeng. Causes Factors of Medical Errors Regional General Hospital of Anwar Makkatutu Bantaeng Regency. Makassar: E-Journal Program Pascasarjana Universitas Hasanuddin; 2018. p. 49-57, 2018.
 16. Sutherland A, Ashcroft DM, Phipps DL. Exploring the human factors of prescribing errors in paediatric intensive care units. *Arch Dis Child* 2019;104:588-95.
 17. Lyell D, Magrabi F, Coiera E. The effect of cognitive load and task complexity on automation bias in electronic prescribing. *Hum Factors* 2018;60:1008-21.
 18. Mohanty S. Awareness of medication error, medication management and prevention among staff nurses in IMS & sum hospital, Odisha. *J Heal Allied Sci NU* 2016;06:18-22.
 19. Guan X, Ni B, Zhang J, Zhu D, Cai Z, Meng W, et al. Association between physicians' workload and prescribing quality in one tertiary hospital in China. *J Patient Saf* 2021;17:e1860-5.
 20. Abdel-Latif MM. Knowledge of healthcare professionals about medication errors in hospitals. *J Basic Clin Pharm* 2016;7:87-92.
 21. Lavan AH, Gallagher PF, O'Mahony D. Methods to reduce prescribing errors in elderly patients with multimorbidity. *Clin Interv Aging* 2016;11:857-66.
 22. Handayani TW. Analisis faktor penyebab medication error di RS Undata Kota Palu. *Farmakol J Farm* 2018;15:168-75.
 23. Farzi S, Irajpour A, Saghaei M, Ravaghi H. Causes of medication errors in intensive care units from the perspective of healthcare professionals. *J Res Pharm Pract* 2017;6:158-65.
 24. Alharbi AI, Gay V, AlGhamdi MJ, Alturki R, Alyamani HJ. Towards an application helping to minimize medication error rate. *Mob Inf Syst* 2021;2021.
 25. Greenwood S. Human factors and safe prescribing. *J Prescr Pract* 2019;1:290-5.
 26. Angkow L, Citraningtyas G, Wiyono WI. Faktor penyebab medication error Di Instalasi Gawat Darurat (Igd) rumah sakit bhayangkara Tk.Iii Manado. *Pharmacon* 2019;8:426.
 27. Gloria L, Yuwono N. Analisis faktor yang mempengaruhi medication error pada pasien kemoterapi Di RSUP DR. Mohammad hoesin palembang. *Maj Kedokt Sriwij* 2017;4:178-84.
 28. Dwi B, Weny WI, Marina M. Pengetahuan dan Sikap Tenaga Kesehatan terhadap Pelaporan dan Sistem Pelaporan Medication Error di RSU Monompia GMIBM Kota Kotamobagu. *Pharmacon* 2019;8:360-70.
 29. Howell AM, Burns EM, Hull L, Mayer E, Sevdalis N, Darzi A. International recommendations for national patient safety incident reporting systems: An expert Delphi consensus-building process. *BMJ Qual Saf* 2017;26:150-63.
 30. Mutair AA, Alhumaid S, Shamsan A, Zaidi AR, Mohaini MA, Al Mutairi A, et al. The effective strategies to avoid medication errors and improving reporting systems. *Medicines (Basel)* 2021;8:46.
 31. Karande S, Marraro GA, Spada C. Minimizing medical errors to improve patient safety: An essential mission ahead. *J Postgrad Med* 2021;67:1-3.
 32. Thiels CA, Ubl DS, Yost KJ, Dowdy SC, Mabry TM, Gazelka HM, et al. Results of a prospective, multicenter initiative aimed at developing opioid-prescribing guidelines after surgery. *Ann Surg* 2018;268:457-68.
 33. Weiner SG, Baker O, Poon SJ, Rodgers AF, Garner C, Nelson LS, et al. The effect of opioid prescribing guidelines on prescriptions by emergency physicians in Ohio. *Ann Emerg Med* 2017;70:799-808.e1.
 34. Lobo F, Río-Álvarez I. Barriers to biosimilar prescribing incentives in the context of clinical governance in Spain. *Pharmaceuticals (Basel)* 2021;14:283.
 35. Purba IE, Rumapea P, Kiyai B. Pengaruh implementasi kebijakan standar operasional prosedur terhadap kinerja pegawai pada sekretariat kota manado. *J Adm Publik UNSRAT* 2014;4:1245.
 36. Brits H, Botha A, Niksch L, Venter K, Terblanché R, Joubert G. Illegible handwriting and other prescription errors on prescriptions at National District Hospital, Bloemfontein. *Prof Nurs Today* 2017;21:53-6.
 37. Shrestha R, Prajapati S. Assessment of prescription pattern and prescription error in outpatient department at tertiary care district hospital, Central Nepal. *J Pharm Policy Pract* 2019;12:16.
 38. Latimer S, Hewitt J, Stanbrough R, McAndrew R. Reducing medication errors: Teaching strategies that increase nursing students' awareness of medication errors and their prevention. Vol. 52. Amsterdam: Nurse Education Today; 2017. 52: p. 7-9.
 39. Sarfati L, Ranchon F, Vantard N, Schwierzt V, Larbre V, Parat S, et al. Human-simulation-based learning to prevent medication error: A systematic review. *J Eval Clin Pract* 2019;25:11-20.
 40. Hassan I. Avoiding medication errors through effective communication in healthcare environment. *Mov Heal Exerc* 2018;7:113-26.
 41. Billstein-Leber M, Carrillo CJ, Cassano AT, Moline K, Robertson JJ. ASHP Guidelines on preventing medication errors in hospitals. *Am J Health Syst Pharm* 2018;75:1493-517.
 42. Wheeler AJ, Scahill S, Hopcroft D, Stapleton H. Reducing medication errors at transitions of care is everyone's business. *Aust Prescr* 2018;41:73-7.
 43. Samadbeik M, Ahmadi M, Sadoughi F, Garavand A. A comparative review of electronic prescription systems: Lessons learned from developed countries. *J Res Pharm Pract* 2017;6:3-11.

T1_22_Completeness_of_Prescription_Contributes

ORIGINALITY REPORT

15%

SIMILARITY INDEX

13%

INTERNET SOURCES

8%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	www.deepdyve.com Internet Source	3%
2	journals.lww.com Internet Source	3%
3	Submitted to Udayana University Student Paper	2%
4	www.researchgate.net Internet Source	2%
5	www.ijhonline.org Internet Source	1%
6	ijehe.mui.ac.ir Internet Source	1%
7	www.ijfmt.com Internet Source	1%
8	Mirna Jayustin Tanjung, Ermi Girsang, Linda Chiuman, Chrismis Novalinda Ginting, Putranto Manalu. "Implementation of Patient Safety in Hospitals: A Qualitative Study", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2021 Publication	1%

9	ejournal.unsri.ac.id Internet Source	1 %
10	ihj.bmj.com Internet Source	1 %
11	Nokky Farra Fazria, Inge Dhamanti. "Knowledge And Attitude Towards Medication Safety: Study From Religion Based Hospital", Jurnal Kesehatan Prima, 2023 Publication	<1 %
12	ir.ymlib.yonsei.ac.kr Internet Source	<1 %
13	wprim.whocc.org.cn Internet Source	<1 %
14	pure.rug.nl Internet Source	<1 %
15	Kathryn Bullen, Nicola Hall, John Sherwood, Nicola Wake, Gemma Donovan. "Prescribing error reporting in primary care: a narrative synthesis systematic review", Integrated Healthcare Journal, 2020 Publication	<1 %

Exclude quotes Off
Exclude bibliography On

Exclude matches < 5 words