

In Silico Analysis of Pongamia pinnata to Inhibit Neuronal Apoptosis after Ischemic Stroke via NMDAR and Caspase-3

by Muhammad Ja'far Shodiq

Submission date: 19-Aug-2023 05:42PM (UTC+0800)

Submission ID: 2147913985

File name: ._In_Silico_Analysis_of_Pongamia_pinnata_to_Inhibit_Neuronal.pdf (705.64K)

Word count: 3356

Character count: 18556



Volume 3 Number 2, July 2023

In Silico Analysis of *Pongamia pinnata* to Inhibit Neuronal Apoptosis after Ischemic Stroke via NMDAR and Caspase-3

Muhammad Ja'far Shodiq¹, Farmindo Hartono¹, Siti Khaerunnisa², Abdulloh Machin³

¹ Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

² Department of Physiology and Medical Biochemistry, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

³ Department of Neurology, Faculty of Medicine, Universitas Airlangga; Dr. Soeotomo General Academic Hospital, Surabaya, Indonesia

Article info

Article History: 22

Received Apr 10, 2023

Revised Jun 2, 2023

Accepted Jun 12, 2023

Published Jul 31, 2023

Keywords:

Cardiovascular disease

In silico study

Ischemic stroke

Neuronal apoptosis

Pongamia pinnata

ABSTRACT

Introduction: One of the cardiovascular diseases with the highest mortality rate is stroke. Stroke is the second-leading cause of death worldwide. Each year, 12.2 million new cases of stroke occur, of which 7.6 million are ischemic strokes. In ischemic stroke, there are several pathways that cause neuronal apoptosis. The activity of NMDAR and caspase-3 is one of the pathways. *Pongamia pinnata* phytochemicals have a neuroprotective function against neurological disorders. However, its use as an inhibitor of apoptosis in ischemic stroke has never been evaluated before. **Objective:** This research was designed to evaluate the phytochemicals of *Pongamia pinnata* as inhibitors of neuronal apoptosis in ischemic stroke using an in silico study. **Methods:** This study used four main phytochemicals of *Pongamia pinnata*, namely Karanjin, Karanjachromene, Pongapin, and Pongachromene. The protein targets for neuronal apoptosis were NMDAR and caspase-3. The molecular docking processes were ligand preparation, protein preparation, grid box determination, molecular docking, and visualized molecular docking. **Results:** In silico results showed that at NMDAR target proteins, Karanjin, Karanjachromene, Pongapin, and Pongachromene have binding energies of -5.12, -5.83, -5.03, and -5.13 kcal/mol. At protein targets, Caspase-3, Karanjin, Karanjachromene, Pongapin, and Pongachromene have binding energies of -4.87, -4.98, -4.88, and -5.08 kcal/mol. **Conclusion:** The phytochemicals of *Pongamia pinnata* have the potential to inhibit neuronal apoptosis via NMDAR and caspase-3 in ischemic stroke. The binding of Karanjachromene to NMDAR demonstrated the compound's best interaction.

Corresponding Author

Siti Khaerunnisa

Department of Physiology and Medical Biochemistry, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

email: st.khaerunnisa@fk.unair.ac.id

Available at <https://e-journal.unair.ac.id/index.php/aksona>



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License

INTRODUCTION

Stroke is one of the cardiovascular diseases with the highest mortality rate. Stroke is the second-leading cause of death in the world. Strokes cause 6.5 million fatalities worldwide. In a year, there are 12.2 million new cases of stroke, of which 7.6 million are ischemic strokes. In ischemic stroke, there are several pathways that cause neuronal apoptosis.¹ One of the pathways is the activity of N-methyl-D-aspartate receptors (NMDAR), the main receptor that plays a role in the intrinsic apoptotic pathway. NMDAR activation causes excess Ca²⁺ influx and triggers calpain activation. Calpain converts the Bcl-2 interacting domain (BID) into a tBID that interacts with Bax. Bax forms homooligomers and then incorporates them into the outer mitochondrial membrane. Consequently, the pores of the mitochondrial permeability transition (mPTP) open and allow the release of apoptogen (Cytc). The released cytc combines with Apaf-1 and pro-caspase-9 and will gradually activate caspase-3. It is this caspase-3 activation that ultimately induces neuronal apoptosis.²

Pongamia pinnata (*P. pinnata*) is a plant originating from India and Southeast Asia. This species' phytochemical studies led to the identification of several compounds from various classes, including flavonoids and terpenoids. These plant phytochemical compounds exhibit various pharmacological activities such as antioxidant, antimicrobial, antiparasitic, anti-inflammatory, anticonvulsant, antidiabetic, cytotoxic, anthelmintic, insecticidal, and immunomodulatory activity.³ *P. pinnata* extract contains many flavonoid compounds, namely Karanjin, Karanjachromene, Pongapin, and Pongachromene.⁴ These compounds have a neuroprotective function in Alzheimer's disease.⁵ However, its use as an apoptosis inhibitor in ischemic stroke has never been evaluated. Therefore, we carried out this in silico research.

OBJECTIVE

This research was designed to evaluate the main phytochemicals of *Pongamia pinnata*, namely karanjin, karanjachromene, pongapin, and pongachromene, as inhibitors of neuronal apoptosis in ischemic stroke using an in silico study.

METHODS

System Configuration

This study used a 32-bit Windows 10 laptop with an Intel Core i3 processor and 2 GB of RAM. Avogadro, AutoDock 4.2, Biovia Discovery Studio 2019, and PyMol were applications employed in this in silico research.

Ligand Preparation

P. pinnata has the main phytochemical compounds, namely Karanjin, Karanjachromene, Pongapin, and Pongachromene (Figure 1).³ The phytochemical compounds of *P. pinnata* were analyzed using the Swiss-Adme website to predict Lipinski's parameters. The compounds were downloaded from NCBI PubChem in.sdf format. All the phytochemicals were optimized using Avogadro software. Every compound's energy was minimized using the MM4 force field. After that, the polar hydrogen was added, the nonpolar hydrogen was merged, and the gasteiger charge was computed using AutoDock 4.2.

Protein Preparation

NMDAR (PDB ID: 5H8Q) and caspase-3 (PDB ID: 3DEI) were chosen as the protein targets in this study because they are key players in the pathophysiology of neuronal apoptosis following ischemic stroke in patients.² The protein target structure was downloaded from the Protein Data Bank (<http://www.rcsb.org>) in.pdb format. NMDAR has two chains, namely chains A and B. We used chain B for molecular docking because the native ligand 5YE is in chain B. Caspase-3 has four chains, namely chains A, B, C, and D. We used chain C for molecular docking because the native ligand RXB is in chain C. The proteins were then optimized by including polar hydrogens, merging nonpolar ones, and adding Kollman charges. The 3D structures of NMDAR and caspase-3 with their native ligands are shown in Figure 2.

Grid Box Determination

Using AutoDock 4.2 software, the native ligand position determined the grid box. In NMDAR, the native ligand grid 5YE was set at 30x30x30 (XYZ) point size, -14.959, -14.301, -25.150 center coordinates, and 0.375 Å spacing. In the caspase-3 protein, the native ligand RXB grid was set at 30x30x30 (XYZ) point size, -46.620, 15.373, -22.195 center coordinates, and 0.375 Å spacing.

Molecular Docking

Molecular docking was performed using AutoDock 4.2 software after proteins and ligands were prepared and grid boxes were defined. The ligand conformation was analyzed using the Lamarckian Genetic Algorithm. The docking parameters were left as defaults. A favorable conformation was selected based on the lowest energy binding (ΔG) and inhibitory constant (K_i). After that, the docking results were visualized using PyMol software and BIOVIA Discover Studio 2019.⁶



RESULTS

P. pinnata has around 70 phytochemical compounds. The main phytochemicals of *P. pinnata* were karanjin, karanjachromene, pongapin, and pongachromene. All these phytochemicals can be found in all parts of *P. pinnata*. All the main phytochemical compounds were screened before docking. All phytochemicals met Lipinski's parameters with no infractions (Table 1). All ligands have the ability to pass the blood-brain barrier to neuronal apoptosis inhibitory receptors in the brain.

Based on the result of molecular docking, karanjachromene has the potential to inhibit neuronal apoptosis via NMDAR inhibitors. Karanjachromene has the strongest binding energy of the other ligand and the native ligand 5YE with NMDAR. Karanjachromene has a binding energy of -5.83 kcal/mol, an inhibition constant of 53.64 μ M, and amino acid bonds with ILE128, PRO141, LYS143, TYR144, SER249, GLY250, and HIS273. While the native ligand 5YE has a binding energy of -5.50 kcal/mol with an inhibition constant of 93.10 μ M. The other ligands have inhibitory activity against NMDAR. It is shown by the negative energy binding value. Karanjachromene (ΔG -5.12 kcal/mol and K_i 176.24 μ M), Pongapin (ΔG -5.03 kcal/mol and K_i 206.90 μ M), and Pongachromene (ΔG -5.13 kcal/mol and K_i 174.27 μ M) have a lower relative energy binding value than the native ligand 5YE (Table 2). The molecular docking visualization of *P. pinnata* phytochemical compounds with NMDAR is shown in Figure 3.

At the caspase-3 receptor, the phytochemical of *P. pinnata* that has the strongest binding energy is Pongachromene, with a binding energy of -5.08 kcal/mol, an inhibition constant of 189.85 μ M, and an amino acid bond with THR166, GLU167, LEU168, CYS170, TYR204, THR255, PHE256, and LYS259. The other ligands, Karanjachromene, Karanjachromene, and Pongapin, have binding energies of -4.87, -4.98, and -4.88 kcal/mol with inhibition constants of 269.40, 223.26, and 266.27 μ M. Meanwhile, the native ligand RXB has a binding energy of -5.82 kcal/mol with inhibition constant of 54.30 μ M (Table 2). The molecular docking visualization of *P. pinnata* phytochemical compounds with caspase-3 is shown in Figure 4.

DISCUSSION

Prior to molecular docking, Lipinski's parameters screened the phytochemical compounds of *P. pinnata*. Lipinski's parameters include molecular weight, log P, H-bond donor, and H-bond acceptor.⁷ All the phytochemicals of *P. pinnata* met Lipinski's parameters. According to the molecular docking results, all the

phytochemicals of *P. pinnata* have negative bonds with NMDAR and caspase-3. Through NMDAR and caspase-3, these phytochemicals can inhibit neuronal apoptosis in ischemic stroke. Karanjachromene inhibits NMDAR better than other ligands and native ligands due to its best binding interaction. Stronger ligand-receptor interactions result from ligands with greater negative binding energy, and those ligands are more effective in inhibiting the receptors.⁸

GluN1 and GluN2 subunits combine to form NMDAR (PDB ID: 5H8Q), a glutamate-gated ion channel. In the CNS, NMDAR functions in both neuronal survival and neuronal death.⁹ NMDAR plays an important role in synaptic plasticity and initiates cellular responses in the CNS, such as learning, brain development, and memory. However, NMDAR also has a harmful side for neurons. When the brain is in ischemia or hypoxia, there is a rapid increase in glutamate levels in the ischemic area, which can destroy neurons.^{10,11} 5H8Q has a structure weight of 65.92 kDa, a modeled residue count of 557, a deposited residue count of 578, an atom count of 4786, two unique protein chains, and a native ligand 5YE that binds to LYS140, PRO141, PHE142, LYS143, TYR144, ARG248, SER249, GLY250, LEU270, and HIS273.¹² Caspase-3 (PDB ID: 3DEI) is a cysteine protease that plays a critical role in human apoptotic cell death. Caspase-3 is a key mediator of neuronal death in the acute stage of ischemic stroke because it involves the final common pathway of apoptosis. Caspase-3 is a major executioner caspase, and it will trigger when it is activated by other caspases.^{13,14,15} 3DEI has a structure weight of 114.63 kDa, 933 modeled residue, 966 deposited residue, 7672 atom count, 1 unique protein chain, and a native ligand RXB that binds to THR166, GLU167, LEU168, CYS170, TYR204, THR255, and LYS259.¹⁶

Karanjin is a bioactive furoflavanoid. This phytochemical was first isolated from *P. pinnata*. Karanjachromene can be isolated from the seed, flower, root, leaf, and stem bark of *P. pinnata*.^{3,17,18} Karanjachromene has multiple health benefits, including anti-diabetic, anti-cancer, antioxidant, anti-colitis, anti-ulcer, gastroprotective, anti-inflammation, and antibacterial. Karanjachromene is a phytochemical with anti-Alzheimer activity and is also a neuroprotective agent.^{5,17,19} Karanjachromene is a bioactive chromenoflavone. Pongachromene is another name for Karanjachromene. This phytochemical is isolated from *P. pinnata* seed.^{3,20} Karanjachromene has some pharmacological activities, such as antibacterial, antioxidant, and anti-aging. However, research on this phytochemical is still limited.^{3,21} Pongapin is a bioactive furanoflavone isolated from the root bark, seed, and stem bark of *P. pinnata*.^{3,22} This phytochemical has not yet been evaluated. According to a recent study, pongapin has anticancer and antidiabetic activity via alpha-glucoside



inhibitors.^{23,24,25} Pongachromene is a bioactive chromene flavone. The first chromene flavone from *P. pinnata* to be reported was this phytochemical. Pongachromene was isolated from the root and stem of *P. pinnata*.^{24,26} The pharmacological activities of pongachromene are still limited. This phytochemical has been reported as antibacterial, antidiabetic, and antioxidant.^{24,27}

P. pinnata with its main phytochemical compounds Karanjin, Karanjachromene, Pongapin, and Pongachromene, has the neuroprotective potential to inhibit neuronal apoptosis after ischemic stroke. It is shown from the molecular docking of the compounds that they can inhibit NMDAR and caspase-3. Through this mechanism, neuronal death can be prevented in ischemic stroke patients.²

CONCLUSION

Molecular docking of *P. pinnata* compounds has the potential to inhibit neuronal apoptosis via NMDAR and caspase-3 in ischemic stroke. The binding of Karanjachromene to NMDAR with a binding energy of -5.83 demonstrated the best interaction of the compound. This bond is slightly stronger than the native ligand to NMDAR (ΔG : -5.50). More *in vitro* and *in vivo* studies are needed to prove the potency of *P. pinnata* in inhibiting neuronal apoptosis after ischemic stroke, especially the Karanjachromene compound in inhibiting NMDAR.

Acknowledgment

We thank to the Department of Physiology and Medical Biochemistry, and the Department of Neurology, Faculty of Medicine, Universitas Airlangga, Indonesia.

Conflict of Interest

All authors have no conflict of interest in this article

Funding

No funding was received for this study

Author Contribution

The first and second authors designed the concept, collected and analyzed the data, and wrote the manuscript. The third and fourth authors provided independent consultations for the review and revision of the manuscript.

REFERENCES

1. Feigin VL, Brainin M, Norrving B, Martins S, Sacco RL, Hacke W, et al. World stroke organization (WSO): Global stroke fact sheet 2022. *Int J Stroke*. 2022;17(1):18–29.
2. Mao R, Zong N, Hu Y, Chen Y, Xu Y. Neuronal death mechanisms and therapeutic strategy in ischemic stroke. *Neurosci Bull*. 2022;38(10):1229–47.
3. Al Muqarrabun LMR, Ahmat N, Ruzaina SAS, Ismail NH, Sahidin I. Medicinal uses, phytochemistry and pharmacology of *Pongamia pinnata* (L.) Pierre: A review. *J Ethnopharmacol*. 2013;150(2):395–420.
4. Das S, Tiwari GJ, Ghosh A. In silico analysis of new flavonoids from *Pongamia pinnata* with a therapeutic potential for age-related macular degeneration. *3 Biotech*. 2020;10(12):536.
5. Gnanaraj C, Sekar M, Fuloria S, Swain SS, Gan SH, Chidambaram K, et al. In silico molecular docking analysis of karanjin against alzheimer's and parkinson's diseases as a potential natural lead molecule for new drug design, development and therapy. *Molecules*. 2022;27(9):2834.
6. Khaerunnisa S, Suhartatu, Awaluddin R. Penelitian in silico untuk pemula. Surabaya: Airlangga University Press; 2020.
7. Benet LZ, Hosey CM, Ursu O, Oprea TI. BDDCS, the Rule of 5 and drugability. *Adv Drug Deliv Rev*. 2016;101:89–98.
8. Pansar T, Poso A. Binding affinity via docking: Fact and fiction. *Molecules*. 2018;23(8):1899.
9. Wu QJ, Tymianski M. Targeting NMDA receptors in stroke: new hope in neuroprotection. *Mol Brain*. 2018;11(1):15.
10. Seillier C, Lesept F, Toutirais O, Potzeha F, Blanc M, Vivien D. Targeting NMDA receptors at the neurovascular unit: Past and future treatments for central nervous system diseases. *Int J Mol Sci*. 2022;23:10336.
11. Shen Z, Xiang M, Chen C, Ding F, Wang Y, Shang C, et al. Glutamate excitotoxicity: Potential therapeutic target for ischemic stroke. *Biomed Pharmacother*. 2022 ;151:113125.
12. Hackos DH, Lupardus PJ, Grand T, Chen Y, Wang T-M, Reynen P, et al. Positive allosteric modulators of GluN2A-containing NMDARs with distinct modes of action and impacts on circuit function. *Neuron*. 2016 ;89(5):983–99.
13. Gill R, Soriano M, Blomgren K, Hagberg H, Wybrecht R, Miss M-T, et al. Role of caspase-3 activation in cerebral ischemia-induced neurodegeneration in adult and neonatal brain. *J Cereb Blood Flow Metab* . 2002 ;22(4):420–30.
14. Fan W, Dai Y, Xu H, Zhu X, Cai P, Wang L, et al. Caspase-3 modulates regenerative response after stroke. *Stem Cells* . 2014 Feb 1;32(2):473–86. 4
15. Machin A, Susilo I, Purwanto DA. Green tea and its active compound epigallocatechin-3-gallate (EGCG) inhibit neuronal apoptosis in a middle cerebral artery occlusion (MCAO) model. *J Basic Clin Physiol Pharmacol*. 2021 ;32(4):319–25.
16. Du J-Q, Wu J, Zhang H-J, Zhang Y-H, Qiu B-Y, Wu F, et al. Isoquinoline-1,3,4-trione derivatives inactivate caspase-3 by generation of reactive oxygen species. *J Biol Chem*. 2008 ;283(44):30205–15.
17. Singh A, Bhatt G, Gujre N, Mitra S, Swaminathan R, Limaye AM, et al. Karanjin. *Phytochemistry*. 2021 ;183:112641.
18. National Center for Biotechnology Information. Karanjin. PubChem Compound Summary for CID 100633. 2023
19. Noor AA, Othman SNN, Lum PT, Mani S, Shaikh MF, Sekar M. Molecules of interest – Karanjin – A review. *Pharmacogn J*. 2020 ;12(4):938–45.
20. National Center for Biotechnology Information. Karanjachromene. PubChem Compound Summary for CID 14033983. 2023
21. Rashid N, Abbasi MSA, Tahir MK, Yusof NM, Yamin BM. Isolation and crystal structure of karanjachromene. *Anal Sci X-ray Struct Anal Online*. 2008;24:X21–2.
22. National Center for Biotechnology Information. Pongapin. PubChem Compound Summary for CID 3083586. 2023
23. Roy R, Pal D, Sur S, Mandal S, Saha P, Panda CK. Pongapin and Karanjin, furanoflavonoids of *Pongamia pinnata*, induce G2/M arrest and apoptosis in cervical cancer cells by differential reactive oxygen species modulation, DNA damage, and nuclear factor kappa-light-chain-enhancer of activated B cell signal. *Phyther Res*. 2019;33(4):1084–94.
24. Sharma R, Williams IS, Gatchie L, Sonawane VR, Chaudhuri



- B, Bharate SB. Furanoflavones pongapin and lanceolatin B blocks the cell cycle and induce senescence in CYP1A1-overexpressing breast cancer cells. *Bioorg Med Chem.* 2018;26(23–24):6076–86.
25. Roy R, Mandal S, Chakrabarti J, Saha P, Panda CK. Downregulation of hyaluronic acid-CD44 signaling pathway in cervical cancer cell by natural polyphenols plumbagin, pongapin and karanjin. *Mol Cell Biochem.* 2021;476(10):3701–9.
26. National Center for Biotechnology Information. Pongachromene. PubChem Compound Summary for CID 14033985. 2023
27. Afrin S, Muhit MA, Sohrab MH, Hasan CM, Ahsan M. Antioxidant, thrombolytic, antimicrobial and cytotoxic activities of flavonoids isolated from the root bark of pongamia pinnata. *Dhaka Univ J Pharm Sci.* 2020;19(1):1–8.

TABLES AND FIGURES

Table 1. ADME Analysis of *P. pinnata* Phytochemical Compounds

Phytochemical Compounds	CID	Lipinski's Rule of Five				Violations	BBB Permeant
		Molecular Weight (g/mol)	Log P	H-bond donor	H-bond acceptor		
Karanjin	100633	292.29	3.43	0	4	0	Yes
Karanjachromene	14033983	334.37	3.94	0	4	0	Yes
Pongapin	3083586	105,6336.37	3.25	0	6	0	Yes
Pongachromene	14033985	378.37	3.80	0	6	0	Yes

Table 2. Molecular Docking Results of *P. pinnata* Phytochemical with Protein Target

Protein Target	Ligand	ΔG (kcal/mol)	K_i (μM)	Amino Acid Bond
NMDAR	Native Ligand (5YE)	-5.50	93.10	LYS140, PRO141, PHE142, LYS143, TYR144, ARG248, SER249, GLY250, LEU270, HIS273
	Karanjin	-5.12	176.24	ILE128, PRO141, PHE142, LYS143, TYR144, GLY250, HIS273
	Karanjachromene	-5.83	53.64	ILE128, PRO141, LYS143, TYR144, SER249, GLY250, HIS273
	Pongapin	-5.03	206.90	ILE128, PRO141, PHE142, LYS143, TYR144, SER249, GLY250, HIS273
	Pongachromene	-5.13	174.27	ILE128, PRO141, TYR144, SER249, GLY250, HIS273
Caspase-3	Native Ligand (RXB)	-5.82	54.30	THR166, GLU167, LEU168, CYS170, TYR204, THR255, LYS259
	Karanjin	-4.87	269.40	THR166, GLU167, LEU168, CYS170, THR255, PHE256, LYS259
	Karanjachromene	-4.98	223.26	THR166, LEU268, TYR204, TRP206, THR255, PHE256
	Pongapin	-4.88	266.27	THR166, GLU167, LEU168, CYS170, TYR204, THR255, LYS259
	Pongachromene	-5.08	189.85	THR166, GLU167, LEU168, CYS170, TYR204, THR255, PHE256, LYS259






Figure 1. Chemical Structure of *P. pinnata* Phytochemical Compounds (i) Karanjin, (ii) Karanjachromene, (iii) Pongapin, and (iv) Pongachromene




Figure 2. 3D Structure of Protein Target (i) NMDAR (5H8Q) with Native Ligand (5YE) and (ii) Caspase-3 (3DEI) with Native Ligand (RXB)




Figure 3. Visualization of *P. pinnata* phytochemical compounds (i) Karanjin, (ii) Karanjachromene, (iii) Pongapin, and (iv) Pongachromene with NMDAR




Figure 4. Visualization of *P. pinnata* phytochemical compounds (i) Karanjin, (ii) Karanjachromene, (iii) Pongapin, and (iv) Pongachromene with Caspase-3

In Silico Analysis of Pongamia pinnata to Inhibit Neuronal Apoptosis after Ischemic Stroke via NMDAR and Caspase-3

ORIGINALITY REPORT

14%

SIMILARITY INDEX

10%

INTERNET SOURCES

11%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| 1 | Submitted to iGroup
Student Paper | 1% |
| 2 | link.springer.com
Internet Source | 1% |
| 3 | www.medicopublication.com
Internet Source | 1% |
| 4 | Zihuan Shen, Mi Xiang, Chen Chen, Fan Ding, Yuling Wang, Chang Shang, Laiyun Xin, Yang Zhang, Xiangning Cui. "Glutamate excitotoxicity: Potential therapeutic target for ischemic stroke", Biomedicine & Pharmacotherapy, 2022
Publication | 1% |
| 5 | Abdulloh Machin, Imam Susilo, Djoko A. Purwanto. "Green tea and its active compound epigallocatechin-3-gallate (EGCG) inhibit neuronal apoptosis in a middle cerebral artery occlusion (MCAO) model", Journal of Basic and Clinical Physiology and Pharmacology, 2021 | 1% |

6	journal.ugm.ac.id Internet Source	1 %
7	Julien Pietrzak, Harold Common, Henri Migaud, Gilles Pasquier, Julien Girard, Sophie Putman. "Have the frequency of and reasons for revision total knee arthroplasty changed since 2000? Comparison of two cohorts from the same hospital: 255 cases (2013–2016) and 68 cases (1991–1998)", <i>Orthopaedics & Traumatology: Surgery & Research</i> , 2019 Publication	1 %
8	L.M.R. Al Muqarrabun, N. Ahmat, S.A.S. Ruzaina, N.H. Ismail, I. Sahidin. "Medicinal uses, phytochemistry and pharmacology of <i>Pongamia pinnata</i> (L.) Pierre: A review", <i>Journal of Ethnopharmacology</i> , 2013 Publication	1 %
9	Submitted to University of Hertfordshire Student Paper	1 %
10	cyberleninka.org Internet Source	1 %
11	Wenying Fan, Yiqin Dai, Haochen Xu, Ximin Zhu, Ping Cai, Lixiang Wang, Chungang Sun, Changlong Hu, Ping Zheng, Bing - Qiao Zhao. "Caspase - 3 Modulates Regenerative Response After Stroke", <i>STEM CELLS</i> , 2014	<1 %

-
- | | | |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 12 | repository.uhamka.ac.id
Internet Source | <1 % |
| 13 | www.discoveryjournals.org
Internet Source | <1 % |
| 14 | www.mdpi.com
Internet Source | <1 % |
| 15 | www.medcraveonline.com
Internet Source | <1 % |
| 16 | Hanna Helander, Henna Aikio, Pigga Keskitalo, Tuija Turunen. "chapter 21 Land-Based Participatory Pedagogical Experiment in Sami Language Distance Teaching", IGI Global, 2022
Publication | <1 % |
| 17 | Ziqing Wei, Yi Xie, Mingze Wei, Huijuan Zhao, Kaidi Ren, Qi Feng, Yuming Xu. "New insights in ferroptosis: Potential therapeutic targets for the treatment of ischemic stroke", <i>Frontiers in Pharmacology</i> , 2022
Publication | <1 % |
| 18 | research.ijcaonline.org
Internet Source | <1 % |
| 19 | bmcneurosci.biomedcentral.com
Internet Source | <1 % |
| 20 | Jais R. Bjelke, Egon Persson, Hanne B. Rasmussen, Birthe B. Kragelund, Ole H. Olsen. | <1 % |

"A loop of coagulation factor VIIa influencing macromolecular substrate specificity", FEBS Letters, 2007

Publication

21

Ping Cai, Haiyu Luo, Haochen Xu, Ximin Zhu et al. "Recombinant ADAMTS 13 Attenuates Brain Injury After Intracerebral Hemorrhage", Stroke, 2015

Publication

22

Rachmaniyah Rachmaniyah, Rusmiati Rusmiati. "Sansevieria trifasciata extract effectively inhibit the growth of Escherichia coli bacteria, in vitro", International Journal of Public Health Science (IJPHS), 2023

Publication

23

ijper.org
Internet Source

<1 %

24

seap.taylors.edu.my
Internet Source

<1 %

25

theecafellc.com
Internet Source

<1 %

26

www.wjgnet.com
Internet Source

<1 %

27

A. Singh, G. Bhatt, N. Gujre, S. Mitra, R. Swaminathan, A.M. Limaye, L. Rangan. "Karanjin", Phytochemistry, 2021

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

In Silico Analysis of Pongamia pinnata to Inhibit Neuronal Apoptosis after Ischemic Stroke via NMDAR and Caspase-3

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
