

Source details

International Journal of Angiology

Scopus coverage years: from 1992 to Present

Publisher: Thieme

ISSN: 1061-1711 E-ISSN: 1615-5939

Subject area: Medicine: Cardiology and Cardiovascular Medicine

Source type: *Journal*

CiteScore 2020

1.6



SJR 2020

0.293



SNIP 2020

0.540



[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology ✕

CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore 2020 ▼

$$1.6 = \frac{259 \text{ Citations } 2017 - 2020}{167 \text{ Documents } 2017 - 2020}$$

Calculated on 05 May, 2021

CiteScoreTracker 2021 📘

$$1.8 = \frac{290 \text{ Citations to date}}{160 \text{ Documents to date}}$$

Last updated on 06 April, 2022 • Updated monthly

CiteScore rank 2020 📘

Category	Rank	Percentile
Medicine		
— Cardiology and Cardiovascular Medicine	#214/317	32nd

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site 🌐](#)



International Journal of Angiology

COUNTRY

United States

Universities and research institutions in United States

SUBJECT AREA AND CATEGORY

Medicine
Cardiology and Cardiovascular
Medicine

PUBLISHER

Thieme
Medical
Publishers
Inc.

H-INDEX

26

PUBLICATION TYPE

Journals

ISSN

10611711, 16155939

COVERAGE

1992-2020

INFORMATION

[Homepage](#)
[How to publish in this journal](#)
john.eleftheriades@yale.edu




SCOPE




To provide a medium for the timely publication of scientific materials in the field of vascular and cardiovascular medicine and surgery. To promote a multi-disciplinary approach to the management and treatment of vascular and cardiovascular disease. To report the latest clinical outcomes of new surgical and clinical techniques in the field of angiology.

Quartiles

FIND SIMILAR JOURNALS

<p>1</p> <p>Vasa - European Journal of Vascular Medicine</p> <p>CHE</p> <p>77% similarity</p>	<p>2</p> <p>Journal of Cardiovascular Surgery</p> <p>ITA</p> <p>76% similarity</p>	<p>3</p> <p>Vascular</p> <p>GBR</p> <p>74% similarity</p>	<p>4</p> <p>Italian Jour and Endova</p> <p>ITA</p> <p>7 s</p>
---	--	---	---





International Journal of Angiology

Q3 Cardiology and Cardiovascular Medicine
best quartile

SJR 2020
0.29

powered by scimagojr.com

Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scima
```

SCImago Graphica

Explore, visually communicate and make sense of data with our new free tool.

Get it



Metrics based on Scopus® data as of April 2021

Leave a comment

Name

Email

(will not be published)

VOLUME 52 NUMBER 1 JULY 2019


ISSN 1091-2688

INTERNATIONAL JOURNAL OF ANGIOLOGY

Official Journal of the International College of Angiology
an ISS of journals of the International College of Angiology
the official journal for interventional vascular surgery

Editor in Chief

John A. Chaterlain, MD, FRCR



Angiology publishes
original research articles, clinical reports,
and reviews in the field of
vascular medicine.



Editorial Board

International Journal of Angiology

Editor-in-Chief

John A. Elefteriades, New Haven, Connecticut
Director, The Aortic Institute
Yale University School of Medicine

Co-Editor-in-Chief

Abeel A. Mangi, Washington, DC
MedStar Hear and Vascular Institute

Founding Editor

John B. Chang, Roslyn, New York

Consulting Editor

Kailash Prasad, Saskatoon, Canada
University of Saskatchewan, College of Medicine

Editorial Advisory Board

Eugene Braunwald, Boston, Massachusetts
Peter P. Liu, Toronto, Canada
Michael F. O'Rourke, Darlinghurst, Australia
Kailash Prasad, Saskatoon, Canada
Stephen Westaby, Oxford, United Kingdom

Senior Editors

Pertti T. Aarnio, Kauniainen, Finland
John D. Corson, Albuquerque, New Mexico
Victor A. Ferraris, Lexington, Kentucky
University of Kentucky Medical Center
Larry H. Hollier, New Orleans, Louisiana
LSU Health Science Center
Frank Manetta, Bay Shore, New York
Northwell Health
Takao Ohki, Tokyo, Japan
Jikei University School of Medicine
Medical University of Innsbruck
Pravin M. Shah, Newport Beach, California
Hoag Heart Institute
Sibu P. Saha, Lexington, Kentucky
University of Kentucky
Bauer E. Sumpio, New Haven, Connecticut
Yale University School of Medicine
Josef Veselka, Prague, Czech Republic
University Hospital Motol, 2nd Medical School, Charles University


Editorial Board

Elie D. Aboulafia, Michigan
Andreia P. Andreev, Bulgaria
Nobuyoshi Azuma, Japan
Semih I. Barlas, Turkey
Jeffrey S. Borer, New York
Anthony J. Comerota, Ohio
Gert J. DeBorst, The Netherlands
Iwan Dakota, Indonesia
Juergen Falkensammer, Austria
Alik Farber, Massachusetts
Glenn R. Faust, New York
Randall W. Franz, Ohio

Gary Giangola, New York
James J. Glazier, Michigan
Petr Hájek, Czech Republic
Michael H. Hall, New York
Hans M. Hoffmeister, Germany
Thomas Jahnke, Germany
Michael C. Kim, New York
Mark Kissin, New York
K.V. Krishnasastri, New York
Sean Kwon, New York
Gregg Landis, New York
Byung-Boong Lee, Virginia
J. Ernesto Molina, Minnesota
Ernesto P. Molmenti, New York
Debabrata Mukherjee, Texas
Stefan Müller-Hülsbeck, Germany
H. Müller-Wiefel, Germany
Vinay Nair, New York
Aravinda Nanjundappa, West Virginia
Alexander N. Orekhov, Russia
Ramdas G. Pai, California
Thomas Panetta, New York
Jeffrey T. Prem, Ohio
Dinker B. Rai, New York
Chand Ramaiah, Tennessee
Anwar Santoso, Indonesia
Darren B. Schneider, New York
Richard W. Schutzer, New York
Nicolas W. Shamma, Iowa
Rajinder P. Sharma, Michigan
Hiroshi Shigematsu, Japan
Ahmed S. Shoker, Canada
Tahir Tak, Minnesota
Eiji Tamiya, Japan
Matti R. Tarkka, Finland
Sanjay Tyagi, India
Padmini Varadarajan, California
Michael A. Winkler, Kentucky
Malka Yahalom, Israel
Mitsuhiro Yamamura, Japan
Wei Zhou, Arizona

Managing Editor

Denise M. Rossignol, Vermont



(<https://adfarm1.adition.com/redi?lid=7084502602900964557&gdpr=1&>

gdpr_consent=CPXLBmzPXLBMzAcABBENCKCsAP AAH AACHQIqtf X b3 j- 5 f t0eY1P9 7 -0zjhfdt-
 8N3f X LBx42M7vF36pq4KuR4Eu3LBIQdIH0HcTUmw6okVrzPsbk2cr7NKJ7PEmnMbO2dyGH9 n93TuZKY7 7z v-v v f 7-3f3 5 3--- e V 99zbv9 39nP 9v-
 9 gimASy5AFZZY4Mm0aRQogRhWEH LAoAKKAYWIKwgdXBTsrqJ9QQsAEaQnAiBBiCIBgEAAgkASERASAHgqEQBEAgABAAQqAI2AQWAFgYBAAKAaFiBFAEIEHbKUErYmBAVIFBZPZUJQd7GmEIZZ4EUC-
 ioQEayRAsDISFg5jgCQEvFkgeYoXyAEYAAAAA.f.gAD.gAAAAA&gdpr_pd=0&userid=7084502602900833485&sid=3412883&kid=2995779&bid=9392579&c=23329&keyword=&sr=0&gk=0&mdev=0&clickurl=
 DOI: 10.1055/s-00000167

International Journal of Angiology

Issue 03 · Volume 25 · September 2016

DOI: 10.1055/s-006-32245

[eFirst \(/products/ejournals/issue/eFirst/10.1055/s-00000167\)](/products/ejournals/issue/eFirst/10.1055/s-00000167)

[Issue \(/products/ejournals/journal/10.1055/s-00000167\)](/products/ejournals/journal/10.1055/s-00000167)

[Most Read \(/products/ejournals/topten/10.1055/s-00000167\)](/products/ejournals/topten/10.1055/s-00000167)

Review Article

139 Al Emam, Abdal Rahman A.; Almomani, Ahmed; Gilani, Syed A.; Khalife, Wissam I.:
**Spontaneous Coronary Artery Dissection: One Disease, Variable Presentations, and Different
 Management Approaches (/products/ejournals/abstract/10.1055/s-0035-1563604)**

Full Text HTML (/products/ejournals/html/10.1055/s-0035-1563604)
 PDF (444 kb) (/products/ejournals/pdf/10.1055/s-0035-1563604.pdf)

Original Article

148 Oren, Ohad; Oren, Michal; Turgeman, Yoav:
**Transradial versus Transfemoral Approach in Peripheral Arterial Interventions (/products/ejournals
 /abstract/10.1055/s-0035-1563607)**

Full Text HTML (/products/ejournals/html/10.1055/s-0035-1563607)
 PDF (92 kb) (/products/ejournals/pdf/10.1055/s-0035-1563607.pdf)

153 Akbarin, Mohammad Mehdi; Ravari, Hassan; Rajabnejad, Ataollah; Valizadeh, Narges; Fazeli, Bahare:
**Investigation of the Etiology of Anemia in Thromboangiitis Obliterans (/products/ejournals/abstract
 /10.1055/s-0035-1571190)**

Full Text HTML (/products/ejournals/html/10.1055/s-0035-1571190)
 PDF (151 kb) (/products/ejournals/pdf/10.1055/s-0035-1571190.pdf)

159 Murray, Gary L.; Colombo, Joseph:
**Ranolazine Therapy Reduces Non-ST-Segment-Elevation Myocardial Infarction and Unstable Angina in
 Coronary Disease Patients with Angina (/products/ejournals/abstract/10.1055/s-0036-1572364)**

Full Text HTML (/products/ejournals/html/10.1055/s-0036-1572364)
 PDF (98 kb) (/products/ejournals/pdf/10.1055/s-0036-1572364.pdf)

165 Shammas, Nicolas W.; Shammas, Gail A.; Harris, Thomas; Voelliger, Cara M.; Shammas, Andrew N.; Jerin, Michael:
**Safety and Effectiveness of Closure Devices Applied to a Stented Common Femoral Artery: A
 Retrospective Analysis (/products/ejournals/abstract/10.1055/s-0036-1572524)**

Full Text HTML (/products/ejournals/html/10.1055/s-0036-1572524)
 PDF (68 kb) (/products/ejournals/pdf/10.1055/s-0036-1572524.pdf)

168 Hitchner, Elizabeth; Morrison, Doug; Liao, Phoebe; Rosen, Allyson; Zhou, Wei:
**Genetic Polymorphisms Influence Cognition in Patients Undergoing Carotid Interventions (/products
 /ejournals/abstract/10.1055/s-0036-1580696)**

Full Text HTML (/products/ejournals/html/10.1055/s-0036-1580696)
 PDF (211 kb) (/products/ejournals/pdf/10.1055/s-0036-1580696.pdf)

174 Kaligis, Rinambaan W. M.; Adiarto, Suko; Erwinanto; Nugroho, Johannes; Pradnyana, Bagus Ari; Lefi, Achmad; Rifqi, Sodiqur:
**Carotid Intima-Media Thickness in Indonesian Subjects with Cardiovascular Disease Risk Factors Who
 Were Not Receiving Lipid-Lowering Agents (/products/ejournals/abstract/10.1055/s-0036-1572365)**

Full Text HTML (/products/ejournals/html/10.1055/s-0036-1572365)
 PDF (135 kb) (/products/ejournals/pdf/10.1055/s-0036-1572365.pdf)

Rapid Communication

181 Parameswaran, Sreejit; Sharma, Rajendra K.:
**Insulin Cannot Induce Adipogenic Differentiation in Primary Cardiac Cultures (/products/ejournals
 /abstract/10.1055/s-0035-1571191)**

Full Text HTML (/products/ejournals/html/10.1055/s-0035-1571191)
 PDF (200 kb) (/products/ejournals/pdf/10.1055/s-0035-1571191.pdf)

Case Report

186 Stechovsky, Cyril; Hajek, Petr; Cipro, Simon; Veselka, Josef:
**Mechanical Chest Compressions in Prolonged Cardiac Arrest due to ST Elevation Myocardial Infarction
 Can Cause Myocardial Contusion (/products/ejournals/abstract/10.1055/s-0034-1373734)**

Full Text HTML (/products/ejournals/html/10.1055/s-0034-1373734)
 PDF (202 kb) (/products/ejournals/pdf/10.1055/s-0034-1373734.pdf)

189 Yoon, Uzung; Marinelli, Laura; Ali, Sayed; Huberfeld, Seymour; Barrera, Rafael; Chang, John B.:

Early Amiodarone-Induced Pulmonary Toxicity after Endovascular Aneurysm Repair: A Case Report
(/products/ejournals/abstract/10.1055/s-0034-1387170)

Full Text [HTML \(/products/ejournals/html/10.1055/s-0034-1387170\)](https://products/ejournals/html/10.1055/s-0034-1387170)
PDF (91 kb) (/products/ejournals/pdf/10.1055/s-0034-1387170.pdf)

193 Palvanov, Arkadiy; Maróer, Ruth Leah; Siegel, David:

Asymptomatic Intrahepatic Portosystemic Venous Shunt: To Treat or Not To Treat? (/products/ejournals/abstract/10.1055/s-0035-1564658)

Full Text [HTML \(/products/ejournals/html/10.1055/s-0035-1564658\)](https://products/ejournals/html/10.1055/s-0035-1564658)
PDF (325 kb) (/products/ejournals/pdf/10.1055/s-0035-1564658.pdf)

Reply to Letter to the Editor

199 Prasad, Kailash:

Is Elevated Levels of Serum Soluble Receptor for Advanced Glycation End Products Harmful in Cigarette Smokers? (/products/ejournals/abstract/10.1055/s-0036-1579691)

Full Text [HTML \(/products/ejournals/html/10.1055/s-0036-1579691\)](https://products/ejournals/html/10.1055/s-0036-1579691)
PDF (69 kb) (/products/ejournals/pdf/10.1055/s-0036-1579691.pdf)

Carotid Intima-Media Thickness in Indonesian Subjects with Cardiovascular Disease Risk Factors Who Were Not Receiving Lipid-Lowering Agents

Rinambaan W. M. Kaligis, MD, FICA¹ Suko Adiarso, MD, PhD, FICA¹ Erwinanto, MD²
 Johannes Nugroho, MD, PhD, FICA³ Bagus Ari Pradnyana, MD, FICA⁴ Achmad Lefi, MD⁵
 Sodikur Rifqi, MD⁶

¹Department of Cardiology and Vascular Medicine, Faculty of Medicine, University of Indonesia, Harapan Kita National Cardiovascular Center, Jakarta, Indonesia

²Department of Cardiology and Vascular Medicine, Faculty of Medicine, Padjajaran University, Hasan Sadikin Hospital, Bandung, Indonesia

³Department of Cardiology and Vascular Medicine, Faculty of Medicine, Airlangga University, Soetomo Hospital, Surabaya, Indonesia

⁴Department of Cardiology and Vascular Medicine, Faculty of Medicine, Udayana University, Sanglah Hospital, Denpasar, Indonesia

⁵Department of Cardiology and Vascular Medicine, Darmo Hospital, Surabaya, Indonesia

⁶Department of Cardiology and Vascular Medicine, Faculty of Medicine, Diponegoro University, Kariadi Hospital, Semarang, Indonesia

Address for correspondence Suko Adiarso, MD, PhD, FICA, Department of Cardiology and Vascular Medicine, Faculty of Medicine, University of Indonesia, Harapan Kita National Cardiovascular Center, Jakarta, Indonesia (e-mail: sukoadiarso@gmail.com).

Int J Angiol 2016;25:174–180.

Abstract

Keywords

- ▶ intima-media thickness
- ▶ atherosclerosis
- ▶ cardiovascular diseases risk factors
- ▶ Indonesia
- ▶ common carotid artery
- ▶ lipid-lowering therapy
- ▶ asymptomatic individuals

Carotid intima-media thickness (CIMT) is frequently utilized for detection of subclinical atherosclerosis. This study aims to investigate the association between the CIMT values and demographic characteristics, cardiovascular disease (CVD) risk factors, lipid biochemistry profiles, and high-sensitivity C-reactive protein (hs-CRP) levels among the Indonesian population. Subjects who had two or more CVD risk factors but were not receiving lipid-lowering therapy were recruited from six hospitals of Indonesia. Measurements of CIMT are obtained by ultrasonography of 12 sites within the common carotid artery. CVD risk factors, lipid and glucose profiles, and hs-CRP values were analyzed with respect to distribution of CIMT. The mean–max CIMT was 0.805 ± 0.190 mm (minimum, 0.268 mm; maximum, 1.652 mm) and the mean–mean CIMT was 0.614 ± 0.190 mm (minimum, 0.127 mm; maximum, 1.388 mm). Multivariate analyses confirmed an independent association between increasing CIMT and increasing age (regression coefficient = 0.004; $p = 0.004$). Our data show normative mean–mean CIMT data for Indonesian subjects with two or more CVD risk factors who are not receiving lipid-lowering therapy, which may guide CVD risk stratification of asymptomatic individuals in Indonesia.

Like many countries undergoing socioeconomic transition, Indonesia is struggling with a fast-growing burden of non-communicable diseases such as diabetes and cardiovascular diseases (CVDs). CVD is one of the main causes of death in Indonesia, contributing for 31.9% of all deaths in the country and 11.2% of all deaths in hospitals.¹ Data from the Indonesia Basic Health Research conducted by the Ministry of Health in 2007 showed that among people older than 18 years the prevalence of hypertension was 29.8% and that of heart diseases was 7.2%.² Risk factors such as smoking, poor diet, and lack of exercise are growing in importance and further contributing to the incidence of these diseases.

Atherosclerosis is the underlying cause of the majority of CVDs. It has been known that atherosclerosis starts in the early years and develops over decades as a clinically silent disease, thus providing the opportunity for identification of high-risk individuals in preclinical stage and early initiation of preventive strategies. Early identification of subclinical atherosclerosis by noninvasive measures such as ultrasonic measurements of intima-media thickness (IMT) is often used as a surrogate end point of CVD and coronary artery disease, with increased carotid IMT (CIMT) having been associated with both the presence and extent of coronary artery disease.^{3,4}

Although extensive studies on the use of CIMT have been conducted in the last two decades, data among Asian population have been growing just recently. One study of atherosclerotic disease involving eight Asian countries has reported the distribution of CIMT values in Asian population with at least two CVD risk factors who were not receiving any lipid-lowering treatment.⁵ The study has established that age, male gender, low high-density lipoprotein cholesterol (HDL-C) levels, and elevated fasting blood glucose levels are associated with the increase of mean-mean CIMT. However, the Asian population consisted of diverse ethnic groups, and the study found a significant variation of mean-mean CIMT across countries.

As the third-largest country in Asia and the largest in the Southeast Asia, the Indonesian population represents an important proportion of Asian people who might have distinct CVD profile, risk associations, and different levels of genetic-environmental interactions.

This study aim to investigate the association between the CIMT values and demographic characteristics, CVD risk factors, lipid biochemistry profiles, and high-sensitivity C-reactive protein (hs-CRP) levels among the Indonesian cohort in the Asian study.

Methods

Study Subjects

The methods of this study have been published elsewhere.⁵ In summary, the study participants were people aged 30 to 69 years with at least two CVD risk factors who had not received lipid-lowering treatment within the 3 months prior to study entry. CVD risk factors included age (males ≥ 45 years and females ≥ 55 years), cigarette smoking (any cigarette smoking in the past month), hypertension (arterial blood pressure ≥ 140 mm Hg systolic and/or ≥ 90 mm Hg diastolic or taking

antihypertensive medication), low HDL-C (< 40 mg/dL, corresponding to 1.0 mmol/L), and family history of premature coronary heart disease (CHD) (clinical CHD or sudden death in the father or another first-degree male relative < 55 years of age; CHD in the mother or another first-degree female relative < 65 years of age). Exclusion criteria were a history of CHD or any condition that affects hs-CRP levels, including malignancy or chronic inflammation.

This study was performed in accordance with the ethical principles of the Declaration of Helsinki, and is consistent with International Conference on Harmonisation (ICH)/Good Clinical Practice (GCP) guidelines and applicable regulatory requirements and the AstraZeneca policy on Bioethics and Human Biological Samples. The study protocol and informed consent forms were approved by the Ethics Committee for Medical Research, Harapan Kita National Cardiovascular Centre.

Data Collection

Data collection for each subject was performed during one or two visits, depending on the availability of a sonographer at the study site. The time between visits was not more than 3 months and eligibility criteria were reassessed at the second visit. Subjects' informed consent was taken, physical examination and measurement of vital signs (including weight, height, waist circumference, and blood pressure) were conducted, and demographic data were collected.

Subjects' fasting blood samples were taken for analyses of total cholesterol, triglyceride, low-density lipoprotein cholesterol, HDL-C, fasting blood glucose, and hs-CRP levels, which were performed at a single central laboratory for consistency.

Subjects were asked to complete a questionnaire which recorded information regarding prevailing knowledge of CVD risk factors, cholesterol management, and physician visit frequency.

CIMT Measurement

Measurements of CIMT in all subjects were conducted using a SonoSite MicroMaxx high-resolution, digital, portable ultrasound system with standard accessories and SonoCalc-embedded IMT software by a certified sonographer. Measurements were taken in the left and right distal common carotid artery (CCA) of both the near and far wall at each location from three positions: anterior, lateral, and posterior. Using all 12 measurements taken, the SonoCalc software calculated for each subject the mean (mean-mean CIMT), minimum, and maximum for the average values of the 12 measurements and the mean (mean-max CIMT), minimum, and maximum for the maximum values of the 12 measurements.

Statistical Analysis

The distribution of CIMT measurements was described in terms of mean, standard deviation (SD), median, minimum, and maximum values. The associations between CIMT measurements and demographic characteristics, CVD risk factors, and lipid and hs-CRP values were analyzed using one-way analysis of variance (ANOVA) models (for

categorical variables) and univariate linear regression models (for continuous variables). The effects of identified explanatory variables were further explored using a general linear model approach by stepwise inclusion of explanatory variables into a multivariate model, based on a significance level of 1%. Log-transformed hs-CRP values were used for statistical analyses except for calculating descriptive statistics. The per protocol analysis set (PPS) from available data was used for analysis. It included subjects who did not violate inclusion and exclusion criteria and whose CIMT values were available. The SAS package, version 9.2 (SAS Institute Inc., Cary, NC), was used for all statistical analysis.

Results

Subject Characteristics

A total of 374 patients from six hospitals in Indonesia enrolled in this study. Of those, 356 were included in the PPS. The subjects included more male (236/356, 66.3%), with the mean (SD) age of the subjects being 56 (7.2) years. ▶ **Table 1** shows the demographic characteristics, CHD risk factors, presence of CHD risk equivalence, and metabolic syndrome of the Indonesian subjects included in the PPS. The most common CHD risk factors were older age (332/356 subjects; 93.3%) and hypertension (259/356 subjects; 72.8%), while the most common metabolic syndrome present was high blood pressure (230/355; 64.8%), followed by abdominal obesity (198/355; 55.9%).

Overall, 287 (80.6%) subjects had never received any lipid-lowering treatment. Of those who had previously received lipid-lowering therapy, statins were the most commonly administered lipid-lowering agent and were taken for a mean of 4.6 months (minimum and maximum treatment duration of 1 and 48 months, respectively). Treatment was stopped at least 3 months prior to study entry.

CIMT Measurements

The overall mean (SD) of the average CIMT measurements (mean–mean CIMT) was 0.614 (0.19) mm (▶ **Table 2**). The minimum and maximum values of mean–mean CIMT measurements were 0.127 and 1.388 mm, respectively. Of the 356 Indonesian subjects in the PPS, 209 (58.71%) had a mean–mean CIMT value of 0.6 or higher (▶ **Fig. 1**).

The mean (SD) of maximum region CIMT values (mean–max CIMT) was 0.805 (0.19) mm (▶ **Table 2**). The minimum and maximum values of mean–max CIMT measurements were 0.268 and 1.652 mm, respectively. Of the 356 Indonesian subjects in the PPS, 186 (52.24%) had a mean–max CIMT value of 0.8 or higher (▶ **Fig. 2**).

We found only two subjects with carotid plaque.

Association of Mean–Mean CIMT with Demographic and CVD Risk Factors

Linear regression analysis (▶ **Table 3**) found that factors which have positive association with mean–mean CIMT were age (regression coefficient = 0.003; $p = 0.026$) and

log (hs-CRP) (regression coefficient = 0.020; $p = 0.029$), while triglyceride level was negatively associated with mean–mean CIMT (regression coefficient = -0.0002 ; $p = 0.022$). Other demographic and CVD risk factors were found to be not associated with mean–mean CIMT based on ANOVA analysis. Further multivariate analysis using general linear model found that no factors could be included in the final model at $\alpha = 0.01$ level.

Association of Mean–Max CIMT with Demographic and CVD Risk Factors

Linear regression analysis (▶ **Table 3**) found that factors that have positive association with mean–max CIMT were age (regression coefficient = 0.004; $p = 0.004$) and log (hs-CRP) (regression coefficient = 0.022; $p = 0.021$), while triglyceride level was negatively associated with mean–max CIMT (regression coefficient = -0.0003 ; $p = 0.012$). Other demographic and CVD risk factors were found to be not associated with mean–mean CIMT based on ANOVA analysis. Further multivariate analysis using general linear model found age as the only independent risk factor for the increase of mean–max CIMT (regression coefficient = 0.004; $p = 0.004$).

Discussion

Our study found that the average mean CIMT and average max CIMT of Indonesian patients with two or more CVD risk factors were 0.614 and 0.805 mm, respectively. The only independent association found was between age and average max CIMT.

This result is quite similar with a study among healthy middle-age population in Korea, which found the average mean CIMT of 0.60 among men and 0.53 among female.⁶ However, our study found no significant difference on the average mean CIMT between male and female. The average mean CIMT of the Indonesian cohort is also lower compare with normotensive middle age and elderly population in southern China with the average mean of 0.71 mm.⁷

Analysis of the healthy individuals younger than 45 years from the six cohorts that are part of the USE-Intima Media Thickness project found higher average mean CIMT of 0.63 mm. This analysis of the USE-IMT project included population from Germany, Canada, Finland, Japan, Norway, and the United States.⁸

The overall mean–mean CIMT of the eight Asian countries in the original study was higher (0.662 ± 0.16 mm).⁵ In this study, the Indonesian cohort has the third lowest mean–mean CIMT after Thailand (0.567 ± 0.120 mm) and the Philippines (0.583 ± 0.117 mm). However, it was also reported that the participants from Thailand and the Philippines were relatively younger compare with Indonesian participants (48.8 ± 12.5 and 52.2 ± 7.6 years vs. $0.56.0 \pm 7.2$ years). A significant correlation between increasing age and mean CIMT has been consistently observed in diverse populations, including the Asian.^{5,6,9,10}

Besides older age, low HDL-C, elevated fasting blood glucose, and male gender were independent predictor of the mean–mean CIMT increase in the overall Asian population. Yet,

Table 1 Subjects' demographic characteristics, CHD risk factors, presence of CHD risk equivalence, and metabolic syndrome ($n = 356$)

Factors		
Age (y)	Mean (SD)	56.0 (7.2)
	Median (min/max)	57 (32/69)
Age	30 to 39 y	5 (1.40)
	40 to 49 y	68 (19.10)
	50 to 59 y	160 (44.94)
	60 to 69 y	123 (34.55)
Gender	Male	236 (66.29)
	Female	120 (33.71)
Weight (kg) ^a	Mean (SD)	63.9 (11.7)
	Median (min/max)	64 (26/111)
Height (cm) ^a	Mean (SD)	160.2 (8.2)
	Median (min/max)	160 (134/180)
BMI (kg/m ²) ^a	Mean (SD)	24.8 (3.9)
	Median (min/max)	25 (12.89/36.59)
BMI (kg/m ²) ^a	Normal weight (<25)	194 (54.80)
	Overweight (25 to 30)	122 (34.46)
	Obese (>30)	38 (10.73)
Waist circumference (cm) ^a	Mean (SD)	87.9 (10.9)
	Median (min/max)	89 (58/117)
Sitting SBP (mm Hg) ^a	Mean (SD)	135.4 (21.4)
	Median (min/max)	130 (100/220)
Sitting DBP (mm Hg) ^a	Mean (SD)	83.4 (12.0)
	Median (min/max)	80 (53/140)
CHD risk factors	Age (men ≥ 45 y; women ≥ 55 y)	332 (93.3)
	Cigarette smoking (any cigarette smoking in the past month)	152 (42.7)
	Hypertension	259 (72.8)
	Low HDL-C	18 (5.1)
	Family history of premature CHD	32 (9.0)
	High HDL-C (negative risk factor if ticked "yes")	2 (0.6)
Presence of CHD risk equivalents ^b	Diabetes	38 (10.7)
	Abdominal aortic aneurysm	1 (0.3)
	Carotid artery disease	7 (2.0)
Presence of metabolic syndrome ^b	Abdominal obesity: (men ≥ 90 cm; women ≥ 80 cm) ^a	198 (55.9)
	Triglyceride (TG ≥ 150 mg/dL)	16 (4.5)
	HDL-C: (men < 40mg/dL; women < 50 mg/dL)	22 (6.2)
	Blood pressure ($\geq 130/85$ mm Hg)	230 (64.8)
	Fasting glucose (≥ 100 mg/dL)	20 (5.6)

Abbreviations: BMI, body mass index; DBP, diastolic blood pressure; HDL-C, high-density lipoprotein cholesterol; SBP, systolic blood pressure.

^a $n = 354$.

^b $n = 355$.

in Indonesian cohort, age is only associated independently with mean-max CIMT and not mean-mean CIMT. One study among Iranian population of 35 to 64 years old without history of cardiovascular event also found that age (odds ratio [OR],

1.11; 95% confidence interval [CI], 0.56–4.34; $p < 0.01$) and male gender (OR [95% CI], 1.14 [0.63–2.23]; $p < 0.05$) were the only significant independent predictors of the mean-max CIMT of ≥ 0.8 mm.¹¹

Table 2 CIMT and hs-CRP values of the Indonesian subjects (n = 356)

	Mean	SD	Median	Minimum	Maximum	95% CI	
						Lower	Upper
Mean–mean CIMT (mm)	0.614	0.1885	0.627	0.127	1.388	0.595	0.634
Mean–max CIMT (mm)	0.805	0.1945	0.805	0.268	1.652	0.785	0.825
	G-mean	CV%					
hs-CRP (mg/dL) ^a	0.172	179.76	0.180	0.020	9.240		

Abbreviation: CIMT, carotid intima-media thickness; hs-CRP, high-sensitivity C-reactive protein.

Note: G-mean is the geometric mean. CV is the coefficient variation calculated as $\text{SQRT}([\exp(s^2) - 1] \times 100)$, where s is a standard deviation on a log-transformed scale.

^an = 349.

Although hs-CRP and triglyceride level were found to be associated during bivariate linear regression analysis with both the mean–mean CIMT and mean–max CIMT in the Indonesian cohort, both failed to stand as independent predictor of CIMT during multivariate analysis. Several stud-

ies have shown that hs-CRP is positively correlated with increased risk of CVD. Yet, in the study among Asian population, hs-CRP level did not correlate directly with CIMT values.⁵

Our study has been the first to show the application of the CIMT measurements among Indonesian subjects, which




Fig. 1 Distribution of CIMT in PPS population of Indonesia (n = 356) based on the average of the mean region values (mean–mean CIMT). CIMT, carotid intima-media thickness; PPS, per protocol analysis set.




Fig. 2 Distribution of CIMT in PPS population of Indonesia (n = 356) based on the average of the maximum region values (mean–max CIMT). CIMT, carotid intima-media thickness; PPS, per protocol analysis set.

Table 3 Association of mean–mean CIMT and mean–max CIMT with demographic and CVD risk factors

Factors	Regression coefficient estimate	SE	95% CI		p-Value
			Lower	Upper	
Mean–mean CIMT					
Age (y)	0.003095	0.001384	0.000372	0.005817	0.0260
Weight (kg)	–0.000840	0.000857	–0.002524	0.000845	0.3277
BMI (kg/m ²)	–0.003017	0.002564	–0.008060	0.002026	0.2402
Waist circumference (cm)	–0.000143	0.000928	–0.001968	0.001681	0.8772
TC (mg/dL)	0.000203	0.000242	–0.000273	0.000678	0.4024
LDL-C (mg/dL)	0.000494	0.000291	–0.000078	0.001065	0.0903
HDL-C (mg/dL)	–0.000876	0.000916	–0.002676	0.000925	0.3395
TG (mg/dL)	–0.000244	0.000106	–0.000452	–0.00036	0.0218
Glucose (mg/dL)	0.000203	0.000278	–0.000343	0.000750	0.4648
Log (hs-CRP)	0.020103	0.009183	0.002042	0.038164	0.0293
Mean–max CIMT					
Age (y)	0.004138	0.001421	0.001343	0.006933	0.0038
Weight (kg)	–0.000822	0.000884	–0.002560	0.000917	0.3531
BMI (kg/m ²)	–0.002635	0.002647	–0.007841	0.002571	0.3202
Waist circumference (cm)	–0.000111	0.000957	–0.001993	0.001771	0.9073
TC (mg/dL)	0.000242	0.000250	–0.000249	0.000733	0.3332
LDL-C (mg/dL)	0.000581	0.000300	–0.000009	0.001171	0.0537
HDL-C (mg/dL)	–0.000714	0.000947	–0.002577	0.001149	0.4516
TG (mg/dL)	–0.000276	0.000109	–0.000491	–0.000061	0.0120
Glucose (mg/dL)	0.000285	0.000287	–0.000280	0.000850	0.3216
Log (hs-CRP)	0.021917	0.009488	0.003257	0.040577	0.0215

Abbreviations: BMI, body mass index; CIMT, carotid intima-media thickness; HDL-C, high-density lipoprotein cholesterol; hs-CRP, high-sensitivity C-reactive protein; LDL-C, high-density lipoprotein cholesterol; TC, total cholesterol; TG, triglyceride.

Note: Only the continuous variables are presented. Analysis was done by linear regression.

shows the values among those with at least two CVD risk factors who were not receiving any lipid-lowering treatment. As the CIMT measurement is often used in the identification of subclinical atherosclerotic disease, information on the baseline value is important.

Limitation of our study includes its cross-sectional nature, which is not the strongest design to prove causal associations. In addition, our examination was limited to the CCA, which may not have detected the presence of atherosclerosis in other vascular beds or more distal segments of the carotid artery. Since our study involved subjects with at least two CVD risk factors which possibly have higher risk of CAD, measurement of healthy subjects is necessary to establish the normative CIMT value among Indonesian subjects. A longitudinal study is further needed for a better evaluation of the relationships between cardiovascular risk factors, subclinical atherosclerosis, and the risk of cardiovascular events in this population.

Conclusion

The findings of this study have important implications for CVD screening and the identification of asymptomatic, high-risk patients and the consequent implementation of CVD prevention strategies in Indonesia.

Acknowledgment

This study and publication was sponsored by AstraZenica.

References

- 1 Ministry of Health Republic of Indonesia. Indonesia's Health Profile 2008. Jakarta: Ministry of Health Republic of Indonesia; 2008
- 2 National Institute of Health Research and Development. Basic Health Research 2007. Jakarta: Ministry of Health of the Republic of Indonesia; 2008

- 3 Zhang Y, Guallar E, Qiao Y, Wasserman BA. Is carotid intima-media thickness as predictive as other noninvasive techniques for the detection of coronary artery disease? *Arterioscler Thromb Vasc Biol* 2014;34(7):1341–1345
- 4 van den Oord SC, Sijbrands EJ, ten Kate GL, et al. Carotid intima-media thickness for cardiovascular risk assessment: systematic review and meta-analysis. *Atherosclerosis* 2013;228(1):1–11
- 5 Oh B-H, Kaligis RWM, Wang Y, et al. Survey of atherosclerotic disease in Asian subjects with cardiovascular disease risk factors who were not receiving lipid-lowering agents. *Int J Cardiol* 2013; 168(3):2761–2766
- 6 Youn YJ, Lee NS, Kim J-Y, et al. Normative values and correlates of mean common carotid intima-media thickness in the Korean rural middle-aged population: the Atherosclerosis Risk of Rural Areas in Korea General Population (ARIRANG) study. *J Korean Med Sci* 2011;26(3):365–371
- 7 Liang Y, Yan Z, Sun B, et al. Cardiovascular risk factor profiles for peripheral artery disease and carotid atherosclerosis among Chinese older people: a population-based study. *PLoS ONE* 2014;9(1):e85927
- 8 Eikendal ALM, Groenewegen KA, Anderson TJ, et al; USE-IMT Project Group. Common carotid intima-media thickness relates to cardiovascular events in adults aged <45 years. *Hypertension* 2015;65(4):707–713
- 9 Pastorius CA, Medina-Lezama J, Corrales-Medina F, et al. Normative values and correlates of carotid artery intima-media thickness and carotid atherosclerosis in Andean-Hispanics: The Prevencion Study. *Atherosclerosis* 2010;211(2):499–505
- 10 Bauer M, Delaney JA, Möhlenkamp S, et al; Multi-Ethnic Study of Atherosclerosis; Investigator Group of the Heinz Nixdorf Recall Study. Comparison of factors associated with carotid intima-media thickness in the Multi-ethnic Study of Atherosclerosis (MESA) and the Heinz Nixdorf Recall Study (HNR). *J Am Soc Echocardiogr* 2013;26(6):667–673
- 11 Azarpazhooh MR, Kazemi-Bajestani SMR, Esmaeili H, et al. Cardiovascular risk factors and nutritional intake are not associated with ultrasound-defined increased Carotid Intima Media Thickness in individuals without a history of cardiovascular events. *Int J Prev Med* 2014;5(11):1412–1421