



Source details

IOP Conference Series: Earth and Environmental Science

Scopus coverage years: from 2010 to Present

ISSN: 1755-1307 E-ISSN: 1755-1315

- Subject area:
- Environmental Science: General Environmental Science
 - Earth and Planetary Sciences: General Earth and Planetary Sciences
 - Physics and Astronomy: General Physics and Astronomy

Source type: Conference Proceeding

CiteScore 2020 **0.5**

SJR 2020 **0.179**

SNIP 2020 **0.436**

[View all documents >](#) [Set document alert](#) [Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology x

CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore **2020** v

0.5 = $\frac{25,463 \text{ Citations } 2017 - 2020}{49,883 \text{ Documents } 2017 - 2020}$

Calculated on 05 May, 2021

CiteScoreTracker 2021 i

0.6 = $\frac{44,677 \text{ Citations to date}}{74,322 \text{ Documents to date}}$

Last updated on 06 April, 2022 - Updated monthly

CiteScore rank 2020 i

Category	Rank	Percentile
Environmental Science		
— General Environmental Science	#183/220	17th
Earth and Planetary Sciences		
— General Earth and Planetary Sciences	#157/186	15th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

ISSN 1741-3337

IOP Conference Series

Earth and Environmental Science

**1st Workshop on Metrology for Agriculture
and Forestry (METROAGRIFOR)**

275

VOLUME 275 – 2019

3–2 October 2018
Asolo, Italy

EDITOR
Enrico Ferra Tosi (ed)

The open access journal for conference proceedings

iopscience.org/jpcs

IOP Publishing

NOTICE: Ukraine: Click here to read IOP Publishing's statement.

Table of contents

Volume 441

2020

◀ Previous issue Next issue ▶

2nd International Conference on Fisheries and Marine Science 26 September 2019, Surabaya, Indonesia

Accepted papers received: 16 January 2020

Published online: 24 February 2020

Open all abstracts

Preface

OPEN ACCESS 011001

Preface

+ Open abstract  View article  PDF

OPEN ACCESS 011002

Peer review statement

+ Open abstract  View article  PDF

Papers

Earth and Environmental Science

OPEN ACCESS 012001

The grow-out of abalone (*Haliotis squamata*) at different shelter shape on growth and survival and its marine environmental influences at Lembongan Bay coastal waters

I Ardi, E Setiadi, Rasidi and W S Pranowo

+ Open abstract  View article  PDF

OPEN ACCESS 012002

Stocking density of Rotifera, *Hexarthra mira* on water quality and production

I Ardi, S Cahyaningsih and E Setiadi

+ Open abstract  View article  PDF

OPEN ACCESS 012003

The effect addition of Kappa carrageenan flour to the level of gel strength and acceptability

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



of dumpling from threadfin bream fish (*Nemipterus nematophorus*) surimi

D M Astutik, L Sulmartiwi, E Saputra and D Y Pujiastuti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012004

Maximizing production of a male offspring in *Moina macrocopa* culture through manipulation of rice bran suspension concentration

A S Mubarak, D Jusadi, M Zairin Jr and M A Suprayudi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012005

The Concept of the Right to Management of Coastal Communities in the Regional Autonomy Era: Experience from Community Assistance to Obtain the Right to Manage Sea Cucumbers in Sunsak Bay, East Lombok.

A Wahyono and M Illiyani

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012006

Selective breeding technique: Pandu and Kunti tilapia (*Oreochromis niloticus*) broodstock candidates at PBIAT Janti, Klaten-Central Java

S H Samara, A W Fathurrozi and Sutarno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012007

Potential of *Caulerpa racemosa* extracts as sunscreen creams

E B Ersalina, A A Abdillah and L Sulmartiwi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012008

Histopathology of the gill of Vaname Shrimp (*Litopenaeus vannamei*) infested by protozoan ectoparasite

S R Pribadi, P D W Sari and S Subekti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012009

The effect of iron powder as oxygen absorber active packaging on fish oil total oxidation value

E N Hidayah, RR J Triastuti and A A Abdillah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012010

The occurrence of endoparasite helminth on Threadfin Bream (*Nemipterus japonicus*) from the fish auction place Banyuwangi, Probolinggo, East Java

D S Octaria, P D W Sari and G Mahasri

We use cookies to enhance your browsing experience, to personalize and improve our site, to analyze our usage, and to help us understand our users' needs. To find out more, see our Privacy and Cookies policy.



Contrast does not Affect Adipose Mesenchymal Stem Cell (AMSC) Proliferation

G R Soekarno, I G R Suryawan, A Andrianto and W M Valentidenta

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012175

The Clinical Benefit of D-ribose in Cardiovascular Ischemic Disease: A New Potential Energy Supplement

I Damanik and D Soemantri

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012176

Association between White Blood Cell to Mean Platelet Volume Ratio (WMR) with Troponin for Refractory Angina Prediction in NSTEMI-ACS patients

I E Hermawati and M Y Assegaf

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012177

Lower Gensini Score in Navy Personnel Compared to Civilians, rather than for Non-Commissioned Officers

I G P G Semita, M J Al-Farabi, M F Huda, B Jovie and I G R Suryawan

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012178

Correlation between Global Longitudinal Strain (GLS)-Left Ventricle and TEI Index (TI) with Seattle Heart Failure Model (SHFM) Score in Chronic Heart Failure Patients with Systolic Dysfunction

I Kartikasari, A Lefi, D Soemantri and R R Juwita

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012179

Atria Score for Antithrombotic Agent Utilization in Ischemic Stroke Patients with Atrial Fibrillation

I P Dewi, K P Dewi and R T Pinzon

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012180

Renal Artery Stenosis: The Challenging in Diagnosis and Therapy

I P Dewi and N Mardiana

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012181

Plasma Histamine Level as a Distinguishing Factor between Stable Coronary Artery

[Disease and Acute Coronary Syndrome](#)
 This site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

I S Prihaliningsih, J N E Putranto, D Soemantri and A S Hadi



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012197

Predictors of Arteriovenous Fistula Early Failure in End-Stage Renal Disease Patients: Real-World Data in Surabaya

M R Amadis, J N E Putranto, I Maghfirah and N Mardiana

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012198

Different Grades of Body Mass Index are Correlated with Left Atrium and Ventricle Structures in Patients with Hypertensive Heart Disease

M S Bhisma, M J Al-Farabi and B S Pikir

[+ Open abstract](#) [View article](#) [PDF](#)

JOURNAL LINKS

[Journal home](#)

[Journal scope](#)

[Information for organizers](#)

[Information for authors](#)

[Contact us](#)

[Reprint services from Curran Associates](#)



The Electrochemical Society
241st ECS Meeting
Early registration pricing
ends April 25
ECS members receive up to 30% off
[Register now!](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



PAPER • OPEN ACCESS

Correlation between Global Longitudinal Strain (GLS)-Left Ventricle and TEI Index (TI) with Seattle Heart Failure Model (SHFM) Score in Chronic Heart Failure Patients with Systolic Dysfunction

To cite this article: I Kartikasari *et al* 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **441** 012178

View the [article online](#) for updates and enhancements.

Correlation between Global Longitudinal Strain (GLS)-Left Ventricle and TEI Index (TI) with Seattle Heart Failure Model (SHFM) Score in Chronic Heart Failure Patients with Systolic Dysfunction

I Kartikasari¹, A Lefi^{1,2*}, D Soemantri^{1,2} and R R Juwita¹

¹Faculty of Medicine, Airlangga University, Surabaya, Indonesia

²Department of Cardiology and Vascular Medicine, Dr. Soetomo General Hospital, Surabaya, Indonesia

*Corresponding author: achmadlefi@gmail.com

Abstract. Assessment of left ventricular function in patients with chronic heart failure is important for determining prognosis and treatment plans. The prognostic values of the myocardium and global ventricular functions remain unclear. This cross-sectional study included 30 subjects obtained through purposive sampling. Global longitudinal strains (GLS) is used to assess the left ventricular systolic function and Tei index (TI) is used to assess both global systolic and diastolic ventricular function. SHFM scoring was calculated based on existing patient data. The mean GLS-left ventricle value was -8.08 ± 3.98 , whereas TI value was 0.65 ± 0.14 . GLS-left ventricle had significant negative correlation with estimated one and five years mortality based on SHFM score ($r = -0.676$ and $p = 0.0001$) whereas TI had a significant positive correlation ($r = 0.745$ and $p = 0.0001$; $r = 0.738$ and $p = 0.0001$). GLS-left ventricle had a significant negative correlation with an estimated one and five years mortality based on SHFM score, whereas TI had a significant positive correlation. Hence it is suggested that GLS-left ventricle and TI can be a prognostic factor.

1. Introduction

Heart failure is a significant cause of morbidity and mortality in various countries. In the last few decades, the incidence and prevalence of chronic heart failure have continuously increased. In the United States, 5.8 million people were suffering from heart failure in 2012, which is expected to increase to 8.5 million by 2030, while in the world the population of heart failure is close to 23 million, and is expected to continue to increase [1-3].

Ejection fraction (EF) of the left ventricle is the frequently used parameter to assess systolic function. Echocardiography, besides being a diagnostic modality, can also have prognostic value, especially parameters that assess left the ventricular function [4-5]. Previous research reported that in a population of patients with EF <45%, mortality decreased almost linearly with an increase in EF (EF <15%, mortality: 51.7%; EF 36% - 45%, mortality: 25.6%; $p < 0.0001$) [6]. Other than being used as



diagnostic tools, left ventricular function can also be used to assess the prognosis, determine treatment plans, make decisions related to the therapy of expensive non-surgical devices and assess the response to treatment. Global Longitudinal Strain (GLS) of the left ventricle assessed through Speckle Tracking Echocardiography (STE) is sensitive in determining complex left ventricular mechanics, by analyzing multidimensional myocardial deformation, which allows GLS to be a more accurate and sensitive parameter than EF in assessing left ventricular systolic function [7-9]. Stanton T *et al.* found that GLS was superior compared to EF and the Wall Motion Scoring Index (WMSI) in predicting the occurrence of all-mortality cause in patients with chronic heart failure [10].

Risk stratification is proven to provide benefits to heart failure patients in various ways as optimal management. The risk model has an essential role in facilitating patients and clinicians in understanding the condition of the disease experienced, especially about the possible outcomes. Prediction can be suboptimal if only based on a holistic assessment by the clinician alone [11]. Currently, there are many risk assessment tools which have been developed to predict the prognosis of patients with heart failure. One of multivariable risk scores that has been validated and recommended by ACCF/AHA Guidelines for the Management of Heart Failure 2013 was the Seattle Heart Failure Model (SHFM). SHFM provides an accurate estimation of mortality in patients with heart failure in 1, 2, and 5 year periods by using clinical parameters that are quickly evaluated, such as pharmacological therapy, medical devices, and laboratory characteristics [5,12].

Currently, the role of left ventricular GLS and TI in assessing mortality estimates in patients with chronic heart failure is not yet known, several studies have been carried out in several countries, but this study has never been conducted in Indonesia. This study aims to assess the correlation between left ventricular and TI GLS values with estimates of mortality within one, and five years assessed from SHFM scores, and evaluate which parameters have a stronger correlation with SHFM scores in outpatients with chronic heart failure with systolic dysfunction at Dr. Hospital Soetomo Surabaya.

2. Methods

This study was conducted at the Cardiac and Echocardiographic Room at the Dr. Soetomo General Hospital Surabaya, from February - May 2016, using purposive sampling, and obtained 30 patients with chronic heart failure who had met the inclusion and exclusion criteria. All samples were taken from the cardiac outpatients' clinic in Dr. Soetomo General Hospital Surabaya and were willing to participate in the study by signing informed consent. The Health Research Ethics Committee of Dr. Soetomo General Hospital Surabaya has agreed and stated ethical conduct for this research.

Each subject was subjected to a standard echocardiography examination and performed a calculation of left Ventricular Global Longitudinal Strain (GLS) and Tei Index (TI), as well as a calculated SHFM score when a patient is in a cardiac clinic for a regular visit. Data on sample characteristics include gender, age, risk factors, disease diagnosis, and type of therapy.

Data was collected, then managed with descriptive statistics, and presented in the form of exposures, tables, and diagrams. Data were analyzed using inferential statistics to see whether there is a relationship between variables with SPSS. The relationship between left ventricular GLS values and IT with SHFM scores was determined using the Pearson correlation test for normal data distribution or Spearman rank for abnormal data distribution. The results are significant if the value of $p < 0.05$.

3. Results

3.1 Characteristic of research subjects

The subjects obtained in this study were 30 people, consisting of 22 men (73.3%) and 8 women (26.7%). The youngest subject is 41 years, and the oldest subject is 75 years. The highest number of subjects according was in the age category 50-59 years, at 12 people (40%), followed by 60-69 years, 11 people (36.7%), then 40-49 years, 5 people (16.7%), and the least was 70-79 years, 2 people (6.7%).

The most common comorbidity was dyslipidemia with 22 people (73.3%), then hypertension and diabetes had the same number, which was found in 13 people (43.3%), smoking was found in 12

people (40%), 9 people had renal insufficiency (30%), and the least common was obesity (BMI > 30) only found in 1 person (3.3%). The most prevalent etiology of heart failure is the presence of coronary or ischemic heart disease, namely in 27 people (90%), and ischemic, dilatative cardiomyopathy was found in 3 people (10%).

Furthermore, the most given medical therapy to patients was ACEI / ARB group 27 people (90%), then statins (83.3%), beta-blockers 24 people (80%). The administration of diuretics was almost the same between furosemide and spironolactone, 18 people (60%) and 19 people (63.3%). The least administered group of drugs is digoxin 2 people (6.7%), and calcium blockers 1 person (3.3%). The degree of heart failure in the subjects was evaluated based on the *New York Heart Association* (NYHA), most of the subjects experienced NYHA class II in 24 people (80%), then NYHA class III 5 people (16.7%), and NYHA class I 1 person (3.3%).

The characteristics of echocardiography in subjects with the lowest EF were 19%, and the highest was 45%, with an average EF of $36.2\% \pm 8.5$. Diastolic function in all subjects was disrupted, and most were found to be abnormal relaxation, of 17 people (56.7%), then restrictive filling 8 people (26.7%), and pseudonormal 5 people (16.7%).

To determine the correlation between left ventricular GLS and SHFM score, a statistical correlation test was carried out, which began with a distribution test to determine the distribution normality of the left ventricular GLS value, and the SHFM score using seven distribution tests.

3.1.1 *Correlation between Left Ventricle GLS and 1-year SHFM Score.*

The Spearman correlation test was carried out between left ventricular GLS and 1-year SHFM score because one of the data was abnormally distributed. The results of the Spearman correlation test showed a significant, negative, and strong correlation between left ventricular GLS and 1-year SHFM score with $p = 0,0001$ ($p < 0.05$) (Table 2).

3.2 *Correlation between the Tei Index and SHFM Score*

3.2.1 *Correlation between the Tei Index and SHFM Score of 1-Year.*

The Spearman correlation test was carried out between Tei Index and 1-year SHFM score because one of the data was abnormally distributed. The results of the Spearman correlation test showed a significant, positive, and strong correlation between Tei Index and 1-year SHFM score with $p = 0,0001$ ($p < 0.05$) (Table 4).

3.2.2 *Correlation between the Tei Index and SHFM Score of 5 Years.*

The Pearson correlation test was carried out between the left ventricular GLS and the 5-year SHFM score because one of the data was abnormally distributed. Test results of the Pearson correlation showed a significant, positive, and strong correlation between left ventricular GLS and 5-year SHFM score with $p = 0,0001$ ($p < 0.05$) (Table 5)

3.3 *Correlation between Left Ventricular GLS and Tei Index and EF*

In this analysis, the Pearson correlation test was used because all data analyzed were normally distributed. The results of the Pearson correlation test showed a significant, negative, and strong correlation between left ventricular GLS and Tei Index with $p = 0,0001$ ($p < 0.05$), while the correlation between left ventricular GLS and EF was found to be significant, positive, and strong with $p = 0,0001$ ($p < 0.05$) (Table 6).

3.4 *Intraobserver and interobserver variability*

Assessment of intraobserver and interobserver variability was conducted using the Bland Altman method, with several (15 subjects) subjected to quite good suitability, both on left ventricular GLS examination, and TI. There were no significant differences between intraobserver and interobserver ($p > 0.05$).

Table 1. Characteristics of Research Subjects.

Variable		N%	Mean(SD ^a)
Age	40-49	5 (16.7)	58 ± 8.12
	50-59	12 (40)	
	60-69	11 (36.7)	
	70-79	2 (6.7)	
Gender	Male	22 (73.3)	
	Female	8 (26.7)	
Comorbidity	Dyslipidemia	22 (73.3)	
	Hypertension	13 (43.3)	
	Diabetes	13 (43.3)	
	Smoking	12 (40)	
	Renal insufficiency	9 (30)	
	Obesity (BMI ^b >30)	1 (3.3)	
	HF ^c etiology	Ischemia	27 (90)
Others		3 (10)	
Pharmacotherapy	ACEI ^d /ARB ^e	27 (90)	
	Beta-blocker	24 (80)	
	Spirolactone	18 (60)	
	Furosemide	19 (63.3)	
	Calcium blocker	1 (3.3)	
	Statin	24 (80)	
	Digoxin	2 (6.7)	
NYHA	Class I	1 (3.3)	
	Class II	24 (80)	
	Class III	5 (16.7)	
LV Function	EF ^f %, mean (SD)	36.2 (8.5)	
Diastolic function	Abnormal relaxation	17 (56,7)	
	Pseudonormal	5 (16,7)	
	Restrictive filling	8 (26,7)	

^aSD: Standard Deviation; ^bBMI: Body Mass Index; ^cHF; Heart failure; ^dACEI: Angiotensin-converting enzyme inhibitor; ^eARB: Angiotensin-receptor blockers; ^fEF: Ejection Fraction.

Table 2. Correlation Analysis Between Left Ventricular GLS and 1-Year SHFM Score.

	Mean ± SD	Min - Max	r	p*
Left Ventricular GLS	-8,1±3,98	0 - -14,7	-0,676	0,0001
1-Year SHFM	7,3±8,69	2 - 48		

* Analyzed with the Spearman Rho Correlation test

Table 3. Correlation Analysis Between Left Ventricular GLS and 5 Years SHFM Score.

	Mean ± SD	Min - Max	r	p*
Left Ventricular GLS	-8,1±3,98	0 -14,7	-0,676	0,0001
5 Years SHFM	29,6±20,4	8 - 98		

* Analyzed with Pearson Rho Correlation test

Table 4. Correlation Analysis Between Tei Index and 1-Year SHFM Score.

	Mean ± SD	Min - Max	r	p*
Tei Index	0,65±0,14	0,43 - 0,95	0,745	0,0001
1 Year SHFM	7,3±8,69	2 - 48		

* Analyzed with the Spearman Rho Correlation test

Table 5. Correlation Analysis Between Tei Index and 5 Years SHFM Score.

	Mean ± SD	Min - Maks	r	p*
Tei Index	0,65±0,14	0,43 - 0,95	0,738	0,0001
5 Years SHFM	29,6±20,4	8 -98		

* Analyzed with the Pearson Correlation test

Table 6. Correlation Analysis Between Left Ventricular GLS and Tei Index.

	Mean ± SD	Min - Maks	r	p*
Left Ventricular GLS	-8,1±3,98	0 -14,7	-0,763	0,0001
Tei Index	0,65±0,14	0,43 - 0,95		

* Analyzed with Pearson Rho Correlation test

Table 7. Correlation Analysis Between Left Ventricular GLS and EF.

	Mean ± SD	Min - Maks	R	p*
Left Ventricular GLS	-8,1±3,98	0 -14,7	0,678	0,0001
EF	36,2±8,51	19 - 45		

* Analyzed with Pearson Rho Correlation test

4. Discussions

Assessment of prognosis in heart disease is strongly related to left ventricular systolic function, which is generally assessed with EF. Previous studies have shown that left ventricular GLS is superior to EF both in assessing left ventricular function and as a predictor of mortality and cardiovascular events [7-10]. In this study, there was a significant, positive, and strong correlation between left ventricular GLS and EF ($r = 0.678$, $p < 0.001$). These results are consistent with the existing literature [13]. Research conducted by Brown in 62 post-infarct patients received a correlation significant between left ventricular GLS and EF calculated by 3D-MRI (Magnetic Resonance Imaging) [14].

MPI or TI is an index that can globally describe both systolic and diastolic ventricular function, and is easy to do, and is not affected by heart rate, blood pressure, and the severity of mitral valve regurgitation. Some studies also mention that TI can have prognostic value for various heart diseases [15]. In this study, the average value of TI in heart failure patients was 0.65 ± 0.14 , with the lowest value of 0.43 and the highest value of 0.95. The data from this study were not much different from the research conducted by Bruch and his colleagues on 81 patients with heart failure with NYHA class ≤ 2 . The results of the average TI value were 0.60 ± 0.18 , other than that the results of the study stated that TI values > 0.47 could be used to identify the presence of congestive heart failure with a sensitivity of 86% and specificity of 82% [16].

In this study, left ventricular GLS was found to have a significant, negative, and strong correlation with TI ($r = -0.763$, $p < 0.001$). This correlation is in line with a recent study by Sørensen, in 1088 outpatient type 1 diabetes patients who did not have a history of previous heart disease, and where GLS was significantly correlated with TI ($p < 0.001$) [15].

SHFM is a model developed to predict mortality, or survival at 1, 2, and 5 years in patients with chronic heart failure. SHFM consists of various clinical, laboratory, and therapeutic variables, where each of these variables has a prognostic value and has been recognized by AHA as one of the predictive scores in heart failure that has been well validated [5,12]. This study is the first study conducted in Indonesia that correlates between myocardial deformity (left ventricular GLS) and cardiac time interval (TI) markers with prognostic indicators obtained from a multivariable risk assessment tool (SHFM score). In this study, the selected SHFM score became the prediction of 1-year and 5-year mortality because it was made more aware of mortality or 1-year survival, and it is essential in clinical practice considerations in determining candidate patients who deserve further therapy such as ICD / CRT-D installation. Or even determine the heart transplant candidate. Five-year mortality prediction is used to assess predictions of mortality over a longer period so that it can be used when educating and explaining disease status in patients.

In this study, there was a significant, negative, and strong correlation between left ventricular GLS and 1-year, and 5-year SHFM mortality scores with both having the same values, with $r = -0.676$, $p < 0.001$. The results of this study are in line with the research conducted by Rangel in 54 chronic heart failure patients with systolic dysfunction ($EF \leq 45\%$), which obtained a significant correlation between left ventricular GLS and SHFM score which estimated the value of life expectancy ($r = -0.41$, $p = 0.002$) [9]. The results of this research also support the cohort study of 1,065 HF rEF patients with an observation period of 22-57 months and a median of 40 months conducted by Sengeløv and colleagues. The results showed that GLS left ventricle was an independent predictor of mortality after adjusting the age, sex, BMI, total cholesterol, mean arterial pressure, heart rate, ischemic cardiomyopathy, percutaneous transluminal coronary angioplasty, coronary artery bypass graft surgery, noninsulin-dependent diabetes mellitus, and conventional echocardiographic parameters (hazard ratio (HR): 1.15, 95% confidence interval (CI)) [17]. In this study, a significant, positive, and strong correlation was found between Tei Index and 1-year SHFM score ($r = 0.745$, $p < 0.001$), and 5-year SHFM score ($r = 0.738$, $p < 0.001$). This research is the first study to analyze the correlation between TI and SHFM scores, which is a model prognostic score consisting of various validated

prognostic variables. There have been several previous studies that analyzed TI correlation with other prognostic variables such as EF, BNP, and NYHA, and all found significant correlations [18,19].

Harjai et al., investigated the prognostic value of TI in heart failure patients with EF <30%, followed by a period of 24 ± 19 months, assessed as the endpoint of death due to any cause, and heart transplantation. During the study, 28 patients died (49%), and two patients (3.5%) underwent a heart transplant. A strong correlation was found between TI values >1.14 and long-term outcomes and proved independent of other clinical and echocardiographic variables that have been shown to have prognostic value [20]. Sørensen et al. researched the general population and got results showing that TI is a significant prognosticator ($p < 0.05$) for Major Adverse Cardiac Event (MACE) [15].

Some limitations of this study are that the results of this study are supported by a model prognostic score without prospective cohort. Hence, further validation is still required. However, despite SHFM being a well-validated prognostic model score in patients with chronic heart failure, outpatient care, based on a variety of variables that have been known to have prognostic value in patients with heart failure. In this study, STE-2D strains evaluated were only assessed retrospectively, and both radial and circumferential strains were not assessed.

5. Conclusion

GLS-left ventricle had a significant negative correlation with an estimated one and five years mortality based on SHFM score, and it is suggested that GLS-left ventricle and TI can be a prognostic factor.

References

- [1] Bui A L, Horwich T B, and Fonarow G C 2011 Epidemiology and risk profile of heart failure *Nat. Rev. Cardiol.* **8** 30–41
- [2] Go A S, et al 2013 Heart disease and stroke statistics—2013 update *Circulation* **127**
- [3] Ponikowski P, et al 2014 Heart failure: preventing disease and death worldwide *ESC Hear. Fail.* **1** 4–25
- [4] McMurray J J V, et al 2012 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: the task force for the diagnosis and treatment of acute and chronic heart failure 2012 of the European society of cardiology. Developed in collaboration with the Heart *Eur. Heart J.* **33** 1787–847
- [5] Yancy C W, et al 2013 ACCF/AHA Guideline for the management of heart failure *Circulation* **128**
- [6] Curtis J P, et al 2003 The association of left ventricular ejection fraction, mortality, and cause of death in stable outpatients with heart failure. *J. Am. Coll. Cardiol.* **42** 736–42
- [7] Motoki H, Borowski A G, Shrestha K, Troughton R W, Tang W H W, Thomas J D and Klein A L 2012 Incremental prognostic value of assessing left ventricular myocardial mechanics in patients with chronic systolic heart failure *J. Am. Coll. Cardiol.* **60** 2074–81
- [8] Nahum J, et al 2010 Impact of longitudinal myocardial deformation on the prognosis of chronic heart failure patients *Circ. Cardiovasc. Imaging* **3** 249–56
- [9] Rangel I, Gonçalves A, de Sousa C, Almeida P B, Rodrigues J, Macedo F, Silva Cardoso J and Maciel M J 2014 Global longitudinal strain as a potential prognostic marker in patients with chronic heart failure and systolic dysfunction *Rev. Port. Cardiol.* **33** 403–9
- [10] Stanton T, Leano R and Marwick T H 2009 Prediction of all-cause mortality from global longitudinal speckle strain *Circ. Cardiovasc. Imaging* **2** 356–64
- [11] Ketchum E S and Levy W C 2011 Multivariate risk scores and patient outcomes in advanced heart failure *Congest. Hear. Fail.* **17** 205–12
- [12] Levy W C, et al 2006 The Seattle Heart Failure Model *Circulation* **113** 1424–33
- [13] Lang R M, et al 2015 Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American society of echocardiography and the European association of cardiovascular imaging *J. Am. Soc. Echocardiogr.* **28** 1–39.e14

- [14] Brown J, Jenkins C and Marwick T H 2009 Use of myocardial strain to assess global left ventricular function: a comparison with cardiac magnetic resonance and 3-dimensional echocardiography *Am. Heart J.* **157** 102.e1-102.e5
- [15] Biering-Sørensen T, Mogelvang R, Schnohr P and Jensen J S 2016 Cardiac time intervals measured by tissue Doppler imaging M-mode: association with hypertension, left ventricular geometry, and future ischemic cardiovascular diseases *J. Am. Heart Assoc.* **5**
- [16] Bruch C 2000 TEI-Index in patients with mild-to-moderate congestive heart failure *Eur. Heart J.* **21** 1888–95
- [17] Sengeløv M, Jørgensen P G, Jensen J S, Bruun N E, Olsen F J, Fritz-Hansen T, Nochioka K, and Biering-Sørensen T 2015 Global longitudinal strain is a superior predictor of all-cause mortality in heart failure with reduced ejection fraction *JACC Cardiovasc. Imaging* **8** 1351–9
- [18] Mikkelsen K V., Møller J E, Bie P, Ryde H, Videbaek L, and Haghfelt T 2006 Tei index and neurohormonal activation in patients with incident heart failure: serial changes and prognostic value *Eur. J. Heart Fail.* **8** 599–608
- [19] Ogunmola O J, Akintomide A O and Olamoyegun A M 2013 Relationship between clinically assessed heart failure severity and the TEI index in Nigerian patients *BMC Res. Notes* **6** 488
- [20] Harjai K J, Scott L, Vivekananthan K, Nunez E, and Edupuganti R 2002 The Tei index: a new prognostic index for patients with symptomatic heart failure. *J. Am. Soc. Echocardiogr.* **15** 864–8