



FOLIA MEDICA INDONESIA

[UNIVERSITAS AIRLANGGA](#)

✳ P-ISSN : 23558393 <> E-ISSN : 2599056X 📁 Subject Area : Education



**1.06667**  
Impact Factor



**1528**  
Google Citations



**Sinta 2**  
Current  
Accreditation

[Google Scholar](#)

[Garuda](#)

[Website](#)

[Editor URL](#)

History Accreditation

2018

2020

2021

2022

2023

2024

2025

[Garuda](#)

[Google Scholar](#)

[Proportions of Group B Streptococcus Isolation from Pregnant Womenâ€™s Vaginal and Rectal Swab Specimens at a Tertiary Hospital in Surabaya, Indonesia](#)

Universitas Airlangga [Folia Medica Indonesiana Vol. 59 No. 1 \(2023\): March 46-50](#)

📅 2023 [DOI: 10.20473/fmi.v59i1.11708](#) [Accred : Sinta 2](#)

[Analysis of Spatial Working Memory Using the Y-Maze on Rodents Treated with High-Calorie Diet and Moderate-Intensity Exercise](#)

Universitas Airlangga [Folia Medica Indonesiana Vol. 59 No. 1 \(2023\): March 40-45](#)

📅 2023 [DOI: 10.20473/fmi.v59i1.32672](#) [Accred : Sinta 2](#)

[The Effect of Binahong \(Anredera cordifolia \(Ten.\) Steenis\) Leaf Ethanolic Extract on the Reduction of Blood Uric Acid Levels in Hyperuricemic Male White Wistar Rats \(Rattus norvegicus\)](#)

Universitas Airlangga [Folia Medica Indonesiana Vol. 59 No. 1 \(2023\): March 20-25](#)

📅 2023 [DOI: 10.20473/fmi.v59i1.36052](#) [Accred : Sinta 2](#)

[Five Single-Nucleotide Polymorphisms in the PITX2 Gene as Risk Factors for Atrial Fibrillation](#)

Universitas Airlangga [Folia Medica Indonesiana Vol. 59 No. 1 \(2023\): March 85-91](#)

📅 2023 [DOI: 10.20473/fmi.v59i1.37126](#) [Accred : Sinta 2](#)

[The Histopathological Features of Syphilis and Its Mimickers](#)

Universitas Airlangga [Folia Medica Indonesiana Vol. 59 No. 1 \(2023\): March 63-69](#)

📅 2023 [DOI: 10.20473/fmi.v59i1.37286](#) [Accred : Sinta 2](#)

[Back Matter Vol.59 No.1 \[3/2023\]](#)

p-ISSN 2355-8393

e-ISSN 2599-056X

# FOLIA MEDICA INDONESIANA

Vol. 55 No. 1 March 2019

**CORRELATION BETWEEN BACTERIOSTATIC EFFECT AND BACTERICIDE WITH ANTIBIOFILM AND ANTICOLONY SPREADING FROM JAVANESE CITRONELLA OIL ON METHICILLIN-RESISTANT *Staphylococcus aureus* (MRSA)**

(Amaliyah Nurul Hidayah, Eddy Bagus Wasito, Kartuti Debora, Achmad Basori, Isnaeni, Budi Utomo)

**ANALYSIS OF ANC LEVELS AFTER FILGRASTIM THERAPY IN ACUTE LEUKEMIA CHILDREN WITH NEUTROPENIA**

(Reta Anggraeni Widya, Susanto Nugroho, Sri Winarsih, Yulistiani)

**EMULGEL FORMULATION OF PURIFIED EXTRACT OF MORINGA (*Moringa oleifera* L.) LEAF**

(Tutik Wuryandari, Nining Sugihartini, Kintoko)

**INCIDENCE OF EMERGENCE AGITATION IN PEDIATRIC PATIENT AFTER GENERAL ANESTHESIA**

(Lucky Andriyanto, Arie Utariani, Elizeus Hanindito, Kohar Hari Santoso, Hamzah, Eka Ari Puspa)

**MICROBIOLOGICAL ASSESSMENT OF FRESH EXPRESSED BREAST MILK AT ROOM TEMPERATURE IN DR. SOETOMO HOSPITAL NEONATAL UNIT**

(Nur Aisyah Widjaja, Kartika Hardiyani, Meta Herdiana Hanindita, Roedi Irawan)

**DISEASE FREE SURVIVAL OF STAGE I ENDOMETRIAL CANCER AFTER SURGERY WITH OR WITHOUT ADJUVANT TREATMENT**

(Woraluk Moradokkasem, Nungrutai Saealb, Tippawan Liabsuetrakul)

**LACTATE LEVEL AS A PREDICTION FACTOR OF REPERFORATION AFTER REPAIRING GASTRIC PERFORATION**

(Vicky Sumarki Budipramana)

**ASSESSMENT OF RAPID IMMUNOCHROMATOGRAPHIC TEST AS A DIAGNOSTIC TOOL FOR NOROVIRUS RELATED DIARRHEA IN CHILDREN**

(Reza Gunadi Ranuh, Alpha Fardah Athiyah, Deanty Ayu PA, Andy Darma, Dadik Raharjo, et al)

**PROFILE OF DRUG HYPERSENSITIVITY PATIENTS HOSPITALIZED IN DR. SOETOMO HOSPITAL, SURABAYA, INDONESIA. PRELIMINARY DATA OF 6 MONTHS OBSERVATION**

(Nur Moya Isyroqiyah, Gatot Soegiarto, Yuan Setiawati)

**CORRELATION OF LOWER LIMB MUSCLES AND BODY MASS INDEX WITH BODY BALANCE IN THE ELDERLY**

(Natasya Valentina, Patricia M Kurniawati, Margarita M Maramis)

**THE EFFECT OF HYPERCHOLESTEROLEMIA ON CORTICAL BONE THICKNESS OF WISTAR RATS (*Rattus norvegicus*)**

(Auliya Dzaqiyatus Sofka, Pudji Lestari, Gadis Melnar Sari)

**Case report :**

**THROMBOCYTOPENIA IN A PATIENT UNDERGOING PRIMARY PERCUTANEOUS CORONARY INTERVENTION**

(Yudi Her Oktaviono, Feranti Meuthia)

**Case Report :**

**BLOWOUT FRACTURE COMPLEX TYPE**

(Fauzi Helmi, Boedy Setya Santoso)

International Online Distribution by ProQuest™  
www.proquest.com

Folia Medica Indonesiana	Vol. 55	No. 1	Page 1-	Surabaya March 2019	p-ISSN: 2355-8393 e-ISSN: 2599-056X
-----------------------------	---------	-------	---------	------------------------	--

# FOLIA MEDICA INDONESIANA

Vol. 55 No. 1 March 2019

**CORRELATION BETWEEN BACTERIOSTATIC EFFECT AND BACTERICIDE EFFECT WITH ANTIBIOFILM AND ANTICOLONY SPREADING FROM JAVANESE CITRONELLA OIL ON METHICILLIN-RESISTANT *Staphylococcus aureus* (MRSA)**

(Amaliyah Nurul Hidayah, Eddy Bagus Wasito, Kartuti Debora, Achmad Basori, Isnaeni, Budi Utomo)

**ANALYSIS OF ANC LEVELS AFTER FILGRASTIM THERAPY IN ACUTE LEUKEMIA CHILDREN WITH NEUTROPENIA**

(Reta Anggraeni Widya, Susanto Nugroho, Sri Winarsih, Yulistiani)

**EMULGEL FORMULATION OF PURIFIED EXTRACT OF MORINGA (*Moringa oleifera* L.) LEAF**

(Tutik Wuryandari, Nining Sugihartini, Kintoko)

**INCIDENCE OF EMERGENCE AGITATION IN PEDIATRIC PATIENT AFTER GENERAL ANESTHESIA**

(Lucky Andriyanto, Arie Utariani, Elizeus Hanindito, Kohar Hari Santoso, Hamzah, Eka Ari Puspita)

**MICROBIOLOGICAL ASSESSMENT OF FRESH EXPRESSED BREAST MILK AT ROOM TEMPERATURE IN DR. SOETOMO HOSPITAL NEONATAL UNIT**

(Nur Aisyah Widjaja, Kartika Hardiyani, Meta Herdiana Hanindita, Roedi Irawan)

**DISEASE FREE SURVIVAL OF STAGE I ENDOMETRIAL CANCER AFTER SURGERY WITH OR WITHOUT ADJUVANT TREATMENT**

(Woraluk Moradokkasem, Nungrutai Saeai, Tippawan Liabsuetrakul)

**LACTATE LEVEL AS A PREDICTION FACTOR OF REPERFORATION AFTER REPAIRING GASTRIC PERFORATION**

(Vicky Sumarki Budipramana)

**ASSESSMENT OF RAPID IMMUNOCHROMATOGRAPHIC TEST AS A DIAGNOSTIC TOOL FOR NOROVIRUS RELATED DIARRHEA IN CHILDREN**

(Reza Gunadi Ranuh, Alpha Fardah Athiyyah, Deanty Ayu PA, Andy Darma, Dadik Raharjo, et al)

**PROFILE OF DRUG HYPERSENSITIVITY PATIENTS HOSPITALIZED IN DR. SOETOMO HOSPITAL, SURABAYA, INDONESIA. PRELIMINARY DATA OF 6 MONTHS OBSERVATION**

(Nur Moya Isyroqiyyah, Gatot Soegiarto, Yuani Setiawati)

**CORRELATION OF LOWER LIMB MUSCLES AND BODY MASS INDEX WITH BODY BALANCE IN THE ELDERLY**

(Natasya Valentina, Patricia M Kurniawati, Margarita M Maramis)

**THE EFFECT OF HYPERCHOLESTEROLEMIA ON CORTICAL BONE THICKNESS OF WISTAR RATS (*Rattus norvegicus*)**

(Auliya Dzaqiyatus Sofka, Pudji Lestari, Gadis Meinar Sari)

**Case report:**

**THROMBOCYTOPENIA IN A PATIENT UNDERGOING PRIMARY PERCUTANEOUS CORONARY INTERVENTION**

(Yudi Her Oktaviono, Feranti Meuthia)

**Case Report:**

**BLOWOUT FRACTURE COMPLEX TYPE**

(Fauzi Helmi, Boedy Setya Santoso)

International Online Distribution by ProQuest™  
www.proquest.com

Folia Medica Indonesiana	Vol. 55	No. 1	Page 1-	Surabaya March 2019	p-ISSN: 2355-8393 e-ISSN: 2599-056X
-----------------------------	---------	-------	---------	------------------------	--

# **FOLIA MEDICA INDONESIA**

**p-ISSN 2355-8393, e-ISSN 2599-056X**

**Vol. 55 no. 1 March 2019**

---

**Medical journal, published by Airlangga University School of Medicine, Surabaya, publishing original basic medical and clinical articles presented as research articles and review articles**

## **EDITOR-IN CHIEF**

**Ni Made Mertaniasih**, Department of Medical Microbiology, Faculty of Medicine,  
Universitas Airlangga, Surabaya, Indonesia

## **DEPUTY EDITORS**

**Muhammad Miftahussurur**, Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga, Indonesia

## **EDITORIAL BOARD**

**Arend Frederik Bos**, Division Neonatology, Faculty of Medical Sciences, University of Groningen, Netherlands

**Hiroaki Kimura**, Department of Physical Medicine and Rehabilitation, Hiroshima University Hospital, Japan

**Jitti Hanprasertpong**, Division of Gynecologic Oncology, Department of Obstetrics and Gynecology,  
Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla, Thailand

**Anucha Thatrimontrichai**, Prince of Songkla University, Thailand

**Surasak Sangkhathat**, Pediatric Surgery Unit, Department of Surgery, Prince of Songkla University,  
Songkhla, Thailand

**Aryati**, Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga, Indonesia

**Viskasari Pintoko Kalanjati**, Department of Anatomy and Histology, Faculty of Medicine,  
Universitas Airlangga, Surabaya, Indonesia

**Irwanto**, Department of Pediatrics, Faculty of Medicine, Universitas Airlangga, Indonesia

**Wihasto Suryaningtyas**, Department of Neurosurgery, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

**Asra Al Fauzi**, Department of Neurosurgery, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

**Azimatul Karimah**, Department of Psychiatry, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

**Lucky Prasetiowati**, Department of Anatomy & Histology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

**Bambang Purwanto**, Department of Medical Physiology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

**Arni Kusuma Dewi**, Department of Anatomy & Histology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

## **MANAGING EDITOR**

**Ahmad Suryawan**, Department of Pediatrics, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

## **PRODUCTION EDITORS**

**Moch. Zuhdy**, Center for Medical Science Community, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

**Athfiyatul Fatati**, Center for Medical Science Community, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

**Alviana Nur Afifah**, Center for Medical Science Community, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

Published by : **GRAMIK (Graha Masyarakat Ilmiah Kedokteran)**  
(Center for Medical Science Community)  
Quarterly (March, June, September, and December)

Address : **Airlangga University School of Medicine**  
Jl. Prof dr Moestopo 47 Surabaya 60131  
Phone: 62-31-5013749, 5020251-3 ext. 135  
Fax : 62-31-5013749, 62-31-5022472  
E-mail: fmi@journal.unair.ac.id, foliamedica@gmail.com

Accredited no. 2/E/KPT/2015

## CONTENTS

<b>CORRELATION BETWEEN BACTERIOSTATIC AND BACTERICIDE EFFECT WITH ANTIBIOFILM AND ANTICOLONY SPREADING FROM JAVANESE CITRONELLA OIL ON METHICILLIN-RESISTANT <i>Staphylococcus aureus</i> (MRSA)</b> (Amaliyah Nurul Hidayah, Eddy Bagus Wasito, Kartuti Debora, Achmad Basori, Isnaeni, Budi Utomo)	1 – 9
<b>ANALYSIS OF ANC LEVELS AFTER FILGRASTIM THERAPY IN ACUTE LEUKEMIA CHILDREN WITH NEUTROPENIA</b> (Reta Anggraeni Widya, Susanto Nugroho, Sri Winarsih, Yulistiani)	10 – 16
<b>EMULGEL FORMULATION OF PURIFIED EXTRACT OF MORINGA (<i>Moringa oleifera</i> L.) LEAF</b> (Tutik Wuryandari, Nining Sugihartini, Kintoko)	17 – 24
<b>INCIDENCE OF EMERGENCE AGITATION IN PEDIATRIC PATIENT AFTER GENERAL ANESTHESIA</b> (Lucky Andriyanto, Arie Utariani, Elizeus Hanindito, Kohar Hari Santoso, Hamzah, Eka Ari Puspita)	25 – 29
<b>MICROBIOLOGICAL ASSESSMENT OF FRESH EXPRESSED BREAST MILK AT ROOM TEMPERATURE IN DR. SOETOMO HOSPITAL NEONATAL UNIT</b> (Nur Aisyah Widjaja, Kartika Hardiyani, Meta Herdiana Hanindita, Roedi Irawan)	30 – 36
<b>DISEASE FREE SURVIVAL OF STAGE I ENDOMETRIAL CANCER AFTER SURGERY WITH OR WITHOUT ADJUVANT TREATMENT</b> (Woraluk Moradokkasem, Nungrutai Saeai, Tippawan Liabsuetrakul)	37 – 42
<b>LACTATE LEVEL AS A PREDICTION FACTOR OF REPERFORATION AFTER REPAIRING GASTRIC PERFORATION</b> (Vicky Sumarki Budipramana)	43 – 47
<b>ASSESSMENT OF RAPID IMMUNOCHROMATOGRAPHIC TEST AS A DIAGNOSTIC TOOL FOR NOROVIRUS RELATED DIARRHEA IN CHILDREN</b> (Reza Gunadi Ranuh, Alpha Fardah Athiyyah, Deanty Ayu PA, Andy Darma, Dadik Raharjo, et al)	48 – 53
<b>PROFILE OF DRUG HYPERSENSITIVITY PATIENTS HOSPITALIZED IN DR. SOETOMO HOSPITAL, SURABAYA, INDONESIA. PRELIMINARY DATA OF 6 MONTHS OBSERVATION</b> (Nur Moya Isyroiyyah, Gatot Soegiarto, Yuani Setiawati)	54 – 57
<b>CORRELATION OF LOWER LIMB MUSCLES AND BODY MASS INDEX WITH BODY BALANCE IN THE ELDERLY</b> (Natasya Valentina, Patricia M Kurniawati, Margarita M Maramis)	58 – 62
<b>THE EFFECT OF HYPERCHOLESTEROLEMIA ON CORTICAL BONE THICKNESS OF WISTAR RATS (<i>Rattus norvegicus</i>)</b> (Auliya Dzaqiyatus Sofka, Pudji Lestari, Gadis Meinar Sari)	63 – 67
<b>Case report: THROMBOCYTOPENIA IN A PATIENT UNDERGOING PRIMARY PERCUTANEOUS CORONARY INTERVENTION</b> (Yudi Her Oktaviono, Feranti Meuthia)	68 – 73
<b>Case Report: BLOWOUT FRACTURE COMPLEX TYPE</b> (Fauzi Helmi, Boedy Setya Santoso)	74 – 81

## INCIDENCE OF EMERGENCE AGITATION IN PEDIATRIC PATIENT AFTER GENERAL ANESTHESIA

**Lucky Andriyanto, Arie Utariani, Elizeus Hanindito, Kohar Hari Santoso, Hamzah, Eka Ari Puspita**

Department of Anesthesiology and Reanimation, Faculty of Medicine, Universitas Airlangga, Dr Soetomo Hospital, Surabaya, Indonesia

### ABSTRACT

*Post anesthesia agitation is common problem in pediatric post anesthesia care unit. The incidences range from 10 to 80%. EA has been described as a dissociated state of consciousness in which the child is inconsolable, irritable, and uncooperative typically thrashing, crying, moaning or incoherent. This study was done to determine the incidence of emergence agitation and associated risk factors in pediatric patients who underwent general anesthesia. This descriptive and analytic study was performed on 105 pediatric patient aged 1-12 years that underwent general anesthesia for various elective diagnostic and surgeries at Dr. Soetomo Hospital between January and February 2016. The presence of emergence agitation was recorded using Pediatric Anesthesia Emergence Delirium (PAED) scale. The factors that linked with Emergence Agitation were recorded in a questionnaire. The data were analyzed using SPSS software with logistic regression. p-values less than 0.05 were considered as significant. Forty two (40%) children had Emergence Agitation. Preoperative anxiety (p=0.006) and Pain (p=0.035) were associated with higher rates of post anesthetic emergence agitation. This study identified preoperative anxiety and pain as risk factors, which are associated with emergence agitation in children. To minimize the incidence of post anesthetic emergence agitation, these risk factors should be considered in the routine care by anesthetist.*

**Keywords:** General anesthesia; pediatric; emergence agitation

### ABSTRAK

*Agitasi pada pediatri paska anestesi umum adalah permasalahan yang banyak terjadi di ruang paska anestesi pada anak. Insiden berkisar antara 10-80 %. EA di deskripsikan sebagai suatu stadium disosiatif di mana anak akan sulit ditenangkan, iritabel, tidak kooperatif, menangis, menendang dan berbicara tidak koheren. Penelitian ini bertujuan untuk menentukan angka kejadian EA dan faktor resiko yang mempengaruhi EA pada pasien pediatri paska anestesi umum. Penelitian ini bersifat deskriptif analitik dengan 105 sampel pasien pediatri usia 1-12 tahun yang menjalani anesthesia umum untuk tindakan pembedahan dan diagnostik di RSUD Dr. Soetomo antara Januari dan Februari 2016. Kejadian EA ditentukan dengan skala PAED. faktor faktor yang dihubungkan dengan EA direkam dalam suatu kuisisioner. Data dianalisa dengan SPSS dengan metode regresi logistik. Faktor yang memiliki nilai p<0,05 dinyatakan signifikan. Empat puluh dua (40%) anak mengalami EA. Kecemasan preoperatif (p=0.006) dan nyeri (p=0.035) secara signifikan berhubungan dengan tingginya kejadian EA. Penelitian ini menemukan kecemasan preoperatif dan nyeri sebagai faktor resiko yang signifikan pada terjadinya EA. Untuk meminimalkan kejadian EA, faktor ini harus diperhatikan oleh setiap ahli anestesi.*

**Kata kunci:** Anestesi umum; pediatri; emergence agitation

**Correspondence:** Lucky Andriyanto, Department of Anesthesiology and Reanimation, Faculty of Medicine, Universitas Airlangga, Dr Soetomo Hospital, Surabaya, Indonesia. E-mail: lky.anest@gmail.com

pISSN:2355-8393 • eISSN: 2599-056x • doi: <http://dx.doi.org/10.20473/fmi.v55i1.12546>  
 • Fol Med Indones. 2019;55:25-29 • Received 21 Nov 2017 • Accepted 22 Mar 2018  
 • Open access under CC-BY-NC-SA license • Available at <https://e-journal.unair.ac.id/FMI/>

## INTRODUCTION

Emergence agitation (EA) is a common problem among postoperative pediatric patients. Emergence agitation, which frequently used interchangeably with emergence delirium (ED) had been defined as a dissociated state of consciousness in which the child is restless,

inconsolable, irritable, and uncooperative typically thrashing, kicking, crying, moaning or incoherent. (Jildenstål et al 2014, Vljakovic & Sindjelic 2007). Factors can affect agitation after surgery, are age, anxiety, pain, anesthetics, surgery type, perioperative drugs or physiologic compromise and the incidences of EA range from 10 to 80%, (Vljakovic & Sindjelic

2007, Voepel-Lewis et al 2003, Mohkamkar et al 2014). Emergence agitation usually occurs within 30 minutes post anesthesia. Although most of EA resolve spontaneously, it can cause harm to the child such as fall, bleed, self-injury, or parent distress. This descriptive analytical study was done to determine the incidence of emergence agitation and associated risk factors in pediatric patients who underwent general anesthesia.

## MATERIALS AND METHODS

With the approval from ethical committee and parental consent, all healthy 105 children (1-12 yrs. old) underwent general anesthesia for elective surgical or diagnostic procedures in one month and met inclusion criteria were included. All patient should categorized in physical status ASA 1-2, did not have preoperative awareness disturbances, the surgery last less than 30 minutes, severe post operative pain, or sedated and ventilated mechanically after anesthesia. There was no drop out sample in this study. All perianesthesia routine care was performed under supervision of an anesthesiologist. At pre anesthesia period, vital signs, and perioperative anxiety measured using Modified Yale Preoperative Anxiety Scale (mYPAS) were recorded. After induction, induction-score was measured using Induction Compliant Checklist (ICC) scale. Duration of anesthesia, time of awakening were recorded. In recovery room emergence agitation was measured using valid Pediatric Anesthesia Emergence Delirium (PAED) scales at 15 minutes and 30 minutes. The Face-Legs-Activity-Cry-Consolability (FLACCS) scale also measured at the same time. All data recorded was presented by descriptive cross table and correlation of factors influence EA incidences was analyzed by logistic regression.

## RESULTS

105 children aged 1-12 years with ASA class I-II were enrolled over one month (January to February 2016). Characteristic of samples / patients then performed by cross tab (table 1). Age of the sample differ to pre school (1 to 7 yrs. old) age and school age. (7 to 12 years old)

Some of the samples (66.7%, n=70) had preoperative anxiety (mYPAS >30). Almost all of the samples (93%, n=98) premedicated with intravenous midazolam. Induction compliance of the samples was good. 80% samples (n=84) had ICC score=5.

Duration of anesthesia of samples varies between 0.5 hour to 4.5 hours and the mean of anesthesia duration was 3.6 hours. Time of awakening was duration since anesthetics stopped until the first time the sample could contact with the anesthetist. Most of the sample awake less than 15 minutes (83.9 %, n=88). The distribution of time of awakening showed in table 3. Post of pain measured using FLACC scale, and differs in 3 categories. Moderate postoperative pain (FLACC 4-6) was found in 21.9% (n=23) samples, mild pain (FLACC 1-3) found in 31.4% (n=33) and no pain (FLACC 0) in 46.7% (n=49).

Table 1. Sample characteristic

Characteristic of samples	Number of Samples	%
Age		
Preschool	67	63.8
School	38	36.2
Sex		
Male	60	57.1
Female	45	42.9
Type of Surgery		
Lower extremity	9	8.6
Upper extremity	3	2.9
Head and neck	26	24.8
Abdominal	12	11.4
Burn	5	4.8
Urogenital	13	12.4
Diagnostic	37	35.2
PS ASA		
1	15	14.3
2	90	85.7

Incidence of emergence agitation (PAED score >10) among samples was high. 40% of samples (n=42) was had EA. Table 4 presents the relationship between perioperative factors and the presence of EA. Using logistic regression analysis, it was founded that perioperative anxiety [(p<0.05 (OR: 4.188, CI 95% (1.5-11.6))] and postoperative pain [(p<0.05 (OR: 1.797, CI 95% (1.0-3.1))] had significant correlation with the presence of emergence agitation among samples.

## DISCUSSION

This study found the incidence of emergence agitation (PAED >10) in pediatric patient who underwent elective surgery was 40 %. PAED scale is reliable and valid tool to measure the presence of emergence agitation, and 10 was the most specific and sensitive cut of point. (Sikich

& Lerman 2004). This incidence was higher than the result found by Voepel-Lewis et al. 2003 and Mohkamkar et al. 2014 that was 18% and 17.9%. But our result was still in range 10-80% as the finding of some studies. (Nasr & Hannalah 2011).

Some studies shows some perioperative factors that correlated to emergence agitation are post-operative pain, short time awaking, anesthetic drugs as well as site of operation. (Mohkamkar et al. 2014; Singh et al. 2012; Voepel-Lewis et al. 2003). Our finding shows perioperative factors that correlated to the incidence of emergence of agitation were perioperative anxiety [(p<0.05 (OR:4.188, CI 95% (1.5-11.6))] and postoperative pain [(p<0.05 (OR : 1.797, CI 95% (1.0-3.1))].

Our study used Modified Yale Preoperative Anxiety Scale (mYPAS) as a tool to assess perioperative anxiety of the patients. It was measured in preparation area by anesthesiologist when the parents was still accompany the patient. mYPAS scale measured by attending anesthesiologist was accurate to assess perioperative e anxiety (Maclaren et al. 2010). Study by (Kain et al. 2004) shown that marked increase in mYPAS score correlated linearly to the increase of emergence agitation.

Pain was proven as a factor correlated to the incidence of EA. Some studies showed that anesthesia for non painful procedures with sevoflurane anesthesia has higher incidence of EA than the same procedures with halothane or propofol. These findings suggest that EA/ED may be a clinical phenomenon that is separate from pain. (Vlajkovic & Sindjelic 2007, Mohkamkar et al 2014) In this study we excluded patients with severe pain (FLACCS >7). Severe pain was the most confounding variable when assessing a child’s behavior upon emergence because of the overlapping clinical picture with EA/ED. (Vlajkovic & Sindjelic 2007) (Mohkamkar et al. 2014) . Adequate pain management would reduce incidence of EA but would not eliminated completely. (Mohkamkar et al 2014).

In our study age, physical status, induction score, type of surgery and time of awakening not statistically significant correlated to emergence agitation.

Aside to concern some predictor factor, it is important to take steps to prevent EA. The role of premedication of midazolam as an EA preventive measure remains a matter of debate. Intravenous administration of midazolam just before the end of surgery proved to reduces emergence agitation (Lee & Ph 2015). Intravenous anesthesia with propofol mixed with ketamin or injection of propofol mixed with ketamin

before completion of anesthesia also showed that EA incidence was lower than control group. (Rizk & Samir 2014; Angheliescu et al. 2011). But other study showed that the concurrent use of midazolam or propofol did not reduce the incidence of emergence agitation but did delay emergence and recovery (Cohen et al. 2002). Fentanyl and dexmedetomidine injection before end of surgery also shown reduces EA. (Abdelhalim & Alarfaj 2013, Makkar et al 2016).

Emergence agitation management includes nonpharmacological and pharmacological approaches. Non-pharmacological approaches include measures to ensure comfort and safety of patients such as motivation, elution, sling that can be done by health workers and parents. The most common pharmacological approach is the administration of midazolam, propofol and opioid such as fentanyl. (Wong & Bailey 2015)

Table 2. Perioperative factors

Perioperative factors	Number of Samples	%
Preoperative Anxiety		
mYPAS <30	35	33.3
mYPAS >30	70	66.7
Induction compliance		
ICC ≤ 5	95	90.5
ICC > 5	10	9.5
Premedication		
Pre medicated	98	93.3
Not pre medicated	7	6.7
Time of awakening (minutes)		
5	17	16.2
10	30	28.6
15	41	39.1
20	13	12.4
25	1	1.0
30	3	2.9
Post operative pain		
No pain	49	46.7
Mild	33	31.4
Moderate	23	21.9

Table 3. Incidence of emergence agitation

Emergence Agitation	Number of Samples	%
No EA	63	60
EA	42	40



Table 4. incidence of emergence agitation relative to perioperative factors

Variable	EA	%	p Value	Relative risk (95% CI)
Perioperative Anxiety			0.006	4.188 (1.5-11.6)
mYPAS <30	36	85.7		
mYPAS >30	6	14.3		
Post operative pain			0.035	1.797 (1.0-3.1)
No pain	14	33.3		
Mild	13	31		
Moderate	15	35.7		
Age			0.868	NA
Preschool	14	33.3		
School	28	66.7		
Physical status			0.134	NA
1	3	7.1		
2	39	92.9		
Induction compliance			0.253	NA
ICC ≤ 5	35	83.3		
ICC > 5	7	16.7		
Type of surgery			0.932	NA
Lower extremity	5	11.9		
Upper extremity	0	0		
Head and neck	11	26.2		
Abdominal	7	16.7		
Burn	3	7.1		
Urogenital	5	11.9		
Diagnostic	11	26.2		
Time of awakening			0.557	NA
5	5	11.9		
10	13	31		
15	16	38.1		
20	5	11.9		
25	0	0		
30	3	7.1		

## CONCLUSION

This study has found the incidence of emergence agitation among pediatric patient who underwent elective surgery was 40%. Perioperative factors that correlated to the presence of emergence agitation were preoperative anxiety and postoperative pain. Thus anesthetist should consider these risk factors in anesthesia routine care.

## REFERENCES

Abdelhalim A, Alarfaj A (2013). The effect of ketamine versus fentanyl on the incidence of emergence agitation after sevoflurane anesthesia in pediatric patients undergoing tonsillectomy with or without adenoidectomy. *Saudi Journal of Anaesthesia* 7, 392

Angheltescu DL, et al (2011). Prevention of emergence agitation in seven children receiving low-dose

ketamine and propofol total intravenous anesthesia. *AANA Journal* 79, 238-242

Cohen IT, Drewsen S, Hannallah RS (2002). Propofol or midazolam do not reduce the incidence of emergence agitation associated with desflurane anaesthesia in children undergoing adenotonsillectomy. *Paediatric Anaesthesia* 12, 604-609

Jildenstål PK, et al (2014). Routines for reducing the occurrence of emergence agitation during awakening in children, a national survey. *SpringerPlus* 3, 572

Kain ZN, et al (2004). Preoperative anxiety and emergence delirium and postoperative maladaptive behaviors. *Anesthesia and Analgesia* 99, 1648-1654

Lee HW, Ph D (2015). Midazolam with placebo on prevention of emergence agitation in children having strabismus surgery. 1354-1361

Maclaren JE, et al (2010). Prediction of preoperative anxiety in children: Who is most accurate? *108*, 1777-1782

- Makkar JK, et al (2016). A comparison of single dose dexmedetomidine with propofol for the prevention of emergence delirium after desflurane anaesthesia in children. *Anaesthesia* 71, 50-57
- Mohkamkar M, et al (2014). Postanesthetic emergence agitation in pediatric patients under general anesthesia. *Iranian journal of pediatrics* 24, 184-90
- Nasr VG, Hannalah RS (2011). Emergence agitation in children a review. *Mid. East J. Anaesthesiol* 21, 175-82
- Rizk SN, Samir EM (2014). Use of ketofol to control emergence agitation in children undergoing adenotonsillectomy. *Egyptian Journal of Anaesthesia* 30, 13-19
- Sikich N, Lerman J (2004). Development and psychometric evaluation of the pediatric anesthesia emergence delirium scale. *Anesthesiology* 100, 1138-1145
- Singh R, et al (2012). Comparative evaluation of incidence of emergence agitation and post-operative recovery profile in paediatric patients after isoflurane, sevoflurane and desflurane anaesthesia. *Indian Journal of Anaesthesia* 56, 156-161
- Vlajkovic GP, Sindjelic RP (2007). Emergence delirium in children: Many questions, few answers. *Anesthesia and Analgesia* 104, 84-91
- Voepel-Lewis T, Malviya S, Tait AR (2003). A prospective cohort study of emergence agitation in the pediatric postanesthesia care unit. *Anesthesia & Analgesia*, 1625-1630
- Wong D, Bailey C (2015). Emergence delirium in children. *Anaesthesia* 70, 375-392