

RJPT

by Lutfi Ashar Mauludin

Submission date: 20-Oct-2023 03:51PM (UTC+0800)

Submission ID: 2192292760

File name: potential_insect_antimicrobial_of_black_soldier_fly_larvae.pdf (256.14K)

Word count: 7270

Character count: 38879

RESEARCH ARTICLE

58

A Potential Insect Antimicrobial of Black Soldier Fly Larvae (*Hermetia illucens*) against Pathogenic Bacteria

Diyantoro^{1,2,3*}, Aliyah Siti Sundari^{1,2}, Rizka Oktarianti Ainun Jariah^{1,2}, Dwi Wahyu Indriati^{1,2},
Dwi Wahyu Indriani⁴

¹Department of Health Science, Faculty of Vocational Studies, Universitas Airlangga, Indonesia.

²Antimicrobial Resource Alternative Studies for Emerging and Re-Emerging Infectious Disease,
Universitas Airlangga, Indonesia.

³Doctoral Program of Medical Science, Faculty of Medicine, Universitas Airlangga, Indonesia.

⁴National Research and Innovation Agency, Indonesia.

*Corresponding Author E-mail: diyantoro_dvm@vokasi.unair.ac.id

ABSTRACT:

54

Antimicrobial activity of black soldier fly larvae extract against gram-positive and gram-negative pathogenic bacteria has been discovered. Water-soluble extracts have the most potential and effectiveness as antimicrobials, with MIC values ranging from 12.5 to 25 mg/mL for both gram-positive and gram-negative bacteria. BSF larvae also create peptides called defensins and cecropins, which have antibacterial properties. Defensins were the best and most effective peptides in inhibiting bacterial growth, inhibiting both gram-positive and gram-negative bacteria (MIC value = > 29.97 μ M), while cecropins were more effective on gram-negative bacteria (MIC value = 0.52 – 2.07 μ M) than gram-positive bacteria (MIC value not detected).

KEYWORDS: Insect antimicrobial, BSF Larvae, pathogenic bacteria.

INTRODUCTION:

Antibiotics have successfully prevented or treated infections, not only saving patients' lives, but also assisting in major medical and surgical advancements. By therapeutically modifying the outcome of bacterial infections, it helped to extend projected life spans. Despite efforts to improve antibiotics, bacteria continue to change in reaction to the medications they are exposed to, and many environmental species already have antimicrobial resistance¹. Several reasons for the promotion of resistant bacteria include the overuse of antibiotics² and the use of inappropriately prescribed antibiotics³. Now, antibiotic resistance is becoming a global public health concern, as is the development of novel antibiotics, which is still in the research stage⁴. Several bacteria that are known to be resistant to antibiotic include MRSA (*Methicillin resistant staphylococcus aureus*)⁵,

Pathogenic ESBL (Extended Spectrum β -lactamases) producing *E. coli*⁶, and Multidrug-resistant (MDR) and extensively drug-resistant (XDR) tuberculosis⁷.

Several research have examined the possibility of antibiotics made from natural substances as an alternative. This is due to the fact that antibiotics derived from materials can prevent resistance to antibiotics developed from chemicals. Tea leaves⁸, *Bryonia seabra*⁹, flowers extract from *Cassia auriculata*¹⁰, *Allium Kurrat*, and *Eruca sativa*¹¹ are some of the natural substances that are commonly used as a source of novel antibiotics, while the antimicrobial from extract of insects are still limited.

Several researchers have been concerned about antimicrobial peptides (AMPs), which are considered natural antibiotics. Several researchers have been studying the potential use of this chemical in a variety of fields, including the pharmaceutical business. In general, this chemical does not cause bacterial resistance to develop. AMPs are abundant in insects, and they have antibacterial activity by destroying the bacterial cell

membrane. Gram-positive bacteria's peptidoglycan contains lipoteichoic and teichoic acids, while Gram-negative bacteria's lipopolysaccharide contains anionic phospholipid and phosphate. When a peptide attaches to the cytoplasmic membrane, it is incorporated into the bilayer phospholipid of the cytoplasmic membrane, causing the membrane structure of bacteria to change. The penetration of AMPs into a cell has the potential to influence nucleic acid and protein synthesis, which could explain their involvement in antibiotic-resistant bacteria¹². Defensins, cecropins, attacins, and dipterocins are the four main groups of insect AMPs, with defensins being the biggest. Many insect species from the orders Diptera, Hymenoptera, Coleoptera, Lepidoptera, Hemiptera, Isoptera, and Odonata have been found to have Defensins¹³.

30 The black soldier fly (*Hermetia illucens*), a non-pest insect that lives in tropical places with mild temperatures that reduce the concentration of animal dung and other biological substances, is one of the Diptera order's species¹⁴⁻¹⁶. This species is also known as an ecological decomposer since it lives in areas with high concentrations of hazardous microbes including bacteria and fungi¹⁷. As a result, the insects' ability to protect themselves from dangerous microbial infections is critical to their survival. BSF larvae have also been found to express a variety of AMPs, including defensins, cecropins, attacins, and dipterocins¹⁸, and are capable of reducing the amount of bacteria in substrates such as *Escherichia coli*¹⁹.

9 The purpose of this review is to discuss the potential use of AMPs of BSF larvae as an alternative to antibiotics for pathogenic bacteria. The discussion covers the knowledge of BSF larvae and its preparation for further examination, the biochemical contents, and the antimicrobial activities against pathogenic bacteria.

53
Black Soldier Fly (BSF) Larvae (*Hermetia illucens*)
Insect description:

Hermetia illucens, also known as the black soldier fly (BSF) larvae, is a member of the Order Diptera, Family Stratiomyidae, Genus *Hermetia*²⁰, and may be found in practically every temperate and tropical region of the world²¹. Adults are medium to large, measuring 15-20 mm long and slender, and resembling bees (wasplike)^{22,23}. They simply require water to survive, having reserves of nutrients for reproduction accumulated during the larvae stage^{24,25}. Female *H. illucens* are commonly observed ovipositing near organic waste, such as decomposing animal or plant waste²⁶. The female *H. illucens* has a blue-black body, whilst the male's abdomen is brown. The tips of the legs are white in both sexes, and the wings are black and gray when resting²². The abdomen is extended at the top

and narrows towards the bottom, with a translucent portion visible in the first two segments²⁷. While the C vein does not entirely surround the wing, the wing venation is tightly packed towards the ribs and more pigmented than the rear²⁷.

Distribution of insect:

These insects are almost universally distributed, with tropical, subtropical, and temperate regions accounting for the majority of their distribution²⁸⁻³⁰. The first documented occurrences from Asia date from the 1940s (for example, Malaysia), but no data from China have been confirmed until 1960³⁰. These insects have been found in Nepal, India, Sri Lanka, Thailand, Malaysia, Vietnam, Japan, Taiwan, Indonesia, and the Philippines, among other Asian countries. *Hermetia illucens* populations have been reported in Indonesia's Java, Sulawesi, and Papua provinces, but the species is now found practically everywhere in the country.

Larvae collection:

25 Egg traps composed of three layers of double-faced corrugated cardboard glued together and cut into 2.5 by 5-cm blocks can be used to gather larvae from the attack colony initially³¹. The egg trap was then placed 2-5cm above the wet Gainesville House Fly Diet, which was made up of 50% wheat bran, 30% alfalfa meal, and 20% maize meal, in a 5 liter plastic bucket³². A oviposition attractant or larval development medium was made by mixing equal parts of this dry media and water. *H. illucens* ovipositing on wet substrates was less appealing³¹. To enhance oviposition in the dry cardboard egg traps, water was added to the substance used as an oviposition attractant until it reached near saturation.

At 30°C and 80% humidity, eggs and larvae were kept in an insectery. Until the eggs hatched, they were placed in a 460-ml plastic cup (MC160 Sweetheart Cup, Chicago, IL) with a cover. In a 460-ml squat plastic cup, neonate larvae were fed 80g of moist Gainesville house fly meal. Fungus was considered to be the reason of low hatch and neonatal survival when eggs were placed on media. High humidity caused by a paper towel kept securely over the cup with a rubber band to contain wandering neonates resulted in fungal development. When larvae were placed on fresh media, fungus formation was infrequent³³.

Larvae process and extraction:

The following procedure was used to obtain BSF larvae extract. 1500 larvae were sterilized and stored at -80°C after being rinsed three times with sterile water. 100g of freeze-dried larvae The mixture was agitated for 12 hours at 4°C and then filtered through filter paper to remove the solid residues. To separate the solution layer containing the larval extract, the mixture was

centrifuged for 20 minutes at 13,000g. After that, a sterilized 0.45m membrane filter was used to filter the separated layer (Advantec MFS, Dublin, CA, USA). The material was kept at -20 degrees Celsius until needed³⁴.

Two-week-old *H. illucens* larvae were prepared. *H. illucens* larvae (600g) were initially rinsed with distilled water three times to remove all impurities. The larvae were immunized with probiotics *Lactobacillus casei* using a needle as previously described³⁵. Small scissors were used to open the rear part of the larva one by one. Afterwards, 10µl of hemolymph from each larva was collected and a total of 2ml from 200 larvae were collected for use in the next step. The hemolymphs were fractionated into seven portions using C18 cartridges (Waters Co., USA), and subsequently evaporated under reduced pressure using a rotary evaporator at 50°C as indicated previously^{36,37}. The seven column fractions were collected and each fraction was tested in vitro for antimicrobial effect before subjecting to further high performance liquid chromatography (HPLC) analysis.

The bacteria were incubated in 10 mL MH agar at 37°C and 30°C (*S. lutea* and *B. subtilis*) for 24 h. 10mL of this bacterial culture was inoculated into a fresh MH agar, which was incubated for 24 h. They were used for the experiment once the optical density (OD) 550 reached 0.5. Cell counts were determined by counting the colonies after Petri dish plates (Ø 87mm) were incubated at 37°C and 30°C for 24 h.

We investigated zone of growth inhibition of bacteria on the antibacterial effects of the extracts of *H. illucens* larvae. The respective bacteria were incubated in 10mL of MH agar at 37°C for 24 h, which were adjusted to a density of 2×10^6 bacteria/mL in fresh LB broth. Petri dishes with nutrient agar were inoculated with 40mL of bacterial culture (a final concentration of 2×10^6 bacteria/mL). The culture was evenly mixed and immediately distributed on Petri plates. Then, 45ml of the extract solution and antibiotics were added to each 6 mm disc, which were dried at room temperature for 30 min. The discs were placed onto the surface of agar plates and the agar plates were incubated at 37°C. Penicillinstreptomycin (10U/mL and 10mg/mL/disc) was used as the antibacterial positive control group. The diameters of bacterial inhibition zone surrounding the discs were measured in 12 and 24 h, respectively. The diameter of the zone of inhibition was measured in mm and the result was recorded. All experiments were performed in triplicate.

Biochemical Studies:

BSF larvae had 17.5% crude protein, 14% fat, 3.5 ash, 3 % acid detergent fiber, 0.8% nitrogen free extract, and 61.2% moisture, according to proximate analysis (Table

1). The amount of crude protein in BSF larvae is relatively significant. BSF larvae protein content varies based on the meal ingested during the growing period. Crude protein content in larvae increased immediately after hatching and then steadily dropped throughout the next 4–12 days of larval development, with a minimum concentration of 38% crude protein (CP) in the larval phase and a further increase of 39.2% in mature larvae on day 14. The concentration of CP increased steadily throughout the development phases, peaking at 46.2% CP in the early pupa stage. CP content was found to be the highest at the post-mortem adult stage (57.6%)³⁸.

BSF larvae are known to include a variety of minerals and vitamins, the most abundant of which are calcium 0.93% and choline 0.11% (Table 2). The fact that the outer layer of the larvae's epidermis releases a deposit of calcium carbonate (CaCO₃), which may lead to the high calcium and ash content, could also explain such diversity^{39,40}. Despite its prominence as a calcium-rich feeder insect, black soldier fly (BSF) larvae have received a lot of attention in the animal feed industry. However, there has been little research on non-production species like reptiles. Calcium serves a variety of functions in the human body. Cell signaling pathways, neurotransmitter release, muscle contractions, cardiac health, bone strength, enzyme function, and blood clotting are all impacted by it⁴¹. When calcium or phosphorus intake is either low or too high, the body attempts to maintain blood homeostasis by diverting or removing calcium that should be stored in bones. Bones become mushy or brittle as a result of this. Long bones are prone to pathological fractures, while the jaw bones deteriorate and become fibrous^{41,42}.

Lauric acid constituted a significant component among the SFAs (5.12%). Furthermore, linoleic acid (1.69%) and palmitic acid (1.61%) accounted for a significant portion of the fatty acids found in the larvae (14%). Lauric acid (C12:0) was found in significant concentrations (13–52%) in all larvae, but only in trace amounts (0–1%) in their meals. As noted by Spranghers *et al.*⁴³, this strongly suggests that it is synthesized by the larvae (2017). The larvae may also produce lauric acid from the carbohydrates in the substrate, according to those authors. This may explain why the largest quantities of lauric acid were seen in BSFL fed on bread (79% carbohydrates) (28% on dry matter basis). There was a positive association between larval weight and % age of lauric acid (C12:0) and total SFA in the larvae, indicating that these fatty acids accumulate the greatest as the larvae gain weight. BSFL at later phases of development had larger % ages of SFA and lauric acid⁴⁴. It has been shown that BSFL differs from other insect larvae due to its high SFA concentration⁴⁵. As adult BSF is not fed by the larvae, fatty acids are most likely a way

of store energy for this later stage of life⁴⁶. The cause of energy storage under SFA in particular could be, as suggested by Ushakova et al.⁴⁷, because these fatty acids are less susceptible to oxidization.

Aspartic acid and glutamic acid were the most abundant amino acids in the larvae. BSFL protein hydrolysate amino acid composition has been analyzed with HPLC instruments. Glutamic acid was the most prevalent

amino acid in BSFL hydrolysate (18.4%). Janssen et al.⁴⁸ and Müller et al.⁴⁹ reported the dominance of glutamic acid in BSFL. Also glutamic acid is the most prevalent amino acid in the prepupa of the BSF sample⁵⁰. Glutamic acid is also the most prevalent amino acid with a composition of up to 19% from other insect species such as Tropical banded grickets (*Grylodes sigillatus*)⁵¹.

Table 1: Proximate Analysis, Mineral, and Vitamin Content of BSF Larvae⁵²

Proximate analysis			Mineral content			Vitamin Content		
Composition	Value	(%) per kg	Composition	Value	(%) per kg	Composition	Value	(%) per kg
Moisture (g/kg)	612	61.20	Calcium (mg/kg)	9,340	0.93	Choline (mg/kg)	1,100	0.11
Crude Protein (g/kg)	175	17.50	Potassium (mg/kg)	4,530	0.45	Camitine (mg/kg)	83.8	0.01
Crude Fat (g/kg)	140	14.00	Phosphorus (mg/kg)	3,560	0.36	Niacin (mg/kg)	71.0	0.01
Ash (g/kg)	35	3.50	Magnesium (mg/kg)	1,740	0.17	Pantothenic acid (mg/kg)	38.5	< 0.01
Acid Detergent Fiber (g/kg)	30	3.00	Chloride (mg/kg)	1,160	0.12	Riboflavin (mg/kg)	16.2	< 0.01
Nitrogen Free Extract (g/kg)	8	0.80	Sodium (mg/kg)	887	0.09	Vitamin C (mg/kg)	< 10.0	< 0.01
			Iron (mg/kg)	66.6	0.01	Thiamin (mg/kg)	7.7	< 0.01
			Manganese (mg/kg)	61.8	0.01	Vitamin E (mg α-tocopherol/kg)	6.2	< 0.01
			Zinc (mg/kg)	56.2	0.01	Pyridoxine (mg/kg)	6.01	< 0.01
			Copper (mg/kg)	4.03	< 0.01	Folic acid (mg/kg)	2.70	< 0.01
			Selenium (mg/kg)	0.32	< 0.01	Biotin (mg/kg)	0.35	< 0.01
			Iodine (mg/kg)	0.26	< 0.01	Vitamin A (µg retinol/kg)	< 300	< 0.01
						Vitamin B12 (µg/kg)	55.8	< 0.01
						Vitamin D3 (IU/kg)	100	< 0.01
						Vitamin D2 (IU/kg)	< 80	< 0.01

Table 2: Amino Acid and Fatty Acid Content of BSF Larvae⁵²

Amino acid content			Fatty acid content		
Composition	Value	(%) per kg	Composition	Value	(%) per kg
Glutamic acid (g/kg)	19.7	1.97	Lauric 12:0 (g/kg)	51.2	5.12
Phenylalanine + tyrosine (g/kg)	19.66	1.97	Linoleic 18:2 (g/kg)	16.9	1.69
Aspartic acid (g/kg)	16.5	1.65	Palmitic 16:0 (g/kg)	16.1	1.61
Valine (g/kg)	12.9	1.29	Oleic 18:1 (g/kg)	15.6	1.56
Arginine (g/kg)	12.3	1.23	Myristic 14:0 (g/kg)	12.0	1.20
Alanine (g/kg)	12.2	1.22	Palmitoleic 16:1 (g/kg)	4.96	0.50
Leucine (g/kg)	12.1	1.21	Stearic 18:0 (g/kg)	2.45	0.25
Tyrosine (g/kg)	12.1	1.21	0.01 Capric 10:0 (g/kg)	0.69	0.07
Lysine (g/kg)	11.9	1.19	Linolenic 18:3 (g/kg)	0.65	0.07
Proline (g/kg)	10.2	1.02	Myristoleic 14:1 (g/kg)	0.50	0.07
Glycine (g/kg)	9.14	0.91	Heptadecanoic 17:0 (g/kg)	0.20	0.05
Isoleucine (g/kg)	7.62	0.76	Arachidic 20:0 (g/kg)	0.16	0.02
Phenylalanine (g/kg)	7.56	0.76	Pentadecanoic 15:0 (g/kg)	0.12	0.01
Serine (g/kg)	7.02	0.70	Eicosadienoic 20:2 (g/kg)	< 0.10	< 0.01
Threonine (g/kg)	6.82	0.68	Benhenic 22:0 (g/kg)	0.09	< 0.01
Histidine (g/kg)	5.94	0.59	Eicosenoic 20:1 (g/kg)	< 0.08	< 0.01
Methionine + cystine (g/kg)	4.39	0.44	Arachidonic 20:4 (g/kg)	< 0.08	< 0.01
Methionine (g/kg)	3.37	0.34	Heptadecenoic 17:1 (g/kg)	< 0.08	< 0.01
Tryptophan (g/kg)	3.00	0.30			
Cystine (g/kg)	1.02	0.10			
Taurine (g/kg)	0.1	0.01			

Antimicrobial Activities:

Antimicrobial activities of several extract fraction of BSF Larva:

Several types of extracts from BSF larvae indicated antibacterial activity against both Gram positive and Gram negative bacteria at certain concentrations. The

results of several studies showed that the water-soluble extract of BSF larvae was more potential compared to other types of extracts. This was indicated by the lower average of MIC value of the extract were in a similar range from 12.5 to 25mg/mL. Several results show that the water soluble fraction of the entire body extract had

considerably stronger antibacterial activity against MRSA and indicate the presence of more than one "antibacterial" product acting in synergy to enhance the action. Furthermore, the aqueous extract of *H. illucens* is highly solid, resilient to a number of freezing cycles and lyophilization and stable, as a frozen preparation, which all are key elements for the pharmaceutical development of a product. It was recently discovered that the presence and generation of a new class of tiny, maybe semi-peptidergic antimicrobial chemicals is not limited to the hemolymph or fat body, and that preparing a whole-body extract could be more efficient when attempting to

isolate those compounds¹⁷.

Furthermore, this sort of water-soluble extraction is highly recommended since the extraction method is simple, quick, and easy to use⁵³, and the protein content in the extraction is not destroyed but reduced by 17.2%. The decreased protein concentration is most likely due to the difficulty of extracting proteins from insects or the hydrophobicity of proteins, which are difficult to dissolve in aqueous buffer systems and remain bonded to the solid cellular components⁵⁴.

Table 3: Antimicrobial activities of several extract fraction of BSF Larvae against pathogenic bacteria

Bacteria Group	Type of extract	Bacteria tested	MIC	References
Gram-positive bacteria	Water-soluble extract	Methicillin-resistant <i>Staphylococcus aureus</i> (MRSA) † (clinically isolated)	25 mg/ml	17
		<i>Staphylococcus aureus</i> KCCM 40881	> 100 mg/ml	
		<i>Staphylococcus aureus</i> KCCM 35494	50 mg/ml	
		<i>Staphylococcus epidermidis</i> KCCM 35494	50 mg/ml	
		<i>Kocuria rhizophila</i> KCCM 11236	25 mg/ml	
		<i>Micrococcus luteus</i> KCCM 11326	25 mg/ml	
		<i>Bacillus subtilis</i> KCCM 11316	12.5 mg/ml	
	Methanol extract	<i>Bacillus subtilis</i> KCTC 1325	ND	14
		<i>Streptococcus mutans</i> KCTC 3289	ND	
		<i>Sarcina lutea</i> ATCC 9341	ND	
	Hexadenolic acid	MRSA	137.37 µg/ml	55
		<i>Staphylococcus aureus</i> (ATCC 25923)	140.38 µg/ml	
	Ethyl acetate	<i>Staphylococcus epidermidis</i> KCCM 35494	25 mg/ml	14
		<i>Staphylococcus aureus</i> KCCM 40881	> 100 mg/ml	
<i>Bacillus subtilis</i> KCCM 11316		25 mg/ml		
Gram-negative bacteria	Water-soluble extract	<i>Escherichia coli</i> KCCM 11234	12.5 mg/ml	17
		<i>Enterococcus aerogenes</i> KCCM 12177	25 mg/ml	
		<i>Pseudomonas aeruginosa</i> KCCM 11328	12.5 mg/ml	
	Methanol extract	<i>Klebsiella pneumoniae</i> KCTC 2253	44.74 mg/ml	14
		<i>Neisseria gonorrhoeae</i>	43.98 mg/ml	
		<i>Shigella sonnei</i> KCTC 2581	43.96 mg/ml	
		<i>Escherichia coli</i>	160 mg/ml	
		<i>Salmonella</i> sp.	160 mg/ml	
		<i>Salmonella typhimurium</i>	325 mg/ml	
		<i>Pseudomonas aeruginosa</i>	325 mg/ml	
	Hexadenolic acid	<i>Klebsiella pneumoniae</i> (ATCC 13883)	139.12 µg/ml	17
		<i>Shigella dysenteriae</i> (ATCC 9750)	139.70 µg/ml	
		<i>Klebsiella pneumoniae</i>	1.25 mg/ 50µL	
	Ethyl acetate	<i>Escherichia coli</i> KCCM 11234	> 100 mg/ml	14

Potential of Antimicrobial Peptides (AMPs) of BSF Larvae:

Due to their minimal toxicity to eukaryotic cells and broad spectrum of action against bacteria, mycobacteria, fungus, viruses, and cancer cells, AMPs are interesting prospects as alternatives to conventional antibiotics⁵⁸. AMPs can kill bacteria in a variety of ways, including disrupting membranes, targeting intracellular components, and interfering with bacterial metabolism⁵⁹⁻⁶¹. Moreover, most AMPs are cationic, with a positive net charge that promotes electrostatic interaction with bacterial membranes which are negatively charged⁶². Every living organism produces AMPs where the high biodiversity and extremely diverse living conditions make insects one of the richest

sources. As it is possible to observe in other species like in *Eristalis tenax*, the *H. illucens* immune system is highly developed because this species feeds on decaying substrates and manure extremely rich in pathogenic microorganisms. In fact, 20 AMPs of the Diptera *E. tenax* have been identified that have been able to adapt to different aquatic habitats with heavy microbial load (sewage tanks and manure pits)⁶³. An essential part of immunological protection is AMPs which synthesize the fat body and hemocytes and then secrete into the hemolymph^{64,65}.

The defensins and cecropins of AMPs from BSF larvae extract have potential as antimicrobials against bacteria. Insect defensins can inhibit the growth of both gram-positive and gram-negative bacteria, but they are more

effective against gram-positive bacteria. In comparison, insect cecropins are more active and effective at inhibiting gram-negative bacteria development. The MIC value of insect defensins against gram-positive bacteria is in the range 0.01 – 14.99 μM, while for gram-negative bacteria is about > 29.97 μM. Six conserved cysteines generate three intramolecular disulfide bonds in the insect defensins, a class of tiny (4 kDa) cysteine-rich cationic peptides. The insect defensin family shares a lot of similarities with mammalian defensins. An N-terminal loop is the most common structure of insect defensins⁶⁶. By interacting with phospholipids, insect defensins target the cytoplasmic membrane of microbes, producing channels that cause membrane permeabilization and, ultimately, microbial cell death⁶⁶⁻⁶⁸.

MIC values of insect cecropins from BSF larvae extract could not be detected in gram-positive bacteria, but only detected in gram-negative bacteria with MIC values of 0.52 – 2.07 μM. The tiny cationic cecropin peptides have antibacterial activity against both Gram-positive and Gram-negative bacteria, as well as fungi⁶⁶. The amidation of the C-terminus of the peptide, which is necessary for the interaction with liposomes of the attacking bacterium, has also been linked to cecropin antibacterial activity^{69,70}. In a hydrophobic environment, cecropins are bactericidal as helical forms (e.g. the bacterial cell membrane or lipopolysaccharides). Cecropins are made up of an N-terminal amphipathic -helix and a C-terminal hydrophobic -helix joined by an intramolecular hinge region⁷¹.

Table 4: Potential of Antimicrobial Peptides (AMPs) of BSF Larvae against Gram-positive Bacteria

Antimicrobial peptide		Antimicrobial peptide	MIC	Preference	
Name	Class				
Defensin-like peptide 2 (DLP2)	Defensins	<i>Staphylococcus aureus</i> ATCC25923	0.01 μM	72	
		<i>Staphylococcus aureus</i> ATCC43300	0.12 μM		
		<i>Staphylococcus aureus</i> ATCC6538	0.12 μM		
		<i>Staphylococcus aureus</i> CICC546	0.23 μM		
		<i>Streptococcus suis</i> CVCC606	0.93 μM		
		<i>Listeria ivanovii</i> ATCC19119	0.12 μM		
Defensin-like peptide 3 (DLP3)	Defensins	MRSA (methicillin-resistant <i>Staphylococcus aureus</i> → clinical isolated)	5 μg/mL	73	
		<i>Staphylococcus aureus</i> KCCM 40881	5 μg/mL		
		<i>Staphylococcus aureus</i> KCCM 12256	10 μg/mL		
		<i>Staphylococcus epidermidis</i> KCCM 35494	10 μg/mL		
Defensin-like peptide 4 (DLP4)	Defensins	MRSA	0.59-1.17 μM	34	
		<i>Staphylococcus aureus</i> KCCM 40881	0.59-1.17 μM		
		<i>Staphylococcus aureus</i> KCCM 12256	1.17-2.34 μM		
		<i>Staphylococcus epidermidis</i> KCCM 35494	0.59-1.17 μM		
		<i>Bacillus subtilis</i> KCCM 11316	0.02-0.04 μM	72	
		<i>Staphylococcus aureus</i> ATCC25923	0.01 μM		
		<i>Staphylococcus aureus</i> ATCC43300	0.23 μM		
		<i>Staphylococcus aureus</i> ATCC6538	0.47 μM		
		<i>Staphylococcus aureus</i> CICC546	0.47 μM		
		<i>Streptococcus suis</i> CVCC606	1.88 μM		
		<i>Listeria ivanovii</i> ATCC19119	0.12 μM		
		<i>Staphylococcus aureus</i> CVCC 546	3.75 μM		74
		<i>Staphylococcus epidermidis</i> ATCC 12228	14.99 μM		
		<i>Streptococcus pneumoniae</i> CVCC 2350	7.50 μM		
<i>Streptococcus suis</i> CVCC 3928	3.75 μM				
ID13	Defensins	<i>Staphylococcus aureus</i> CVCC 546	0.95 μM	74	
		<i>S. epidermidis</i> ATCC 12228	1.91 μM		
		<i>Streptococcus pneumoniae</i> CVCC 2350	0.05 μM		
		<i>Streptococcus suis</i> CVCC 3928	0.95 μM		
Cecropin-like peptide 1 (CLP1)	Cecropins	<i>Staphylococcus aureus</i> KCCM 40881	ND	75	
		<i>Staphylococcus aureus</i> KCCM 12256	ND		
		<i>Staphylococcus epidermidis</i> KCCM 35494	ND		
Trx-stomoxynZH1	Cecropins	<i>Staphylococcus aureus</i>	27-54 μg/ml	76	

Table 5: Potential of Antimicrobial Peptides (AMPs) of BSF Larvae against Gram-negative Bacteria

Antimicrobial peptide		Antimicrobial peptide	MIC	Preference
Name	Class			
Defensin-like peptide 2 (DLP2)	Defensins	<i>Escherichia coli</i> CVCC1515	>29.97 μM	72
		<i>Escherichia coli</i> CICC21530 (serotype O157:H7)	>29.97 μM	
		<i>Salmonella typhimurium</i> ATCC14028	>29.97 μM	
		<i>Salmonella enteritidis</i> CMCC50336	>29.97 μM	
Defensin-like peptide 3 (DLP3)	Defensins	<i>Escherichia coli</i> KCCM 11234	10 μg/mL	73
		<i>Pseudomonas aeruginosa</i> KCCM 11328	40 μg/ml	

Defensin-like peptide 4 (DLP4)	Defensins	<i>Escherichia coli</i> CVCC1515	>29.98 μM	72	
		<i>Escherichia coli</i> CICC21530 (serotype O157:H7)	>29.98 μM		
		<i>Salmonella typhimurium</i> ATCC14028	>29.98 μM		
		<i>Salmonella enteritidis</i> CMCC50336	>29.98 μM		
		<i>Escherichia coli</i> ATCC 25922	>29.98 μM		74
		<i>Escherichia coli</i> K88	>29.98 μM		
		<i>Salmonella pullorum</i> CVCC 533	>29.98 μM		
ID13	Defensins	<i>Salmonella enteritidis</i> CVCC 3377	>29.98 μM	74	
		<i>Escherichia coli</i> ATCC 25922	> 30.50 μM		
		<i>Escherichia coli</i> K88	> 30.50 μM		
		<i>Salmonella pullorum</i> CVCC 533	> 30.50 μM		
Cecropin-like peptide 1 (CLP1)	Cecropins	<i>Salmonella enteritidis</i> CVCC 3377	> 30.50 μM	75	
		<i>Escherichia coli</i> KCCM 11234	0.52–1.03 μM		
		<i>Enterobacter aerogenes</i> KCCM 12177	1.03–2.07 μM		
Trx-stomoxynZH1	Cecropins	<i>Pseudomonas aeruginosa</i> KCCM 11328	1.03–2.07 μM	76	
		<i>Escherichia coli</i>	15-30 $\mu\text{g/ml}$		

ACKNOWLEDGEMENT:

The authors are grateful to my supervisor for his valuable and constructive suggestions during the planning and development of this review work. His willingness to give his time so generously has been very much appreciated.

12

CONFLICT OF INTEREST:

The authors declare no conflict of interest.

REFERENCES:

- Sreeja MK, Gowrishankar NL, Adisha S, Divya KC. Antibiotic Resistance-Reasons and the Most Common Resistant Pathogens – A Review. *Research Journal of Pharmacy and Technology* 2017; 10(6): 1886-1890. doi: 10.5958/0974-360X.2017.00331.6
- Shivani MD, Vaishali RU. Antibiotic Overuse and Resistance: An Awareness Study. *Research Journal of Pharmacy and Technology* 2019; 12(6): 2794-2798. doi: 10.5958/0974-360X.2019.00470.0
- Aishwarya JR. History of Antibiotics and Evolution of Resistance. *Research Journal of Pharmacy and Technology* 2015; 8(12): 1719-1724. doi: 10.5958/0974-360X.2015.00309.1
- Maria JD, Vijey AM. An Overview on Antibiotic use and resistance. *Research Journal of Pharmacy and Technology* 2017; 10(8): 2793-2796. doi: 10.5958/0974-360X.2017.00494.2
- Sharma S, Upadhayay UM, Mistry S. Antimicrobial Activity of Extracts against MRSA. *Research Journal of Pharmacy and Technology* 2016; 9(12):2283-2286. doi: 10.5958/0974-360X.2016.00460.1
- Nanda A, Dhamodharan S, Nayak BK. Antibiotic Resistance Pattern Exhibited by ESBL (Extended Spectrum β -Lactamases) in Multidrug Resistant Strains, *Escherichia coli*. *Research Journal of Pharmacy and Technology* 2017; 10(11): 3705-3708. doi: 10.5958/0974-360X.2017.00672.2
- Sandip Z, Shweta AP, Sushmita SR. Rise of Antibiotic Resistance in Tuberculosis. *Research Journal of Pharmacy and Technology* 2018; 11(7): 3201-3204. doi: 10.5958/0974-360X.2018.00588.7
- Hindi NKK, Abdul-Husin IF, Al-Mahdi ZKA, Ewadh RMJ, Hossain AO, Kadhim MJ, Alnasraui AHF, Al-Yaseri AA. Effectiveness of Aqueous extract of Green, Black and Red Tea Leaves against some types of Gram positive and negative bacteria. *Research Journal of Pharmacy and Technology* 2017; 10(6): 1957-1962. doi: 10.5958/0974-360X.2017.00343.2
- Elayaraja A, Rao GD. Anti-Bacterial Activity of Various Crude Extracts of *Bryonia seabra*. *Research Journal of Pharmacy and Technology*. 2008; 1(3): 283-284.
- Bargah RK, Kushwaha A, Tirkey A, Hariwanshi B. In Vitro Antioxidant and Antibacterial Screening of flowers Extract from *Cassia auriculata* Linn. *Research Journal of Pharmacy and Technology* 2020; 13(6):2624-2628. doi: 10.5958/0974-360X.2020.00466.7
- El-Sayed MA, Kamel MM, El-Raei MA, Osman SM, Gamil L, Abbas HA. Study of Antibacterial Activity of Some Plant Extracts Against Enterohemorrhagic *Escherichia coli* O157:H7. *Research Journal of Pharmacy and Technology* 2013; 6(8): 916-919.
- Żyłowska M, Wyszynska A, Jagusztyn-Krynicka EK. Antimicrobial peptides – defensins (in Polish). *Post. Mikrobiol.* 2011; 50: 223–234.
- Yi HY, Chowdhury M, Huang YD, Yu XQ. Insect antimicrobial peptides and their applications. *Applied Microbiology and Biotechnology* 2014; 98(13): 5807–5822. doi: 10.1007/s00253-014-5792-6
- Choi WH, Yun JH, Chu JP, Chu KB. Antibacterial effect of extracts of *Hermetia illucens* (Diptera: Stratiomyidae) larvae against Gram-negative bacteria. *Entomological Research* 2012; 42(5): 219–226. doi: 10.1111/j.1748-5967.2012.00465.x
- Kim W, Bae S, Park K, Lee S, Choi Y, Han S, Koh Y. Biochemical characterization of digestive enzymes in the black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae). *Journal of Asia-Pacific Research* 2011; 14(1): 11–14. doi: 10.1016/j.aspen.2010.11.003
- Diener S, Solano NMS, Gutierrez FR, Zurbrugg C, Tockner K. Biological Treatment of Municipal Organic Waste using Black Soldier Fly Larvae. *Waste and Biomass Valorization* 2011; 2(4): 357–363. doi: 10.1007/s12649-011-9079-1
- Park SI, Chang BS, Yoe SM. Detection of antimicrobial substances from larvae of the black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae). *Entomological Research* 2014; 44(2): 58–64. doi: 10.1111/1748-5967.12050
- Müller A, Wolf D, Gutzeit HO. The black soldier fly, *Hermetia illucens* – a promising source for sustainable production of proteins, lipids and bioactive substances. *Zeitschrift für Naturforschung* 2017; 72(9–10): 351–363. doi: 10.1515/znc-2017-0030
- Liu Q, Tomberlin JK, Brday JA, Sanford MR, Yu Z. Black soldier fly (Diptera: Stratiomyidae) larvae reduce *Escherichia coli* in dairy manure. *Environmental Entomology* 2008; 37(6): 1525–1530. doi: 10.1603/0046-225X-37.6.1525
- Hem S. Final Report: Maggot – Bioconversion Research Program in Indonesia, Concept of New Food Resources Result and Applications 2005-2011. Perancis: Institut de Recherche pour le Développement. 2011.
- Tomberlin JK, Adler PH, Myers HM. Development of the black soldier fly (Diptera: Stratiomyidae) in relation to temperature. *Environmental Entomology*. 2009; 38(3): 930-4. doi: 10.1603/022.038.0347
- Byrd JH, Castner JL. Insects of Forensic Importance. In: Byrd JH, Castner JL, editors. *Forensic Entomology: The Utility of Arthropods in Legal Investigations* (Second Edition). Boca Raton: CRC Press. 2010; 2nd ed: pp. 39-122.

23. Triplehorn CA, Johnson NF. Borror and deLong's Introduction to the Study of Insects (Seventh Edition). Belmont: Thomson Brooks/Cole. 2005.
24. Tomberlin JK, Sheppard DC, Joice JA. Selected life-history traits of black soldier flies (Diptera: Stratiomyidae) reared on three artificial diets. *Annals of the Entomological Society of America* 2002; 95(3):379-86. doi: 10.1603/0013-8746(2002)095[0379:SLHTOB]2.0.CO;2
25. Sheppard DC, Tomberlin JK, Joice JA, Kiser BC, Somner SM. Rearing methods for the black soldier fly (Diptera: Stratiomyidae). *Journal of Medical Entomology* 2002; 39(4):695-8. doi: 10.1603/0022-2585-39.4.695
26. Newton GL, Sheppard DC, Watson DW, Burtle GJ, Dove CR, Tomberlin JK, Thelen EE. The black soldier fly, *Hermetia illucens*, as a manure management / resource recovery tool. *Symposium on the State of the Science of Animal Manure and Waste Management* 2005; 5-7
27. DuPont MW, Larish LB. Soldier fly: Livestock Management Insect Pests, LM-10.7. Cooperative Extension Service, College of Tropical Agriculture and Human Resources. University of Hawaii, Manoa. 2003.
28. Rozkosný R. A biosystematic study of the European Stratiomyidae (Diptera). In: Spencer, K.A. (Ed.), Clitellariinae, Hermetiinae, Pachygasteriinae and Bibliography. *Series Entomologica* 1983; 2:172-176.
29. Leclercq M. A propos de *Hermetia illucens* (Linnaeus, 1758) ("soldier fly") (Diptera: Stratiomyidae: Hermetiinae). *Bulletin et Annales de la Société Belge d'entomologie* 1997; 133:275-282.
30. Marshall SA, Woodley NE, Hauser M. The historical spread of the Black Soldier Fly, *Hermetia illucens* (L.) (Diptera, Stratiomyidae, Hermetiinae), and its establishment in Canada. *Journal of the Entomological Society of Ontario* 2015; 146:51-54.
31. Booth DC, Sheppard C. Oviposition of the black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae): eggs, masses, timing, and site characteristics. *Environmental Entomology* 1984; 13(2): 421-423. doi: 10.1093/ee/13.2.421
32. Hogsette J. New diets for production of house-flies and stable flies (Diptera, Muscidae) in the Laboratory. *Journal of Economic Entomology* 1992; 85: 2291-2294. <https://digitalcommons.unl.edu/usdaarsfacpub/1005>
33. Beskin KV, Holcomb CD, Cammack JA, Crippen TL, Knap AH, Sweet ST, Tomberlin JK. Larval digestion of different manure types by the black soldier fly (Diptera: Stratiomyidae) impacts associated volatile emissions. *Waste Management* 2018; 74: 213-220. doi: 10.1016/j.wasman.2018.01.019.
34. Park SI, Kim JW, Yoe SM. Purification and characterization of a novel antibacterial peptide from black soldier fly (*Hermetia illucens*) larvae. *Developmental and Comparative Immunology* 2015; 52: 98-106. doi: 10.1016/j.dci.2015.04.018
35. Robertson M, Postlethwait JH. The humoral antibacterial response of *Drosophila* adults. *Developmental and Comparative Immunology* 1986; 10(2): 167-179. doi: 10.1016/0145-305x(86)90001-7
36. Chu K, Jeon G, Quan F. Hexanedioic acid from *Hermetia illucens* larvae (Diptera: Stratiomyidae) protects mice against *Klebsiella pneumoniae* infection. *Entomological Research* 2014; 44(1): 1-8. doi: 10.1111/1748-5967.12043
37. Choi WH, Choi H, Goo TW, Quan F. Novel antibacterial peptides induced by probiotics in *Hermetia illucens* (Diptera: Stratiomyidae) larvae. *Entomological Research* 2018; 48(4): 237-247. doi: 10.1111/1748-5967.12259
38. Liu X, Chen X, Wang H, Yang Q, ur Rehman K, Li W, Cai M, Li Q, Mazza L, Zhang J. Dynamic changes of nutrient composition throughout the entire life cycle of black soldier fly. *PLoS ONE*. 2017; 12: e0182601. doi: 10.1371/journal.pone.0182601
39. Barragan-Fonseca KB, Dicke M, van Loon JJA. Nutritional value of the black soldier fly (*Hermetia illucens* L.) and its suitability as animal feed – a review. *Journal of Insects as Food and Feed* 2017; 3(2): 105-120. doi: 10.3920/JIFF2016.0055
40. Newton GL, Booram CV, Barker RW, Hale OM. Dried *Hermetia illucens* larvae meal as a supplement for swine. *Journal of Animal Science* 1977; 44(3): 395-400. doi: 10.2527/jas1977.443395x
41. Raiti P. Endocrinology. In Divers SJ and Stahl SJ (eds.): *Mader's Reptile and Amphibian Medicine and Surgery*. Elsevier, St. Louis, MO. 2019; 835-848.
42. Boyer TH, Scott PW. Nutritional Secondary Hyperparathyroidism. In Divers SJ and Stahl SJ (eds.): *Mader's Reptile and Amphibian Medicine and Surgery*. Elsevier, St. Louis, MO. 2019; 1326-1327.
43. Spranghers T, Ottoboni M, Klootwijk C, Owyn A, Debrouere S, De Meulenaer B, De Smet S. Nutritional composition of black soldier fly (*Hermetia illucens*) prepupae reared on different organic waste substrates. *Journal of the Science of Food and Agriculture* 2017; 97(8):2594-2600. doi: 10.1002/jsfa.8081
44. Liu X, Chen X, Wang H, Yang QQ, Rehman KU, Li W, Cai MM, Li Q, Mazza L, Zhang JB, Yu ZN, Zheng LY. Dynamic changes of nutrient composition throughout the entire life cycle of black soldier fly. *PLoS ONE*. 2017; 12: 21. doi: 10.1371/journal.pone.0182601
45. Ramos-Bueno RP, González-Fernández MJ, Sánchez-Muros-Lozano MJ, García-Barroso F, Guil-Guerrero JL. Fatty acid profiles and cholesterol content of seven insect species assessed by several extraction systems. *European Food Research and Technology* 2016; 242: 1471-1477. doi: 10.1007/s00217-016-2647-7
46. Tomberlin JK, Sheppard DC. Factors influencing mating and oviposition of black soldier flies (Diptera: Stratiomyidae) in a colony. *Journal of Entomological Science* 2002; 37: 345-352. doi: 10.18474/0749-8004-37.4.345
47. Ushakova NA, Brodskii ES, Kovalenko AA, Bastrakov AI, Kozlova AA, Pavlov DS. Characteristics of lipid fractions of larvae of the black soldier fly *Hermetia illucens*. *Doklady Biochemistry and Biophysics* 2016; 468(1): 209-212. doi: 10.1134/S1607672916030145
48. Janssen RH, Vincken JP, van den Broek LAM, Fogliano V, Lakemond CMM. Nitrogen-to-Protein Conversion Factors for Three Edible Insects: *Tenebrio molitor*, *Alphitobius diaperinus*, and *Hermetia illucens*. *Journal of Agricultural and Food Chemistry* 2017; 65(11): 2275-2278. doi: 10.1021/acs.jafc.7b00471
49. Muller A, Wolf D, Gutzeit HO. The black soldier fly, *Hermetia illucens* – a promising source for sustainable production of proteins, lipids and bioactive substances. *Zeitschrift für Naturforschung - Section C Journal of Biosciences* 2017; 72(9-10): 351-363. doi: 10.1515/znc-2017-0030.
50. Surendra KC, Olivier R, Tomberlin JK, Jha R, Khanal SK. Bioconversion of organic wastes into biodiesel and animal feed via insect farming. *Renewable Energy* 2016; 98:197-202. doi: 10.1016/j.renene.2016.03.022
51. Hall FG, Jones OG, O'Haire ME, Liceaga AM. Functional properties of tropical banded cricket (*Gryllobates sigillatus*) protein hydrolysates. *Food Chemistry* 2017; 224: 414-422. doi: 10.1016/j.foodchem.2016.11.138
52. Finke MD. Complete nutrient content of four species of feeder insects. *Zoo Biology* 2013; 32(1):27-36. doi: 10.1002/zoo.21012
53. Rammo RNN. Bactericidal and Anti-biofilm Formation of Aqueous Plant Extracts against Pathogenic Bacteria. *Asian Journal of Pharmaceutical Research* 2017; 7(1): 25-29. doi: 10.5958/2231-5691.2017.00005.3
54. Meylaers K, Cerstiaens A, Vierstraete E, Baggerman G, Michiels CW, De Loof A, Schoofs L. Antimicrobial compounds of low molecular mass are constitutively present in insects: characterisation of β -alanyl-tyrosine. *Current Pharmaceutical Design* 2003;9(2):159-74. doi: 10.2174/1381612033392279.
55. Choi WH, Jiang MH. Evaluation of antibacterial activity of hexanedioic acid isolated from *Hermetia illucens* larvae. *Journal of Applied Biomedicine* 2014; 12(3): 179-189. doi: 10.1016/j.jab.2014.01.003
56. Harlystiarini, Mutia R, Wibawan IWT, Astuti DA. In vitro antibacterial activity of black soldier fly (*Hermetia illucens*) larva

- extracts against Gram-negative bacteria. *Buletin Peternakan* 2019; 43(2): 125-129. doi: 10.21059/buletinpeternak.v43i2.42833
57. Auza FA, Purwanti S, Syamsu JA, Natsir A. Antibacterial activities of black soldier flies (*Hermetia illucens*, L.) extract towards the growth of *Salmonella typhimurium*, *E. coli*, and *Pseudomonas aeruginosa*. *IOP Conference, Series: Earth and Environmental Science* 2020; 492: 012024. doi:10.1088/1755-1315/492/1/012024
 58. Hu H, Wang C, Guo X, Li W, Wang Y, He Q. Broad activity against porcine bacterial pathogens displayed by two insect antimicrobial peptides moricin and cecropin B. *Molecules and Cells* 2013; 35(2): 106-114. doi: 10.1007/s10059-013-2132-0
 59. Jozefak A, Engberg RM. Insect proteins as a potential source of antimicrobial peptides in livestock production: A review. *Journal of Animal and Feed Sciences* 2017; 26(2): 87-99. doi: 10.22358/jafs/69998/2017
 60. Bechinger B, Gorr SU. Antimicrobial peptides: Mechanisms of action and resistance. *Journal of Dental Research* 2017; 96(3): 254-260. doi: 10.1177/0022034516679973
 61. Oñate-Garzón J, Manrique-Moreno M, Trier S, Leidy C, Torres R, Patino E. Antimicrobial activity and interactions of cationic peptides derived from *Galleria mellonella* cecropin D-like peptide with model membranes. *Journal of Antibiotics* 2017; 70(3): 238-245. doi: 10.1038/ja.2016.134
 62. Yeaman MR, Yount NY. Mechanisms of antimicrobial peptide action and resistance. *Pharmacological Review* 2003; 55(1): 27-55. doi: 10.1124/pr.55.1.2.
 63. Hirsch R, Wiesner J, Bauer A, Marker A, Vogel H, Hammann PE, Vilcinskas A. Antimicrobial peptides from rat-tailed maggots of the drone fly *Eristalis tenax* show potent activity against multidrug-resistant gram-negative bacteria. *Microorganisms* 2020; 8(5): 626. doi: 10.3390/microorganisms8050626.
 64. Bulet P, Hetru C, Dimarcq JL, Hoffman D. Antimicrobial peptides in insects: Structure and function. *Developmental and Comparative Immunology* 1999; 23(4-5): 329-344. doi: 10.1016/s0145-305x(99)00015-4.
 65. Hofmann JA, Reichhart JM. *Drosophila* innate immunity: An evolutionary perspective. *Nature Immunology* 2002; 3(2): 121-126. doi: 10.1038/ni0202-121.
 66. Yi HY, Chowdhury M, Huang YD, Yu XQ. Insect antimicrobial peptides and their applications. *Applied Microbiology and Biotechnology* 2014; 98(13): 5807-22. doi: 10.1007/s00253-014-5792-6.
 67. Cociancich S, Ghazi A, Hetru C, Hoffmann JA, Letellier L. Insect defensin, an inducible antibacterial peptide, forms voltage-dependent channels in *Micrococcus luteus*. *Journal of Biological Chemistry* 1993; 268(26): 19239-45. PMID: 7690029
 68. Malanovic N, Lohner K. Antimicrobial peptides targeting Gram-positive bacteria. *Pharmaceuticals (Basel)* 2016; 9(3): 59. doi: 10.3390/ph9030059
 69. Li ZQ, Merrifield RB, Boman IA, Boman HG. Effects on electrophoretic mobility and antibacterial spectrum of removal of two residues from synthetic sarcotoxin IA and addition of the same residues to cecropin B. *FEBS Letters* 1988; 231(2): 299-302. doi: 10.1016/0014-5793(88)80837-8.
 70. Nakajima Y, Qu XM, Natori S. Interaction between liposomes and sarcotoxin IA, a potent antibacterial protein of *Sarcophaga peregrina* (flesh fly). *Journal of Biological Chemistry* 1987; 262(4): 1665-9. PMID: 3543008
 71. Landon C, Meudal H, Boulanger N, Bulet P, Vovelle F. Solution structures of stomoxyn and spinigerin, two insect antimicrobial peptides with an alpha-helical conformation. *Biopolymers* 2006; 81(2): 92-103. doi: 10.1002/bip.20370.
 72. Li Z, Mao R, Teng D, Hao Y, Chen H, Wang X, Wang X, Yang N, Wang J. Antibacterial and immunomodulatory activities of insect defensins-DLP2 and DLP4 against multidrug-resistant *Staphylococcus aureus*. *Scientific Reports* 2017; 7: 12124. doi:10.1038/s41598-017-10839-4.
 73. Park SI, Yoe SM. Defensin-like peptide3 from black soldier fly: Identification, characterization, and key amino acids for anti-Gram-negative bacteria. *Entomological Research* 2017; 47(1): 41-47. doi: 10.1111/1748-5967.12214
 74. Li B, Yang N, Wang X, Hao Y, Mao R, Li Z, Wang Z, Teng D, Wang J. An Enhanced Variant Designed From DLP4 Cationic Peptide Against *Staphylococcus aureus* CVCC 546. *Frontier in Microbiology* 2020; 11: 1057. doi: 10.3389/fmicb.2020.01057
 75. Park SI, Yoe SM. A novel cecropin-like peptide from black soldier fly, *Hermetia illucens*: Isolation, structural and functional characterization. *Entomological Research* 2017(2); 47: 115-124. doi: 10.1111/1748-5967.12226
 76. Elhag O, Zhou D, Song Q, Soomro AA, Ci M, Zheng L, Yu Z, Zhang J. Screening, Expression, Purification and Functional Characterization of Novel Antimicrobial Peptide Genes from *Hermetia illucens* (L.). *Plos One* 2017. doi:10.1371/journal.pone.0169582.

ORIGINALITY REPORT

18%

SIMILARITY INDEX

11%

INTERNET SOURCES

14%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Victoria University Student Paper	1%
2	d-nb.info Internet Source	1%
3	Azza M. Khedre, Tarek G. Ismail, Gehad A. Hashem, Islam M. Zakaria. "In vitro antibacterial activity and synergetic effect of crude extract of the Wohlfahrtia nuba (Diptera: Sarcophagidae) flesh fly larvae", Brazilian Journal of Microbiology, 2023 Publication	1%
4	coek.info Internet Source	1%
5	pdffox.com Internet Source	1%
6	pdfs.semanticscholar.org Internet Source	1%
7	docplayer.net Internet Source	<1%

8	www.info.pmda.go.jp Internet Source	<1 %
9	A. Józefiak, R. Engberg. "Insect proteins as a potential source of antimicrobial peptides in livestock production. A review", <i>Journal of Animal and Feed Sciences</i> , 2017 Publication	<1 %
10	Lucian T. Miron, Richard P. Postma, Guido Bosch, Michel H.M. Eppink. " Preliminary Evaluation of Aqueous Protein Extraction from Black Soldier Fly Larvae (L.) ", <i>Industrial Biotechnology</i> , 2019 Publication	<1 %
11	citeseerx.ist.psu.edu Internet Source	<1 %
12	link.springer.com Internet Source	<1 %
13	www.cell.com Internet Source	<1 %
14	www.mdpi.com Internet Source	<1 %
15	www.qb.uson.mx Internet Source	<1 %
16	Omnia Abass, Asmaa Haggag, Rawda Badawy, Eman Essa. "Microbial Stress Resistance of <i>Eristalis tenax</i> Rat-Tailed	<1 %

Maggots", Egyptian Academic Journal of Biological Sciences. A, Entomology, 2020

Publication

17

patentimages.storage.googleapis.com

Internet Source

<1 %

18

Carmen Scieuzo, Fabiana Giglio, Roberta Rinaldi, Marilena E. Lekka et al. "In Vitro Evaluation of the Antibacterial Activity of the Peptide Fractions Extracted from the Hemolymph of *Hermetia illucens* (Diptera: Stratiomyidae)", *Insects*, 2023

Publication

<1 %

19

Submitted to University of Warwick

Student Paper

<1 %

20

Hueih Min Chen, Wei Wang, David Smith, Siu Chiu Chan. "Effects of the anti-bacterial peptide cecropin B and its analogs, cecropins B-1 and B-2, on liposomes, bacteria, and cancer cells", *Biochimica et Biophysica Acta (BBA) - General Subjects*, 1997

Publication

<1 %

21

Bexfield, A.. "Detection and partial characterisation of two antibacterial factors from the excretions/secretions of the medicinal maggot *Lucilia sericata* and their activity against methicillin-resistant *Staphylococcus aureus* (MRSA)", *Microbes and Infection*, 200411

<1 %

22

Submitted to National University of Singapore

Student Paper

<1 %

23

epub.ub.uni-greifswald.de

Internet Source

<1 %

24

patents.justia.com

Internet Source

<1 %

25

Submitted to University of Sheffield

Student Paper

<1 %

26

Allan Berezcki. "Ressonadores dinamicamente estáveis de com meio ativo birrefringente de Nd:YAG", Universidade de Sao Paulo, Agencia USP de Gestao da Informacao Academica (AGUIA), 2021

Publication

<1 %

27

Submitted to Far Eastern University

Student Paper

<1 %

28

Submitted to J S S University

Student Paper

<1 %

29

Kieran Magee, Joe Halstead, Richard Small, Iain Young. "Valorisation of Organic Waste By-Products Using Black Soldier Fly (*Hermetia illucens*) as a Bio-Convertor", Sustainability, 2021

Publication

<1 %

Submitted to University of Melbourne

30

Student Paper

<1 %

31

www.agrobiology.ru

Internet Source

<1 %

32

www.esbjergpaints.dk

Internet Source

<1 %

33

www.scpscience.com

Internet Source

<1 %

34

Submitted to Afyon Kocatepe University

Student Paper

<1 %

35

Zechao Hu, Handong Li, Sha Liu, Rongrong Xue, Jian Sun, Hong Ji. "Assessment of black soldier fly (*Hermetia illucens*) larvae meal as a potential substitute for soybean meal on growth performance and flesh quality of grass carp *Ctenopharyngodon idellus*", *Animal Nutrition*, 2023

Publication

<1 %

36

www.iwmi.cgiar.org

Internet Source

<1 %

37

Mochamad Firmansyah, Muhammad Yusuf Abduh. "Production of protein hydrolysate containing antioxidant activity from *Hermetia illucens*", *Heliyon*, 2019

Publication

<1 %

38

Internet Source

<1 %

39

nlist.inflibnet.ac.in

Internet Source

<1 %

40

psasir.upm.edu.my

Internet Source

<1 %

41

Diane M. O'Brien, Carol L. Boggs, Marilyn L. Fogel. "Making eggs from nectar: the role of life history and dietary carbon turnover in butterfly reproductive resource allocation", *Oikos*, 2004

Publication

<1 %

42

uploads.consultaremedios.com.br

Internet Source

<1 %

43

www.phytopharmajournal.com

Internet Source

<1 %

44

Ülkü Yılmaz, Hasan Küçükbay, Nihat Şireci, Mehmet Akkurt, Selami Günal, Rıza Durmaz, M. Nawaz Tahir. "Synthesis, microwave-promoted catalytic activity in Suzuki-Miyaura cross-coupling reactions and antimicrobial properties of novel benzimidazole salts bearing trimethylsilyl group", *Applied Organometallic Chemistry*, 2011

Publication

<1 %

45

"Recent Advances in Edible Fats and Oils Technology", Springer Science and Business Media LLC, 2022

Publication

<1 %

46

Bizimenyera, S.E., G.E. Swan, H. Chikoto, and J.N. Eloff. "Rationale for using *Peltophorum africanum* (Fabaceae) extracts in veterinary medicine", Journal of the South African Veterinary Association, 2012.

Publication

<1 %

47

Julia Depta, Paulina Małkowska, Monika Wysokińska, Karolina Todorska et al. "Therapeutic Role of Antimicrobial Peptides in Diabetes Mellitus", Biologics, 2022

Publication

<1 %

48

K.C. Surendra, Jeffery K. Tomberlin, Arnold van Huis, Jonathan A. Cammack, Lars-Henrik L. Heckmann, Samir Kumar Khanal. "Rethinking organic wastes bioconversion: Evaluating the potential of the black soldier fly (*Hermetia illucens* (L.)) (Diptera: Stratiomyidae) (BSF)", Waste Management, 2020

Publication

<1 %

49

Lingyan Kong, Gang Li, Xiangsheng Wang. "Mild oxidation of thiophene over TS-1/H₂O₂", Catalysis Today, 2004

Publication

<1 %

50

Mahmoud Hussein, Viju V. Pillai, Joshua M. Goddard, Hui G. Park, Kumar S. Kothapalli, Deborah A. Ross, Quirine M. Ketterings, J. Thomas Brenna, Mark B. Milstein, Helene Marquis, Patricia A. Johnson, Jan P. Nyrop, Vimal Selvaraj. "Sustainable production of housefly (*Musca domestica*) larvae as a protein-rich feed ingredient by utilizing cattle manure", PLOS ONE, 2017

Publication

<1 %

51

Matthias Leippe. "Amoebapores, a family of membranolytic peptides from cytoplasmic granules of *Entamoeba histolytica*: isolation, primary structure, and pore bacterial cytoplasmic membranes", Molecular Microbiology, 12/1994

Publication

<1 %

52

Monika Kaczor, Piotr Bulak, Kinga Proc-Pietrycha, Marina Kirichenko-Babko, Andrzej Bieganski. "The Variety of Applications of *Hermetia illucens* in Industrial and Agricultural Areas—Review", Biology, 2022

Publication

<1 %

53

Submitted to Universiti Teknologi Malaysia

Student Paper

<1 %

54

edepot.wur.nl

Internet Source

<1 %

55	journalarrb.com Internet Source	<1 %
56	www.dovepress.com Internet Source	<1 %
57	www.i-scholar.in Internet Source	<1 %
58	www.juriskes.com Internet Source	<1 %
59	www.koreascience.or.kr Internet Source	<1 %
60	www.smujo.id Internet Source	<1 %
61	Afsaneh Moslemi Mehni, Mohammad Rahnamaeian, Nader Hassanzadeh, Hamid Forootan Far. "Pseudomonas tolaasii, the causal agent of mushroom brown blotch, is susceptible to insect cecropins", Journal of Plant Pathology, 2023 Publication	<1 %
62	Eunice C. Y. Li-Chan, Shih-Li Hunag, Chia-Ling Jao, Kit-Pan Ho, Kuo-Chiang Hsu. "Peptides Derived from Atlantic Salmon Skin Gelatin as Dipeptidyl-peptidase IV Inhibitors", Journal of Agricultural and Food Chemistry, 2012 Publication	<1 %

63

Osama Elhag, Dingzhong Zhou, Qi Song, Abdul Aziz Soomro, Minmin Cai, Longyu Zheng, Ziniu Yu, Jibin Zhang. "Screening, Expression, Purification and Functional Characterization of Novel Antimicrobial Peptide Genes from *Hermetia illucens* (L.)", PLOS ONE, 2017

Publication

<1 %

64

B.A. Rumpold, H. Speckmann, O. Schlüter, W. Kloas, A. Prochnow. " Potentials of a biogenic residue-based production of as fish meal replacement in aquafeed for in Germany ", Journal of Insects as Food and Feed, 2018

Publication

<1 %

65

Damian Józefiak, Agata Józefiak, Bartosz Kierończyk, Mateusz Rawski et al. "1. Insects – A Natural Nutrient Source for Poultry – A Review", Annals of Animal Science, 2016

Publication

<1 %

66

Foroumadi, A.. "Synthesis and antibacterial activity of N-[2-(5-bromothiophen-2-yl)-2-oxoethyl] and N-[(2-5-bromothiophen-2-yl)-2-oximinoethyl] derivatives of piperazinyl quinolones", Bioorganic & Medicinal Chemistry Letters, 20051015

Publication

<1 %

67

Waldemar Tejchman, Bartosz Orwat, Izabela Korona-Główniak, Anna Barbasz et al. "Highly

<1 %

efficient microwave synthesis of rhodanine and 2-thiohydantoin derivatives and determination of relationships between their chemical structures and antibacterial activity", RSC Advances, 2019

Publication

68

Yongqi Meng, Xiuxia Zhang, Zelong Zhang, Jiajun Li, Peihua Zheng, Juntao Li, Jiarui Xu, Jianan Xian, Yaopeng Lu. "Effects of Microorganisms on Growth Performance, Body Composition, Digestive Enzyme Activity, Intestinal Bacteria Flora and Antimicrobial Peptide (AMP) Content of Black Soldier Fly Larvae (*Hermetia illucens*)", Animals, 2023

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On

RJPT

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/0

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
