

# Demographic diversity of covid-19 patients

*by* Diah Wahyuningsih

---

**Submission date:** 22-Sep-2023 09:32AM (UTC+0800)

**Submission ID:** 2173156515

**File name:** Demographic\_diversity\_of\_covid-19\_patients.pdf (487.43K)

**Word count:** 2991

**Character count:** 16015

## Demographic diversity of COVID-19 patients treated at ICU Special Hospital for Infections Surabaya

Diah Wahyuningsih<sup>1</sup>, Anna Surgean Veterini<sup>1</sup>, Hamzah Hamzah<sup>2</sup>, Lucky Andriyanto<sup>1</sup>

14

### Abstract

**Background:** The diversity demography of coronavirus disease 2019 (COVID-19) has prompted us to collect data in our workplace. These data are essential because they can serve as a data bank of demographics in one big hospital in East Java. We analyzed patients' characteristics with COVID-19 admitted to the Intensive Care Unit (ICU) Special Hospital for Infections, Airlangga University, Surabaya.

**Methods:** Retrospective study from medical record of 180 patients with confirmed COVID-19 admitted to the ICU Special Hospital for Infections, Airlangga University of Surabaya between April - September 2020.

**Results:** Most of the patients were male (67.2%), with median age was 55 (22-83) years. The body mass index (BMI) data consisted of normal (48.9%), overweight (39.4%), obesity class I 7.8%, obesity class II 1.1%, obesity class III 0.6%, and underweight 2.2%. We found severe acute respiratory distress syndrome (ARDS) in 63.9%. The use of invasive mechanical ventilation was 74% of the total patients. Most of the

patients (85%) had comorbidity: hypertension, diabetes mellitus, and geriatrics. The average length of stay in the ICU was 8.5 days. We transferred 29% of patients to a low-care ward, and 66.7% died. We identified gastrointestinal symptoms on admission to the ICU were 43.3%, predominantly by nausea and vomiting. Forty-six point seven percent of patients with gastrointestinal symptoms during hospitalization consisted of gastric retention and diarrhea, some with hematin.

**Conclusion:** The demographic data we present above are limited in our area. The demographic data of COVID-19 patients in other places may be different from the information we obtained. However, data like this may represent the patient's condition in areas similar to ours. Besides, this data can warn that the patient's condition, as in our data, requires special attention. It is necessary to add data from all corners of Indonesia to represent the demographic data of COVID-19 patients in Indonesia.

**Key words:** COVID-19, patient's characteristic, clinical complaints, demography.

32

<sup>1</sup>Department of Anesthesiology and Intensive Therapy, Special Hospital for Infections, Universitas Airlangga, Surabaya, Indonesia

<sup>2</sup>Anesthesiology and Intensive Care Department, Faculty of Medicine, Universitas Airlangga-Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

### Address for correspondence:

Anna Surgean Veterini  
Department of Anesthesiology and Intensive Therapy, Special Hospital for Infections, Universitas Airlangga, Surabaya, Indonesia  
Campus C Mulyorejo, Universitas Airlangga  
Tel: +6281802080000  
Email: annasurgeonveterini@gmail.com, annasurgeon@fk.un-air.ac.id

### Introduction

The cause of acute respiratory distress syndrome (ARDS) in severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection is a cytokine storm, an uncontrolled systemic inflammatory response caused by the release of large amounts of proinflammatory cytokines and chemokines. (1,2) This excessive immune response can cause lung damage and fibrosis, resulting in functional disability. (3) Some literature say that certain demographic conditions have a predisposition to being more easily infected with coronavirus disease 2019 (COVID-19). (4,5) Therefore, we collected demographic data on COVID-19 patients in Intensive Care Unit (ICU) Special Hospital for Infections

to strengthen existing data, even though the size of the study subjects we took was relatively small. The demographic data we collected were gender, age, body mass index (BMI), comorbid, ARDS type, length of stay (LOS), and outcome.

Previous studies have stated that COVID-19 does not always start with respiratory complaints but can also manifest in other organs, for example, gastrointestinal tract (GIT). The study by Xiao et al. (2020) reported that they could detect the virus in feces, even though it was clear in airway samples. (6) The explanation is that angiotensin converting enzyme 2 (ACE-2) protein is also highly expressed in intestinal epithelial cells, which functions as a co-receptor to enter nutrients into the intestine. (7) The theory mentioned before explains why the symptoms of COVID-19 are not only in the respiratory tract but also in the gastrointestinal tract. Many patients experience gastrointestinal disorders such as diarrhea, abdominal pain, etc., before being confirmed as COVID-19. Our retrospective descriptive study showed complaints of GIT before being admitted to the ICU Special Hospital for Infections, as well as criticisms of GIT while in the ICU Special Hospital for Infections. These findings expect to add data on the characteristics of COVID-19 patients that may have the advantage for the development of handling COVID-19.

## 12 Materials and methods

Our study was a retrospective study by taking the medical record of COVID-19 patients in the Intensive Care Unit of the Special Hospital for Infections, Airlangga University Surabaya, between April-September 2020. According to Berlin criteria, all cases included in this study were male or female adults aged  $\geq 18$  years, confirmed positive for SARS-CoV-2 infection, diagnosed with ARDS. Exclusion criteria were patients who finished ICU care before 3 days (discharge, moved to another room, died, or were referred to another hospital) because there was a periodic laboratory examination every 3 days to evaluate the patient's condition. We took the data regarding the gender, age, BMI, comorbidities, ARDS category, use of invasive mechanical ventilation, length of stay, outcome, and gastrointestinal symptoms. No conflict of interest regarding the publication. The author funded this research. We got ethical clearance from the Ethical Committee at Airlangga University Hospital, Surabaya, Indonesia (179/KEP/2020).

## Results

A total of 180 medical records met the criteria. **Table 1**

**Table 1** shows that subjects were dominated by male at 67.2%, while female was 32.8%. This study shows a minimum age of 22 years and a maximum age of 83 years, with a median age of 55 years of research subjects. BMI was normal at 48.9%, followed by overweight at 39.4%, obesity class I 7.8%, obesity class II 1.1%, obesity class III 0.6%, and underweight 2.2%. We found severe ARDS in 115 patients (63.9%), moderate ARDS in 58 patients (32.2%), mild ARDS in 7 patients (3.9%). In this study, 133 patients (74%) were under invasive mechanical ventilation. One hundred fifty-three patients (85%) had comorbidities; the top three were hypertension in 91 patients, diabetes mellitus in 89 patients, and geriatrics in 40 patients. The mean length of stay in the Intensive Care Unit was 8.5 (3-34) days. Fifty-two subjects (29%) were transferred to low-care rooms, and 120 subjects (66.7%) died. Overall data on gastrointestinal symptoms at ICU were identified in 78 patients (43.3%), predominantly nausea (48 patients) and vomiting (32 patients); the rest were decreased appetite, abdominal pain, diarrhea, bloating, and constipation. Meanwhile, gastrointestinal symptoms experienced during hospitalization were gastric retention in 76 patients ( $>100$  ml, some patients with hematin) and diarrhea in 14 patients.

## Discussion

In this study, male gender dominated as much as 67.2%. Previous research supported that diabetes mellitus, hypertension, male gender, and active smoking are risk factors for SARS-CoV-2 infection. This phenomenon is because of the increasing number of ACE-2 receptor expressions in this population. (8) Women are said to have more robust innate and humoral immune responses than men, giving them a better response to many viral, fungal, bacterial, and parasitic infections. One meta-analysis study said that the need for ICU care was almost three times higher in male patients, and the likelihood of dying was higher than that of women. Socio-cultural differences and gender-based behavior also contribute to the sex differences seen in the severity of COVID-19. (9)

In this study, the patients' mean age was 55 years, with the youngest 22 years, and the oldest 83 years. There is significant variation in cases of the severity of COVID-19 across countries. However, the main focus of COVID-19 is predominantly on older people; the study by Sudharsanan et al. (2020) showed that people of all ages were equally susceptible to severe acute metabolic syndrome infection. (10) Most of the patients (48.9%) had normal body mass index, 39.4% were overweight, and 9.5% obese. If

we compare the data we got, there was no difference in the risk of COVID-19 between normal BMI, overweight, and obesity. Several things that may explain this phenomenon are that BMI cannot represent fat-free adipose tissue mass. So, people on a normal BMI scale can experience an increase in body fat percentage, especially in the visceral area; this is called normal weight obesity. (11) Visceral adiposity, regardless of BMI, is associated with a higher risk for critical illness in patients with COVID-19. (12,13)

We found severe ARDS in 115 patients (63.9%), moderate ARDS in 58 patients (32.2%), and mild ARDS in 7 patients (3.9%). In this study, 133 patients (73.9%) used invasive mechanical ventilation. Based on the previous literature, the reported mortality rate in COVID-19 ranged from 50 to 97% in those requiring mechanical ventilation. (14,15) Patients with moderate to severe ARDS required invasive mechanical ventilation and had a poor prognosis.

Total subjects with comorbidity were 153 patients (85%), mainly hypertension and diabetes mellitus, while geriatric was the third most common comorbid. Comorbidity can exacerbate the manifestations of COVID-19. In smokers, hypertension, and diabetes mellitus, it is suspected that there is an increase in ACE-2 receptor expression. (8) Older age is associated with decreased acquired immunity. It is also important to remember that in the elderly, body weight and muscle mass begin to fall. Still, the relative fat mass increases, especially in those with comorbidities such as cardiovascular and respiratory conditions. Older age is also associated with more hypertension and diabetes mellitus due to stiff blood vessels and impaired metabolic efficiency. Older people also have fewer cardiorespiratory reserves to fight COVID-19 infection. (16)

Our study presents the median length of stay in ICU was 8.5 days (3-34 days). The length of stay in our research is longer when we compare with the study by Rees and colleagues, which analyzed the length of stay in the ICU based on eight studies, 4 studies within China and 4 studies outside China, had a median of 8 (5-13) days for China, and 7 (4-11) days outside of China. (17) Understanding how long the hospitalization of COVID-19 patients is critical for planning and predicting bed capacity and staff and associated medical equipment needs.

Based on data analysis of 180 study subjects, 52 subjects (29%) were transferred to low-care rooms, and 120 subjects (66.7%) died. Although experi-

ence with COVID-19 continues to develop, the reported mortality rate was 50 to 97% in those requiring mechanical ventilation. (14) The mortality rate in our study was significantly higher than the published mortality rate for intubated patients with H1N1 influenza (35-46%) and other causes of ARDS. (18)

Gastrointestinal complaints were the main complaints of patients before being hospitalized, in addition to respiratory complaints. Gastrointestinal complaints before hospitalization were in 78 patients (43.3%), predominantly nausea (48 patients) and vomiting (32 patients); the rest were decreased appetite, stomachache, diarrhea, bloating, and constipation. Gastrointestinal symptoms during hospitalization included gastric retention in 76 patients (>100 ml, some patients had hematin) and diarrhea in 14 patients. Critically ill patients are prone to diffuse mucosal injury of the stomach, producing upper gastrointestinal bleeding associated with increased morbidity and mortality. The main risk factors for stress-related mucosal bleeding are prolonged mechanical ventilation and coagulopathy. (19) During critical illness, changes occur in the gut microflora, including circulating stress hormones, intestinal ischemia, immunosuppression, antibiotics, and nutritional deficiencies. Another reason for changes in the gut microbiota is the massive use of antibiotics. (20) It can cause dysbiosis and increase susceptibility to new infections and inflammatory disorders and cause antibiotic-related diarrhea.

### Conclusion

The demographic data we present above are limited in our area. The demographic data of COVID-19 patients in other places may be different from the information we obtained. However, data like this may represent the patient's condition in areas similar to ours. Besides, this data can warn that the patient's condition, as in our data, requires special attention. It is necessary to add data from all corners of Indonesia to represent the demographic data of COVID-19 patients in Indonesia.

### Funding source

This research did not receive any specific grant from funding agencies in the public, commercial, or non-profit sectors.

### Conflict of interests

All authors have no conflict of interest.

**Table 1.** Characteristics of COVID-19 patients in Intensive Care Unit (ICU)

| Characteristic                         |                               | Total, n (%) | Mean±SD     | Median | Range     |
|--|-------------------------------|--------------|-------------|--------|-----------|
| Gender                                 | Male                          | 121 (67.2%)  |             |        |           |
|  | Female                        | 59 (32.8%)   |             |        |           |
| Age                                    |                               | 180 (100%)   | 55.17±11.14 | 55     | 22-83     |
| BMI                                    |                               | 180 (100%)   | 25.6±3.90   | 25     | 16.6-45.8 |
| BMI category (kg/m <sup>2</sup> )      | Underweight (<18.5)           | 4 (2.2%)     |             |        |           |
|  | Normal (18.5-24.9)            | 88 (48.9%)   |             |        |           |
|  | Overweight (25.0-29.9)        | 71 (39.4%)   |             |        |           |
|  | Obesity class I (30.0-34.9)   | 14 (7.8%)    |             |        |           |
|  | Obesity class II (35.0-39.9)  | 2 (1.1%)     |             |        |           |
|  | Obesity class III (>40)       | 1 (0.6%)     |             |        |           |
| ARDS category                          | Mild                          | 7 (3.9%)     |             |        |           |
|  | Moderate                      | 58 (32.2%)   |             |        |           |
|  | Severe                        | 115 (63.9%)  |             |        |           |
| Use of invasive mechanical ventilation | Yes                           | 133 (74%)    |             |        |           |
|  | No                            | 47 (26%)     |             |        |           |
| Comorbid                               | Yes                           | 153 (85%)    |             |        |           |
|  | No                            | 27 (15%)     |             |        |           |
| Distribution of comorbid               | Hypertension                  | 91           |             |        |           |
|  | DM type 2                     | 89           |             |        |           |
|  | Geriatrics                    | 40           |             |        |           |
|  | CKD (regular HD)              | 13           |             |        |           |
|  | Obesity                       | 10           |             |        |           |
|  | AKI                           | 7            |             |        |           |
|  | Post surgery                  | 5            |             |        |           |
|  | Asthma                        | 3            |             |        |           |
|  | CVA history                   | 3            |             |        |           |
|  | Pregnancy                     | 2            |             |        |           |
| Others                                 | 27                            |              |             |        |           |
| Length of stay                         |                               | 180 (100)    | 10.4±6.96   | 8.5    | 3-34      |
| Outcome                                | Discharged from hospital      | 7 (3.9%)     |             |        |           |
|  | Transferred to low-care rooms | 52 (29%)     |             |        |           |
|  | Referred to another hospital  | 1 (0.6%)     |             |        |           |
|  | Died                          | 120 (66.7%)  |             |        |           |
| Gastrointestinal symptoms              | When entering the ICU:        | 78 (43.3%)   |             |        |           |
|  | - Nausea                      | 48           |             |        |           |
|  | - Vomiting                    | 32           |             |        |           |
|  | - Decreased appetite          | 18           |             |        |           |
|  | - Stomachache                 | 14           |             |        |           |
|  | - Diarrhea                    | 12           |             |        |           |
|  | - Bloating                    | 3            |             |        |           |
|  | - Constipation                | 2            |             |        |           |
|  | While in the ICU:             | 84 (46.7%)   |             |        |           |
|  | - Gastric retention*          | 76           |             |        |           |
| - Diarrhea                             | 14                            |              |             |        |           |

Legend: COVID-19=coronavirus disease 2019; SD=standard deviation; BMI=body mass index; ARDS=acute respiratory distress syndrome; DM=diabetes mellitus; CKD=chronic kidney disease; HD=hemodialysis; AKI=acute kidney injury; CVA=cerebrovascular accident. \*(>100 ml, some patients with hematin).

## References

1. Li L, Huang Q, Wang DC, Ingbar DH, Wang X. Acute lung injury in patients with COVID-19 infection. *Clin Transl Med* 2020;10:20-7.
2. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497-506.
3. Zumla A, Hui DS, Azhar EI, Memish ZA, Maeurer M. Reducing mortality from 2019-nCoV: host-directed therapies should be an option. *Lancet (London, England)* 2020;395:e35-6.
4. Karmakar M, Lantz PM, Tipimani R. Association of Social and Demographic Factors With COVID-19 Incidence and Death Rates in the US. *JAMA Netw Open* 2021;4:e2036462.
5. Monod M, Blenkinsop A, Xi X, Hebert D, Bershan S, Tietze S, et al. Age groups that sustain resurging COVID-19 epidemics in the United States. *Science* 2021;371:1-12.
6. Xiao F, Tang M, Zheng X, Liu Y, Li X, Shan H. Evidence for Gastrointestinal Infection of SARS-CoV-2. *Gastroenterology* 2020;158:1831-3.e3.
7. Hashimoto T, Perlot T, Rehman A, Trichereau J, Ishiguro H, Paolino M, et al. ACE2 links amino acid malnutrition to microbial ecology and intestinal inflammation. *Nature* 2012;487:477-81.
8. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med* 2020;8:e21.
9. Peckham H, de Grijter NM, Raine C, Radziszewska A, Ciurtin C, Wedderburn LR, et al. Male sex identified by global COVID-19 meta-analysis as a risk factor for death and ITU admission. *Nat Commun* 2020;11:6317.
10. Sudharsanan N, Didzun O, Bämighausen T, Geldsetzer P. The Contribution of the Age Distribution of Cases to COVID-19 Case Fatality Across Countries: A Nine-Country Demographic Study. *Ann Intern Med* 2020;173:714-20.
11. Kapoor N, Furler J, Paul TV, Thomas N, Oldenburg B. Normal Weight Obesity: An Underrecognized Problem in Individuals of South Asian Descent. *Clin Ther* 2019;41:1638-42.
12. Battisti S, Pedone C, Napoli N, Russo E, Agnoletti V, Nigra SG, et al. Computed Tomography Highlights Increased Visceral Adiposity Associated With Critical Illness in COVID-19. *Diabetes Care* 2020;43:e129-30.
13. Petersen A, Bressemer K, Albrecht J, Thieß H-M, Vahldiek J, Hamm B, et al. The role of visceral adiposity in the severity of COVID-19: Highlights from a unicenter cross-sectional pilot study in Germany. *Metabolism* 2020;110:154317.
14. Zou X, Chen K, Zou J, Han P, Hao J, Han Z. Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front Med* 2020;14:185-92.
15. Wu D, Yang XO. TH17 responses in cytokine storm of COVID-19: An emerging target of JAK2 inhibitor Fedratinib. *J Microbiol Immunol Infect* 2020;53:368-70.
16. Sattar N, McInnes IB, McMurray JJV. Obesity Is a Risk Factor for Severe COVID-19 Infection: Multiple Potential Mechanisms. *Circulation* 2020;142:4-6.
17. Rees EM, Nightingale ES, Jafari Y, Waterlow N, Clifford S, Pearson CAB, et al. COVID-19 length of hospital stay: a systematic review and data synthesis. *BMC Med* 2020;18:270.
18. Zambon M, Vincent J-L. Mortality Rates for Patients With Acute Lung Injury/ARDS Have Decreased Over Time. *Chest* 2008;133:1120-7.
19. Duerksen DR. Stress-related mucosal disease in critically ill patients. *Best Pract Res Clin Gastroenterol* 2003;17:327-44.
20. Guan W-J, Ni Z-Y, Hu Y, Liang W-H, Ou C-Q, He J-X, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020;382:1708-20.

Copyright of Critical Care & Shock is the property of Critical Care & Shock Journal and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.

# Demographic diversity of covid-19 patients

## ORIGINALITY REPORT

15%

SIMILARITY INDEX

11%

INTERNET SOURCES

12%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

|   |  |    |
|---|--|----|
| 1 | <a href="http://rcastoragev2.blob.core.windows.net">rcastoragev2.blob.core.windows.net</a><br>Internet Source  | 1% |
| 2 | <a href="http://www.biorxiv.org">www.biorxiv.org</a><br>Internet Source  | 1% |
| 3 | T. S. Hettiarachchi, A. Askari, E. Rudge, L. T. Hao, S. Sarwar, D. Dowsett, A. El Hadi, Irshad Shaikh. "Comparison of robotic vs laparoscopic left-sided colorectal cancer resections", Journal of Robotic Surgery, 2022<br>Publication                    | 1% |
| 4 | <a href="http://asjo.in">asjo.in</a><br>Internet Source  | 1% |
| 5 | <a href="http://researchonline.jcu.edu.au">researchonline.jcu.edu.au</a><br>Internet Source  | 1% |
| 6 | Po-Ren Hsueh, Sung-Jung Ho, Po-Chuen Hsieh, I-Min Liu, Shio-Shin Jean. "Use of Multiple Doses of Intravenous Infusion of Umbilical Cord-Mesenchymal Stem Cells for the Treatment of Adult Patients with Severe COVID-19-Related Acute Respiratory Distress | 1% |



# Syndrome: Literature Review", Stem Cells International, 2023

Publication

---

|       |   |      |
|-------|---|------|
| 7     | Premananda Bharati, Susmita Bharati, Manoranjan Pal, Suman Chakraborty, Ranjan Gupta. "Chronic Energy Deficiency Among Indian Women by Residential Status", Ecology of Food and Nutrition, 2008               | 1 %  |
| <hr/> |   |      |
| 8     | Stefano Puliatti, Ahmed Eissa, Radwa Eissa, Marco Amato et al. "COVID-19 and Urology: A Comprehensive Review of the Literature", BJU International, 2020  | 1 %  |
| <hr/> |   |      |
| 9     | Leonel Nunes, Catarina Meireles, Carlos Gomes, Nuno Ribeiro. "Acacia dealbata Link. Aboveground Biomass Assessment: Sustainability of Control and Eradication Actions to Reduce Rural Fires Risk", Fire, 2022 | 1 %  |
| <hr/> |   |      |
| 10    | <a href="http://search.ricest.ac.ir">search.ricest.ac.ir</a>  | 1 %  |
| <hr/> |   |      |
| 11    | <a href="http://www.dovepress.com">www.dovepress.com</a>  | <1 % |
| <hr/> |   |      |
| 12    | Sunil Kumar Dodani, Asma Nasim, Tahir Aziz, Anwar Naqvi. "The efficacy of isoniazid prophylaxis in renal transplant recipients in a   | <1 % |

# high tuberculosis burden country", Transplant Infectious Disease, 2021

Publication

13

Chiyong Cha, Gumhee Baek. "Symptoms and management of long COVID: A scoping review", Journal of Clinical Nursing, 2021

Publication

<1 %

14

[ijms.sums.ac.ir](http://ijms.sums.ac.ir)

Internet Source

<1 %

15

[pubmed.ncbi.nlm.nih.gov](http://pubmed.ncbi.nlm.nih.gov)

Internet Source

<1 %

16

Kentaro Shimizu, Haruhiko Hirata, Daijiro Kabata, Natsuko Tokuhira et al. "Ivermectin administration is associated with lower gastrointestinal complications and greater ventilator-free days in ventilated patients with COVID-19: A propensity score analysis", Journal of Infection and Chemotherapy, 2021

Publication

<1 %

17

Umair Mallick. "Chapter 19 Gender Disparity in CoViD19", Springer Science and Business Media LLC, 2022

Publication

<1 %

18

[cms.hasekidergisi.com](http://cms.hasekidergisi.com)

Internet Source

<1 %

19

[danke-mutasd.com](http://danke-mutasd.com)

Internet Source

<1 %

|    |  |      |
|----|--|------|
| 20 | <a href="https://journals.lww.com">journals.lww.com</a><br>Internet Source   | <1 % |
| 21 | <a href="http://www.wjgnet.com">www.wjgnet.com</a><br>Internet Source  | <1 % |
| 22 | Ennio Giulio Favalli, Francesca Ingegnoli, Orazio De Lucia, Gilberto Cincinelli, Rolando Cimaz, Roberto Caporali. "COVID-19 infection and rheumatoid arthritis: Faraway, so close!", <i>Autoimmunity Reviews</i> , 2020<br>Publication | <1 % |
| 23 | Jefferson M. Sesler. "Stress-related Mucosal Disease in the Intensive Care Unit : An Update on Prophylaxis", <i>AACN Advanced Critical Care</i> , 04/2007<br>Publication   | <1 % |
| 24 | Jürgen Schölmerich. "Therapy Insight: prophylaxis of stress-induced gastrointestinal bleeding in critically ill patients", <i>Nature Clinical Practice Gastroenterology &amp; Hepatology</i> , 10/2007<br>Publication                  | <1 % |
| 25 | <a href="http://bmcmmedresmethodol.biomedcentral.com">bmcmmedresmethodol.biomedcentral.com</a><br>Internet Source  | <1 % |
| 26 | <a href="http://doaj.org">doaj.org</a><br>Internet Source  | <1 % |
| 27 | <a href="http://docksci.com">docksci.com</a><br>Internet Source  | <1 % |

---

|    |   |      |
|----|---|------|
| 28 | <a href="http://e-journal.unair.ac.id">e-journal.unair.ac.id</a><br>Internet Source   | <1 % |
| 29 | <a href="http://www.accjournal.org">www.accjournal.org</a><br>Internet Source   | <1 % |
| 30 | <a href="http://www.medrxiv.org">www.medrxiv.org</a><br>Internet Source   | <1 % |
| 31 | <a href="http://www.researchsquare.com">www.researchsquare.com</a><br>Internet Source   | <1 % |
| 32 | "Euroanaesthesia 2004: Joint Meeting of the European Society of Anaesthesiologists and European Academy of Anaesthesiology Lisbon, Portugal, 5–8 June 2004", European Journal of Anaesthesiology, 09/14/2005<br>Publication   | <1 % |
| 33 | Alexandre de Fátima Cobre, Monica Surek, Raquel de Oliveira Vilhena, Beatriz Böger et al. "Influence of foods and nutrients on COVID-19 recovery: A multivariate analysis of data from 170 countries using a generalized linear model", Clinical Nutrition, 2021<br>Publication | <1 % |
| 34 | Laura Timelli, Giuseppina Liuzzi, Alessandro Cannavacciuolo, Nicola Petrosillo, Vincenzo Puro, Enrico Girardi. "Association of COVID-19 case fatality rate with disease burden: an ecological analysis in Italy during the first  | <1 % |

---

# wave", International Journal of Infectious Diseases, 2021

Publication

---

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On

# Demographic diversity of covid-19 patients

---

GRADEMARK REPORT

---

FINAL GRADE

GENERAL COMMENTS

**/100**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---