

# Association between anthropometric

*by Nur Aisiyah Widjaja*

---

**Submission date:** 16-Oct-2023 04:13PM (UTC+0800)

**Submission ID:** 2197288654

**File name:** Association\_between\_anthropometric.pdf (237.57K)

**Word count:** 2098

**Character count:** 11792

## Association between anthropometric parameters and carotid intima-media thickness in obese adolescents

\*Nur Aisiyah Widjaja<sup>1</sup>, Rendi Aji Prihaningtyas<sup>1</sup>, Meta Herdiana Hanindita<sup>1</sup>, Roedi Irawan<sup>1</sup>, Retno Handajani<sup>1</sup>, IDG Ugrasena<sup>1</sup>

*Sri Lanka Journal of Child Health*, 2021; **50**(1): 28-31

### Abstract

**Introduction:** Prevalence of obesity among adolescents is increasing in developing countries. Inflammation in obesity causes atherosclerosis which could develop into cardiovascular disease. Carotid intima-media thickness (CIMT) is a non-invasive subclinical marker of atherosclerosis. Determining the association between body size and atherosclerosis may allow early detection of atherosclerosis risk.

**Objectives:** To assess the association between anthropometric parameters and CIMT in obese adolescents.

**Method:** A cross sectional study was conducted on adolescents with central obesity who visited the paediatric clinic of Dr. Soetomo General Hospital, Surabaya, Indonesia. Eating habits were obtained through food recall. Weight, height, waist circumference and thigh circumference were measured for calculation of body mass index (BMI) and waist to hip ratio (WHR). Obesity was defined as BMI >P<sub>95</sub> according to age and sex. CIMT was obtained through B mode ultrasonography on the neck. Analysis was done using Spearman rho' to analyse association between BMI, waist circumference, WHR and CIMT.

**Results:** There were 59 obese adolescents, comprising 32 (54.2%) males and 27 (45.8%) females. BMI, waist circumference and WHR were not significantly associated with CIMT in obese adolescents (p>0.05).

<sup>1</sup>Airlangga University, Surabaya, Indonesia

\*Correspondence: nuril08@yahoo.com

 <https://orcid.org/0000-0002-4253-8760>

(Received on 13 February 2020: Accepted after revision on 20 March 2020)

The authors declare that there are no conflicts of interest

This research received a grant from Airlangga University, Surabaya, Indonesia.

Open Access Article published under the [Creative](#)

[Commons Attribution CC-BY](#)  License

**Conclusions:** In this study, BMI, waist circumference and WHR were not significantly associated with CIMT in obese adolescents.

<sup>12</sup> DOI: <http://dx.doi.org/10.4038/sljch.v50i1.9398>

(Key words: Obesity, CIMT, BMI, waist circumference, waist to hip ratio)

### Introduction

In Indonesia, the prevalence of obesity is increasing among all age groups, including adolescents<sup>1</sup>. The high level of body fat in obesity, is a risk factor for cardiovascular disease<sup>2</sup>. Atherosclerosis starts in obese children and adolescents<sup>3</sup>. Central obesity correlates positively with atherosclerosis progressivity, although the correlation is still inconsistent<sup>4</sup>. Carotid intima-media thickness (CIMT) is a subclinical atherosclerosis marker and a risk factor for cardiovascular disease that could be measured non-invasively and easily<sup>5,6</sup>. However, most studies on this subject have been done in adults in developed countries.

### Objectives

To assess the association between body mass index (BMI), waist circumference, waist to hip ratio (WHR) and CIMT in obese adolescents.

### Method

A cross sectional study was carried out on obese adolescents, 13-16 years of age, at the paediatric clinic of Dr. Soetomo General Hospital, Surabaya, Indonesia. Exclusion criteria comprised consumption of steroids within 6 months before study, dyslipidaemia drugs within 3 months before study, hormonal therapy, alcohol consumption, smoking, and endocrine disorders. Eating habits were obtained through food recall.

**Anthropometric measurements:** Weight (kg) measurement used a digital scale with subjects wearing light clothes, without footwear or other accessories (Seca, Germany). Height (cm) measurement used a stadiometer, with subjects in erect position with no footwear or headwear (Seca, Germany). For calculation of BMI, the following formula was used. BMI = body weight (kg)/ body height (m<sup>2</sup>). Obesity was defined as BMI > P<sub>95</sub> based on age and gender according to CDC 2000

curve. Waist circumference (cm) was measured parallel to the floor using metlin from upper part of iliac crest and bottom border of ribs on midaxillary line at end of expiration. Thigh circumference was measured from the biggest circumference on the buttocks with parallel position to the floor. WHR was the result of waist circumference (cm) divided by thigh circumference (cm).

**CIMT measurement:** This was done using B-mode ultrasonography (Toshiba, Japan) by cardiologists. Examination was done in the supine position with the neck extended and bent to the right.

**Ethical issues:** Ethical approval was obtained from the Ethics Committee of Dr. Soetomo General Hospital, Surabaya, Indonesia (No. 0698/KEPK/X/2018). Written informed consent was obtained from parents of the subjects before commencing study. All data obtained from the subjects were anonymised.

**Statistical analysis:** Quantitative parameters are presented as mean ± standard deviation. Kolmogorov–Smirnov test was utilised to evaluate data normality. Associations between BMI, waist circumference, WHR and CIMT were analysed

using Spearman rho' SPSS with significant p value of <0.05.

**Results**

There were 59 obese adolescents, comprising 32 (54.2%) males and 27 (45.8%) females. The subjects had a mean BMI of 31.99 ± 3.67 kg/m<sup>2</sup>, mean waist circumference of 100.18 ± 10.63 cm, mean thigh circumference of 105.32 ± 8.27 cm, and a mean CIMT of 0.51 ± 0.10 mm (Table 1).

**Table 1: Study subjects' characteristics**

Variable	Mean (SD)
Age (years)	13.89 ± 0.82
Body weight (kg)	80.77 ± 13.35
Body height (cm)	158.76 ± 7.12
Body mass index (kg/m <sup>2</sup> )	31.99 ± 3.67
Waist circumference (cm)	100.18 ± 10.63
Thigh circumference (cm)	105.32 ± 8.27
WHR	0.95 ± 0.6
CIMT (mm)	0.51 ± 0.10

No association between BMI, waist circumference, or WHR and CIMT was found in obese adolescents (p>0.05) (Table 2)

**Table 2: Association between variables**

CIMT	BMI	Waist circumference	WHR
R	0.019	-0.163	-0.032
P	0.886	0.217	0.812

**Discussion**

Obesity gives rise to a chronic inflammation because of an imbalance between pro-inflammatory and anti-inflammatory cytokines<sup>7</sup>. The high body fat level at stomach causes cell dysfunction and increases risk of cardiovascular disorder in adulthood<sup>2,8,9</sup>. Waist circumference can assess risk factors for cardiovascular disease in obese adolescents<sup>10</sup>.

Cardiovascular disease starts with atherosclerosis process in obese children and adolescents<sup>3</sup>. Long obesity duration and central obesity are associated with subclinical heart disease<sup>11</sup>. Obese adolescents have increased CIMT in comparison with adolescents having normal BMI<sup>10,12</sup>. The presence of dyslipidaemia, hypertension, and diabetes mellitus increases CIMT<sup>13</sup>. A previous study showed that left CIMT is better associated with cardiovascular risk compared to right CIMT<sup>14</sup>. In children, no association was found between CIMT and BMI or body fat<sup>15</sup>. However, CIMT tends to increase after the age of 10 due to hormonal changes<sup>16,17</sup>.

This study shows no association between CIMT and BMI, waist circumference, or WHR. This is in accordance with past studies which showed no association between CIMT and BMI or body fat. However, results from a study in a developed country showed that adiposity is associated with CIMT in adolescents<sup>18</sup>. CIMT is associated with waist circumference and WHR in healthy adolescents<sup>14</sup>. In obese adolescents, CIMT is associated with BMI, waist circumference, and body fat percentage<sup>10</sup>.

This study has a number of limitations. Being a cross sectional study, conclusions regarding causal relation could not be drawn. High-resolution B-mode ultrasonography also has low sensitivity and is operator dependent. This could affect the results of CIMT measurements due to undetected small differences<sup>17</sup>. Puberty data were also not available which could have affected the results of this study<sup>14</sup>.

### Conclusions <sup>1</sup>

In this study, BMI, waist circumference and WHR were not significantly associated with CIMT in obese adolescents

### Acknowledgements

The authors thank the Universitas Airlangga, Surabaya, Indonesia for supporting this study and Dr. Soetomo General Hospital, Surabaya for permitting this study.

### References

1. Rachmi CN, Li M, Alison BL. Overweight and obesity in Indonesia: prevalence and risk factors—a literature review. *Public Health* 2017; **147**:20–9. <https://doi.org/10.1016/j.puhe.2017.02.002> PMID: 28404492
2. Going SB, Lohman TG, Cussler EC, Williams DP, Morrison JA, Horn PS. Percent body fat and chronic disease risk factors in U.S. Children and Youth. *American Journal of Preventive Medicine* 2011; **41**:S77–S86. <https://doi.org/10.1016/j.amepre.2011.07.006> PMID: 21961616
3. Fang J, Zhang JP, Luo CX, Yu XM, Lv LQ. Carotid Intima-media thickness in childhood and adolescent obesity relations to abdominal obesity, high triglyceride level and insulin resistance. *International Journal of Medical Sciences* 2011; **7**:278–83. <https://doi.org/10.7150/ijms.7.278> PMID: 20827427 PMID: PMC2934726
4. Chagas P, Caramori P, Barcellos C, Galdino TP, Gomes I, Schwanke CHA. Association of different anthropometric measures and indices with coronary atherosclerotic burden. *Arq. Bras. Cardiol.* 2011; **97**:397–401. <https://doi.org/10.1590/S0066782X2011005000093> PMID: 21971634
5. van den Oord SCH, Sijbrands EJG, ten Kate GL, van Klaveren D, van Domburg RT, van der Steen AFW, Schinkel AFL. Carotid intima-media thickness for cardiovascular risk assessment: Systematic review and meta-analysis. *Atherosclerosis* 2013; **228**:1–11. <https://doi.org/10.1016/j.atherosclerosis.2013.01.025> PMID: 23395523
6. Gao Z, Khoury PR, McCoy CE, Shah AS, Kimball TR, Dolan LM, et al. Adiposity has no direct effect on carotid intima-media thickness in adolescents and young adults: Use of structural equation modeling to elucidate indirect & direct pathways. *Atherosclerosis* 2016; **246**:29–35. <https://doi.org/10.1016/j.atherosclerosis.2015.11.033> PMID: 26752690 PMID: PMC4764416
7. Todendi PF, Possuelo LG, Klinger EI, Reuter CP, Burgos MS, Moura DJ, et al. Low-grade inflammation markers in children and adolescents: Influence of anthropometric characteristics and CRP and IL6 polymorphisms. *Cytokine* 2016; **88**:177–83. <https://doi.org/10.1016/j.cyto.2016.09.007> PMID: 27643980
8. Staiano AE, Katzmarzyk PT. Ethnic and sex differences in body fat and visceral and subcutaneous adiposity in children and adolescents. *International Journal of Obesity* 2012; **36**:1261–9. <https://doi.org/10.1038/ijo.2012.95> PMID: 22710928 PMID: PMC4129655
9. Weber DR, Levitt KLE, Zemel BS, Gallagher PR, Murphy KM, Dumser SM, et al. Anthropometric measures of abdominal adiposity for the identification of cardio-metabolic risk factors in adolescents. *Diabetes Research and Clinical Practice* 2014; **103**:e14–e17. <https://doi.org/10.1016/j.diabres.2013.12.050> PMID: 24552682 PMID: PMC4384445
10. Elkiran O, Yilmaz E, Koc M, Kamanli A, Ustundag B, Ilhan N. The association between intima media thickness, central obesity and diastolic blood pressure in obese and overweight children: A cross-sectional school-based study. *International Journal of Cardiology* 2013; **165**: 528–32. <https://doi.org/10.1016/j.ijcard.2011.09.080> PMID: 22014414

11. Reis JP, Loria CM, Lewis CE, Powell-Wiley TM, Wei GS, Carr JJ, et al. Association between duration of overall and abdominal obesity beginning in young adulthood and coronary artery calcification in middle age. *Journal of the American Medical Association* 2013; **310**:280.  
<https://doi.org/10.1001/jama.2013.7833>  
PMid: 23860986 PMCID: PMC4226407
12. Stabouli S, Kotsis V, Karagianni C, Zakopoulos N, Konstantopoulos A. Blood pressure and carotid artery intima-media thickness in children and adolescents: the role of obesity. *Hellenic Journal of Cardiology* 2012; **53**:41–7.
13. Le J, Zhang D, Menees S, Chen J, Raghuvver G. “Vascular Age” is advanced in children with atherosclerosis-promoting risk factors. *Circulation: Cardiovascular Imaging* 2010; **3**: 8–14.  
<https://doi.org/10.1161/CIRCIMAGING.109.880070>  
PMid: 19920030
14. Kollias A, Psilopatis I, Karagiaouri E, Glaraki M, Grammatikos E, Grammatikos EE, et al. Adiposity, blood pressure, and carotid intima-media thickness in Greek adolescents. *Obesity* 2013; **21**:1013–7.  
<https://doi.org/10.1002/oby.20194>  
PMid: 23784905
15. Osiniri I, Sitjar C, Soriano-Rodríguez P, Prats-Puig A, Casas-Satre C, Mayol L, et al. Carotid intima-media thickness at 7 years of age: Relationship to C-reactive protein rather than adiposity. *The Journal of Pediatrics* 2012; **160**:276-80.e1.  
<https://doi.org/10.1016/j.jpeds.2011.07.020>  
PMid: 21875718
16. Böhm B, Hartmann K, Buck M, Oberhoffer R. Sex differences of carotid intima-media thickness in healthy children and adolescents. *Atherosclerosis* 2009; **206**:458–63.  
<https://doi.org/10.1016/j.atherosclerosis.2009.03.016>  
PMid: 19389672
17. Baroncini LA, Sylvestre L de C, Pecoits Filho R. Assessment of intima-media thickness in healthy children aged 1 to 15 years. *Arquivos Brasileiros de Cardiologia* 2016; **106**(4):327-32.  
<https://doi.org/10.5935/abc.20160030>  
PMid: 26959401 PMCID: PMC4845706
18. Park MH, Skow A, De Matteis S, Kessel AS, Saxena S, Viner RM, et al. Adiposity and carotid-intima media thickness in children and adolescents: a systematic review. *BMC Pediatrics* 2015; **15**:161-5.  
<https://doi.org/10.1186/s12887-015-0478-5>  
PMid: 26475608 PMCID: PMC4609088

# Association between anthropometric

## ORIGINALITY REPORT

16%

SIMILARITY INDEX

13%

INTERNET SOURCES

9%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

1	Impact of Energy Balance on Cancer Disparities, 2014. Publication	1%
2	pdfs.semanticscholar.org Internet Source	1%
3	worldwidescience.org Internet Source	1%
4	www.ijrrjournal.com Internet Source	1%
5	www.cureus.com Internet Source	1%
6	www.mdpi.com Internet Source	1%
7	V. Strugala, P. W. Dettmar, J. P. Pearson. "Thickness and continuity of the adherent colonic mucus barrier in active and quiescent ulcerative colitis and Crohn's disease", International Journal of Clinical Practice, 2008 Publication	1%

8

Georgios Karapostolakis, Marina Vakaki, Achilleas Attilakos, Antonios Marmarinos et al. "The Effect of Long-Term Atorvastatin Therapy on Carotid Intima-Media Thickness of Children With Dyslipidemia", *Angiology*, 2020

Publication

1 %

9

[mdpi-res.com](https://mdpi-res.com)

Internet Source

1 %

10

[www.cambridge.org](https://www.cambridge.org)

Internet Source

1 %

11

[topsecretapiaccess.dovepress.com](https://topsecretapiaccess.dovepress.com)

Internet Source

1 %

12

[www.masimo.com](https://www.masimo.com)

Internet Source

1 %

13

[www.researchgate.net](https://www.researchgate.net)

Internet Source

1 %

14

Suziane Ungari Cayres, Ricardo Ribeiro Agostinete, Barbara de Moura Mello Antunes, Fabio Santos Lira, Rômulo Araújo Fernandes. "Impact of physical exercise/activity on vascular structure and inflammation in pediatric populations: A literature review", *Journal for Specialists in Pediatric Nursing*, 2016

Publication

1 %

15

[documentserver.uhasselt.be](https://documentserver.uhasselt.be)

1 %

16

Gu, W., Y. Huang, Y. Zhang, J. Hong, Y. Liu, W. Zhan, G. Ning, and W. Wang. "Adolescents and young adults with newly diagnosed Type2 diabetes demonstrate greater carotid intima-media thickness than those with Type1 diabetes", *Diabetic Medicine*, 2014.

Publication

1 %

17

Emelia J. Benjamin, Paul Muntner, Alvaro Alonso, Marcio S. Bittencourt et al. "Heart Disease and Stroke Statistics—2019 Update: A Report From the American Heart Association", *Circulation*, 2019

Publication

&lt;1 %

18

Seref Alpsoy, Aydın Akyuz, Dursun Cayan Akkoyun, Burçin Nalbantoglu, Birol Topcu, Mustafa Metin Donma. "Effect of Obesity on Endothelial Function and Subclinical Atherosclerosis in Children", *Electronic Journal of General Medicine*, 2014

Publication

&lt;1 %

Exclude quotes  OnExclude matches  OffExclude bibliography  On

# Association between anthropometric

---

GRADEMARK REPORT

---

FINAL GRADE

GENERAL COMMENTS

**/100**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---