

Cut-off value of Waist

by Nur Aisyah Widjaja

Submission date: 13-Oct-2023 09:15PM (UTC+0800)

Submission ID: 2194578037

File name: Cut-off_value_of_Waist.pdf (92.88K)

Word count: 3254

Character count: 16452

Cut-off value of Waist-to-Hip Ratio as a predictor of metabolic syndrome in adolescents with obesity

Nur Aisiyah Widjaja, Rizka Arifani, Roedi Irawan

Department of Child Health, University of Airlangga, Surabaya, Indonesia

Abstract. *Background and aim:* Metabolic syndrome is present in 7% of adolescents and 19 to 35% in obese adolescents, and the cause of this condition has not been fully understood. Early identification of the risk involved could be an initial step in preventing a metabolic syndrome. Increased waist circumference, which is a parameter of central obesity, is also risk factor for this condition. This study aims to determine the possibility cut-off value of waist-to-hip (WHR) as a predictor of metabolic syndrome. *Methods:* We studied 208 obese adolescents between the ages of 13 to 18 years from junior and senior high schools who living in rural dan urban areas in East Java. These obese adolescents were divided into two groups, namely with and without a metabolic syndrome. Anthropometric measurements, including waist-to-hip ratio (WHR), were carried out to analyze the cut-off values between the two groups. *Results:* 208 obese adolescents (51.4% males and 48.6% females) without the metabolic syndrome and 104 obese adolescents with metabolic syndrome were evaluated. There was a significant relationship between WHR and metabolic syndrome in obese adolescents ($r = 0.203$, $P = 0.003$). Those adolescents with a high WHR value > 0.891 had twice risk to develop metabolic syndrome compared to those with lower (OR 2.033; 95% CI = 1.165-3.545). *Conclusions:* Increased waist-to-hip ratio > 0.89 in adolescents was associated with higher risk to develop metabolic syndrome and can be proposed as a predictor for metabolic syndrome in obese adolescents. (www.actabiomedica.it)

Key words: Waist-to-hip ratio, Metabolic Syndrome, Adolescence, Risk Factors

Introduction

Metabolic syndrome (MetS), also known as “Insulin Resistance Syndrome” or “Syndrome X”, has an increasing prevalence in children and adolescents. The cause of this condition has not been fully understood (1). Therefore, early identification of MetS risk factors and adopting suitable preventive measures is an essential step in preventing its occurrence (2,3). One of the components of MetS is central obesity, which can be measured using various anthropometric parameters, such as waist and hip circumferences, and waist-hip ratio (WHR). Furthermore, the periodic measurement of these parameters can be used for early identification of MetS (4). The cut-off value of WHR can be used as a predicting indicator of

metabolic syndrome in obese children (5). However, this value varies among different countries (6). This study aims to determine the cut-off value of WHR as a predictor of metabolic syndrome in obese adolescents in Indonesia.

Materials and methods

Study population and design

This is an observational case-control study performed between January to May 2020 on obese adolescents with and without the metabolic syndrome. The cut-off value that discriminates those with versus those without the metabolic syndrome was searched.

The correlation of WHR with the metabolic syndrome in all obese adolescents aged 13 to 18 years in Surabaya and Sidoarjo was measured. This study was approved by the Ethical Committee of the Faculty of Medicine, Airlangga University (No. 65/EC/KEPK/FKUA/2020).

Diagnosis of obesity and metabolic syndrome

Obesity was diagnosed based on the CDC 2000 criteria and body mass index (BMI) for age and gender above the 95th percentile. The metabolic syndrome was diagnosed based on the IDF criteria for diagnosing this condition in children and adolescents.

Physical and laboratory examination

Body weight was measured using a digital weight scale (Seca, Germany No ref. 224 1714009) with a precision of 0.1kg. The subjects were standing barefoot and putting on thin clothes during the measurement. Furthermore, the heights were measured using stadiometers (Seca, Germany No ref. 224 1714009), with an accuracy of 0.1cm. The stadiometer was used to measure the height from heel to vertex and the results were presented in 'meter'. Waist circumference was measured midway between the lowest rib and the iliac crest at the end of expiration. The measurements were made on the skin and recorded to the nearest 0.1cm. Furthermore, the waist-hip ratio (WHR) was calculated using the Bjorntorp formula (7). The subjects had fasted for 12 hours before the blood samples were taken. Fasting blood sample was collected for measuring triglyceride, HDL, and blood glucose levels. The blood samples were centrifuged and the serum was stored at -70°C for further analysis.

Data analysis

The correlation between waist-to-hip ratio and metabolic syndrome in obese adolescents was analyzed using the Mann-Whitney comparative test. The cut-off value of WHR was based on the receiver operating characteristic (ROC). The correlation between WHR (WHR>0.891 vs. WHR<0.891) with the component of metabolic syndrome (central obesity, low level HDL-c, hypertriglyceridemia, hyperglycemia, and hypertension) was calculated with chi square test.

Results

An examination was conducted on adolescents with obesity in junior and senior high schools in Surabaya and Sidoarjo. The characteristics of the subjects are listed in the table below (Table 1).

The Mann-Whitney test was used to show the correlation between WHR and metabolic syndrome in obese adolescents with a P-value of 0.003 (95% CI = 0.689-1.071). There was a significant correlation between WHR and metabolic syndrome in obese adolescents as seen in Table 2. The backward multivariate regression analysis showed that the WHR was the most influential parameter on metabolic syndrome compared to other central obesity parameters, such as waist and hip circumference. Based on table 2, waist circumference has stronger correlation with metabolic syndrome compared to WHR ($r = 0.296$ vs. 0.203 respectively, $P < 0.05$).

A diagnostic power test of the WHR for metabolic syndrome in obese adolescents was performed using ROC to determine the cut-off value (Figure 1). The cut-off value of WHR 0.891 on the ROC curve showed a sensitivity value of 63.72% and a specificity of 52.99%. The analysis showed that adolescents with WHR > 0.891 would be x 2.033 times higher risk to develop metabolic syndrome than a low WHR (< 0.891). Furthermore, a chi-square analysis was performed based on the mean waist-to-hip ratio of 0.891 for the components of the metabolic syndrome, including decreased HDL-c, increased triglyceride, and increased fasting blood sugar levels, and increased blood pressure. A high WHR (> 0.891) had a significant correlation with increased blood pressure in obese adolescents, ($P: 0.027$). It was found that in obese adolescents with WHR > 0.891, the WHR was not correlated with the other components of metabolic syndrome (increased triglycerides, decreased in HDL-c, and increased fasting blood sugar levels ($P: >0.05$)).

Discussion

Gender is one of the factors that play a major role in adolescents suffering from metabolic syndrome. The metabolic syndrome was more common in obese male adolescents compared to females. Similar results were also obtained from studies in Malaysia (8)

Table 1. Demographic characteristics, biochemical and diagnostic parameters.

Characteristic	Metabolic syndrome + SD	Non metabolic syndrome	Total (n=208)	P
Sex				
Male, n (%)	64	37	101 (48.5%)	<0.0001 ¹
Female, n (%)	40	67	107 (51.4%)	
Age, month	181.13 + 18.96	179.23 + 18.98	180.18 + 18.95	0.470 ²
≤ 16 years	80	80	160 (76.9%)	0.565 ¹
> 16 years	24	24	48 (23.1%)	
Triglyceride level	150.70 + 68.27	92.21 + 47.87	121.46 + 65.72	<0.0001 ²
≥ 150 mg/dl	70	20	90 (43.3%)	<0.0001 ¹
< 150 mg/dl	34	84	118 (56.7%)	
HDL level	38.72 + 6.79	45.74 + 7.29	42.23 + 7.86	<0.0001 ²
< 40 mg/dl	98	60	158 (75.9%)	<0.0001 ¹
≥ 40 mg/dl	6	44	50 (24.1%)	
Fasting glucose level	86.74 + 9.42	87.31 + 19.46	87.02 + 15.26	0.789 ²
≥ 100 mg/dl	18	4	22 (10.5%)	<0.0001 ¹
< 100 mg/dl	86	100	186 (89.5%)	
Systolic blood pressure	130.87 + 11.44	119.52 + 12.50	125.19 + 13.24	<0.0001 ²
Diastolic blood pressure	87.11 + 9.65	78.65 + 8.47	82.88 + 9.998	<0.0001 ²
≥ 130/85 mmHg	72	21	93 (44.7%)	<0.0001 ¹
< 130/85 mmHg	32	83	115 (55.3%)	

¹Fischer's exact test; ²Independent sample T-test; Significant of P: <0.05

Table 2. Regression analysis of central obesity parameters in correlation to the metabolic syndrome in adolescents with obesity.

Variable	r	P	OR
Waist circumference	0.296	0.001	0.825 ¹
Hip circumference	0.181	0.014	1.238 ¹
WHR	0.203	0.003	1.248 ¹
Hip circumference	0.181	0.014	1.045 ¹
WHR	0.203	0.003	762.115 ²

¹bivariate regression analysis; ²multivariate regression analysis. Significant of P: <0.05

and Norway (9) which showed that male adolescents had a greater prevalence of metabolic syndrome. This relationship was not significant for ages above 18 years (10).

The different components of the metabolic syndrome had variable prevalence in our obese adolescents. A decrease in HDL levels was detected in (75.9%), while an increase in blood pressure occurred

in (44.7%). Similar results were reported from a study in Virginia on 78 children aged 3 to 18 years with obesity. In their study, high blood pressure was the most prevalent component, followed by a decrease in HDL levels (11).

This study documented a significant correlation between WHR and metabolic syndrome in obese adolescents. Similarly, a cross-sectional studies conducted




Figure 1. ROC curve for the value of WHR as a predictor of metabolic syndrome.

in Brazil (109 children), Turkey (1194 children, mean age: 9 years), and USA (764 children, aged 5 to 12 years) found significant relationship between the waist-hip ratio and the metabolic syndrome in obese children (2). Another study showed that WHR had a significant relationship with components of the metabolic syndrome in obese children (2,12,13). Furthermore, Moore et al. (13), retrospectively showed that the waist-to-hip ratio was associated with the occurrence of metabolic syndrome.

Central obesity has also been associated with the risk of metabolic syndrome in children along with other variables such as waist (WC) and hip circumference (HC), and waist-to-hip ratio (WHR). Furthermore, according to the multivariate regression analysis, the most influential variable in obese adolescents is the waist-to-hip ratio. A study in Spain involving 1001 adolescents with an average age of 13.1 years also showed that WHR was the best predictor of this condition compared to WC and BMI. Furthermore, a cross-sectional study that was conducted in 18 secondary schools showed that WHR had the highest predictive value compared to WC and BMI (14). The results of a retrospective study in Iran which was conducted

on 2064 children and adolescents aged 6 to 18 years showed that WHR has the weakest association with metabolic syndrome when compared to other parameters such as WC, and BMI (15).

Currently, there is no consensus of the best cut-off value for WHR to the risk of metabolic syndrome in obese adolescents. The cut-off value of WHR 0.891 on the ROC curve showed a sensitivity value of 47.1% and a specificity of 35.6% in relation to metabolic syndrome, which was in contrast with this result.

A study involving 958 adolescents in Brazil showed a limiting WHR value of 0.87 on the ROC curve with an AUC of 0.43 and suggested that a WHR value of 0.87 can be used as a predictor of metabolic syndrome in adolescents (sensitivity value of 46.1% and a specificity of 51.2%) (2). A study from Spain on 1001 adolescents (median age of 13 years) suggested a cut-off value of WHR - 0.89 to be a good predictor of metabolic syndrome (14). Other studies in Asia propose a mean value of 0.8 to indicate higher risk of metabolic disorders.

There was a decrease on WHR value at the age of 9 to 15 years, and then stagnant (flat) at the age of 16, which was occurred on males and females in this study. A large cohort cross-sectional study involving 21,111 children aged 8 to 15 years, it was found that the WHR curve declined and then came to a plateau at the age of 15 years (16). In addition, the cut-off value associated with the metabolic syndrome also decreased above 15 years of age with a value of 0.84 in females and 0.91 in males representing the 95th percentile (17).

To our knowledge, no previous research in Indonesia has searched the WHR cut-off value associated with the metabolic syndrome in adolescents. However, Istiqomah et al. (18) analyzed the WHR cut-off value associated with hypertension, a component of the metabolic syndrome, in a large cohort ($n = 325$) and suggested that $WHR > 0.94$ could be used for screening hypertension in children, (AUC value of 0.724 (95% CI 0.672-0.72; $P < 0.001$)). In their study, the WHR value > 0.94 had a sensitivity of 59.6% and a specificity of 77.0% (18). In our study whereby the high WHR (> 0.891) had a significant correlation to the occurrence of increased blood pressure (a component of the metabolic syndrome), in obese adolescents

with a $P < 0.05$. In support to our findings, Widjaja et al. (19) studied 59 Indonesian obese adolescents and reported a significant correlation between WHR and insulin resistance, another component of the metabolic syndrome (p value 0.008 and r value 0.343).

Conclusion

The waist-to-hip ratio significantly correlated with the occurrence of metabolic syndrome in obese adolescents compared to the other central obesity parameters. Therefore, the WHR values of more than 0.891 can be used as a predictor of metabolic syndrome in obese adolescents.

Acknowledgements: The authors are grateful to the 12 high schools in Surabaya and Sidoarjo that volunteered to become respondents. We also express our gratitude to all headmasters and counselling teachers for bringing together the respondents along with their parents before and during the data collection. The authors disclosed receipt of the following financial support for the research, and/or publication of this article.

Conflict of Interest: The authors declare that he or she has no commercial associations that might pose a conflict of interest in connection with the submitted article

Author Contributions: Conceptualization: Nur Aisiyah Widjaja; methodology: Roedi Irawan; data curation: Rizka Arifani; writing—original draft preparation: Nur Aisiyah Widjaja; writing—review and editing: Roedi Irawan; All authors have read and agreed to the published version of the manuscript.

References

- Singh N, Scmhrd. Metabolic Syndrome: Practice Essentials, Background, Pathophysiology. *J Hear Stroke*. Jan 2018;3(1):1044
- Ferreira AP, Ferreira CB, Brito CJ, et al. Prediction of Metabolic Syndrome in Children through Anthropometric Indicators. *Arq Bras Cardiol*. 2011 Feb;96(2):121–5. doi: 10.1590/s0066-782x2011005000005.
- Seo JY, Kim JH. Validation of surrogate markers for metabolic syndrome and cardiometabolic risk factor clustering in children and adolescents: A nationwide population-based study. *PLoS One*. 2017 Oct; 12(10):e0186050. doi: 10.1371/journal.pone.0186050.
- Vincent J, Mohn A. Early diagnosis of metabolic syndrome in children. *Lancet Child Adolesc Heal*. 2017 Oct;1(2):86–8. doi: 10.1016/S2352-4642(17)30043-3.
- Mushtaq MU, Gull S, Abdullah HM, Shahid U, Shad MA, Akram J. Waist circumference, waist-hip ratio and waist-height ratio percentiles and central obesity among Pakistani children aged five to twelve years. *BMC Pediatr*. 2011 Nov 21;11:105. doi: 10.1186/1471-2431-11-105.
- Pacheco LS, Blanco E, Burrows R, Reyes M, Lozoff B, Gahagan S. Early onset obesity and risk of metabolic syndrome among Chilean adolescents. *Prev Chronic Dis*. 2017 Oct 12;14:E93. doi: 10.5888/pcd14.170132.
- Skruze G, Kažoka D. Use of waist to hip ratio in the determination of the body composition in preschool children in Latvian population. *Pap Anthropol*. 2017 Sept;26(2):126. doi:10.12697/poa.2017.26.2.13.
- Fadzlina AA, Harun F, Nurul Haniza MY, et al. Metabolic syndrome among 13 year old adolescents: Prevalence and risk factors. *BMC Public Health*. 2014;14(Suppl 3):S7. doi: 10.1186/1471-2458-14-S3-S7.
- Barstad LH, Júlíusson PB, Johnson LK, Hertel JK, Lekhal S, Hjelmesth J. Gender-related differences in cardio-metabolic risk factors and lifestyle behaviors in treatment-seeking adolescents with severe obesity. *BMC Pediatr*. 2018 Feb;18(1):61. doi: 10.1186/s12887-018-1057-3
- Wittcopp C, Conroy R. Metabolic syndrome in children and adolescents. *Pediatr*. 2016 May;37(5):193–202. doi: 10.1542/pir.2014-0095.
- Co J, Jeffrey J, Emmett M, Modak A, Sondike SB. Obesity, Hypertension and Metabolic Syndrome in Children in West Virginia. *W V Med J*. 2015 Jul-Aug;111(4):20–2, 24. PMID: 26242028.
- Agirbasli M, Agaoglu NB, Ergonul O, et al. Comparison of anthropometric indices in predicting metabolic syndrome components in children. *Metab Syndr Relat Disord*. 2011 Dec;9(6):453–9. doi: 10.1089/met.2011.0018.
- Moore LM, Fals AM, Jennelle PJ, Green JF, Pepe J, Richard T. Analysis of Pediatric Waist to Hip Ratio Relationship to Metabolic Syndrome Markers. *J Pediatr Health Care*. 2015 Jul-Aug;29(4):319–24. doi: 10.1016/j.pedhc.2014.12.003.
- Perona JS, Schmidt-RioValle J, Rueda-Medina B, Correa-Rodríguez M, González-Jiménez E. Waist circumference shows the highest predictive value for metabolic syndrome, and waist-to-hip ratio for its components, in Spanish adolescents. *Nutr Res*. 2017 Sep;45:38–45. doi: 10.1016/j.nutres.2017.06.007.
- Hashemipour M, Soghrati M, Malek Ahmadi M, Soghrati M. Anthropometric indices associated with dyslipidemia in obese children and adolescents: a retrospective study in isfahan. *ARYA Atheroscler*. 2011 Spring;7(1):31–9. PMID: 22577442; PMCID: PMC3347839
- Kelishadi R, Gouya MM, Ardalan G, et al. First reference curves of waist and hip circumferences in an Asian population of youths: CASPIAN study. *J Trop Pediatr*. 2007 Jun;53(3):158–64. doi: 10.1093/tropej/fml090.
- Mederico M, Paoli M, Zerpa Y, et al. Valores de referencia de la circunferencia de la cintura e índice de la cintura/cadera en

- escolares y adolescentes de Mérida, Venezuela: Comparación con referencias internacionales. *Endocrinol y Nutr.* 2013 May;60(5):235–242. doi: 10.1016/j.endonu.2012.12.003.
18. Istiqomah E, Gurnida DA, Hilmanto D, Djalil D, Hakim L, Fauziah PN. Waist circumference and waist-hip ratio as screening tools for hypertension in children aged 6–11 years. *Paediatr Indones.* 2019 Sep;59(5):265–270. doi: 10.14238/pi59.5.2019.265-70.
19. Widjaja NA, Prihaningtyas RA, Hanindita MH, Irawan R. Adiponectin, anthropometric measurements and insulin resistance in adolescence with obesity. *Malays J Nutr.* 2020 Jun;26(2):165–72. doi: 10.31246/mjn-2019-0100.

Correspondence:

Received: 30 September 2022

Accepted: 30 November 2022

Nur Aisiyah Widjaja, MD

Nutrition and Metabolic Disease Unit,

Child Health Departemen Airlangga University, Surabaya

Dharmahasada Mas XIII Blok AK no. 1-2

Surabaya, 60285 Indonesia

Phone: +62 8123073379

E-mail: nurilwidjaja69@gmail.com

Orcid ID: 0000-0002-4253-8760

Cut-off value of Waist

ORIGINALITY REPORT

17%

SIMILARITY INDEX

13%

INTERNET SOURCES

12%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | repository.unair.ac.id
Internet Source | 1% |
| 2 | repo.stikesicme-jbg.ac.id
Internet Source | 1% |
| 3 | boris.unibe.ch
Internet Source | 1% |
| 4 | Anuj Jaiswal, Robin E. Gilbert, Karadi H. Sunil Kumar, Andrew D. Carrothers, Jan-Herman Kuiper, James B. Richardson. "Function and Survival after Revision of Hip Resurfacing", HIP International, 2011
Publication | 1% |
| 5 | mattioli1885journals.com
Internet Source | 1% |
| 6 | Anne S. Biong, Hege M. Rebnord, Ragnhild L. Fimreite, Kerstin U. Trygg, Jetmund Ringstad, Dag S. Thelle, Jan I. Pedersen. "Intake of dairy fat and dairy products, and risk of myocardial infarction: A case-control study", | 1% |

International Journal of Food Sciences and Nutrition, 2009

Publication

7	Budi Santoso, Nanda Yuli Rahmawati, Ashon Sa'adi, Sri Ratna Dwiningsih et al. "Elevated peritoneal soluble endoglin and GDF-15 in infertile women with severe endometriosis and pelvic adhesion", Journal of Reproductive Immunology, 2021 Publication	1 %
8	academic.oup.com Internet Source	1 %
9	brenda-enzymes.org Internet Source	1 %
10	www.jpmmh.org Internet Source	1 %
11	www.mdpi.com Internet Source	1 %
12	dspace.knust.edu.gh Internet Source	1 %
13	Alynne Christian Ribeiro Andaki, Adelson Luiz Araújo Tinôco, Edmar Lacerda Mendes, Roberto Andaki Júnior et al. "Anthropometry and physical activity level in the prediction of metabolic syndrome in children", Public Health Nutrition, 2013 Publication	<1 %

14 Qianyuan Yang, Yalan Liu, Zhaofeng Jin, Leilei Liu, Zhiping Yuan, Degan Xu, Feng Hong. "Evaluation of anthropometric indices as a predictor of diabetes in Dong and Miao ethnicities in China: A cross-sectional analysis of China Multi-Ethnic Cohort Study", PLOS ONE, 2022
Publication

15 bmcpediatr.biomedcentral.com
Internet Source

16 discovery.ucl.ac.uk
Internet Source

17 Sajithya Perera, Vitool Lohsoonthorn, Wiroj Jiamjarasrangi, Somrat Lertmaharit, Michelle A. Williams. "Association between elevated liver enzymes and metabolic syndrome among Thai adults", Diabetes & Metabolic Syndrome: Clinical Research & Reviews, 2008
Publication

18 jped.elsevier.es
Internet Source

19 www.longdom.org
Internet Source

20 Kuba, Valesca, Claudio Leone, and Durval Damiani. "Is waist-to-height ratio a useful indicator of cardio-metabolic risk in 6-10-year-old children?", BMC Pediatrics, 2013.

21

bsdwebstorage.blob.core.windows.net

Internet Source

<1 %

22

Esfarjani, Fatemeh, Marjan Khalafi, Fatemeh Mohammadi, Negar Zamani-Nour, and Roya Kelishad. "Metabolic Syndrome and its determinants in a sample of young Iranian children with obesity", *Pakistan Journal of Medical Sciences*, 2013.

Publication

<1 %

23

Fatemeh Abbaszadeh, Nahid Sarafranz, Mahboobeh Kafeei Atrian, Zohreh Sadat, Azam Bagheri, Alireza Moravveji. "Anthropometric Indices in the Prediction of Hypertension in Female Adolescents", *Iranian Red Crescent Medical Journal*, 2017

Publication

<1 %

24

Javier S. Perona, Jacqueline Schmidt-RioValle, Blanca Rueda-Medina, María Correa-Rodríguez, Emilio González-Jiménez. "Waist circumference shows the highest predictive value for metabolic syndrome, and waist-to-hip ratio for its components, in Spanish adolescents", *Nutrition Research*, 2017

Publication

<1 %

25

Yao Duan, Jiahong Sun, Mingming Wang, Min Zhao, Costan G. Magnussen, Bo Xi. "Association between short sleep duration

<1 %

and metabolic syndrome in Chinese children and adolescents", Sleep Medicine, 2020

Publication

26

ikee.lib.auth.gr

Internet Source

<1 %

27

onlinelibrary.wiley.com

Internet Source

<1 %

28

www.pjms.org.pk

Internet Source

<1 %

29

www.whri.qmul.ac.uk

Internet Source

<1 %

30

Anna Springwald, Patrycja Różana-Kowalska, Piotr Gibała, Olimpia Zajdel-Cwynar, Ewa Małecka-Tendera, Paweł Matusik. "Usefulness of the metabolic syndrome diagnosis in obese children in clinical practice", Pediatric Endocrinology Diabetes and Metabolism, 2019

Publication

<1 %

31

B. Isomaa, M. Henricsson, P. Almgren, T. Tuomi, M-R. Taskinen, L. Groop. "The metabolic syndrome influences the risk of chronic complications in patients with Type II diabetes", Diabetologia, 2001

Publication

<1 %

32

Miharu Tamaoki, Ikumi Honda, Keisuke Nakanishi, Sophathya Cheam, Manabu

<1 %

Okawada, Hisataka Sakakibara. "Prevalence of Metabolic Syndrome and Its Components in Urban Cambodia: A Cross-Sectional Study", *Journal of Epidemiology and Global Health*, 2022

Publication

33

www.pubmedcentral.nih.gov

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Cut-off value of Waist

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
