

Fruit development of groundcherry (*Physalis* *angulata* L.) in dryland

by Sucipto Hariyanto

Submission date: 13-Dec-2022 05:31PM (UTC+0800)

Submission ID: 1980074835

File name: b5-AJCS.pdf (469.23K)

Word count: 4488

Character count: 23324

Fruit development of groundcherry (*Physalis angulata* L.) in dryland

Diana Nurus Sholehah^{1,3}, Sucipto Hariyanto², Hery Purnobasuki*²

¹Program of Mathematics and Natural Science, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia

²Department of Biology, Faculty of Sciences and Technology, Airlangga University, Surabaya, Indonesia

³Program of Agroecotechnology, Faculty of Agriculture, University of Trunojoyo Madura, Bangkalan, Indonesia

*Correspondence: hery-p@fst.unair.ac.id² dianans@trunojoyo.ac.id^{1,3}

Abstract

Physalis angulata are widespread in the tropics with high bioactive compounds and pharmacological effects. This paper aims to identify the stages of fruit development and changes in physicochemical properties to provide comprehensive data on optimal fruit utilization. This study used CM 1 accession found in Madura Island with purplish stems and fruit peels. After 30 days of cultivation, the plant was transplanted into a polybag containing red Mediterranean soil and manure (5:1). It is grown in a greenhouse for 90 days to observe. The experimental design was completely randomized. The stages were selected as treatments: 1, 3, 6, 9, 12, 15, 18, and 21 Day After Anthesis. The study used 15 replications with a total of 15 plants used, except for chemical analysis which used five replication with three plants for each replication. Observations were carried out during the reproductive phase and the parameters included fruit phenology, physical character, and chemical characteristics. The result showed that the growth stage of *P. angulata* consists of fully opened flower, fruit development (fruit set and fruit expansion), and ripening fruit (starting ripe, advanced ripe, and eating ripe). The optimal fruit stage for consumption is 21 days after anthesis, which has a diameter of 13.58 mm, a weight of 1.69 g and firmness of 3.40 N, total soluble solids of 14.96 Brix, total titratable acid of 0.0011%, and vitamin C 20.91 mg/100 g.

Keywords: BBCH; calyx; chemical characteristics; physical characteristics; phenology

Abbreviations: BBCH_Biologische Bundesanstalt, Bundesortenamt und Chemische Industrie; DAA_day after anthesis; TTA_total titratable acidity; TSS_total soluble solid.

Introduction

Physalis is a genus of the Solanaceae Family with the highest number of phytochemical-rich species out of 2300 species belonging to this family (Medina-Medrano et al., 2015). This species easily adapts to different environmental conditions, such as different types of climate. It can grow in a wide range of environments, evidenced by the anatomical plasticity of leaves favouring more excellent radiation resistance (Rodrigues et al., 2014). It causes the species to spread from the subtropics such as Asia, Columbia, Brazil, Peru, and Africa to the tropics (Muniz et al., 2014). Ozaslan et al. (2017) reported on an invasive plant in Turkey.

This species is a monoecious plant with pale yellow single flowers on the axillary leaves (Figueiredo et al., 2020). It bears flowers and fruits throughout the year (Muniz et al., 2014). *Physalis* fruit has interesting physical properties with a distinctive taste, aroma, juicy and crunchy texture and attractive colour (Oliveira et al., 2016).

P. angulata is also known as groundcherry. It has become the centre of attention because of the nutritional and bioactive compounds that provide important body health benefits (Kusumaningtyas et al., 2015; Saavedra et al., 2019). The primary components are physalin and withanolide groups. The three groups of physalin are physalin B, F, and G. A form of withanolide is withanolatin. Anti-inflammatory, anti-

tumour, cytotoxic, anti-carcinoma and trypanocidal properties are all present in both (Sun et al., 2017; Meng et al., 2019). Another important content is antioxidant compounds from phenolic groups such as flavonoids, phenolic acids, and coumarin (Lima et al., 2014). These compounds have also been shown to function as immunomodulators (Kusumaningtyas et al., 2015). According to the study conducted by Medina-Medrano et al. (2015), antioxidant compounds in this species have the highest DPPH scavenging and total antioxidant capacity compared to 5 other *Physalis* species. The fruit showed the highest antioxidant activity (Ferreira et al., 2019) and cytotoxic effect (Iwansyah et al., 2019). Demand for this fruit has recently risen, followed by the rise in production and commercialization of *Physalis* to other regions of the world (Muniz et al., 2015; Oliveira et al., 2016).

Despite numerous research, it is essential to comprehend fruit growth and development and their impact on fruit quality, from seed to harvest. This data will be used to determine maturity index, which is currently unavailable. Determining ripeness is critical for deciding the best harvest time for fresh markets and processing, such as nutritional foods and traditional medicine (Neves et al., 2017).

Growing environments also significantly influence the rate

and period of plant growth and development, which ultimately affects the physicochemical characteristics of fruits, such as their weight, size, and soluble solid (Barroso et al., 2018). The studies on the impact of the growing environment on fruit characteristics lead to understand the possibility of growth of the species in various areas, such as drylands. There is a critical need to consider drylands adapt to climate change, as they will change the phenology of plant growth and senescence. A core aspect of global change studies is the study of biological life cycles, such as phenology and fruit production (Walker et al., 2015). Understanding plant phenological behaviour is critical for improving production and determining proper cultivation management (Ghrab et al., 2016).

This study aims to identify the stages of fruit development and changes in physicochemical properties to provide comprehensive data on optimal fruit utilization in dryland. In this study, there was a comprehensive presentation of fruit development analysis from anthesis until ripe. The aspects of fruit development consist of the phenology and physicochemical change based on BBCH scale

Result and Discussion

Fruiting phenology

The growth stage of groundcherry fruit based on the BBCH scale consists of stages 65, 70, and 80 (Table 1). Fully opened flowers mark stage 65, abscised petals indicate expansion fruit at stage 70, while the ripening stage occurs at stage 80. Observations were carried until eating ripe at 21 DAA because the fruit fell off over that age.

The plant in this study begins to produce flowers after the branches formed, and the formation of other flowers follows this. The transformation from the vegetative to the generative stage is represented by this occurrence (Ramirez et al., 2013). Groundcherry flowered during the reproductive phase until senescence (Figueiredo et al., 2020). The first flower emerged time was 12 days after transplanting. That time was earlier than the result of Sadiyah et al. (2020).

Stage 65 is characterized by a fully opened flower with perpendicular petals to the pedicel. The open flowers had an average diameter of 12,67 mm. The flowers begin to open randomly from 06:00 to 12:00 and start to close at 15:00 and close entirely at 17:00 daily, in line with the petal movement time of Figueiredo et al. (2020). In this study, the flowers opened for 8-9 hours. Flower that open less than that will return to open the next day until they meet the opening time. Anther opened and released pollen for the fertilization process as soon as the flower opened. Flower opening rhythmicities are generally related to circadian rhythms and pollination activity with pollinators (Fenske et al., 2018). Environmental signals such as light, temperature, and nutritional status may influence the circadian clock of flowers (Horibe, 2018, Inoue et al., 2018).

The anthesis flowers are closed and dried together with the anthers the next day (Figueiredo et al., 2020). The dried petals fall or remain between the tips of the calyx, which turn into calyx that cover the fruit. The process occurred for three days and indicated the beginning of stage 70. The early part of fruit development is the growth of ovaries known as fruit sets, followed by cell division and fruit expansion (Handa et al., 2011; McAtee et al., 2013). The success of the fruit set determines the fruit yield. Temperature and humidity were the predominant environmental factors in tomato fruit sets

(Shamshiri et al., 2018). The size and weight of the fruit increase rapidly, and the stage lasted for 12 days.

Stage 81 represents the beginning of the ripening process that occurs at 12 DAA. Harvest was performed at 21 DAA (stage 89). A previous study stated that the fruit ripe based on optimal seed maturity occurred 28 days after anthesis at an average daily temperature of 24-27°C (Santiago et al., 2019). According to Hatfield and Prueger (2015), temperature improves phenological growth by increasing senescence. The reproductive phase is more sensitive to temperature increases than the vegetative phase. The higher daily average temperature of this study (40.27°C) is believed to accelerate ripening. Physical and chemical changes characterize the maturation process.

Physical characteristics

Physical changes include size, weight, firmness, and colour change. The size and weight increase toward the end of the measurement period, but the rate decreased at 12 - 21 DAA (fig 2). The weight and diameter of the fruit at 21 DAA are similar to the values obtained by Santiago et al. at the ideal harvest fruit stage (2019). The weight and size increase shows the accumulation of dry matter in the pulp and seeds (Liguori et al., 2014). The variations may be attributed to the ripening state, in which the fruit, climate, and soil properties were planted along with the genotype (Oliveira et al., 2016, Purquerio et al., 2019).

Texture modifying occurs during the ripening process. The activated ethylene-induced significantly changed the shape and composition of the cell wall and leads to the softening of pericarp tissue (Tucker et al., 2017). Firmness decrease began to occur 12 DAA, and the lowest firmness was 3.39 N in 21 DAA. This value was higher than the ripe *P. peruviana* fruit firmness in Oliveira et al. 2016 with 2.94 N. According to Sadiyah et al. (2020), calyx accession from Madura has a higher firmness than other regions in East Java.

The chromatic coordinate values are shown in Fig 3. The value of *a** was negative at the beginning of the fruit growth and showed a green colour on calyx and fruit peeled up to 12 DAA and 6 DAA, respectively. Yellow discolouration of the calyx and peel occurred during maturation. This is supported by the increasing positive *b** value at 15-18 DAA ranging from 30.06 - 31.07. The intensity was below the *P. peruviana* with *b** value of 57.94 (Oliveira et al., 2016). The value of *b* decreases at 18 DAA, while the *L** value increases up to 21 DAA. A positive *L** value represents increased redness and the reddish-purple colour on the calyx and peel. It is different from the latest study in Madura by Sadiyah et al. (2020), which reported that ripe fruits have green skin. The appearance of purple pigments in many fruits is associated with the presence of anthocyanins. In certain fruits, the purple colour was associated with fruit maturity, and the levels reach maximum in the final stages of fruit development (Khoo et al., 2017).

Chemical characteristic

According to Souza et al. (2017), *Physalis* has a high carbohydrate level at the fruit growth phase. This level will decrease when the carbohydrates are broken down into sugar at the fruit ripening phase, increasing the soluble solid levels. The content of organic acids, sugars, and other dissolved solids is a soluble solid component that affects fruit taste. The soluble solid levels increased rapidly after entering the 12 DAA, but the increase begins to slow down at the end of fruit life (Fig 3). The highest soluble solid levels were 21

Table. 1 Growth stage of *Physalis angulata* fruit.

Stage	BBCH Description	BBCH <i>Physalis angulata</i>	DAA
60	Flowering		
65	Fully open flower	Fully open flower	0
70	Fruit Development		
71	Fruit set	Fruit set; Petal dries or abscised; calyx closed; fruit about to grow	1
72	20 % fruit growth	Fruit reached 20% final weight	3
75	50 % fruit growth	Fruit reached 50% final weight	6
77	70 % fruit growth	Fruit reached 50% final weight	9
80	Fruit Maturity		
81	Starting of ripening	Start of rapid increase soluble sugar and softening	12
85	Advanced ripening	Calyx and fruit peel turned to yellow, enhanced softening	15
88		Slow softening, calyx turned to slightly brownish, and fruit peels started to have purple tinge and seed maturation	18
89	Eating ripe	Completely mature: fruits have typical fully ripe, yellow with an intensive purple tinge, starting abscission of the fruit	21




Fig 1. Monthly rainfall during the development of groundcherry in the years 2019 and 2020.




Fig 2. Fruit Development of *Physalis angulata* Base on BBCH Scale from Open Flower to Eating Ripe. Bar = 2 mm.
Averages followed by the same letter in the same bar do not differ by DMRT test ($P < 0.05$).




Fig 3. Colour of *P. angulata* fruit and calyx indicated by L*¹², a* and b* value. (a) L* (lightness) (b) a* (red / green value) (c) b* (Yellow / blue value). Averages followed by the same letter in the same bar do not differ by DMRT test ($P < 0.05$).




Fig 4. Some important chemical characteristics of *P. angulata* fruit aligned to BBCH scale . Averages followed by the same letter in the same bar do not differ by DMRT test ($P < 0.05$).

DAA with 14.9°Brix, higher than the study by Curi et al. (2018) and Souza et al. (2017) with 12 and 11.3°Brix, respectively. In contrast, the total titratable acid decreased to a minimum level at 21 DAA with 0.0011%. The acid decrease affects consumer preferences for fruit selection. Sweet fruit and low acidity are preferred by consumers and suitable for fresh consumption (Souza et al., 2012). The vitamin C level was 20.91 mg / 100 g, below the study results conducted by Curi et al. (2018), reaching 75.44%.

Material and Methods

Plant material and location of the experiment

The experiment was carried out from October 2019 to February 2020. The plants are cultivated in Bangkalan Madura (112°E 44'9,176"E and -72°47'652"S) which was a dryland

under a wet climate (Dwiratna et al., 2018) with low-moderate monthly rainfall (Fig 1) and annual rainfall of 1662 mm, an average daily maximum temperature of 40.27°C at an altitude of 5 m above sea level. This study used CM 1 accession found in Madura Island. It was characterized by a purplish stem and fruit peel, according to Sholehah and Setiawan (2019). The plant growth medium was a red mediterranean soil:manure (5:1) in a polybag. The plants were transplanted 30 days after sowing and cultivated for 90 days in the greenhouse for observation. The experimental design was completely randomized with development stages of fruit as treatments. The stages were selected as treatments: 1 DAA, 3 DAA, 6 DAA, 9 DAA, 12 DAA, 15 DAA, 18 DAA and 21 DAA (Fig 1). The study used 15 replications with a total of 15 plants, except for chemical analysis which used five replication with three plants for each replication. Five fruits

of each plant were collected for each stage and taken to the laboratory.

The data obtained were analyzed with analysis of variance (ANOVA). Then The means were compared with Duncan Multiple Range Test (DMRT) at 5 % significance.

Phenological monitoring

Phenological observations were carried out on flowers and fruits. Information about flower initiation, longevity, and degree of opening of the petals was systematically collected. At the beginning of the study, the flowers were checked every four hours. The flowers were only observed from 0500 hours to 2100 hours if it was not open overnight. The fruiting phenology is presented based on physicochemical changes from anthesis flowers through ripe fruit according to the BBCH scale (Hack et al., 2013, Ramirez et al., 2013). These data were used to characterize the fruit development study.

Physical characteristics

The physical characteristics include the diameter, weight, colour, and firmness of the fruit. Digital callipers measured the diameter with a sensitivity of 0.01 mm. The weight was measured using a digital balance with a sensitivity of 0.01 mg. The fruit colour was measured using a colour reader (Konica Minolta - CR -10) by measuring L*, a*, and b*. L* represents the value of dark or light, a* represents the variation value from red to green (a positive value = red, a negative value = green), while b* represents the variation value from blue to yellow (b positive value = yellow, b negative value = blue) (Yildiz et al., 2015). Its colour parameter is widely used in agroindustry and effectively determines slight colour variation (Odjo et al., 2012). The fruit firmness was measured using the TA.XT texture analyzer tool. The tool's setting includes a 2 mm probe, a test speed of 1.00 mm/s, and a distance of 6 mm from the observation. The firmness value is obtained in newtons (N).

Chemical characteristics

some critical chemical characteristics of the fruit, including TSS, TTA and vitamin C. The TTA was determined using titration according to the Association of Official Analytical Chemists (AOAC, 2000). Two grams of fruit juice were dissolved in 50 ml of purified water. Then 25 ml filtrate of the mixture was added to three drops of phenolphthalein 1 %. The mixture was titrated with 0.1 N sodium hydroxide (NaOH) until it turned to a stable pink colour. The total titratable acidity was expressed in terms of TTA (%) as the amount of 0.1 N NaOH per 100 mg fruit, according to the following equation:

$$TTA (\%) = \frac{(V \times 0.1 \times 100)}{10 \text{ m}}$$

Where: 0.1 is the normality of NaOH (N), V is the volume of NaOH required (ml), and m is the mass of the fruit juice sample used (mg).

Vitamin C was determined using a modified method of redox titration by Ciancaglini et al. (2001). Two grams of fruit juice were dissolved in 50 ml of purified water. The mixture of 25 ml filtrate was then added to 1 ml of starch indicator drops and then titrated with 0.01 N iodine until it turned to a stable blue colour. The vitamin C content can be calculated according to the following equation:

$$\text{Vitamin C (mg / 100 g)} = \frac{V \times 0.88 \times P \times 100}{W_s}$$

6
V is the volume of iod 0.01 N (ml), P is the dilution factor of the fruit sample, 0.88 is a converted factor of 1 ml for 1 ml iod 0.01 N ascorbic acid milligram, and Ws is the sample weight (g).

Conclusion

In conclusion, the growth stage of *P. angulata* includes fully opened flower, fruit development (fruit set and fruit expansion), and ripening fruit (starting ripe, advanced ripe, and eating ripe). Based on growth conditions, the ripe fruit has a diameter of 13.58 mm, weight 1.69 g, firmness 3.40 N, soluble solid 14.96, total titratable acid 0.0011%, and vitamin C content of 20.91 mg / 100 g. Eating ripe fruit is the optimal condition for consumption at 21 DAA with maximum fruit size, yellow with purple tinge fruit, soft juicy texture, and sweet taste. The sugar level and the level of titratable acid influenced the unique taste of the fruit.

Acknowledgements

We would like to thank Dr. Gita Pawana, M.Si, head of Agriculture Science and Technology Laboratory, to support the laboratory facilities during the research.

References

- Association of Official Analytical Chemistry (1995) Official Methods of Analysis 932.12 Solids (Soluble) in Fruits and Fruit Products. Virginia.
- Association of Official Analytical Chemists (2000). Official method 942.15 Acidity (Titrable) of fruit products read with AOAC official method 920.149. Preparation of test sample. 17th edition. USA : AOAC International.
- Barroso NS, Fonseca JST, Nascimento MN, Pelacani CR (2018). Physalis development according to the growing season in the semi-arid region of the Bahia state, Brazil. Pesqui Agropecu Trop. 48(4):429-435.
- Ciancaglini P, Santos HL, Daghastanli KRP, Thedei Jr G (2001) Using a classical method of vitamin C quantification as a tool for discussion of its role in the body. Biochem Mol Biol Edu. 29:110-114.
- Curi PN, Carvalho CS, Salgado DL, Pio R, Silva DFD, Pinheiro ACM, Souza PR (2018) Characterization of different native american *Physalis* species and evaluation of their processing potential as jelly in combination with brie-type cheese. Food Sci Technol (Campinas). 38(1):112-119.
- Dwiratna S, Bafdal N, Asdak C, Carsono, N (2018) Study of runoff farming system to improve dryland cropping index in Indonesia. Int J Adv Sci Eng Inf Technol. 8(2):390-396.
- Hack H, Klose FR, Stauss R, Buhr FL, Lancashire PD, Boom T, Wicke MHH, Feller C, Weber HBFE (2001) The extended BBCH-scale. In : Meier U (Ed) Growth stages of mono-and dicotyledonous plants BBCH monograph, 2nd edn. Berlin: Federal Biological Research Centre for Agriculture and Forestry.
- Fenske MP, Nguyen LP, Horn EK, Riffell JA, Imaizumi T (2018) Circadian clocks of both plants and pollinators influence flower seeking behavior of the pollinator hawkmoth *Manduca sexta*. Sci Rep. 8(2842):1-13.
- Ferreira LM, do Vale AE, de Souza AJ, Leite KB, Sacramento C, Moreno MLV, Araujo TH, Soares MBP, Grassi MFR (2019) Anatomical and phytochemical characterization of *Physalis angulata* L.: A plant with therapeutic potential. Pharmacogn Res. 11:171-7.

- Figueiredo MCC, Passos AR, Hughes FM, Santos KS, Silva AL, Soares TL (2020) Reproductive biology of *Physalis angulata* L. (Solanaceae). *Sci Hortic.* 267:1-10.
- Ghrab M, Zitouna R, Masmoudi MM, Mechlia NB (2016) Phenology and yield efficiency of early, mid-, and late-maturing cultivars of peach in irrigated orchards under mediterranean climate. *Int J Fruit Sci.* 16(3):323-334.
- Handa AK, Tiznado-Hernández M, Mattock AK (2011) Fruit development and ripening: a molecular perspective. In: Altman A, Hasegawa PM (eds) *Plant Biotechnology and Agriculture*. 1st edn. Oxford: Academic Press.
- Hatfield JL, Prueger JH (2015) Temperature extremes: Effect on plant growth and development. *Weather Clim Extremes.* 10:4-10.
- Horibe T (2018) Approach towards the control of rose flower opening by light environment. *Hortic Int J.* 2(4):197-199.
- Inoue K, Araki T, Endo M (2018) Circadian clock during plant development. *J Plant Res.* 131(1):59-66.
- Iwansyah AC, Juliani WP, Luthfiyanti R (2019) Characterization of nutrition, antioxidant properties and toxicity of *Physalis angulata* plant extract. *A J Pharm Clin Res.* 12(11):95-99.
- Khoo HE, Azlan A, Tang ST, Lim SM (2017) Anthocyanidins and anthocyanins: colored pigments as food, pharmaceutical ingredients, and the potential health benefits. *Food Nutr Res.* 61(1):1-21.
- Kusumaningtyas RW, Laili N, Limandha P (2015) Potential of ciplukan (*Physalis angulata* L.) as source of functional ingredient. In: Setyobudi RH, Scheer H, Limantara L, Shioi Y, Fiedor L, Brotosudarmo SHP, Prihastyanti MNU (eds) HK-ICONS 2014. Proceeding of the Meeting 2nd Humboldt Kolleg in Conjunction with International Conference on Natural Sciences, Procedia Chemistry 14, Malang, March 2015.
- Liguori G, Inglese P, Sortino G, Inglese G (2014) Dry matter accumulation and seasonal partitioning in mature *Opuntia ficus-indica* (L.) Mill. fruiting trees. *Ital J Agron.* 9(537):44-47.
- Lima MS, Evangelista AF, Santos GGL, Ribeiro IM, Tomassini TCB, Soares MBP, Villareal CF (2014) Antinociceptive properties of physalin from *Physalis angulata*. *J Nat Pro.* 77(11):2397-2404.
- McAtee P, Karim S, Schaffer R, David K (2013) A dynamic interplay between phytohormones is required for fruit development, maturation, and ripening. *Front Plant Sci.* 4(79):1-7.
- Medina-Medrano JR, Almaraz-Abarca N, González-Elizondo MS, Uribe-Soto JN, González-Valdes LS, Herrera-Arrieta Y (2015) Phenolic Constituents and antioxidant properties of five wild species of *Physalis* (Solanaceae). *Bot Stud.* 56:24.
- Meng Q, Fan J, Liu Z, Li X, Zhang F, Zhang Y, Sun Y, Li L, Liu X, Hua E (2019) Cytotoxic withanolides from the whole herb of *Physalis angulata* L. *Molecules.* 24(8):1-9.
- Muniz J, Kretzschmar AA, Rufato L, Pelizza TR, Rufato ADR, Macedo TA (2014) General aspect of *Physalis* cultivation. *Cienc Rural.* 44(6):964-970.
- Neves LC, Campos AJ, Cisneros-Zevallos L, Colombo RC, Roberto SR (2017) Postharvest behavior of camu-camu fruits based on harvesting time and nutraceutical properties. *Sci Hortic.* 217:276-284.
- Odjo S, Malumba P, Dossou J, Janas S, Béra F (2012) Influence of drying and hydrothermal treatment of corn on the denaturation of salt-soluble proteins and color parameters. *J Food Eng.* 109:562-570.
- Oliveira SF, Gonçalves FJA, Correia PMR, Guiné RPF (2016) Physical properties of *Physalis peruviana* L. *Open Agric.* 1:55-59.
- Ozaslan C, Farooq S, Onen H, Ozcan S, Bukun B, Gunal H (2017) Germination biology of two invasive *Physalis* species and implication for their management in arid and semi-arid region. *Sci Rep.* 7(16960):1-10.
- Purquerio LPV, Mattar GS, Duart AM, Moraes CC, Araújo HS, Santos FF (2019) Growth, yield, nutrient accumulation and export and thermal sum of Italian zucchini. *Hortic Bras.* 37(2):221-227.
- Ramírez F, Fischer G, Davenport TL, Pinzón JCA, Ulrichs C (2013) Short communication cape gooseberry (*Physalis peruviana* L.) phenology according to the BBCH phenological scale. *Sci Hort.* 162:39-42.
- Rodrigues FA, Soares JDR, Silva RAL, Penoni ES, Pasqual M, Pereira FJ, Castro EM (2014) Anatomy of vegetative organs and seed histochemistry of *Physalis peruviana* L. *Aust J Crop Sci.* 8(6):895-900.
- Saavedra JCM, Zaragoza FAR, Toledo DC, Hernández VCS, Vargas-Ponce O (2019) Agromorphological characterization of wild and weedy populations of *Physalis angulata* in Mexico. *Sci Hort.* 26:86-94.
- Sadiyah H, Soegianto A, Waluyo B, Ashari S (2020) Short communication: Preliminary characterization of groundcherry (*Physalis angulata*) from East Java Province, Indonesia based on morpho-agronomic traits. *Biodiversitas.* 21(2):759-769.
- Shamshiri RR, Jones JW, Thorp KR, Ahmad D, Man HC, Taheri S (2018) Review of optimum temperature, humidity, and vapour pressure deficit for microclimate evaluation and control in greenhouse cultivation of tomato: a review. *Int Agrophys.* 32:287-302.
- Santiago WR, Gama JSN, Torres SB, Bacchetta G (2019) Physiological maturity of *Physalis angulata* L. seeds. *Rev Ciênc Agron.* 5(3):431-438.
- Sholehah DN and Setiawan E (2019). Report of *Physalis angulata* L. from Madura: Quality profile. IOP Conference Series: Earth and Environmental Science 276:1-7.
- Souza VR, Pereira PAP, Queiroz F, Borges SV, Carneiro JDS (2012) Determination of bioactive compounds, antioxidant activity and chemical composition of Cerrado Brazilian fruits. *Food Chem.* 134(1):381-386.
- Souza CLM, Souza MO, Oliveira RS, Nascimento MN, Pelacani CR (2017) Biometric characteristics of fruits and physiological characterization of seeds of *Physalis* species (Solanaceae). *Rev Bras Ciênc Agr.* 12(3):
- Sun C, Qiu C, Zhao F, Kang N, Chen L, Qiu F (2017) Physalins V-IX, 16,24-cyclo-13,14-seco withanolides from *Physalis angulata* and their antiproliferative and anti-inflammatory activities. *Sci Rep.* 7(1):1-10.
- Tucker G, Yin X, Zhang A, Wang MM, Zhu Q, Liu X, Xie X, Chen K, Grierson D (2017) Ethylene and fruit softening. *Food Qual Saf.* 1(4):253-267.
- Walker J, Beurs K, Wynne RH (2015) Phenological response of an Arizona dryland forest to short-term climatic extremes. *Remote Sens.* 7:10832-10855.
- Yıldız G, İzli N, Halil U, Uylâşer V (2015) Physical and chemical characteristics of goldenberry fruit (*Physalis peruviana* L.). *J Food Sci Tech.* 52(4):2320-2327.

Fruit development of groundcherry (*Physalis angulata* L.) in dryland

ORIGINALITY REPORT



PRIMARY SOURCES

- | | | |
|---|---|-----|
| 1 | periodicos.ifap.edu.br
Internet Source | 1 % |
| 2 | www.pubmedcentral.nih.gov
Internet Source | 1 % |
| 3 | www.agriculturejournals.cz
Internet Source | 1 % |
| 4 | iopscience.iop.org
Internet Source | 1 % |
| 5 | STACEY D. SMITH, SANDRA KNAPP. "The natural history of reproduction in <i>Solanum</i> and <i>Lycianthes</i> (Solanaceae) in a subtropical moist forest", Bulletin of The Natural History Museum. Botany Series, 2002
Publication | 1 % |
| 6 | link.springer.com
Internet Source | 1 % |
| 7 | cyberleninka.org
Internet Source | 1 % |
-

8	assets.researchsquare.com Internet Source	1 %
9	www.derpharmacemica.com Internet Source	1 %
10	repositorio.upn.edu.pe Internet Source	<1 %
11	www.rbherbicidas.com.br Internet Source	<1 %
12	docplayer.net Internet Source	<1 %
13	Manik Chandra Roy, Majbaul Alam, Abu Saeid, Bijoy Chandra Das, Md. Biplob Mia, Md. Atikur Rahman, Jong Bang Eun, Maruf Ahmed. "Extraction and characterization of pectin from pomelo peel and its impact on nutritional properties of carrot jam during storage", Journal of Food Processing and Preservation, 2018 Publication	<1 %
14	philippinetambayan.com Internet Source	<1 %
15	issuu.com Internet Source	<1 %
16	www.alice.cnptia.embrapa.br Internet Source	<1 %

- 17 Pattison, Richard J., Fabiana Csukasi, and Carmen Catalá. "Mechanisms regulating auxin action during fruit development", *Physiologia Plantarum*, 2014.
Publication <1 %
- 18 Diana N. Sholehah, Sucipto Hariyanto, Hery Purnobasuki. "Effect of salinity on growth, physiology, and production of groundcherry (*Physalis angulata* L.)", *AIMS Agriculture and Food*, 2022
Publication <1 %
- 19 pharmacia.pensoft.net <1 %
Internet Source
- 20 www.locus.ufv.br <1 %
Internet Source
- 21 Dewi Desnilasari, Nok Afifah, Novita Indrianti. "Physicochemical, baking quality, and sensory evaluation of gluten free bread made from modified sweet potato flour with addition of nuts flour", *AIP Publishing*, 2019
Publication <1 %
- 22 Diana Nurus Sholehah, Dini Ermavitalini, Miratul Khasanah, Edy Setiti Wida Utami, Hery Purnobasuki. "Effect of foliar nutrients application to the growth and yield of *Physalis peruviana* and *Physalis alkekengi*", *IOP* <1 %

Conference Series: Earth and Environmental Science, 2022

Publication

- 23 R Mastuti, B Waluyo, J Batoro. "Response of seed germination and seedling growth of Physalis accession from East Java", Journal of Physics: Conference Series, 2020 <1 %
Publication
-
- 24 Rui Sun, Lei Sun, Kai-fang Wang. "Partial Least Squares Analysis of the Relationship Between Chemical Composition and Phenolic Compounds in Blueberry Cultivars", Food Analytical Methods, 2016 <1 %
Publication
-
- 25 Shahid Farooq, Huseyin Onen, Cumali Ozaslan, Ahmed M. El-Shehawi, Mona M. Elseehy. "Characteristics and methods to release seed dormancy of two ground cherry (Physalis) species", Journal of Applied Research on Medicinal and Aromatic Plants, 2021 <1 %
Publication
-
- 26 biodiversitas.mipa.uns.ac.id <1 %
Internet Source
-
- 27 cigrjournal.org <1 %
Internet Source
-
- 28 ejournal.unib.ac.id <1 %
Internet Source

<1 %

29 ijpp.gau.ac.ir <1 %
Internet Source

30 smujo.id <1 %
Internet Source

31 www.isnff-jfb.com <1 %
Internet Source

32 www.rgv.uefs.br <1 %
Internet Source

33 www.scielo.br <1 %
Internet Source

34 C.-W. Shao, H.-L. Chiu, S.-K. Chang. "A Study <1 %
on Generic Representation of Skeletal
Remains Replication of Prehistoric Burial",
ISPRS - International Archives of the
Photogrammetry, Remote Sensing and
Spatial Information Sciences, 2015
Publication

35 Mariana Carvalho Chaves Figueiredo, Adriana <1 %
Rodrigues Passos, Frederic Mendes Hughes,
Keylla Souza dos Santos et al. "Reproductive
biology of Physalis angulata L. (Solanaceae)",
Scientia Horticulturae, 2020
Publication

36

Ruiz Carrasco, Karina Beatriz <1972>
(Torrigiani, Patrizia). "Expression profiling in developing peach seed and mesocarp as affected by growth regulators", Alma Mater Studiorum - Università di Bologna, 2012.

<1 %

Publication

37

Yanan Hu, Zhenyun Han, Yaqiang Sun, Shuai Wang, Ting Wang, Yi Wang, Kenong Xu, Xinzhou Zhang, Xuefeng Xu, Zhenhai Han, Ting Wu. "ERF4 affects fruit firmness through TPL4 by reducing ethylene production", The Plant Journal, 2020

<1 %

Publication

38

Naseer Ahmad Bhat, Adil Gani, Faiza Jan, Khalid Muzaffar. "Process Standardization and Characterization of Chhurpi- A Himalayan Homemade Hard Cheese", Applied Food Research, 2022

<1 %

Publication

39

Samuel Luiz Fioreze, João Paulo Costa Carneiro, Daniela Dias Pinto, João Domingos Rodrigues. "Foliar application of calcium and kinetin on soybean at reproductive stage", Journal of Plant Nutrition, 2017

<1 %

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Fruit development of groundcherry (*Physalis angulata* L.) in dryland

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
