

View all documents >

Source details



0.144

Bali Medical Journal	0.2	(i)
Scopus coverage years: from 2019 to Present	0.2	
Publisher: Sanglah General Hospital		
ISSN: 2089-1180 E-ISSN: 2302-2914	SJR 2022 0.117	(i)
Subject area: (Medicine: General Medicine)	0.117	
Source type: Journal		
	SNIP 2022	\odot

☐ Save to source list Source Homepage

CiteScore CiteScore rank & trend Scopus content coverage

Set document alert

i Improved CiteScore methodology

CiteScore 2022 counts the citations received in 2019-2022 to articles, reviews, conference papers, book chapters and data papers published in 2019-2022, and divides this by the number of publications published in 2019-2022. Learn more >



CiteScore rank 2022 ①

Category	Rank	Percentile
Medicine General Medicine	#767/830	7th

View CiteScore methodology \gt CiteScore FAQ \gt Add CiteScore to your site c°

9/4/23, 10:46 AM Bali Medical Journal






Bali Medical Journal 8

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
Indonesia Universities and	Medicine Medicine	Sanglah General Hospital	4
research institutions in Indonesia	(miscellaneous)		
Media Ranking in Indonesia			
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	20891180, 23022914	2020-2022	Homepage
			How to publish in this journal
			srimaliawan@unud.ac.id

SCOPE

Bali Med. J. is open access, peer-reviewed journal aiming to communicate high-quality research articles, reviews, and general articles in the field. Bali Med. J. publish articles that encompass all aspects of basic research/clinical studies related to medical sciences. The Journal aims to bridge and integrate the intellectual, methodological, and substantive medical scholarship diversity and encourage a vigorous dialogue between medical scholars and practitioners.

Q Join the conversation about this journal



communicate and make sense of data with our **new data visualization**

tool.

Metrics based on Scopus® data as of April 2023

R. Sutomo 4 months ago

Dear SCImago Team,

I am an Indonesian, and I know exactly that the home base of this journal is in Bali, Indonesia. Would you kindly revise the information on the SCImago website? Thank you

reply



Melanie Ortiz 4 months ago

Dear Sutomo, thank you for contacting us. We will proceed to analyze your request as soon as possible. Greetings from Spain and thank you for using the SCImago products, SCImago Team

A Ahmad Al-Sarabbi 1 year ago

Dear Scimago,

On this page, you mentioned that the country of origin is Italy, but the journal's website says Indonesia. And if I am not mistaken, Bali is indeed a very famous, beautiful island in Indonesia.

Which one is accurate? Does your website post misleading information? Are other information regarding other journals can be trusted?

reply

Melanie Ortiz 1 year ago

Dear Ahmad,

Thank you for contacting us. We will revise that information based on Scopus as soon as

https://www.scimagojr.com/journalsearch.php?q=21101024217&tip=sid&clean=0

SCImago Team

SCImago Team





(https://balimedicaljournal.org)

Open Access & Peer Reviewed Multidisciplinary Journal of Medical Sciences

Search)

Advanced Search (/index.php/bmj/search/search)

(II)

Dr Barrie Tan - SG ENT Expert **Submit Your Paper** Ad Dr Barrie Tan ENT Ad Dove Medical Press Distributor ACP seven Leader in Shipping & Logistics Ad Alex58 Ad MSC Mana Mana Fine Living with Ascott Ad Mana Mana Ad DiscoverASR 34 Data Centres Including Asia JinPeng Mining Ad OVHcloud Ad JinPeng Mining Opening Offer 25% Off Coohom Fresh Sale for Pro Ad Ying'nFlo Admiralty Ad Coohom 3D Space Design

 $Home\ (https://balimedicaljournal.org/index.php/bmj/index) > Editorial\ Board\ \&\ Reviewer$

Editor-in-Chief

 $\textbf{Prof. Dr. Sri Maliawan, SpBS} \ (http://www.baliroyalhospital.co.id/halaman_staff.php?ditail=229)$

(Scopus ID (https://www.scopus.com/authid/detail.uri?authorId=15738530400)), (Google scholar (https://scholar.google.co.id/citations?user=qVJ57aYAAAAJ&hl=id)) srimaliawan@unud.ac.id / maliawans@yahoo.com
Department of Neuro Surgery, Udayana University

Sanglah General Hospital

Bali - Indonesia

Associate Editor

Prof. Putra Manuaba, M.Phil (http://profpuma.weebly.com/)

 $(Scopus ID) (https://www.scopus.com/authid/detail.uri?authorld=8412278400), (Google Scholar (https://scholar.google.com/citations?user=jnmT14kPWNcC&hl=en)) \\ putramanuaba@unud.ac.id / putramanuaba28@yahoo.com$

Biomedicine Postgraduate Program, Udayana University

Bali - Indonesia

4

Prof. Ketut Suwiyoga, SpOG (http://www.scopus.com/results/authorNamesList.url?sort=count-f&src=al&sid=01CAC4E9A2FB056A0A90221C03EC65FE.FZg2ODcJC9ArCe8WOZPv. NAME%28EQUALS%28Suwiyoga%29%29&st1=Suwiyoga&orcidId=&selectionPageSearch=anl&reselectAuthor=false&activeFlag=false&showDocument=false&resultsPerPage=20 (Scopus ID (https://www.scopus.com/authid/detail.uri?authorId=54080784800))

suwiyoga@unud.ac.id

Faculty of Medicine, Udayana University, Sanglah Hospital Denpasar, Bali-Indonesia

Editorial Board for Regional America

Ankit Sakhuja, M.B.B.S., F.A.C.P., F.A.S.N. (http://www.med.umich.edu/intmed/nephrology/STAFF/sakhuja_a1.htm)

(Scopus ID (http://www.scopus.com/authid/detail.url?authorld=16744977200))

asakhuja@med.umich.edu

Nephrology and Hypertension Cleveland Clinic (United States)

Editorial Board for Regional Australia

Professor John Svigos, AM

MBBS; DRCOG; CBioEth; FRCOG; FRANZCOG

(http://www.womenshealthspecialists.com.au/jsvigos.html)

(Scopus ID) (https://www.scopus.com/authid/detail.uri?authorId=6603773825)

john@svigos.com.au (mailto:john@svigos.com.au)

Discipline of Obstetrics & Gynaecology

Faculty of Health & Medical Sciences

University of Adelaide, South Australia

dr Deasy Ayuningtyas Tandio MPH-MBA (http://orcid.org/0000-0001-7847-2831).

(OrcidID) (https://orcid.org/0000-0001-7847-2831)

deasytandio@yahoo.com

James Cook University Australia Master of Public Health Master Of Business Administration, Indonesia

Editorial Board for Regional Europa

Prof. Harald Hoekstra

(Scopus ID) (https://www.scopus.com/authid/detail.uri?authorId=36038081900)

jsvigos@iprimus.com.au

Universitair Medisch Centrum Groningen, Division of Surgical Oncology, Groningen the Netherland

Editorial Board for Regional Asia

Prof Huang Qin (http://accu.cqu.edu.cn/web/eallprof/559.jhtml)

(Scopus ID) (https://www.scopus.com/authid/detail.uri?authorId=7409535321)

qhuang@cqu.edu.cn

Chairman Dept. of Neurosurgery, Guangdong 999 Hospital Guangzhou China

Assoc. Prof. Mohammad Amin Bahrami

(Scopus ID) (https://www.scopus.com/authid/detail.uri?authorId=55524082200)

aminbahrami1359@gmail.com

 $Head\ of\ healthcare\ management\ department,\ Shahid\ Sadoughi\ University\ of\ Medical\ Sciences,\ Yazd, Iran$

Dr. Tanveer Beg, PhD

(Scopus ID) (https://www.scopus.com/authid/detail.uri?authorId=6505772852)

tbmirza@jazanu.edu.sa

Assistant Professor, Department of Biology, Faculty of Science, Jazan University, Jazan, Saudi Arabia.

Editorial Board Members

Prof. Andi Asadul Islam

(Scopus ID) (https://www.scopus.com/authid/detail.uri?authorld=55504893500), (Google Scholar) (https://scholar.google.co.id/citations?user=vWs1RdMAAAAJ&hl=id&oi=sra) undee@med.unhas.ac.id

Faculty of Medicine Hasanudin University, Makasar-Indonesia



Prof. Dr. dr. Abdul Hafid Bajamal, Sp.BS

(Scopus ID) (https://www.scopus.com/authid/detail.uri?authorId=57192378862)

hfbajamal@gmail.com

Faculty of Medicine Airlangga University, Surabaya-Indonesia

Dr. dr. I Wayan Sudarsa, Sp.B(K) Onk, FINACS, FICS.

(Scopus ID (https://www.scopus.com/authid/detail.uri?authorId=57205145862)), (Google Scholar (https://scholar.google.co.id/citations?hl=id&user=SdInHKwAAAAJ)), (Researchg

$\textbf{dr. I.B. Amertha P. Manuaba, SKed, MBiomed.} \ (\text{https://scholar.google.co.} id/citations?user=KzCQgA0AAAAJ\&hl=en)$

AmerthaManuaba@gmail.com / Amertha_Manuaba@unud.ac.id

Faculty of Medicine, Universitas Udayana, Indonesia

Editorial inquiries to be addressed to:

email 1: editorbalimedicaljournal@gmail.com (mailto:editor@balimedicaljournal.org)

email 2: editor@balimedicaljournal.org (mailto:editor@balimedicaljournal.org)



WEB OF SCIENCE** (https://mjl.clarivate.com/search-results?issn=2089-1180&hide_exact_match_fl=true&utm_source=mjl&utm_medium=share-bylink&utm_campaign=search-results-share-this-journal)



(https://www.scopus.com/sourceid/21101024217)



(https://doaj.org/toc/2302-2914)



(https://sinta3.kemdikbud.go.id/journals/profile/2513)

Full Indexing List (https://balimedicaljournal.org/index.php/bmj/pages/view/indexing)

In Press (https://balimedicaljournal.org/index.php/bmj/issue/view/30)

Submit An Article (https://balimedicaljournal.org/index.php/bmj/login)

Scopus Citedness (https://balimedicaljournal.org/index.php/bmj/pages/view/scopus)



(https://balimedicaljournal.org/index.php/bmj/)

Published by: (http://www.discoversys.ca/)

For Indonesian Physician Forum and Indonesia College of Surgeons, Indonesia	
Bali Medical Journal, Bali-Indonesia	
☐ 62 (0369) 225206	
♣ 62 (0369) 225206	
■ administrator@balimedicaljournal.org (mailto:administrator@balimedicaljournal.org)	
Contact (/index.php/bmj/pages/view/contact)	
Journal Information (/index.php/bmj/pages/view/journalinfo)	
Editorial Board (/index.php/bmj/pages/view/editorialboard)	
Abstracting & Indexing (/index.php/bmj/pages/view/indexing)	
Privacy Statement (http://discoversys.ca/privacy.html)	
Home (/index.php/bmj/index)	
Last Issue (/index.php/bmj/issue/current)	
Archive (/index.php/bmj/issue/archive)	
Author Guidelines (/index.php/bmj/pages/view/authorguidlines)	
Open-Access Licence (/index.php/bmj/pages/view/OAlicence)	
Copyright © 2008-2022 DiscoverSys Inc (http://discoversys.ca/). All rights reserved. [CO] SYN DEFINITION (http://creativecommons.org/licenses/by-nc-nd/4.0/) (http://www.crossref.org/citedby/index.html) (http://creativecommons.org/licenses/by-nc-nd/4.0/) (http://www.crossref.org/citedby/index.html) (http://creativecommons.org/licenses/by-nc-nd/4.0/) (http://discoversys.ca/privacy.html) No Fee (http://discoversys.ca	





(https://www.balimedicaliournal.org)

Open Access & Peer Reviewed Multidisciplinary Journal of Medical Sciences

Search		
(

Advanced Search (/index.php/bmj/search/search)

Free ChatGPT Browser Extension	Ramen Tonkotsu di Fukuoka Ikian Ali Nippon Airways
Write With Confidence	New Body is Waiting
Articles and Info	Claim Your Free Trial Iklan American Journal Experts
Ship your Produces with MSC	Be a VTuber with nizima LIVE
Annual Cybersecurity Report Iklan CrowdStrike®	Create an Event on Facebook
No Setup Fee	Online interactive learning Relan LIGS University
Sprei California by My Love	Hapuskan Risiko di Proyek Anda
Create an Event on Facebook	•

Home (https://www.balimedicaljournal.org/index.php/bmj/index) > Archives (https://www.balimedicaljournal.org/index.php/bmj/issue/archive) > Vol. 11 No. 1 (2022): (Available online: 1 April 2022)

Vol. 11 No. 1 (2022): (Available online: 1 April 2022)


ORIGINAL ARTICLE


Evaluating Low Values of Early Diastolic Velocity (e') as a Predictor of Major Cardiovascular **Events in Patients with Acute** Myocardial Infarction


ORIGINAL ARTICLE

The prevalence and characteristics of perineal rupture during vaginal delivery at Sanglah General Hospital and Regional Hospitals in Bali from January (https://www.balimedicaljournal.org/index.php/bmj/article/view/3360)




























Olivia Desty Sabunga, Cahyono Kaelan, Upik Anderiani Miskad, Andi Alfian Zainuddin, Ni Ketut Sungowati, Muhammad Husni Cangara Online First: Apr 8, 2022	Malayana Rahmita Nasution, Putri Chadijah Tampubolon, Irma Sari Nasution Online First: Apr 30, 2022
P pdf Abstract (https://www.balimedicaljournal.org/i	Abstract (https://www.balimedicaljournal.org/index.php/bmj/article/view/3185/ndex.php/bmj/article/view/3078/2006)
ORIGINAL ARTICLE Relationship between CD4 levels and mucocutaneous manifestations in HIV-AIDS patients at Dr. Soetomo General Academic Teaching Hospital, Surabaya, Indonesia (https://www.balimedicaljournal.org/inc	ORIGINAL ARTICLE Correlation between sociodemographic and attitude of Malang citizens about self medication on urticaria (https://www.balimedicaljournal.org/index.php/bmj/article/vialdex.php/bmj/ar
Citra Dwi Harningtyas, Damayanti, Maylita Sari, Muhammad Yulianto Listiawan, Diah Mira Indramaya, Linda Astari, Budi Utomo, Dwi Murtiastutik, Setyana Widyantari, Astindari, Afif Nurul Hidayati Online First: Apr 13, 2022	Online First: Apr 17, 2022 Podf
Abstract (https://www.balimedicaljournal.org/ii ORIGINAL ARTICLE The effect of metformin on	ORIGINAL ARTICLE Antiproliferation and Apoptosis
autophagy by LC3 expression in Type 2 Diabetes Mellitus (T2DM) human skeletal muscle cell culture (https://www.balimedicaljournal.org/inc Jongky Hendro Prajitno, Agung Pranoto, Robert Dwitama Adiwinoto, Soebagijo Adi Soelistijo Online First: Apr 30, 2022	Effect of Cisplatin and Nanocurcumin on Ovarian Cancer SKOV3 Cell (https://www.balimedicaljournal.org/index.php/bmj/article/violdex.php/bmj/article/vi
Abstract (https://www.balimedicaljournal.org/i	
ORIGINAL ARTICLE	ORIGINAL ARTICLE





















WEB OF SCIENCE** (https://mjl.clarivate.com/search-results?issn=2089-1180&hide_exact_match_fl=true&utm_source=mjl&utm_medium=share-bylink&utm_campaign=search-results-share-this-journal)





(https://doaj.org/toc/2302-2914)



Full Indexing List (https://balimedicaljournal.org/index.php/bmj/pages/view/indexing)

In Press (https://balimedicaljournal.org/index.php/bmj/issue/view/30)

Submit An Article (https://balimedicaljournal.org/index.php/bmj/login)

Scopus Citedness (https://balimedicaljournal.org/index.php/bmj/pages/view/scopus)

(//clustrmaps.com/site/1a4xh? utm_source=globe)



(https://balimedicaljournal.org/index.php/bmj/)

Published by Pali Medical Journal, Bali-Indonesia (http://www.discoversys.ca/) 62 (0369) 225206 **4** 62 (0369) 225206 For Indonesian Physician Forum and Indonesia College of Surgeons, Indonesia ▲ administrator@balimedicaljournal.org (mailto:administrator@balimedicaljournal.org) Home (/index.php/bmj/index) Contact (/index.php/bmj/pages/view/contact) Last Issue Iournal Information (/index.php/bmj/issue/current) (/index.php/bmj/pages/view/journalinfo) Archive Editorial Board (/index.php/bmj/issue/archive) (/index.php/bmj/pages/view/editorialboard) Author Guidelines Abstracting & Indexing (/index.php/bmj/pages/view/authorguidlines) (/index.php/bmj/pages/view/indexing) Open-Access Licence Privacy Statement (/index.php/bmj/pages/view/OAlicence) (http://discoversys.ca/privacy.html)

Copyright © 2008-2022 DiscoverSys Inc (http://discoversys.ca/). All rights reserved.

(http://creativecommons.org/licenses/by-nc-nd/4.0/) (http://www.crossref.org/citedby/index.html) (http://www.sherpa.ac.uk/romeo/pub/1931/) (http://discoversys.ca/privacy.html) (http://discoversys.ca/privacy.html) (http://discoversys.ca/privacy.html) (http://discoversys.ca/privacy.html) (http://discoversys.ca/privacy.html) (http://discoversys.ca/privacy.html) (http://discoversys.ca/privacy.html) (http://discoversys.ca/privacy.html) (http://the-acap.org/acap-enabled.php)

Bali Medical Journal (*Bali MedJ*) 2022, Volume 11, Number 1: 349-355 P-ISSN.2089-1180, E-ISSN: 2302-2914



The effect of metformin on autophagy by LC3 expression in Type 2 Diabetes Mellitus (T2DM) human skeletal muscle cell culture



Jongky Hendro Prajitno¹, Agung Pranoto^{2*,} Robert Dwitama Adiwinoto³, Soebagijo Adi Soelistijo²

¹Doctoral Program Student, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia;
²Division of Endocrinology, Diabetes, and Metabolism, Department of Internal Medicine, Dr. Soetomo Teaching Hospital, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia;
³Division of Endocrinology, Diabetes, and Metabolism, Department of Internal Medicine, Dr. Ramelan Naval Hospital, Surabaya, Indonesia;

*Corresponding author:
Agung Pranoto;
Division of Endocrinology, Diabetes,
and Metabolism, Department of
Internal Medicine, Dr. Soetomo Teaching
Hospital, Faculty of Medicine, Universitas
Airlangga, Surabaya, Indonesia;
pranotoagung@yahoo.com

Received: 2022-02-13 Accepted: 2022-04-20 Published: 2022-04-30

ABSTRACT

Background: Skeletal muscle is an essential tissue in glucose metabolism. Reduced autophagic capacity to remove damaged contractile proteins in skeletal muscle cells will contribute to the loss of muscle mass. Metformin is the first-line agent for treating type 2 diabetes mellitus (T2DM) patients. This study aims to investigate the effects of metformin on autophagy through LC3 expression in human skeletal muscle cell culture (SkMC).

Methods: The T2DM human SkMC was obtained from T2DM treatment naïve patients, purchased from AcceGen Biotech®. Fully differentiated myotubes were randomized into the control and treatment groups. The treatment group was given Metformin in three doses (1 mM, 2 mM, and 3 mM). An immunoblotting assay of AMPKα and LC3 was performed using electrochemiluminescence (ECL). The quantitative expression of AMPKα and LC3 were measured at baseline, after 24-hour, 48-hour, and 72-hour. Data were analyzed using SPSS version 22 for Windows. Results: AMPKα and LC3 expression were higher in the treatment group compared to the control group. The levels of AMPKα and LC3 expression in the treatment group increased dose-dependent. Linear regression analysis demonstrated a significant correlation between metformin administration and LC3 expression levels (p<0.0001).

Conclusion: Metformin administration on T2DM human SkMC resulted in increased autophagic activity, marked by increased LC3 expression.

Keywords: type 2 diabetes mellitus, skeletal muscle, autophagy, Metformin.

Cite This Article: Prajitno, J.H., Pranoto, A., Adiwinoto, R.D., Soelistijo, S.A. 2022. The effect of metformin on autophagy by LC3 expression in Type 2 Diabetes Mellitus (T2DM) human skeletal muscle cell culture. *Bali Medical Journal* 11(1): 349-355. DOI: 10.15562/bmj.v11i1.3203

INTRODUCTION

Skeletal muscle plays an important role in glucose metabolism.1 During muscle contraction, the blood glucose levels in the peripheral blood vessels decrease. This is evidenced by a previous study that indicated an association between muscle strength and blood glucose control.2 Insulin triggers glucose uptake by muscle cells through the Glucose transporter (GLUT) Sodium dependent glucose transporter-2 (SGLT-2).3 A greater muscle mass and GLUT4 expression were found in the trained muscles and associated with greater glucose uptake into the muscle. Meanwhile, a decrease in muscle mass and GLUT4 expression in unhealthy muscles diminish glucose uptake.4 Older patients with T2DM have a two-fold muscle mass loss compared to the elderly without T2DM.5 Due to this important role in

glucose utilization, loss of skeletal muscle mass (sarcopenia) will worsen insulin resistance (IR) in T2DM.6

Macroautophagy (autophagy) is a process in a cell characterized by developing a closed double-membrane vesicle called the autophagosome. The autophagosome then fused with the lysosome to degrade its content. Autophagy largely engulfs a portion of the cytoplasm non-selectively (bulk). However, it can also selectively target dysfunctional organelles or harmful proteins through specific adaptor proteins.7 This selectivity may preserve cellular function in many tissues during aging. Previously, autophagy was assumed to increase protein degradation; therefore, it might result in sarcopenia. Recent studies have indicated that autophagy maintains muscle mass. Reduced autophagic capacity to remove damaged contractile proteins and dysfunctional organelles contribute to age-related decline in myofibril function and muscle strength in humans.8

Type 2 Diabetes Mellitus (T2DM) is a metabolic disorder that developed due to pancreatic islet β-cells' failure to sustain the hyperinsulinemia required to compensate for IR.9,10 Insulin resistance diminishes autophagy signaling in skeletal muscle cells. Decreased autophagy will affect the function of insulin-sensitive tissues, including skeletal muscle, and autophagy-deficient skeletal displays similar characteristics as insulinresistant muscle, such as mitochondrial dysfunction. The mechanism by which IR affects autophagy might be due to an intrinsic defect in autophagy or a response to hyperinsulinemia.11 Previous study has demonstrated reduced autophagy response to metabolic stress in human muscle precursor cells from humans with T2DM.12 Insulin has been linked to

inhibition of autophagy through activation of mammalian target of rapamycin (mTOR) which is a negative regulator of autophagy.¹³

Metformin, a derivate of biguanide, is the first-line agent for treating T2DM patients. Metformin can lower blood sugar levels by inhibiting gluconeogenesis adenosine monophosphatethrough activated protein kinase (AMPK) activation.14 A previous study indicated that metformin treatment positively impacted skeletal muscle IR levels.15 It also lowers fat mass, increases muscle mass which is atrophied due to obesity, and attenuates muscle cells oxidative stress in mice models. 16,17 However, some studies also reported that metformin might cause mitochondrial dysfunction and inhibit satellite cell proliferation, impeding mass muscle gain.18-20 Microtubule-associated protein 1A/1B-light chain 3 (LC3) immunoblotting is commonly used to evaluate autophagy in intact muscle tissue. It is an autophagy-related gene that is essential for forming autophagosomes in muscle cells.21

Although several studies have examined the effects of Metformin on skeletal muscle in mice models, to our knowledge, the data regarding the effects of Metformin on human skeletal muscle is limited. Therefore, our present study aimed to investigate the effects of Metformin on autophagy through LC3 expression in human SkMC.

METHODS

The present study was an experimental, in vitro laboratory research, with a posttestonly control group design, carried out at the laboratory of the physiology of Universitas Brawijava. The T2DM human SkMC originated from Human Skeletal Muscle Myoblasts (HSMM) ABC-TC3957 T2DM cell line, which was obtained from T2DM treatment naïve patients, purchased from AcceGen Biotech®. The SkMC were grown in a 36-mm petri dish with glucose-rich DMEM (Dulbecco's Modified Eagle's medium containing 4.5 grams of glucose) and 20% Fetal Bovine Serum (FBS) and 1% antibiotic solution (penicillin-streptomycin-glutamine, PSG), incubated at 37°C with 5% CO2 for three days. To fully differentiate the myotubes,

dishes with a density of 65 - 75% cells were cultured in DMEM with 2% Horse Serum and 1% PSG solution, incubated at 37°C with 5% CO2 until the cell density reached 80% for a day. The media was changed every day during the fusion process, and the process was monitored with a contrast microscope using a 10 - 20 times magnification. Fully differentiated myotubes were randomized using the simple random sampling technique into control and treatment groups. The treatment group was given Metformin in three different doses (1 mM, 2 mM, and 3 mM), with 4 replications in each group, and then incubated for 2 days before the further examination. 15,22,23

All myotubes were lysed, and then the lysates were rinsed with G-agarose for 20 minutes and incubated with a specific antibody overnight at 4°C. Protein G particles were precipitated for an hour at 4°C and then washed 4 times with lysis buffer solution before being inserted into polyacrylamide gel. After being inserted into 15% SDS-polyacrylamide gel, electrophoresis was performed on all samples. The nitrocellulose membrane was blocked with Tris buffer solution and then incubated overnight at 4°C with antibodies: anti-AMPKa2 (Ser173) and anti-LC3. Anti-AMPKa2 (Ser173) was diluted 1: 2000.24 The nitrocellulose membrane was then rinsed 3 times using PBS-T solution and incubated for an hour with

Horseradish Peroxidase (HRP) secondary conjugated antibody. 25,26 Immunoblotting assay of AMPK α and LC3 was performed using ECL according to the manufacturer protocol based on the previous study. 27,28 The quantitative expression of AMPK α and LC3 were measured at baseline, after 24-hour, 48-hour, and 72-hour.

The collected data are presented as the mean ± standard error (SE). Analysis using one-way analysis of variance (ANOVA) followed by Tukey's post-hoc test was performed to detect a significant difference between groups. Kruskal-Wallis test was used instead for data that were not normally distributed. Differences were considered statistically significant at p < 0.05. Analysis using linear regression was performed to determine the correlation between the dose of metformin treatment, the timing of observation, and the expression of AMPKa and LC3. The analysis was performed using SPSS for Windows version 22.

RESULTS

Since Metformin is a drug that acts through AMPK activation, the treatment group expressed AMPK α higher than the control group in this present study. This observation became more evident after 72-hour of metformin administration which displayed a significant difference in AMPK α levels from the control




Figure 1. Effects of Metformin on AMPKα expression. Immunoblotting quantification (mean ± SE) was measured at 72-hour after Metformin was administered. Analysis of variance (ANOVA) followed by Tukey's post-hoc test results: ^a*p* < 0.05 when compared to control group, ^b*p* < 0.05 when compared to Met 1 mM group, ^c*p* < 0.05 when compared to Met 2 mM group, ^d*p* < 0.05 when compared to Met 3 mM group. Met 1 mM: group treated with 1 mM of Metformin; Met 2 mM: group treated with 2 mM of Metformin; Met 3 mM: group treated with 3 mM of Metformin.

group and each increment of metformin concentration (Figure 1). The level of AMPK α expression at baseline, after 24-hour, and 48-hour of metformin administration are represented in Figures 2, 3, and 4. Figure 5 depicts the visualization of AMPK α expression under

the fluorescent microscope for 72-hour.

Treatment with Metformin significantly increased the levels of LC3 expression in T2DM human SkMC. The treatment group with the concentration of 3 mM displayed the highest expression of LC3 levels both at baseline and after 24 48-hour of metformin




Figure 2. Effects of Metformin on AMPKα expression. Immunoblotting quantification (mean \pm SE) was measured at baseline. Analysis of variance (ANOVA) followed by Tukey's post-hoc test results: ${}^ap < 0.05$ when compared to control group, ${}^bp < 0.05$ when compared to Met 1 mM group, ${}^cp < 0.05$ when compared to Met 2 mM group, ${}^dp < 0.05$ when compared to Met 3 mM group. Met 1 mM: group treated with 1 mM of Metformin; Met 2 mM: group treated with 2 mM of Metformin; Met 3 mM: group treated with 3 mM of Metformin.




Figure 3. Effects of Metformin on AMPKα expression. Immunoblotting quantification (mean \pm SE) was measured at 24-hour after Metformin was administered. Analysis of variance (ANOVA) followed by Tukey's post-hoc test results: ap < 0.05 when compared to control group, bp < 0.05 when compared to Met 1 mM group, cp < 0.05 when compared to Met 2 mM group, dp < 0.05 when compared to Met 3 mM group. Met 1 mM: group treated with 1 mM of Metformin; Met 2 mM: group treated with 2 mM of Metformin; Met 3 mM: group treated with 3 mM of Metformin.

administration (Figures 6, 7, and 8). After 72-hour of metformin administration, LC3 expression in the group treated with 1 mM and 2 mM metformin raised significantly, displaying a statistically significant difference from the 3 mM group (Figure 9). Figure 10 represents the visualization of LC3 expression under the fluorescent microscope for 72-hour.

By using linear regression analysis, a significant correlation between metformin administration at increased concentrations and LC3 expression levels was found (p<0.0001). However, no significant correlation was established between the expression of AMPK α and LC3. These findings suggested that metformininduced autophagy in T2DM human SkMC is somewhat AMPK α independent.

DISCUSSION

Metformin is the first-line agent for treating hyperglycemia in T2DM patients. There is abundant evidence that the pharmacological action of Metformin is from the activation of AMPK and its downstream target. Considering AMPK as the main sensor of the cell's energy status, it also plays an important role in regulating autophagy. Previously, autophagy was presumed to promote sarcopenia of skeletal muscle. However, a line of evidence has accumulated, demonstrating that autophagy is important in skeletal muscle homeostasis. Skeletal muscle is an essential tissue in glucose metabolism and LC3 is widely used to monitor autophagy in skeletal muscle. Therefore, we are interested in investigating the effect of metformin administration on LC3 expression inT2DM human SkMC.

The process of autophagy is tightly regulated, involving numerous autophagyrelated genes (ATG). In general, autophagy can be broken down into the following steps: initiation, vesicle (autophagosome) expansion, lysosome fusion, degradation. As the sensor of cellular energy status, AMPK incites autophagy initiation, especially during starvation.29 Numerous pharmacological agents are also known to target AMPK activation, directly and indirectly, for example, The Metformin.30 ULK-1 complex activation is the central mechanism in autophagy initiation. Upon activation,




Figure 4. Effects of Metformin on AMPKα expression. Immunoblotting quantification (mean \pm SE) measured at 48-hour after Metformin was administered. Analysis of variance (ANOVA) followed by Tukey's post-hoc test results: ap < 0.05 when compared to control group, bp < 0.05 when compared to Met 1 mM group, cp < 0.05 when compared to Met 2 mM group, dp < 0.05 when compared to Met 3 mM group. Met 1 mM: group treated with 1 mM of Metformin; Met 2 mM: group treated with 2 mM of Metformin; Met 3 mM: group treated with 3 mM of Metformin.




Figure 5. Representative images of AMPKα expression with immunofluorescence at 72-hour after Metformin was administered. Met 1 mM: group treated with 1 mM of Metformin; Met 2 mM: group treated with 2 mM of Metformin; Met 3 mM: group treated with 3 mM of Metformin.

the ULK-1 complex then assembles the autophagosome with the Beclin 1-Vps34 complex in the next step. The autophagosome formation is completed once the two ubiquitin-like conjugation systems, namely ATG12-ATG5-ATG16L and ATG8, form the lipid conjugated form of LC3. The autophagosome is then fused with the lysosome to form the autolysosome, followed by degradation of its content.³¹

The main result of our present study was the significant correlation between metformin administration and LC3 expression. Metformin raised LC3 expression levels in the T2DM human SkMC dose-dependent. This finding is

similar to the study by Kanamori H et al., which demonstrated an increase in LC3 and autophagic activity in cardiomyocytes treated with Metformin in the long term.³² Those findings might be due to inhibition p62 activity, which suppresses autophagy. During starvation, the increase in AMPKa and FoxO3 expression will be followed by increased LC3 activity.25 In contrast, Nwadike C et al., stated that AMPK activation might inhibit early and late phases of autophagy.33 Autophagy regulation in skeletal muscle is also dependent on FoxO3 activity. In a starved skeletal muscle model, deletion of FoxO3 activity will impede the LC3 lipidation process.34 This finding is also backed by

the study of Sanchez AM et al., on AICAR treated muscle cells which removed AMPK α and FoxO3 activity, still resulting in LC3 expression due to AMPK α and FoxO3 residual activity.²⁵

Our present study demonstrated that Metformin raised the expression of AMPKalevels in T2DM human SkMC. This finding is supported by the study of Li DI et al., in which metformin administration with a dose of 2 mM in vitro for 48hour raised AMPKa expression levels significantly in skeletal muscle cells.22 The elevation of AMPKα activity is related to the increase of Thr-172 phosphorylation and Acetyl-CoA Carboxylase (ACC) inactivation.35,36 This increase in AMPKa activity is independent of the cellular ADP/ATP ratio.35 However, some in vivo studies reported different results. The studies by Suwa M et al., in humans and Kristensen JM et al., in mice reported that even though the increase in AMPKa expression and ACC phosphorylation can be observed within 5 - 6 hours after metformin administration, after days, these findings were negligible.37,38 Administration of Metformin with the dose of 0.4 mM resulted in a dramatic increase in AMPKa expression within 24-hour without inhibition in mTOR expression.39 Queiroz EA et al., also reported no significant difference in the AMPKa expression after 10 mM metformin administration in breast cancer cells, although notable phosphorylation of AMPKa and Thr-172 were observed after 48 and 72 hours. 40

The present study failed to demonstrate a significant correlation between AMPKa and LC3 expression in the SkMC in the context of autophagic activity. This finding, however, is in contrast with several previous studies on autophagy in which the majority stated that AMPK is associated with autophagic flux. Autophagy is a complex process involving an interconnected web of key regulators rather than a linear pathway. Many of these regulators can induce autophagy through different requirements depending on the type and length of induction signals.41 Aside from the AMPK-dependent Regulation of autophagic degradation, an AMPK-independent pathway triggers Ca²⁺-mediated autophagy, involving the PI(3)P-effector protein WIPI-1 and LC3.⁴² A study by Tomic T et al., regarding the autophagic and apoptotic properties of Metformin on melanoma cells, revealed that the increase in LC3 activity occurred by either AMPK-dependent or AMPK-independent pathways.⁴³

Our study has several limitations

to be acknowledged. First, the doses of Metformin used in this present study were relatively small (1 mM, 2 mM, and 3 mM). The effect of a higher dose (≥ 5 mM) of Metformin on the autophagic activity of human skeletal muscle cells still needs further investigation. Second, the HSMM ABC-TC3957 T2DM cell




Figure 6. Effects of Metformin on LC3 expression. Immunoblotting quantification (mean \pm SE) was measured at baseline. Analysis of variance (ANOVA) followed by Tukey's post-hoc test results: ${}^{a}p < 0.05$ when compared to control group, ${}^{b}p < 0.05$ when compared to Met 1 mM group, ${}^{c}p < 0.05$ when compared to Met 2 mM group, ${}^{d}p < 0.05$ when compared to Met 3 mM group. Met 1 mM: group treated with 1 mM of Metformin; Met 2 mM: group treated with 2 mM of Metformin; Met 3 mM: group treated with 3 mM of Metformin.




Figure 7. Effects of Metformin on LC3 expression. Immunoblotting quantification (mean \pm SE) was measured at 24-hour after Metformin was administered. Analysis of variance (ANOVA) followed by Tukey's post-hoc test results: ap < 0.05 when compared to control group, bp < 0.05 when compared to Met 1 mM group, cp < 0.05 when compared to Met 2 mM group, dp < 0.05 when compared to Met 3 mM group. Met 1 mM: group treated with 1 mM of Metformin; Met 2 mM: group treated with 2 mM of Metformin; Met 3 mM: group treated with 3 mM of Metformin.

lines used in this study were derived from Caucasians. Therefore, our results should be interpreted carefully, and future studies with more T2DM muscle cell lines are required to validate these findings. Third, our study only observed some of the autophagy-related proteins. Other downstream protein targets of AMPK (FoxO3, ULK-1, MURF-1) and their relation to autophagy in T2DM muscle cells still need to be investigated.

CONCLUSION

This present study has demonstrated that metformin administration on T2DM human SkMC resulted in increased autophagic activity, marked by increased LC3 expression in a dose-dependent manner. Our study could not establish the correlation between the increase in AMPKα expression and LC 3 expression, which suggests that increased autophagy in the skeletal muscle might have occurred in AMPK-independent pathways.

ACKNOWLEDGMENTS

None.

CONFLICT OF INTEREST

The authors declared no conflict of interests.

ETHICS CONSIDERATION

This research was reviewed and approved by the Health Research Ethics Committee of the Faculty of Medicine, Universitas Airlangga, Surabaya, with reference number: 51/EC/KEPK/FKUA/2021.

FUNDING

None.

AUTHOR CONTRIBUTION

All authors contribute to the study from the conceptual framework, data acquisition, and data analysis until reporting the study results through publication.

REFERENCES

 Jaiswal N, Gavin MG, Quinn WJ 3rd, Luongo TS, Gelfer RG, Baur JA, et al. The role of skeletal muscle Akt in the regulation of muscle mass and glucose homeostasis. Mol Metab. 2019;28:1-13.

- Bawadi H, Alkhatib D, Abu-Hijleh H, Alalwani J, Majed L, Shi Z. Muscle Strength and Glycaemic Control among Patients with Type 2 Diabetes. Nutrients. 2020;12(3):771.
- Evans PL, McMillin SL, Weyrauch LA, Witczak CA. Regulation of Skeletal Muscle Glucose
- Transport and Glucose Metabolism by Exercise Training. *Nutrients*. 2019;11(10):2432.
- Sinacore DR, Gulve EA. The role of skeletal muscle in glucose transport, glucose homeostasis. and insulin resistance:




Figure 8. Effects of Metformin on LC3 expression. Immunoblotting quantification (mean \pm SE) measured at 48-hour after Metformin was administered. Analysis of variance (ANOVA) followed by Tukey's post-hoc test results: ap < 0.05 when compared to control group, bp < 0.05 when compared to Met 1 mM group, cp < 0.05 when compared to Met 2 mM group, dp < 0.05 when compared to Met 3 mM group. Met 1 mM: group treated with 1 mM of Metformin; Met 2 mM: group treated with 2 mM of Metformin; Met 3 mM: group treated with 3 mM of Metformin.




Figure 9. Effects of Metformin on LC3 expression. Immunoblotting quantification (mean \pm SE) was measured at 72-hour after Metformin was administered. Analysis of variance (ANOVA) followed by Tukey's post-hoc test results: ap < 0.05 when compared to control group, bp < 0.05 when compared to Met 1 mM group, cp < 0.05 when compared to Met 2 mM group, dp < 0.05 when compared to Met 3 mM group. Met 1 mM: group treated with 1 mM of Metformin; Met 2 mM: group treated with 2 mM of Metformin; Met 3 mM: group treated with 3 mM of Metformin.

- implications for physical therapy. *Phys Ther*. 1993;73(12):878-891.
- Park SW, Goodpaster BH, Lee JS, Kuller LH, Boudreau R, de Rekeneire N, et al. Excessive loss of skeletal muscle mass in older adults with type 2 diabetes. Diabetes Care. 2009 Nov;32(11):1993-7.
- Abidin Öztürk ZA, Türkbeyler İH, Demir Z, Bilici M, Kepekçi Y. The effect of blood glucose regulation on sarcopenia parameters in obese and diabetic patients. *Turk J Phys Med Rehabil*. 2017;64(1):72-79.
- Mizushima N, Komatsu M. Autophagy: renovation of cells and tissues. Cell. 2011;147(4):728-741.
- Jiao J, Demontis F. Skeletal muscle autophagy and its role in sarcopenia and organismal aging. Curr Opin Pharmacol. 2017;34:1-6.
- Prentki M, Nolan CJ. Islet beta cell failure in type 2 diabetes. J Clin Invest. 2006;116(7):1802-1812.
- Kahn SE, Cooper ME, Del Prato S. Pathophysiology and treatment of type 2 diabetes: perspectives on the past, present, and future. *Lancet*. 2014;383(9922):1068-1083.
- Dudani S, Poodury S, Mangalesh S. Study of neutrophil-lymphocyte ratio (NLR) in recent onset type 2 diabetes mellitus. Bali Medical Journal. 2021;10(1):11-16.
- Henriksen TI, Wigge LV, Nielsen J, Pedersen BK, Sandri M, Scheele C. Dysregulated autophagy in muscle precursor cells from humans with type 2 diabetes. Sci Rep. 2019;9(1):8169.
- Settembre C, Di Malta C, Polito VA, Garcia Arencibia M, Vetrini F, Erdin S, et al. TFEB links autophagy to lysosomal biogenesis. Science. 2011;332(6036):1429-33.
- Zhou G, Myers R, Li Y, Chen Y, Shen X, Fenyk-Melody J, et al. Role of AMP-activated protein kinase in mechanism of metformin action. J Clin Invest. 2001;108(8):1167-74.
- Yuan H, Hu Y, Zhu Y, Zhang Y, Luo C, Li Z, et al. Metformin ameliorates high uric acid-induced insulin resistance in skeletal muscle cells. Mol Cell Endocrinol. 2017;443:138-145.
- Hasan MM, Shalaby SM, El-Gendy J, Abdelghany EMA. Beneficial effects of metformin on muscle atrophy induced by obesity in rats. J Cell Biochem. 2019;120(4):5677-5686.
- Diniz Vilela D, Gomes Peixoto L, Teixeira RR, Belele Baptista N, Carvalho Caixeta D, Vieira de Souza A, et al. The Role of Metformin in Controlling Oxidative Stress in Muscle of Diabetic Rats. Oxid Med Cell Longev. 2016;2016:6978625.
- Wessels B, Ciapaite J, van den Broek NM, Nicolay K, Prompers JJ. Metformin impairs mitochondrial function in skeletal muscle of both lean and diabetic rats in a dose-dependent manner. PLoS One. 2014;9(6):e100525.
- Mulyani WRW, Sanjiwani MID, Sandra, Prabawa IPY, Lestari AAW, Wihandani DM, et al. Chaperone-Based Therapeutic Target Innovation: Heat Shock Protein 70 (HSP70) for Type 2 Diabetes Mellitus. Diabetes Metab Syndr Obes. 2020;13:559-568.
- Walton RG, Dungan CM, Long DE, Tuggle SC, Kosmac K, Peck BD, et al. Metformin blunts




Figure 10. Representative images of LC3 expression with immunofluorescence at 72-hour after Metformin was administered. Met 1 mM: group treated with 1 mM of Metformin; Met 2 mM: group treated with 2 mM of Metformin; Met 3 mM: group treated with 3 mM of Metformin.

- muscle hypertrophy in response to progressive resistance exercise training in older adults: A randomized, double-blind, placebo-controlled, multicenter trial: The MASTERS trial. Aging Cell. 2019;18(6):e13039.
- Fritzen AM, Madsen AB, Kleinert M, Treebak JT, Lundsgaard AM, Jensen TE, et al. Regulation of autophagy in human skeletal muscle: effects of exercise, exercise training and insulin stimulation. J Physiol. 2016 Feb 1;594(3):745-61
- Li DJ, Huang F, Lu WJ, Jiang GJ, Deng YP, Shen FM. Metformin promotes irisin release from murine skeletal muscle independently of AMPactivated protein kinase activation. *Acta Physiol* (Oxf). 2015;213(3):711-721.
- Jadhav KS, Dungan CM, Williamson DL. Metformin limits ceramide-induced senescence in C2C12 myoblasts. Mech Ageing Dev. 2013;134(11-12):548-559.
- Brown AE, Dibnah B, Fisher E, Newton JL, Walker M. Pharmacological activation of AMPK and glucose uptake in cultured human skeletal muscle cells from patients with ME/ CFS. Biosci Rep. 2018;38(3):BSR20180242.
- Sanchez AM, Csibi A, Raibon A, Cornille K, Gay S, Bernardi H, et al. AMPK promotes skeletal muscle autophagy through activation of forkhead FoxO3a and interaction with Ulk1. J Cell Biochem. 2012;113(2):695-710.
- Clavel S, Siffroi-Fernandez S, Coldefy AS, Boulukos K, Pisani DF, Dérijard B. Regulation of the intracellular localization of Foxo3a by stress-activated protein kinase signaling pathways in skeletal muscle cells. *Mol Cell Biol*. 2010;30(2):470-480.

- Thomson DM. The Role of AMPK in the Regulation of Skeletal Muscle Size, Hypertrophy, and Regeneration. *Int J Mol Sci.* 2018;19(10):3125.
- Mizushima N, Yoshimori T. How to interpret LC3 immunoblotting. Autophagy. 2007;3(6):542-545.
- Xia Q, Huang X, Huang J, Zheng Y, March ME, Li J, et al. The Role of Autophagy in Skeletal Muscle Diseases. Front Physiol. 2021;12:638983.
- Coughlan KA, Valentine RJ, Ruderman NB, Saha AK. AMPK activation: a therapeutic target for type 2 diabetes?. *Diabetes Metab Syndr Obes*. 2014;7:241-253.
- Sebastián D, Zorzano A. Self-Eating for Muscle Fitness: Autophagy in the Control of Energy Metabolism. Dev Cell. 2020;54(2):268-281.
- Kanamori H, Naruse G, Yoshida A, Minatoguchi S, Watanabe T, Kawaguchi T, et al. Metformin Enhances Autophagy and Provides Cardioprotection in δ-Sarcoglycan Deficiency-Induced Dilated Cardiomyopathy. Circ Heart Fail. 2019;12(4):e005418.
- Nwadike C, Williamson LE, Gallagher LE, Guan JL, Chan EYW. AMPK Inhibits ULK1-Dependent Autophagosome Formation and Lysosomal Acidification via Distinct Mechanisms. Mol Cell Biol. 2018;38(10):e00023-18.
- Milan G, Romanello V, Pescatore F, Armani A, Paik JH, Frasson L, et al. Regulation of autophagy and the ubiquitin-proteasome system by the FoxO transcriptional network during muscle atrophy. Nat Commun. 2015;6:6670.
- Hawley SA, Gadalla AE, Olsen GS, Hardie DG. The antidiabetic drug metformin activates

- the AMP-activated protein kinase cascade via an adenine nucleotide-independent mechanism. *Diabetes*. 2002;51(8):2420-2425.
- Musi N, Hirshman MF, Nygren J, Svanfeldt M, Bavenholm P, Rooyackers O, et al. Metformin increases AMP-activated protein kinase activity in skeletal muscle of subjects with type 2 diabetes. Diabetes. 2002;51(7):2074-81.
- Suwa M, Egashira T, Nakano H, Sasaki H, Kumagai S. Metformin increases the PGC-1alpha protein and oxidative enzyme activities possibly via AMPK phosphorylation in skeletal muscle in vivo. J Appl Physiol (1985). 2006;101(6):1685-1692.
- Kristensen JM, Treebak JT, Schjerling P, Goodyear L, Wojtaszewski JF. Two weeks of metformin treatment induces AMPKdependent enhancement of insulinstimulated glucose uptake in mouse soleus muscle. Am J Physiol Endocrinol Metab. 2014;306(10):E1099-E1109.
- Langone F, Cannata S, Fuoco C, Lettieri Barbato D, Testa S, Nardozza AP, et al. Metformin protects skeletal muscle from cardiotoxin induced degeneration. PLoS One. 2014;9(12):e114018.
- Queiroz EA, Puukila S, Eichler R, Sampaio SC, Forsyth HL, Lees SJ, et al. Metformin induces apoptosis and cell cycle arrest mediated by oxidative stress, AMPK and FOXO3a in MCF-7 breast cancer cells. PLoS One. 2014;9(5):e98207.
- Corona Velazquez AF, Jackson WT. So Many Roads: the Multifaceted Regulation of Autophagy Induction. Mol Cell Biol. 2018;38(21):e00303-18.
- Grotemeier A, Alers S, Pfisterer SG, Paasch F, Daubrawa M, Dieterle A, et al. AMPK-independent induction of autophagy by cytosolic Ca2+ increase. Cell Signal. 2010;22(6):914-25.
- Tomic T, Botton T, Cerezo M, Robert G, Luciano F, Puissant A, et al. Metformin inhibits melanoma development through autophagy and apoptosis mechanisms. Cell Death Dis. 2011;2(9):e199.



This work is licensed under a Creative Commons Attribution