



# Source details

## Bali Medical Journal

Scopus coverage years: from 2019 to Present

Publisher: Sanglah General Hospital

ISSN: 2089-1180 E-ISSN: 2302-2914

Subject area: Medicine: General Medicine

Source type: Journal

CiteScore 2022

0.2



SJR 2022

0.117



SNIP 2022

0.144



[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

### i Improved CiteScore methodology



CiteScore 2022 counts the citations received in 2019-2022 to articles, reviews, conference papers, book chapters and data papers published in 2019-2022, and divides this by the number of publications published in 2019-2022. [Learn more >](#)

CiteScore 2022 ▼

$$0.2 = \frac{180 \text{ Citations 2019 - 2022}}{889 \text{ Documents 2019 - 2022}}$$

Calculated on 05 May, 2023

CiteScoreTracker 2023 ⓘ

$$0.4 = \frac{428 \text{ Citations to date}}{1,080 \text{ Documents to date}}$$

Last updated on 05 September, 2023 • Updated monthly

## CiteScore rank 2022 ⓘ

Category	Rank	Percentile
Medicine		
General Medicine	#767/830	7th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)



# Bali Medical Journal

## COUNTRY

Indonesia



Universities and research institutions in Indonesia



Media Ranking in Indonesia

## SUBJECT AREA AND CATEGORY

Medicine  
Medicine  
(miscellaneous)

## PUBLISHER

Sanglah General Hospital

## H-INDEX

4

## PUBLICATION TYPE

Journals

## ISSN

20891 180, 23022914

## COVERAGE


2020-2022






## INFORMATION

- [Homepage](#)
- [How to publish in this journal](#)
- [srimaliawan@unud.ac.id](mailto:srimaliawan@unud.ac.id)




## SCOPE

Bali Med. J. is open access, peer-reviewed journal aiming to communicate high-quality research articles, reviews, and general articles in the field. Bali Med. J. publish articles that encompass all aspects of basic research/clinical studies related to medical sciences. The Journal aims to bridge and integrate the intellectual, methodological, and substantive medical scholarship diversity and encourage a vigorous dialogue between medical scholars and practitioners.

 Join the conversation about this journal



- Cites / Doc. (4 years)
- Cites / Doc. (3 years)
- Cites / Doc. (2 years)



Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com/journalsearch.php?q=21101024217&tip=sid&clean=0"
```


600

300



**SCImago Graphica**

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of April 2023

R **R. Sutomo** 4 months ago

Dear SCImago Team,

I am an Indonesian, and I know exactly that the home base of this journal is in Bali, Indonesia.

Would you kindly revise the information on the SCImago website? Thank you

reply

SCImago Team



**Melanie Ortiz** 4 months ago

Dear Sutomo, thank you for contacting us. We will proceed to analyze your request as soon as possible. Greetings from Spain and thank you for using the SCImago products,  
SCImago Team

A **Ahmad Al-Sarabbi** 1 year ago

Dear Scimago,

On this page, you mentioned that the country of origin is Italy, but the journal's website says Indonesia. And if I am not mistaken, Bali is indeed a very famous, beautiful island in Indonesia.

Which one is accurate? Does your website post misleading information?

Are other information regarding other journals can be trusted?

reply

SCImago Team



**Melanie Ortiz** 1 year ago

Dear Ahmad,

Thank you for contacting us. We will revise that information based on Scopus as soon as

**Bali**  
**Medical Journal**



INDONESIAN PHYSICIAN FORUM &  
INDONESIA COLLEGE OF SURGEONS,  
INDONESIA



<https://balimedicaljournal.org>

Open Access & Peer Reviewed Multidisciplinary  
Journal of Medical Sciences

[Advanced Search \(/index.php/bmj/search/search\)](/index.php/bmj/search/search)

**Dr Barrie Tan - SG ENT Expert**

Ad Dr Barrie Tan ENT

**Submit Your Paper**

Ad Dove Medical Press



**Distributor ACP seven**

Ad Alex58

**Leader in Shipping & Logistics**

Ad MSC

**Mana Mana**

Ad Mana Mana

**Fine Living with Ascott**

Ad DiscoverASR

**34 Data Centres Including Asia**

Ad OVHeloud

**JinPeng Mining**

Ad JinPeng Mining

**Opening Offer 25% Off**

Ad Ying'nFlo Admiralty

**Coohom Fresh Sale for Pro**

Ad Coohom 3D Space Design

Home (<https://balimedicaljournal.org/index.php/bmj/index>) > Editorial Board & Reviewer

**Editor-in-Chief**

**Prof. Dr. Sri Maliawan, SpBS** ([http://www.baliroyalhospital.co.id/halaman\\_staff.php?ditail=229](http://www.baliroyalhospital.co.id/halaman_staff.php?ditail=229))

(Scopus ID (<https://www.scopus.com/authid/detail.uri?authorId=15738530400>)), (Google scholar (<https://scholar.google.co.id/citations?user=qVJ57aYAAAAJ&hl=id>))

[srimaliawan@unud.ac.id](mailto:srimaliawan@unud.ac.id) / [maliawans@yahoo.com](mailto:maliawans@yahoo.com)

Department of Neuro Surgery, Udayana University

Sanglah General Hospital

Bali - Indonesia

**Associate Editor**

**Prof. Putra Manuaba, M.Phil** (<http://profpuma.weebly.com/>)

(Scopus ID) (<https://www.scopus.com/authid/detail.uri?authorId=8412278400>), (Google Scholar (<https://scholar.google.com/citations?user=jnmT14kPWNcC&hl=en>))

[putramanuaba@unud.ac.id](mailto:putramanuaba@unud.ac.id) / [putramanuaba28@yahoo.com](mailto:putramanuaba28@yahoo.com)

Biomedicine Postgraduate Program, Udayana University

Bali - Indonesia



**Prof. Ketut Suwiyoga, SpOG** (<http://www.scopus.com/results/authorNamesList.uri?sort=count-f&src=al&sid=01CAC4E9A2FB056A0A90221C03EC65FE.FZg2ODcJC9ArCe8WOZPv.NAME%28EQUALS%28Suwiyoga%29%29&st1=Suwiyoga&orcidId=&selectionPageSearch=anl&reselectAuthor=false&activeFlag=false&showDocument=false&resultsPerPage=20>)  
 (Scopus ID (<https://www.scopus.com/authid/detail.uri?authorId=54080784800>))  
 suwiyoga@unud.ac.id  
 Faculty of Medicine, Udayana University, Sanglah Hospital Denpasar, Bali-Indonesia

### Editorial Board for Regional America

**Ankit Sakhuja, M.B.B.S., F.A.C.P., F.A.S.N.** ([http://www.med.umich.edu/intmed/nephrology/STAFF/sakhuja\\_a1.htm](http://www.med.umich.edu/intmed/nephrology/STAFF/sakhuja_a1.htm))  
 (Scopus ID (<http://www.scopus.com/authid/detail.uri?authorId=16744977200>))  
 asakhuja@med.umich.edu  
 Nephrology and Hypertension Cleveland Clinic (United States)

### Editorial Board for Regional Australia

**Professor John Svigos, AM**  
**MBBS; DRCOG; CBioEth; FRCOG; FRANZCOG**  
 (<http://www.womenshealthspecialists.com.au/jsvigos.html>)  
 (Scopus ID) (<https://www.scopus.com/authid/detail.uri?authorId=6603773825>)  
 john@svigos.com.au (mailto:john@svigos.com.au)  
 Discipline of Obstetrics & Gynaecology  
 Faculty of Health & Medical Sciences  
 University of Adelaide, South Australia

**dr Deasy Ayuningtyas Tandio MPH-MBA** (<http://orcid.org/0000-0001-7847-2831>).  
 (OrcidID) (<https://orcid.org/0000-0001-7847-2831>)  
 deasytandio@yahoo.com  
 James Cook University Australia Master of Public Health Master Of Business Administration, Indonesia

### Editorial Board for Regional Europa

**Prof. Harald Hoekstra**  
 (Scopus ID) (<https://www.scopus.com/authid/detail.uri?authorId=36038081900>)  
 jsvigos@iprimus.com.au  
 Universitair Medisch Centrum Groningen, Division of Surgical Oncology, Groningen the Netherland

### Editorial Board for Regional Asia

**Prof Huang Qin** (<http://accu.cqu.edu.cn/web/eallprof/559.jhtml>)  
 (Scopus ID) (<https://www.scopus.com/authid/detail.uri?authorId=7409535321>)  
 qhuang@cqu.edu.cn  
 Chairman Dept. of Neurosurgery, Guangdong 999 Hospital Guangzhou China

**Assoc. Prof. Mohammad Amin Bahrami**  
 (Scopus ID) (<https://www.scopus.com/authid/detail.uri?authorId=55524082200>)  
 aminbahrami1359@gmail.com  
 Head of healthcare management department, Shahid Sadoughi University of Medical Sciences, Yazd,Iran

**Dr. Tanveer Beg, PhD**  
 (Scopus ID) (<https://www.scopus.com/authid/detail.uri?authorId=6505772852>)  
 tbmirza@jazanu.edu.sa  
 Assistant Professor, Department of Biology, Faculty of Science, Jazan University, Jazan, Saudi Arabia.

### Editorial Board Members

**Prof. Andi Asadul Islam**  
 (Scopus ID) (<https://www.scopus.com/authid/detail.uri?authorId=55504893500>), (Google Scholar) (<https://scholar.google.co.id/citations?user=vWs1RdMAAAJ&hl=id&oi=sra>)  
 undee@med.unhas.ac.id  
 Faculty of Medicine Hasanudin University, Makasar-Indonesia



**Prof. Dr. dr. Abdul Hafid Bajamal, Sp.BS**

(Scopus ID) (<https://www.scopus.com/authid/detail.uri?authorId=57192378862>)

[hfbajamal@gmail.com](mailto:hfbajamal@gmail.com)

Faculty of Medicine Airlangga University, Surabaya-Indonesia

**Dr. dr. I Wayan Sudarsa, Sp.B(K) Onk, FINACS, FICS.**

(Scopus ID (<https://www.scopus.com/authid/detail.uri?authorId=57205145862>)), (Google Scholar (<https://scholar.google.co.id/citations?hl=id&user=SdInHKwAAAAJ>))), (Researchg

**dr. I.B. Amertha P. Manuaba, SKed, MBIomed.** (<https://scholar.google.co.id/citations?user=KzCQgA0AAAAJ&hl=en>)

(Scopus ID) (<https://www.scopus.com/authid/detail.uri?authorId=57195520004>), (Google Scholar) (<https://scholar.google.co.id/citations?user=KzCQgA0AAAAJ&hl=en>), (ORCID) (h

[AmerthaManuaba@gmail.com](mailto:AmerthaManuaba@gmail.com) / [Amertha\\_Manuaba@unud.ac.id](mailto:Amertha_Manuaba@unud.ac.id)

Faculty of Medicine, Universitas Udayana, Indonesia

**Editorial inquiries to be addressed to:**

email 1: [editorbalimedicaljournal@gmail.com](mailto:editorbalimedicaljournal@gmail.com) (<mailto:editor@balimedicaljournal.org>)

email 2: [editor@balimedicaljournal.org](mailto:editor@balimedicaljournal.org) (<mailto:editor@balimedicaljournal.org>)

-----



WEB OF SCIENCE™ ([https://mjl.clarivate.com/search-results?issn=2089-1180&hide\\_exact\\_match\\_fl=true&utm\\_source=mjl&utm\\_medium=share-by-link&utm\\_campaign=search-results-share-this-journal](https://mjl.clarivate.com/search-results?issn=2089-1180&hide_exact_match_fl=true&utm_source=mjl&utm_medium=share-by-link&utm_campaign=search-results-share-this-journal))



ELSEVIER  
Scopus

(<https://www.scopus.com/sourceid/21101024217>)

DOAJ DIRECTORY OF  
OPEN ACCESS  
JOURNALS

(<https://doaj.org/toC/2302-2914>)



(<https://sinta3.kemdikbud.go.id/journals/profile/2513>)

Full Indexing List (<https://balimedicaljournal.org/index.php/bmj/pages/view/indexing>)

In Press (<https://balimedicaljournal.org/index.php/bmj/issue/view/30>)

Submit An Article (<https://balimedicaljournal.org/index.php/bmj/login>)

Scopus Citedness (<https://balimedicaljournal.org/index.php/bmj/pages/view/scopus>)





**Claim Your Free Trial**

Ad American Journal Experts

**Evolutionary Human Sciences - Read the latest article** <sup>①</sup>

Ad cambridge.org

**Publish Open Access**

Ad Taylor & Francis

**Door and Window**

Ad Topp

**Generously simple and free**

Ad New Relic

**No Setup Fee**

Ad OVHcloud

**Online Universities**

Ad Prorateer

**Be a VTuber with nizima LIVE**

Ad Live2D Inc.

**Web & File Instant Translation**

Ad Sider

**Download disini**

Ad zbook

**research paper publication**

Ad IJSDR Research Journal

**Peer review, Referred Journal**

Ad IJCRT Research Journal



Home (<https://www.balimedicaljournal.org/index.php/bmj/index>) > Archives (<https://www.balimedicaljournal.org/index.php/bmj/issue/archive>) > Vol. 11 No. 2 (2022):  
(Available online : 1 August 2022)

**Vol. 11 No. 2 (2022): (Available online : 1 August 2022)**

ORIGINAL ARTICLE

**Stress associated with hypertension in middle-age and elderly in Binong, Tangerang**  
(<https://www.balimedicaljournal.org/index.php/bmj/article/view/3356>)

Martina Pakpahan, Ni Gusti Ayu Eka, Maria Veronika Ayu Florensa

Online First: Jul 4, 2022 |

 Abstract

 pdf (<https://www.balimedicaljournal.org/index.php/bmj/article/view/3356/2085>)

Online First: Aug 8, 2022 |

 Abstract


 pdf (<https://www.balimedicaljournal.org/index.php/bmj/article/view/3407/2137>)

REVIEW

**The The outcome of sternum healing among diabetic patients undergoing open heart surgery: a literature review**  
(<https://www.balimedicaljournal.org/index.php/bmj/article/view/3513>)

Taufik Nur Yahya, Yan Efrata Sembiring, Soebagijo Adi Soelistijo

Online First: Aug 15, 2022 |

 Abstract


 pdf (<https://www.balimedicaljournal.org/index.php/bmj/article/view/3513/2153>)

REVIEW

**Current surgical interventions of Charcot neuroarthropathy of the foot and ankle: a systematic review**  
(<https://www.balimedicaljournal.org/index.php/bmj/article/view/3619>)

I Wayan Subawa, I Gusti Ngurah Bagus Andhika Pramana, Adiet Wahyu Kristian, Putu Bihan Surya Kinanta

Online First: Aug 5, 2022 |

 Abstract

 pdf (<https://www.balimedicaljournal.org/index.php/bmj/article/view/3619/2123>)



CASE REPORT

**Complications of repeated percutaneous transhepatic biliary drainage (PTBD) for palliation of obstructive jaundice in cholangiocarcinoma patient** (<https://www.balimedicaljournal.org/index.php/bmj/article/view/3465>)

Radhitya Farizky Deta Juniawan, Ummi Maimunah, Titong Sugihartono, Ulfa Kholili, Budi Widodo, Husin Thamrin, Muhammad Miftahussurur, Amie Vidyani

Online First: Aug 16, 2022 |

 Abstract

 pdf (<https://www.balimedicaljournal.org/index.php/bmj/article/view/3465/2155>)

CASE REPORT

**A fatal case of Norwegian scabies in a patient with diabetes mellitus**  
(<https://www.balimedicaljournal.org/index.php/bmj/article/view/3500>)

## The outcome of sternum healing among diabetic patients undergoing open heart surgery: a literature review



Taufik Nur Yahya<sup>1\*</sup>, Yan Efrata Sembiring<sup>2</sup>, Soebagijo Adi Soelistijo<sup>3</sup>

### ABSTRACT

**Background:** As part of the surgery, sternotomy is a common technique to expose the underlying organs, allowing surgeons to explore the targeted organ. Despite the benefits offered, it has several considerable complications; one of them is wound healing defect. The impairment of sternal wound healing after sternotomy could be superficial and/or deep. This condition is influenced by many factors, including diabetes mellitus, obesity, and other comorbidities. These anomalies will affect the normal healing process of the bone and other connective tissue, particularly after a major invasive event, such as in sternotomy. This study aimed to review the outcome of sternum healing among diabetics that underwent heart surgery with a sternotomy approach.

**Methods:** Works of literature reviewed in this study were obtained from Pubmed and Google Scholar databases starting from 1992 until April 2022. The keywords used were 'sternal wound healing', 'cardiac surgery', and 'diabetes mellitus'. An advanced search based on the exact phrases was conducted on Google Scholar. Gathered kinds of literature were then selected based on relevancy.

**Results:** The majority of articles reviewed were observational, and most of them had CABG as a part of open heart surgery, followed by valvular and aortic surgeries and others. The incidence of impaired sternal healing and other complications (superficial and/or deep) was more commonly seen among patients with diabetes. Most studies also reported a significant correlation between DM and the incidence of impaired sternal healing, suggesting that DM was a significant predictor of it.

**Conclusion:** Diabetes mellitus is a medical condition that must be taken into account among the candidates for open heart surgery, particularly if it is poorly controlled. The Hyperglycaemic state experienced by the patients will lead to multiorgan damage and immunological dysfunction that could affect the healing process of the sternum.

**Keywords:** cardiac surgery, diabetes mellitus, sternal wound healing.

**Cite This Article:** Yahya, T.N., Sembiring, Y.E., Soelistijo, S.A. 2022. The outcome of sternum healing among diabetic patients undergoing open heart surgery: a literature review. *Bali Medical Journal* 11(2): 818-826. DOI: 10.15562/bmj.v11i2.3513

<sup>1</sup>Resident of Thoracic, Cardiac, and Vascular Surgery Department, Universitas Airlangga, Dr. Soetomo Academic General Hospital, Surabaya, East Java, Indonesia;

<sup>2</sup>Senior Attending of Thoracic, Cardiac, and Vascular Surgery Department, Universitas Airlangga, Dr. Soetomo Academic General Hospital, Surabaya, East Java, Indonesia;

<sup>3</sup>Senior Attending of Internal Medicine Department, Universitas Airlangga, Dr. Soetomo Academic General Hospital, Surabaya, East Java, Indonesia;

\*Corresponding author:

Taufik Nur Yahya;  
Resident of Thoracic, Cardiac, and Vascular Surgery Department, Universitas Airlangga, Dr. Soetomo Academic General Hospital, Surabaya, East Java, Indonesia;  
taufiknuryahya.md@gmail.com

Received: 2022-06-12

Accepted: 2022-08-01

Published: 2022-08-15

### INTRODUCTION

Diabetes mellitus (DM) is a chronic metabolic disease that is characterized by the elevation of blood glucose levels, also referred to as hyperglycemia. World Health Organization (WHO) stated in 2014, approximately 422 million people around the globe had diabetes mellitus; and in 2012, nearly 1.5 million people died due to this debilitating disease.<sup>1</sup> According to American Diabetes Association (ADA), diabetes mellitus can be classified into several groups based on the etiology, which are as follows: type I, type II, and gestational.<sup>2</sup> There is a significant association between diabetes with cardiovascular disease manifestation.<sup>3</sup> Among the other comorbidities associated with diabetes, atherosclerosis is a

significant cause of mortality in humans, particularly if it occurs within the coronary vessels, because this could result in myocardial infarction – the leading cause of death worldwide.<sup>4,5</sup> Irreversibility of atherosclerosis and the need for adequate, continuous perfusion to the myocardial tissue made several attempts created to get rid of this problem. One of them is coronary artery bypass graft (CABG), by which the occluded vessel is bypassed by the new vessel in order to improve myocardial perfusion and viability, hence, reducing the incidence of myocardial infarction in the future.<sup>6</sup> CABG requires sternotomy – a procedure to cut and 'damage' the sternum, which must be done to allow the surgeons to explore the underlying structures. A sternotomy

is a standard approach performed in almost every surgical procedure on the heart and mediastinum.<sup>7</sup> Sternotomy is also indicated in heart transplantation, valvular reconstruction, and pulmonary surgeries.<sup>8,9</sup> Immunological dysregulation and microangiopathy are the other comorbidities associated with the hyperglycemic state, and these conditions can impair cellular regeneration and vascular perfusion to the periphery sites, such as in wound healing after surgery. It has long been known that adequate oxygenation and nutrient supply to the site of injury are needed to achieve optimal healing and tissue renewal; otherwise, it could be halted, and several disorders could ensue, for example, in the case of malunion, osteonecrosis, and wound

dehiscence.<sup>10</sup> Considering the explanations above, this review aimed to describe the outcome of sternal healing among diabetic patients undergoing cardiac/other opened chest surgery.

## METHODS

The pieces of literature reviewed in this study were obtained from PubMed and Google Scholar databases, with the publication years starting from 1992 – 2022. The advanced search was applied in Google Scholar with; all of the words: 'sternal healing after cardiac surgery among diabetic patients'; exact phrase: 'sternal healing'; and at least one word: 'diabetes mellitus.' These words were meant to be included anywhere in the article. The advanced search was not done on PubMed, with the keywords used: 'sternal wound healing', 'cardiac surgery', and 'diabetes mellitus.' This strategy resulted in 44 and 252 kinds of literature shown from PubMed and Google Scholar, respectively. The next step was a selection based on the relevancy, accessibility, and the languages of the articles, where the articles reviewed were only in English. Furthermore, related references in the articles were also reviewed in order to broaden the search. This strategy resulted in 24 eligible articles from the past 30 years.

## RESULTS

The data gathered from the articles consisted of the author (year of publication), study design, age and gender of the study population (for the age we used: mean  $\pm$  SD or median [IQR]), indication for the cardiac/open chest surgery, and the outcome (sternal healing).

Of the 24 articles cited to be reviewed, most of the studies ( $n = 19$ ) were observational, and the rest were literature reviews ( $n = 3$ ) and systematic reviews/meta-analyses ( $n = 2$ ). Almost all papers showed that CABG was the most common cardiac surgery performed, followed by valve reconstruction, aortic/other vessel surgeries, emergency surgeries, heart transplantation, and others (unmentioned); one study by Zahiri *et al.* that investigated the sternum reconstruction using the autologous flap.<sup>28</sup>

## DISCUSSION

### Comorbidities Associated with Diabetes Mellitus

In general, the complications exerted by a person with chronic diabetes mellitus are classified into microvascular and macrovascular involvements. The diabetic microvascular pathophysiology is associated with many chronic inflammatory processes triggered by some cytokines and growth factors.<sup>35-43</sup> Increased levels of glucose will further increase the production of advanced glycation end product (AGE); this substance has several deleterious effects, one of which is to directly cross-link the collagenous portion beneath the vascular wall. This non-enzymatic reaction, if it occurs for a long period, will result in microangiopathy—characterized by the thickening and infiltration of immunological cells of the small blood vessels that can be found in the retina, skin, and glomerulus. AGE will also bind to its receptor (RAGE), expressed mainly on osteoclasts and osteocytes; the activation of RAGE will lead to osteoclastogenesis. Thickened and occluded vascular lumen will lead to poor tissue and connective tissue perfusion, including the tissue of the skin, bone, and peripheral nerves.<sup>44</sup>

A hyperglycemic state will also promote the formation of intracellular fructose and sorbitol via the sorbitol (polyol) pathway, particularly in the non-insulin-dependent tissues (endothelial cells, neurons and peripheral nerve cells, pancreatic  $\beta$ -cells, and kidney). Excess glucose will be first converted to sorbitol by an enzyme called aldose reductase. Next, the sorbitol is oxidized by sorbitol dehydrogenase into fructose as the final product. Accumulation of these substances inside the cells will result in an osmotic pressure gradient that eventually damages the cell membrane; moreover, this enzymatic reaction utilizes nicotinamide adenine dinucleotide phosphate (NADPH) that is also needed to make GSH (glutathione), a potent antioxidant; hence, increased polyol reaction as seen in diabetes will promote the excess formation of reactive oxygen species (ROS) while decreasing the rate of its degeneration, making the cells previously mentioned injured and damaged.<sup>45</sup>

Immune disruption is also seen in diabetic patients. Several experimental and *in vitro* studies have shown that the excess blood glucose level will impair cytokine formation, defect in leukocytes recruitment and activation at the injured site, and wound healing along with tissue regeneration. Diabetic patients also reflect poor immunological performance in terms of eradicating infection, which mainly involves the activation and effective signaling of macrophages and natural killer cells. This also could explain the common incidence of both superficial and deep wound infections and poorer prognosis of diabetics in facing severe infectious agents compared with normoglycemic individuals.<sup>11,14,46</sup>

### Sternal Healing

The bone healing process consists of four stages: inflammation, soft callus formation, complex callus formation, and bone remodeling.<sup>47</sup> Bony compartments can achieve optimal healing in several ways after getting injured or in the case of fracture, namely intramembranous and endochondral ossification. Intramembranous ossification is one of the healing processes that mainly requires the direct differentiation of mesenchymal precursor cells (MPC) to osteoblasts without going through the cartilaginous phase – which is not the case of endochondral ossification, which undergoes cartilaginous phase (the proliferation of MPCs to chondrocyte first). Intramembranous ossification occurs notably in flat bones (e.g., sternum and cranial bone). Meanwhile, endochondral ossification usually takes place at the long (diaphyseal) bones (femur and tibia).<sup>48,49</sup>

Several factors that influence the normal bone healing process are proper oxygenation, adequate supply of various nutrients consisting of glucose & other mineral components, and activity of the nearby precursor cells that secrete growth factors needed to assemble the new tissue. These events are orchestrated well only if good perfusion exists, which is not the case with chronic diabetes.<sup>49,50</sup>

Besides biological factors, physical/mechanical factors also influence the outcome of a bone healing process. The sternum is anatomically located at the


anterior, midline chest and is also a crucial part of the rib cage. Furthermore, it has an important barrier function towards the underlying vital organs that comprise the heart, great vessels, and pulmonary tissues. The rib cage itself is slightly mobile since it moves on several occasions: breathing, coughing, and when one individual lift a heavy object. This makes the sternum tends to be strained and stretched most of the time, leading to lateral distraction.<sup>49,51</sup>

**Diabetes Mellitus and Sternal Wound Impairment**

Normally, circulating insulin exerts its anabolic effect on the bone by binding with INSR/IRS-1 expressed on osteoblasts. After binding, the subsequent event will be the activation of Akt and PI3K with appropriate suppression of the FOXO1 gene. Decreased level of insulin, either because of the lack of production or ineffective utilization, will lead to the impairment of normal bone growth that is characterized by abnormal mineralization and matrix deposition along with the abnormality of bone architecture. These abnormalities are prominently seen among T1DM patients since this population totally lacks insulin production.<sup>42,52</sup>

After an injury, the bone will develop an intramedullary hematoma, by which this event is purposed to effectively recruit inflammatory cells and adjacent fibroblasts. Bone mineralization, along with inorganic matrix deposition, occurs with the help of these cells and requires high content of collagen, notably type I. AGE, a protein glycosylated with the aldose sugar, directly cross-linked the exposed collagen and halted its utilization by fibroblasts and macrophages, decreasing the collagen's availability within the injured bony tissue.<sup>52,53</sup>

The abnormally high level of proinflammatory cytokines (IL-6, TNF $\alpha$ ), AGE, along with the accumulation of intracellular ROS, will disrupt the differentiation of mesenchymal stem cells (MSC) and survivability of osteoblasts. High levels of cytokines and alterations in growth hormone secretion also will shift the differentiation from osteoblastic to the osteoclastic direction, worsening the performance of the bone healing process even further. Osteoclastogenesis



**Figure 1.** Diabetes affects cellular components of the sternal bone healing.

occurs due to the activation of RAGE, and as a consequence, the production of Receptor Activator of Nuclear Factor  $\kappa$ -B (RANK) by nearby osteocytes and osteoblasts is increased. RANK binds to its ligand (RANKL), which is located along the osteoclast cellular membrane. RAGE also down-regulates the expression of osteoprotegerin (OPG) to enhance osteoclastogenesis and bone resorption.<sup>52,54,55</sup>

**Previous Studies Related to Sternal Healing Outcome**

Out of 24 articles reviewed, as shown in Table 1, most of the works of literature found that diabetes, regardless of the type, was significantly associated with the incidence of sternal wound complications that comprise sternal dehiscence and infection (superficial and deep), and malunion.<sup>11,13-15,18-23,25,28-31,34</sup> This corresponds with the previously postulated theory regarding the multiorgan and immunological dysfunction associated with a hyperglycemic state. Some studies, however, stated that the type of diabetes was more significant in T1DM rather than T2DM. This suggests that lacking insulin condition was more severe among

T1 diabetes patients compared to T2 diabetics.<sup>16,21,42,52</sup> Other contradictory results were reported, where significance between the incidence of sternal wound complication was not associated with diabetes mellitus.<sup>10,17,24,26,27,32,33</sup> Some studies mentioned that besides diabetes, the outcome of sternal healing among patients receiving CABG procedure also depended on the grafting technique, particularly if internal mammary arteries, which is the first choice of conduit, were used as the vessel source since the major vascularization of the sternum was derived from these arteries.<sup>20,24,34,55</sup>

**Evaluation for Sternal Healing**

Several parameters were set to evaluate the outcome of sternal healing, comprised of sternal wound infection (superficial and/or deep), sternal dehiscence, and sternal instability. According to National Nosocomial Infections Surveillance (NNIS), superficial infection is defined as an infection that involves the skin (epidermis and dermis) and the subcutaneous layer that is clinically assessed by at least 1 (one) finding as follows: 1) purulent discharge, 2) positive culture result from the discharge, 3) erythema, pain/tenderness and warm

sensation that surrounds the incision site, and 4) surgeon's decision.<sup>23,39,31</sup> Deep sternal infection based, on the Centers for Disease Control and Prevention (CDC), is defined where the subcutaneous tissue and the space underneath it are involved, with at least one of the following criteria: 1) positive culture/organism isolation from mediastinal tissue, 2) clinical signs of mediastinitis, and 3) systemic manifestation such as fever (body temperature of  $\geq 38^{\circ}\text{C}$ ), sternal/chest pain, and/or sternal instability. Almost all studies used these criteria to diagnose sternal wound infection after open cardiac surgery.<sup>11-14,16-22,25-34</sup> CT scan was used by Shin *et al.* to investigate the sternal healing that was proved by the union of the sternal body and manubrium.<sup>15</sup> Culture and pathogen identification from the mediastinal and the incisional discharge was conducted by Heilmann *et al.*, Zacharias *et al.*, and Fakhri *et al.*, with most of the pathogens isolated, were methicillin-susceptible *S.aureus* (MSSA), coagulase-negative staphylococci, gram-negative aerobes, *Corynebacterium sp.*, and *Candida sp.*<sup>16,23,31</sup>

## CONCLUSION

Open heart surgery is a major invasive procedure that has several complications that are broadly categorized into local and systemic. The most common local involvement is impaired sternal healing. It can be caused by many factors. One of them is diabetes mellitus with chronic hyperglycemia. The literature review results showed that chronic hyperglycemia could lead to several organ damages and immunological dysregulation, resulting in defective bone repair, perfusion, and ineffective remodeling.

## CONFLICT OF INTEREST

The authors declare that there is no conflict of interest regarding the publication of this article.

## FUNDING

The authors are responsible for all research funding without obtaining financial support.

## ETHICS APPROVAL

Not applicable.

## AUTHOR CONTRIBUTION

Taufik Nur Yahya, Yan Efrata Sembiring, and Soebagijo Adi Soelistijo conceptualized and designed the study. All authors analyzed and interpreted the study results and revised the manuscript.

## REFERENCES

- Silva JAD, Souza ECF, Echazú Böschemeier AG, Costa CCMD, Bezerra HS, Feitosa EELC. Diagnosis of diabetes mellitus and living with a chronic condition: participatory study. *BMC Public Health*. 2018;18(1):699. Published 2018 Jun 5. doi:10.1186/s12889-018-5637-9.
- Kharroubi AT, Darwish HM. Diabetes mellitus: The epidemic of the century. *World J Diabetes*. 2015;6(6):850-867. doi:10.4239/wjcd.v6.i6.850.
- Marchelia LZ, Purwati P, Wironegoro R. High blood glucose level increase cardiovascular disease risk in type 2 diabetes mellitus. *Folia Medica Indonesiana*. 2016;52(2):127-30.
- Artha IMJR, Bhargah A, Dharmawan NK, Pande UW, Triyana KA, Mahariski PA, Yuwono J, Bhargah V, Prabawa IPY, Manuaba IBAP, Rina IK. High level of individual lipid profile and lipid ratio as a predictive marker of poor glycemic control in type-2 diabetes mellitus. *Vasc Health Risk Manag*. 2019 Jun 5;15:149-157. doi: 10.2147/VHRM.S209830.
- Ioacara S, Popescu AC, Tenenbaum J, et al. Acute Myocardial Infarction Mortality Rates and Trends in Romania between 1994 and 2017. *Int J Environ Res Public Health*. 2019;17(1):285. Published 2019 Dec 31. doi:10.3390/ijerph17010285.
- Bachar BJ, Manna B. Coronary Artery Bypass Graft [Internet]. *StatPearls*. 2021. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK507836/>
- Amal I, Soebroto H, Puruhito. Comparison of bone wax and chitosan usage on post-sternotomy bone healing. *Asian Cardiovasc Thorac Ann*. 2021;29(3):203-207. doi:10.1177/0218492320984097.
- Pezzella AT. Global aspects of cardiothoracic surgery with focus on developing countries. *Asian Cardiovasc Thorac Ann*. 2010;18(3):299-310. doi:10.1177/0218492310370060.
- Boudoulas KD, Ravi Y, Garcia D, et al. Type of Valvular Heart Disease Requiring Surgery in the 21st Century: Mortality and Length-of-Stay Related to Surgery. *Open Cardiovasc Med J*. 2013;7:104-109. Published 2013 Sep 4. doi:10.2174/1874192420130902001.
- Cope G. The effects of smoking on wound healing. *Wounds UK*. 2014;10(2):10-8.
- Lemaigen A, Birgand G, Ghodhbane W, et al. Sternal wound infection after cardiac surgery: incidence and risk factors according to clinical presentation. *Clin Microbiol Infect*. 2015;21(7):674.e11-674.e6.74E18. doi:10.1016/j.cmi.2015.03.025.

- Sofer D, Gurevitch J, Shapira I, et al. Sternal wound infections in patients after coronary artery bypass grafting using bilateral skeletonized internal mammary arteries. *Ann Surg*. 1999;229(4):585-590. doi:10.1097/00000658-199904000-00020.
- Oswald I, Boening A, Pons-Kuehnemann J, Grieshaber P. Wound Infection after CABG Using Internal Mammary Artery Grafts: A Meta-Analysis. *Thorac Cardiovasc Surg*. 2021;69(7):639-648. doi:10.1055/s-0040-1713662.
- Balachandran S, Lee A, Denehy L, et al. Risk Factors for Sternal Complications After Cardiac Operations: A Systematic Review. *Ann Thorac Surg*. 2016;102(6):2109-2117. doi:10.1016/j.athoracsur.2016.05.047.
- Shin YC, Kim SH, Kim DJ, et al. Sternal healing after coronary artery bypass grafting using bilateral internal thoracic arteries: assessment by computed tomography scan. *Korean J Thorac Cardiovasc Surg*. 2015;48(1):33-39. doi:10.5090/kjctcs.2015.48.1.33.
- Heilmann C, Stahl R, Schneider C, et al. Wound complications after median sternotomy: a single-centre study. *Interact Cardiovasc Thorac Surg*. 2013;16(5):643-648. doi:10.1093/icvts/ivs554.
- Gansera B, Delalic A, Eszlari E, Eichinger W. 14-Year Results of Bilateral versus Single Internal Thoracic Artery Grafts for Left-Sided Myocardial Revascularization in Young Diabetic Patients. *Thorac Cardiovasc Surg*. 2017;65(4):272-277. doi:10.1055/s-0036-1593864.
- Zalewska-Adamic M, Bachorzewska-Gajewska H, Malyszko J, et al. Impact of diabetes on mortality and complications after coronary artery by-pass graft operation in patients with left main coronary artery disease. *Adv Med Sci*. 2014;59(2):250-255. doi:10.1016/j.advm.2014.02.006.
- Nakano J, Okabayashi H, Hanyu M, et al. Risk factors for wound infection after off-pump coronary artery bypass grafting: should bilateral internal thoracic arteries be harvested in patients with diabetes? *J Thorac Cardiovasc Surg*. 2008;135(3):540-545. doi:10.1016/j.jtcvs.2007.11.008.
- Peterson MD, Borger MA, Rao V, Peniston CM, Feindel CM. Skeletonization of bilateral internal thoracic artery grafts lowers the risk of sternal infection in patients with diabetes. *J Thorac Cardiovasc Surg*. 2003;126(5):1314-1319. doi:10.1016/s0022-5223(03)00808-0.
- Zacharias A, Habib RH. Factors predisposing to median sternotomy complications. Deep vs superficial infection. *Chest*. 1996;110(5):1173-1178. doi:10.1378/chest.110.5.1173.
- Pevni D, Uretzky G, Mohr A, et al. Routine use of bilateral skeletonized internal thoracic artery grafting: long-term results. *Circulation*. 2008;118(7):705-712. doi:10.1161/CIRCULATIONAHA.107.756676.
- Zuckermann A, Barten MJ. Surgical wound complications after heart transplantation. *Transpl Int*. 2011;24(7):627-636. doi:10.1111/j.1432-2277.2011.01247.x.
- Lazar HL. The risk of mediastinitis and deep sternal wound infections with single

- and bilateral, pedicled and skeletonized internal thoracic arteries. *Ann Cardiothorac Surg.* 2018;7(5):663-672. doi:10.21037/acs.2018.06.11.
25. Fu RH, Weinstein AL, Chang MM, Argenziano M, Ascherman JA, Rohde CH. Risk factors of infected sternal wounds versus sterile wound dehiscence. *J Surg Res.* 2016;200(1):400-407. doi:10.1016/j.jss.2015.07.045.
  26. Gurevitch J, Paz Y, Shapira I, et al. Routine use of bilateral skeletonized internal mammary arteries for myocardial revascularization. *Ann Thorac Surg.* 1999;68(2):406-412. doi:10.1016/s0003-4975(99)00460-9.
  27. Shaheen Y, Kasab I, Galal M. Comparative Study Between the effect of Skeletonized and Pedicled Internal Thoracic Artery on sternum healing in Patient Undergoing CABG. *Benha Med J.* 2020;0(0):0-0.
  28. Zahiri HR, Lumpkins K, Kelishadi SS, et al. Significant predictors of complications after sternal wound reconstruction: a 21-year experience. *Ann Plast Surg.* 2012;69(4):439-441. doi:10.1097/SAP.0b013e318231d1ef.
  29. Lenz K, Brandt M, Fraund-Cremer S, Cremer J. Coronary artery bypass surgery in diabetic patients - risk factors for sternal wound infections. *GMS Interdiscip Plast Reconstr Surg DGPW.* 2016;5:Doc18. Published 2016 Jul 28. doi:10.3205/ipsr000097.
  30. Savage EB, Grab JD, O'Brien SM, et al. Use of both internal thoracic arteries in diabetic patients increases deep sternal wound infection. *Ann Thorac Surg.* 2007;83(3):1002-1006. doi:10.1016/j.athoracsur.2006.09.094.
  31. Fakhri MG, Sharma M, Khatib R, et al. Increase in the rate of sternal surgical site infection after coronary artery bypass graft: a marker of higher severity of illness. *Infect Control Hosp Epidemiol.* 2007;28(6):655-660. doi:10.1086/518347.
  32. Sakamoto H, Fukuda I, Oosaka M, Nakata H. Risk factors and treatment of deep sternal wound infection after cardiac operation. *Ann Thorac Cardiovasc Surg.* 2003;9(4):226-232.
  33. Kieser TM, Rose MS, Aluthman U, Montgomery M, Louie T, Belenkie I. Toward zero: deep sternal wound infection after 1001 consecutive coronary artery bypass procedures using arterial grafts: implications for diabetic patients. *J Thorac Cardiovasc Surg.* 2014;148(5):1887-1895. doi:10.1016/j.jtcvs.2014.02.022.
  34. Dai C, Lu Z, Zhu H, Xue S, Lian F. Bilateral internal mammary artery grafting and risk of sternal wound infection: evidence from observational studies. *Ann Thorac Surg.* 2013;95(6):1938-1945. doi:10.1016/j.athoracsur.2012.12.038.
  35. Liu M, Weiss MA, Arunagiri A, et al. Biosynthesis, structure, and folding of the insulin precursor protein. *Diabetes Obes Metab.* 2018;20 Suppl 2(Suppl 2):28-50. doi:10.1111/dom.13378.
  36. Petersen MC, Shulman GI. Mechanisms of Insulin Action and Insulin Resistance. *Physiol Rev.* 2018;98(4):2133-2223. doi:10.1152/physrev.00063.2017.
  37. Yee LD, Mortimer JE, Natarajan R, Dietze EC, Seewaldt VL. Metabolic Health, Insulin, and Breast Cancer: Why Oncologists Should Care About Insulin. *Front Endocrinol (Lausanne).* 2020;11:58. Published 2020 Feb 20. doi:10.3389/fendo.2020.00058.
  38. DiMeglio LA, Evans-Molina C, Oram RA. Type 1 diabetes. *Lancet.* 2018;391(10138):2449-2462. doi:10.1016/S0140-6736(18)31320-5.
  39. Lucier J, Weinstock RS. Diabetes Mellitus Type 1. In: StatPearls. Treasure Island (FL): StatPearls Publishing; May 11, 2022.
  40. Bowden S. Partial Remission (honeymoon phase) in Type 1 Diabetes Mellitus. 2017. 10.2174/9781681089348121070001.
  41. Pratley RE. The early treatment of type 2 diabetes. *Am J Med.* 2013;126(9 Suppl 1):S2-S9. doi:10.1016/j.amjmed.2013.06.007.
  42. Saini V. Molecular mechanisms of insulin resistance in type 2 diabetes mellitus. *World J Diabetes.* 2010;1(3):68-75. doi:10.4239/wjdv.1i3.68.
  43. Romadhon PZ, Sutjahjo A, Novida H, Soelistijo SA, Wibisono S, Prajitno JH et al. HBA1C and plasma transforming growth factor-beta 1 in type-2 diabetes mellitus patients. *New Armenian Medical Journal.* 2019;13(1):69-73.
  44. Baltzis D, Eleftheriadou I, Veves A. Pathogenesis and treatment of impaired wound healing in diabetes mellitus: new insights. *Adv Ther.* 2014;31(8):817-836. doi:10.1007/s12325-014-0140-x.
  45. Moemen, L.A., Abdel Hamid, M.A., Wahab, S.A. et al. Role of advanced glycation end products and sorbitol dehydrogenase in the pathogenesis of diabetic retinopathy. *Bull Natl Res Cent* 44. 2020;58. <https://doi.org/10.1186/s42269-020-00304-0>.
  46. Berbudi A, Rahmadika N, Tjahjadi AI, Ruslami R. Type 2 Diabetes and its Impact on the Immune System. *Curr Diabetes Rev.* 2020;16(5):442-449. doi:10.2174/1573399815666191024085838.
  47. Wibowo H, Widiyanti P. The Effect of Diclofenac Sodium on Callus Formation in White Male Rat (*Rattus norvegicus*) Cruris Fracture Healing. *Folia Medica Indonesiana (FMI).* 2022;58(2):108-12.
  48. Oryan A, Monazzah S, Bigham-Sadegh A. Bone injury and fracture healing biology. *Biomed Environ Sci.* 2015;28(1):57-71. doi:10.3967/bes2015.006.
  49. Bigham-Sadegh A, Oryan A. Basic concepts regarding fracture healing and the current options and future directions in managing bone fractures. *Int Wound J.* 2015;12(3):238-247. doi:10.1111/iwj.12231.
  50. Kemmler J, Bindl R, McCook O, et al. Exposure to 100% Oxygen Abolishes the Impairment of Fracture Healing after Thoracic Trauma. *PLoS One.* 2015;10(7):e0131194. Published 2015 Jul 6. doi:10.1371/journal.pone.0131194.
  51. Parker R, Adams JL, Ogola G, et al. Current activity guidelines for CABG patients are too restrictive: comparison of the forces exerted on the median sternotomy during a cough vs. lifting activities combined with valsalva maneuver. *Thorac Cardiovasc Surg.* 2008;56(4):190-194. doi:10.1055/s-2008-1038470.
  52. Jiao H, Xiao E, Graves DT. Diabetes and Its Effect on Bone and Fracture Healing. *Curr Osteoporos Rep.* 2015;13(5):327-335. doi:10.1007/s11914-015-0286-8.
  53. Yao D, Brownlee M. Hyperglycemia-induced reactive oxygen species increase expression of the receptor for advanced glycation end products (RAGE) and RAGE ligands. *Diabetes.* 2010;59(1):249-255. doi:10.2337/db09-0801.
  54. Marin C, Luyten FP, Van der Schueren B, Kerckhofs G, Vandamme K. The Impact of Type 2 Diabetes on Bone Fracture Healing. *Front Endocrinol (Lausanne).* 2018;9:6. Published 2018 Jan 24. doi:10.3389/fendo.2018.00006.
  55. Roysse, Alistair & Roysse, Colin & Boggett, Stuart & Clarke-Errey, Sandy & Pawanis, Zulfayandi. Why and how to achieve total arterial revascularisation in coronary surgery. *Vessel Plus.* 2020. [10.20517/2574-1209.2019.34](https://doi.org/10.20517/2574-1209.2019.34).



This work is licensed under a Creative Commons Attribution

**Table 1.** Clinical characteristics of populations included in this study.

No	Author (year)	Design	Age (year)	Gender (year)	Indication(s) of Surgery	DM Type	Outcome
1	Lemaigen <i>et al.</i> <sup>11</sup> (2015)	Observational	65.6 [55.4–74.9]	Male & Female	Cardiac surgeries: surgical emergency, isolated valve surgery, isolated CABG, valve + CABG & others.	Type I (insulin-dependent) and Type II	Diabetes mellitus (both non-insulin and insulin-dependent) were significantly correlated with sternal wound infection and wound healing impairment (p<0.001)
2	Sofer <i>et al.</i> <sup>12</sup> (1999)	Observational	≥65	Female	CABG	NS	Advanced age and the presence of diabetes mellitus were not significant risk factors for sternal wound infection (SWI).
3	Oswald <i>et al.</i> <sup>13</sup> (2012)	Meta-analysis	>65	Female	CABG	NS	CABG in patients with diabetes mellitus was strongly associated with sternal wound healing impairment.
4	Balachandran <i>et al.</i> <sup>14</sup> (2016)	Systematic Review	≥18	Male & Female	Cardiac surgeries: valve surgery, CABG, valve + CABG	Type I (insulin-dependent) and Type II	Impaired sternal healing and wound infection were significantly correlated with diabetes mellitus, both non-insulin and insulin-dependent
5	Shin <i>et al.</i> <sup>15</sup> (2015)	Observational	64.4	Male & Female	CABG	NS	Diabetes mellitus was strongly associated (p<0.05) with poor sternal healing.
6	Heilmann <i>et al.</i> <sup>16</sup> (2013)	Observational	67.0 ± 12.7	Male & Female	Cardiac surgeries: surgical emergency, isolated valve surgery, isolated CABG, valve + CABG & aortic operations	Type I (insulin-dependent) and Type II	Type I diabetes mellitus but not type II was associated with both postoperative superficial SWI and deep SWI. The other contributing factors were obesity and COPD.
7	Gansera <i>et al.</i> <sup>17</sup> (2016)	Observational	60.1 ± 5.3	Male	CABG	Type I (insulin-dependent) and Type II	Even though both types (I&II) DM patients had a higher risk of developing deep SWI after sternotomy, it did not reach clinical significance. There was also no significant correlation between BIMA (bilateral IMA grafting) and SIMA grafting method with long-term outcomes among diabetics who underwent CABG.



No	Author (year)	Design	Age (year)	Gender (year)	Indication(s) of Surgery	DM Type	Outcome
8	Zalewska-Adamiec <i>et al.</i> <sup>18</sup> (2014)	Observational	69.7 ± 6.75	Male	CABG	Type I (insulin-dependent) and Type II	Wound healing defect and sternal dehiscence were significantly higher among diabetics (both types) compared with non-diabetics.
9	Nakano <i>et al.</i> <sup>19</sup> (2008)	Observational	69.3 ± 8.1	Female	Cardiac surgeries: surgical emergency, isolated valve surgery, isolated CABG, valve + CABG & aortic operations	Type I (insulin-dependent) and Type II	Sternal wound infection and healing impairment were significantly higher among diabetics than non-diabetics.
10	Peterson <i>et al.</i> <sup>20</sup> (2003)	Observational	60.1 ± 9.3	Male	CABG	Type I (insulin-dependent) and Type II	Sternal healing impairment and healing complications were significantly lowered among diabetics that received skeletonized IMA grafts rather than non-skeletonized.
11	Zacharias <i>et al.</i> <sup>21</sup> (1996)	Observational	63	Male & Female	Cardiac surgeries: CABG only, the valve only, others	Type I (insulin-dependent) & Type II	Diabetes mellitus, mainly insulin-dependent (type I), was strongly associated with the incidence of SWI after cardiac surgery.
12	Pevni <i>et al.</i> <sup>22</sup> (2008)	Observational	>70	Male & Female	CABG	Type I (insulin-dependent) & Type II	Diabetic patients had a strong association with the incidence of SWI.
13	Zuckermann <i>et al.</i> <sup>23</sup> (2011)	Review	NS	Male	Heart transplantation	NS	Diabetic patients undergoing heart transplantation have a higher incidence of deep SWI and also a higher risk (Odds ratio/OR; 2.1 – 5.5)
14	Lazar <i>et al.</i> <sup>24</sup> (2018)	Review	NS	Male & Female	CABG	NS	There was no significant difference between diabetic and non-diabetic in deep sternal wound healing. This was suggested to be correlated with the method used – skeletonization, by which it preserved collateral circulation to the sternum.
15	Fu <i>et al.</i> <sup>25</sup> (2015)	Observational	65.6 ± 15.7	Male & Female	Cardiac surgeries: elective & emergency surgeries, IMA grafting	NS	Diabetic patients that require medication, BMI > 30, respiratory disorders, and emergency surgery were significant predictors of sternal wound dehiscence.

No	Author (year)	Design	Age (year)	Gender (year)	Indication(s) of Surgery	DM Type	Outcome
16	Gurevitch <i>et al.</i> <sup>26</sup> (1999)	Observational	65	Female	CABG	NS	Sternal wound infection and healing impairment were not significantly correlated with diabetes mellitus and advanced age (>65 years old). Healing impairment was only encountered in 30% of patients.
17	Shaheen <i>et al.</i> <sup>27</sup> (2020)	Observational	60 ± 1.25	Male & Female	CABG	NS	Superficial and deep sternal wound infections along with impaired healing were seen more commonly among diabetic patients, although P-value didn't reach a significant value (P>0.05)
18	Zahiri <i>et al.</i> <sup>28</sup> (2012)	Observational	61.8	Male & Female	Sternal Wound Reconstruction	NS	Sternal wound complication after flap reconstruction was 2.6 times more frequently to be encountered among diabetics.
19	Lenz <i>et al.</i> <sup>29</sup> (2016)	Observational	NS	NS	CABG	Type I (insulin-dependent) & Type II	Sternal wound infections (superficial and deep) were significantly higher in diabetic patients, mainly insulin-dependent diabetes mellitus
20	Savage <i>et al.</i> <sup>30</sup> (2007)	Observational	64.6 ± 10.2	Male & Female	CABG	Type I (insulin-dependent) & Type II	The incidence of deep sternal wound complications was significantly higher in patients with diabetes mellitus, especially in insulin-dependent diabetics.
21	Fakih <i>et al.</i> <sup>31</sup> (2007)	Observational	67.2 ± 11	Male	CABG	NS	Deep and superficial SWI were encountered more frequently in candidates with diabetes mellitus (p<0.005).
22	Sakamoto <i>et al.</i> <sup>32</sup> (2003)	Observational	62.0 ± 11.9	Male	Cardiac surgeries: surgical emergency, isolated valve surgery, isolated CABG, valve + CABG & aortic operations	NS	This study showed no significant correlation between DM and deep sternal wound infection
23	Kieser <i>et al.</i> <sup>33</sup> (2014)	Observational	65 ± 10.4	Male & Female	CABG	Type I insulin-dependent) & Type II	There was no significant impact of diabetes mellitus on the incidence of deep SWI both in male and female diabetics.

No	Author (year)	Design	Age (year)	Gender (year)	Indication(s) of Surgery	DM Type	Outcome
24	Dai <i>et al.</i> <sup>34</sup> (2013)	Review	>65	NS	CABG (BIMA & SIMA)	NS	Patients with diabetes mellitus had more adverse and extended sternal wound complications after undergoing CABG surgery,