



Source details

Annals of Medicine and Surgery

Scopus coverage years: from 2012 to Present

Publisher: Elsevier

E-ISSN: 2049-0801

Subject area: Medicine: Surgery

Source type: Journal

CiteScore 2022

1.7



SJR 2022

0.387



SNIP 2022

0.765



[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

Improved CiteScore methodology

CiteScore 2022 counts the citations received in 2019-2022 to articles, reviews, conference papers, book chapters and data papers published in 2019-2022, and divides this by the number of publications published in 2019-2022. [Learn more >](#)

CiteScore 2022

$$1.7 = \frac{5,570 \text{ Citations 2019 - 2022}}{3,335 \text{ Documents 2019 - 2022}}$$

Calculated on 05 May, 2023

CiteScoreTracker 2023

$$2.4 = \frac{7,967 \text{ Citations to date}}{3,380 \text{ Documents to date}}$$

Last updated on 05 September, 2023 • Updated monthly

CiteScore rank 2022

Category	Rank	Percentile
Medicine		
Surgery	#269/495	45th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

Annals of Medicine and Surgery

COUNTRY

Netherlands



Universities and research institutions in Netherlands



Media Ranking in Netherlands

SUBJECT AREA AND CATEGORY

Medicine

Medicine (miscellaneous)
Surgery

PUBLISHER

Elsevier BV

H-INDEX

36

PUBLICATION TYPE

Journals

ISSN

20490801

COVERAGE

2012-2022

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

annalsjournal@elsevier.com

SCOPE







Annals of Medicine and Surgery is an online-only, peer-reviewed open access journal with a global outlook and focus on those training in medicine and surgery (postgraduate and undergraduate). AMS contains a mix of original clinical and basic science research, reviews, editorials, commentary, perspectives, debate, opinion, case reports and journal club reports. The journal crosses the whole disease spectrum of medicine and surgery but does also have a special focus on the following areas: -Patient safety- Human factors, teamwork, communication and professionalism- Quality improvement science and practice- Evidence based medicine- Implementation science – clinical evidence into practice- Leadership and management- Medical education, teaching, and training- Public and global health- Healthcare policy, delivery, commissioning, and resource management- Use of technology and health informatics- Clinical ethics and medical law- Research and innovation – bedside to bench and back again – including applications like personalised medicine.




Join the conversation about this journal

FIND SIMILAR JOURNALS ?


<p>1 International Journal of Surgery NLD</p> <p>80% similarity</p>	<p>2 International Journal of Surgery Open GBR</p> <p>73% similarity</p>	<p>3 International Journal of Surgery Case Reports NLD</p> <p>67% similarity</p>	<p>4 BMC Surgery GBR</p> <p>61% similarity</p>
---	--	--	--





SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of April 2023

M Muhammad Arsham Javed 2 months ago

Hi there,
Can you please tell what is the acceptance rate of annals of medicine and Surgery?

reply



Melanie Ortiz 1 month ago

SCImago Team

Dear Muhammad,
Thank you for contacting us.
We suggest you visit the journal's homepage or contact the journal's editorial staff , so they could inform you more deeply.
Best Regards, SCImago Team

C Chia 4 months ago

Would like to understand whether this AMS journal is in 2023 SCI list?

reply

update of the data. This update is send to us around April / May every year. Thus, the indicators for 2019 will be available in June 2020 and before that we can't know what will happen with this journal. Best Regards, SCImago Team

Leave a comment

Name

Email

(will not be published)

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS
Horatio (Satire 1, 1, 106)

[Cookie settings](#)[Cookie policy](#)



ISSN 2090-0695

ANNALS OF MEDICINE & SURGERY



www.annalsjournal.com

ScienceDirect

Editorial Board : Annals of Medicine and Surgery

Advertisement

Editorial Board

Editor-In-Chief

Riaz Agha, London, UK

Statistical Editor

L. C. See, Taoyuan, Taiwan

Associate Editors

H. Ahmed, Sulaimani, Iraq

S. Al-Hakkak, Najaf, Iraq

G. Alsfasser, Rostock, Germany

F. C. Campanile, Civita Castellana, Italy

O. P. Choudhary, Punjab, India

D. Chowdhury, Manchester, UK

A. Coppola, Rome, Italy

A. Hasan, Cairo, Egypt

A. Jha, Delhi, India

K. Kopeva, Tomsk, Russia

M. Mitchell, New York, USA

R. Moshref, Jeddah, Saudi Arabia

E. Ntouvali, Athens, Greece

M. Oley, Manado, Indonesia

C. Paasch, Brandenburg, Germany

R. Rosyidi, Mataram, Indonesia

P. Schemmer, Graz, Austria

C. Sohrabi, London, UK

D. A. Syahputra, Aceh, Indonesia

N. Tenreiro, Vila Real, Portugal

Editorial Board

X-P. Chen, Wuhan, China

M. Cohen, Newark, USA

S. Hamid, Karachi, Pakistan

P. Hutchinson, Cambridge, UK

P. Littlejohns, London, UK

A. Mahapatra, Bhubaneswar, India

J. McKimm, Swansea, UK

D. Muzumdar, Mumbai, India

F. Nicoli, London, UK

D. Sokol, London, UK

J. Talati, Karachi, Pakistan

J. V. Thomas, Birmingham, USA

M. C. Warlé, Nijmegen, Netherlands

T. G. Williams, Tunbridge Wells, UK

Advertisement

[^Back to Top](#)**Never Miss an Issue**

Get new journal Tables of Contents sent right to your email inbox

Browse Journal Content

- [Register on the website](#)

Customer Service

Contact us at:

- Support:
[Submit a Service Request](#)
- TEL:
800-638-3030 (within the USA)
301-223-2300 (outside of the USA)

-
- [Privacy Policy \(Updated June 29, 2023\)](#)
- [Legal Disclaimer](#)
- [Terms of Use](#)
- [Open Access Policy](#)
- Copyright © 2023
- [Wolters Kluwer Health, Inc. and/or its subsidiaries. All rights reserved.](#)

Advertisement



March 2022 - Volume 75 - Issue : Annals of Medicine and Surgery

[Home](#) > March 2022 - Volume 75 - Issue

< [Previous Issue](#) | [Next Issue](#) >



March 2022 - Volume 75 - Issue

[Table of Contents Outline](#)

[eTOC Alerts](#)

[Contributor Index](#)

Case Report

[Immune thrombocytopenia purpura flare post COVID-19 vaccine](#)

Ali, Elrazi; Al-Maharmeh, Qusai; Rozi, Waail Mohammed; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[A very rare case of chylomicronemia revealed by cerebral thrombophlebitis in a 4-month-old infant](#)

Onci-Es-Saad, ; Lamzouri, Oussama; Bouchlarhem, Amine; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-

- [Permissions](#)
- [Free](#)

[**Primary diffuse B-cell non-Hodgkin's breast lymphoma; A case report with a brief literature review**](#)

Hammood, Zuhair D.; Salih, Abdulwahid M.; Latif, Shaban; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[**Giant symptomatic adrenal myelolipoma: A case report**](#)

Ramdani, Abdelbassir; Aissaoui, Asmae; Bouhout, Tariq; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[**Hemi-mandibulectomy without bony reconstruction: A case report of mandibular metastasis from a silent differentiated papillary thyroid cancer**](#)

Wahyono, Artanto; Dwianingsih, Ery Kus; Avanti, Widya Surya; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[**Capillary hyperpermeability syndrome: A fatal complication of acute leukaemia: Case report and review**](#)

Benzekri, Hajar; Bouchlarhem, Amine; lamassab, Nour el houda; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[**A promising contribution to negative pressure wound therapy in treatment of prosthetic joint infection. Discussion based on case report**](#)

Tsvetkov, Vitaly O.; Ivkov, Alexey V.; Mikaelyan, Liana S.; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)

- [Free](#)

[Magnitude and factors associated with preoperative depression among elective surgical patients at University of Gondar comprehensive specialized hospital, North West Ethiopia: A cross-sectional study](#)

Kebede, Yeneneh Negesse; Denu, Zewditu Abdissa; Aytolign, Habtu Adane; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[Residual symptoms and the quality of life in individuals recovered from COVID-19 infection: A survey from Pakistan](#)

Qamar, Mohammad Aadil; Martins, Russell Seth; Dhillon, Rubaid Azhar; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[Extraintestinal manifestation of inflammatory bowel disease and associated factors in pediatric patients](#)

Rahmani, Parisa; Rasti, Ghazaleh; Gorgi, Mojtaba; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[Profile of cardiovascular disease risk in type 2 diabetes mellitus patients receiving statin therapy: A cross-sectional study](#)

Ikhsan, Yasmin Karimah; Soelistijo, Soebagijo Adi; Putranto, Johannes Nugroho Eko

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[Assessment of post-operative nausea and vomiting prophylaxis usage for cesarean section, 2021: A cross sectional study](#)

Admass, Biruk Adie; Tawye, Hailu Yimer; Endalew, Nigussie Simeneh; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)

Kang, Ajaipal S.; Kang, Kevin S.

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[Obstructive oligo-anuria revealing pelvic gynecological cancers, analysis of a series of 102 cases](#)

Kbirou, A.; Sayah, M.; Sounni, F.; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[Acute orchiepididymitis: Epidemiological and clinical aspects: An analysis of 152 cases](#)

Kbirou, A.; Alafifi, M.; Sayah, M.; More

Annals of Medicine & Surgery. 75:, March 2022.

- [⊕ Abstract](#)
-
-
- [Permissions](#)
- [Free](#)

[Table of Contents Outline](#) | [Back to Top](#)

- [1](#)
- [2](#)
-

Show 100 results per page



[^Back to Top](#)



Never Miss an Issue

Get new journal Tables of Contents sent right to your email inbox

Browse Journal Content


- [Register on the website](#)

Customer Service

- Contact us at:
 - Support:
[Submit a Service Request](#)
 - TEL:
800-638-3030 (within the USA)
301-223-2300 (outside of the USA)

[Manage Cookie Preferences](#)

- [Privacy Policy \(Updated June 29, 2023\)](#)
- [Legal Disclaimer](#)
- [Terms of Use](#)
- [Open Access Policy](#)
- Copyright © 2023
- [Wolters Kluwer Health, Inc. and/or its subsidiaries. All rights reserved.](#)



Cross-sectional Study

Profile of cardiovascular disease risk in type 2 diabetes mellitus patients receiving statin therapy: A cross-sectional study

Yasmin Karimah Ikhsan^a, Soebagijo Adi Soelistijo^{b,*}, Johannes Nugroho Eko Putranto^c

^a Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

^b Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

^c Department of Vascular and Cardiology Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

ARTICLE INFO

Keywords:

Atherosclerosis vascular disease
Framingham risk score
Type 2 diabetes mellitus

ABSTRACT

Background: Cardiovascular disease is still the number 1 cause of death globally. Meanwhile, type 2 diabetes mellitus (T2DM) is a risk factor for atherosclerosis vascular disease (ASCVD), so an assessment using Framingham Risk Score (FRS) is needed to predict the risk of ASCVD in the future.

Objective: Analyzing the risk factor of ASCVD using the Framingham Risk Score (FRS) in T2DM patients.

Methods: This study was conducted from July 2020 to July 2021, which the participants were measured for FRS including age, gender, current smoking, diabetes, blood pressure (systolic), high-density lipoprotein (HDL) cholesterol, total cholesterol (TC), and ASCVD risk score. The analysis employed multiple linear tests and ANOVA tests with $p < 0.05$.

Results: Several ASCVD risk factors in T2DM patients were found, including gender ($t = 6.015$; $p < 0.001$), age ($t = 6.901$; $p < 0.001$), HDL level ($t = 2.287$; $p = 0.024$), CT level ($t = 5.273$; $p < 0.001$), blood pressure ($t = 5.850$; $p < 0.001$), and current smoking ($t = 2.638$; $p = 0.009$). The results of analysis between ASCVD risk factor and level of ASCVD risk obtained a significant association ($F = 36,642$; $p < 0.001$).

Conclusion: Risk factors of ASCVD in T2DM patients such as gender, age, HDL level, CT level, blood pressure, and current smoking.

1. Introduction

Cardiovascular disease (CVD) is the number one death for non-communicable diseases each year globally. Around 17.9 million people in the world die from this disease, and it is 31% of the causes of death worldwide [1]. Data from WHO shows that in Indonesia in 2012, ischemic heart disease (IHD) placed second as the disease that can cause death in sufferers by 8.9% or around 138.4 thousand population, where one of the main causes is atherosclerosis and diabetes [2]. Overall, an estimated 2.6 million deaths are caused by elevated cholesterol and another 29.7 million people experience disability each year. The prevalence of coronary heart disease is 1.5% that increases with age, where the highest group is at the age of 65–74 years [3].

Diabetes mellitus is a metabolic disorder caused by genetic and environmental factors. Type 2 diabetes mellitus (T2DM) is a major risk factor affecting coronary artery disease (CAD) which approximately 75% of T2DM patients die as a consequence of atherosclerosis vascular

disease (ASCVD) [4]. In T2DM patients there is insulin resistance which triggers hyperglycemia which causes multiple metabolic abnormalities that increase the risk of ASCVD [5,6]. The gold-standard treatment for the prevention of primary and secondary ASCVD used statins [7] which statins have been shown to reduce the risk of CVD in T2DM [8]. This study aimed to analyze the profile of the risk factors for ASCVD in T2DM patients.

2. Methods

2.1. Participants

Participants in this study were patients diagnosed with T2DM [9,10]. Participant inclusion criteria included patients diagnosed with T2DM (HbA1c of $\geq 6.5\%$, fasting plasma glucose of ≥ 126 mg/dL, 2-h plasma glucose & random plasma glucose ≥ 200 mg/dL) [11,12], and consuming statins. Meanwhile, participant exclusion criteria included

* Corresponding author. Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Jl. Mayjend Prof. Dr. Moestopo No. 6-8, Airlangga, Gubeng, Surabaya, East Java, 60286, Indonesia.

E-mail address: soebagijo1121@gmail.com (S.A. Soelistijo).

<https://doi.org/10.1016/j.amsu.2022.103368>

Received 17 December 2021; Received in revised form 4 February 2022; Accepted 10 February 2022

Available online 15 February 2022

2049-0801/© 2022 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

patients with a history of CVD, patients who had received nicotinic acid therapy, bile acid sequestrants, fibrates, ezetimibe, and PCSK9 inhibitors. Participants were required to fill out an informed consent before starting the study.

2.2. Study design

A cross-sectional study was conducted using purposive sampling from July 2020 to June 2021 with 124 T2DM patients. This study was conducted based on Strengthening the Reporting of Cohort Studies in Surgery (STROCSS) 2021 Guideline [13]. Measurement of ASCVD risk score was calculated based on the Framingham score (FRS), of which the calculation was based on ASCVD risk factors including age, TC, HDL, blood pressure (systolic), current smoking, and diabetes [14]. The measurement results were categorized into 3, namely low risk (<10%), middle risk (11–19%), and high risk ($\geq 20\%$). This measurement was used to predict that participants are likely to experience ASCVD disorders during the next 10 years [15].

2.3. Blood pressure examination

Blood pressure is a vital sign measured by a tensimeter that is used to make a clinical decision. It is divided into two, systolic and diastolic, with a measurement unit of mmHg. Blood pressure used in this study was only systolic, which are categorized into hypertension (>140 mmHg), normal (100–140 mmHg), and hypotension (systolic <100 mmHg) [16]. The categorization of blood pressure used in this study was <120 mmHg, 120–29 mmHg, 130–139 mmHg, 140–149 mmHg, 150–159 mmHg, and ≥ 160 mmHg.

3. Cholesterol examination

Cholesterol levels in this study were measured using a chemical auto-analyzer (Toshiba, Japan) [17], in which the cholesterol used in this study included TC and HDL cholesterol [18]. Measurement of TC in this study was in mg/dL, which was categorized into <160 mg/dL, 160–199 mg/dL, 200–239 mg/dL, 240–279 mg/dL, and ≥ 280 mg/dL. Meanwhile, HDL level measurements were categorized as follows <35 mg/dL, 35–44 mg/dL, 45–49 mg/dL, 50–59 dL, and 60 mg/dL.

3.1. Statistical analysis

The measurement data were collected and analyzed using the statistical package for the social science (SPSS) version 24.0 software (IBM Corp., Armonk, NY, USA). The data were presented in the form of tables or figures. Analysis to determine the risk factors for ASCVD in patients with diabetes mellitus used the ANOVA test which was declared significant if $p < 0.05$. In addition, this study used multiple linear regression with $p < 0.05$.

4. Results

4.1. Characteristic of participant

Most participants were female as many as 85 participants (65.8%), and all participants had T2DM. Most participants were non-smokers as many as 89 participants (71.8%). The participants' average age was 56.15 ± 9.01 , with a median value of 57.00 (51.00–62.25) years. The youngest and oldest participants were 30 years old and 75 years old, respectively. Most participants were in the age range of 50–54 years as many as 27 participants (21.7%). The participants' average HDL levels were 56.43 ± 17.42 mg/dL, with a median value of 52.00 (45.00–61.00) mg/dL. The lowest and highest HDL levels were 34.00 mg/dL and 124.00 mg/dL, respectively. Meanwhile, the participants' HDL levels ranged from 50.00 to 59.00 mg/dL and 60.00 mg/dL, each of which has a total of 37 participants (29.8%). The participants' average CT value

was 205.19 ± 36.47 mg/dL, with a median value of 199.00 (178.00–225.25) mg/dL. The highest and lowest CT scores were 303.00 mg/dL and 129 mg/dL, respectively. Most participants had CT values ranging from 160.00 to 199.00 mg/dL as many as 53 participants (42.7%). The participants' average blood pressure was 123.98 ± 11.36 mmHg, with a median value of 120.00 (120.00–130.00) mmHg. The participants' lowest and highest blood pressures were 100.00 mmHg and 181 mmHg, respectively. Most participants had blood pressure ranging from 120.00 to 129.00 mmHg as many as 53 participants (42.7%; Table 1).

4.2. Risk factor of atherosclerotic cardiovascular disease in type 2 diabetes mellitus patient

The FRS participant measurement showed that 26 participants (21%) had low risk of ASCVD, 52 participants (42%) had middle risk of ASCVD, and 46 participants (37%) had high risk of ASCVD. Based on the analysis, several risk factors for ASCVD in T2DM patients were found, including gender ($t = 6.015$; $p < 0.001$), age ($t = 6.901$; $p < 0.001$), HDL level ($t = 2.287$; $p = 0.024$), CT level ($t = 5.273$; $p < 0.001$), blood pressure ($t = 5.850$; $p < 0.001$), and current smoking ($t = 2.638$; $p = 0.009$; Table 2). There was a significant relationship between ASCVD risk factor and ASCVD risk level ($F = 36,642$; $p < 0.001$).

5. Discussion

Age is a risk factor for ASCVD because increasing age will increase the risk of atherosclerotic cardiovascular disease [19]. Age is a risk

Table 1
Characteristic of participant.

Variable	n (%)
Gender	
Male	39 (31.4)
Female	85 (68.5)
Age (years old)	
30-34	3 (2.4)
35-39	3 (2.4)
40-44	6 (4.8)
45-49	14 (11.2)
50-54	27 (21.7)
55-59	23 (18.5)
60-64	22 (17.7)
65-69	22 (17.7)
70-74	3 (2.4)
>75	1 (0.8)
HDL level (mg/dL)	
<35	1 (0.8)
35-44	25 (20.1)
45-49	24 (19.3)
50-59	37 (29.8)
≥ 60	37 (29.8)
Cholesterol total (mg/dL)	
<160	10 (8.0)
160-199	53 (42.7)
200-239	37 (29.8)
240-279	22 (17.7)
≥ 280	2 (1.6)
Blood pressure (mmHg)	
<120	23 (18.5)
120-129	53 (42.7)
130-139	41 (33.0)
140-149	1 (0.8)
150-159	4 (3.2)
≥ 160	2 (1.6)
Diabetes mellitus	
Yes	124 (100.0)
No	0 (0.0)
Current smoking	
Yes	35 (28.2)
No	89 (71.8)

Table 2
Risk factor of atherosclerotic cardiovascular disease in type 2 diabetes mellitus.

	FRS			t	p
	Low	Middle	High		
Gender				6.015	<0.001**
Male	4 (10.2)	6 (15.3)	29 (74.3)		
Female	22 (25.8)	46 (54.1)	17 (20.0)		
Age (years old)				6.901	<0.001**
30–34	3 (100.0)	0 (0.0)	0 (0.0)		
35–39	1 (33.3)	2 (66.7)	0 (0.0)		
40–44	4 (66.7)	2 (33.3)	0 (0.0)		
45–49	8 (57.1)	3 (21.4)	3 (21.4)		
50–54	5 (18.5)	15 (55.5)	7 (26.0)		
55–59	4 (17.4)	12 (52.1)	7 (30.4)		
60–64	1 (4.5)	8 (36.5)	13 (59.0)		
65–69	0 (0.0)	10 (45.5)	12 (54.5)		
70–74	0 (0.0)	0 (0.0)	3 (100.0)		
>75	0 (0.0)	0 (0.0)	1 (100.0)		
HDL level (mg/dL)				2.287	0.024*
<35	0 (0.0)	0 (0.0)	1 (100)		
35–44	3 (12.0)	7 (28.0)	15 (60.0)		
45–49	5 (20.8)	10 (41.6)	9 (37.5)		
50–59	9 (24.3)	16 (43.2)	12 (32.4)		
≥60	9 (24.3)	19 (51.3)	9 (24.3)		
Cholesterol total (mg/dL)				5.273	<0.001**
<160	3 (30.0)	3 (30.0)	4 (40)		
160–199	17 (32.0)	22 (41.6)	14 (26.4)		
200–239	6 (16.2)	17 (46.0)	14 (37.8)		
240–279	0 (0.0)	9 (41.0)	13 (59.0)		
≥280	0 (0.0)	1 (50.0)	1 (50.0)		
Blood pressure (mmHg)				5.850	<0.001**
<120	13 (56.5)	7 (30.5)	3 (13.0)		
120–129	8 (15.0)	25 (47.1)	20 (37.7)		
130–139	5 (21.1)	20 (48.7)	16 (39.0)		
140–149	0 (0.0)	0 (0.0)	1 (100.0)		
150–159	0 (0.0)	0 (0.0)	4 (100.0)		
≥160	0 (0.0)	0 (0.0)	2 (100.0)		
Diabetes mellitus				–	–
Yes	26 (21.0)	52 (42.0)	46 (37.0)		
No	0 (0.0)	0 (0.0)	0 (0.0)		
Current smoking				2.638	0.009*
Yes	5 (14.3)	7 (20.0)	23 (65.7)		
No	21 (23.5)	45 (50.5)	23 (26.0)		

Note: FRS = Framingham risk score; HDL = high-density lipoprotein; *significant <0.05; **significant <0.001.

factor for CAD as increasing age will increase the risk of CVD. The older the age, the greater the possibility of atheroma plaques sticking to the walls and disrupting blood flow to the tissues [20]. ASCVD is more common in men than women as supported by previous research, stating that the majority of patients with ASCVD are men [21]. The latest study in Indonesia stated that ASCVD was found in men as much as 61% with a prevalence ratio of 2:1 between men and women [3].

Lower HDL cholesterol levels are associated with a higher risk of cardiovascular disease. Several studies have also shown that people with HDL levels of <35 mg/dL are more likely to experience ASCVD [18,22]. HDL will carry cholesterol from tissues including coronary arteries to be catabolized in the liver. The newly formed HDL from its precursors, namely apo a-1, apo a-2, and apo a-milano proteins, will attract cholesterol with the help of the ATP binding cassette transporter A1 (ABCT A1) enzyme. This HDL cholesterol will later increase in size and can attract more fat and cholesterol in the body for catabolism [23]. In addition, high CT levels lead to deposition, narrowing, and plaque in blood vessels and increase the risk of ASCVD [18,24]. Hypertension is a major risk factor contributing to ASCVD, which has been mentioned in several kinds of literature and studies [25]. Recent studies have also stated that patients with hypertension have a 10 times risk of developing ASCVD including coronary heart disease [26].

Meanwhile, current smoking increases the risk of ASCVD which mechanisms that cause atherosclerosis are direct endothelial injury due to agents in cigarettes such as carbon monoxide and nicotine that cause blebs on the lumen surface, microphile formation, and release of endothelial cells (endothelial damage), changes in platelets, increased fibrinogen and C-reactive protein levels and induce proinflammatory cytokines [27]. Type 2 DM is a disease related to glucose in the blood, in which when hyperglycemia occurs, the blood will experience a viscosity so that the blood supply to the tissue is blocked and has a high risk of ASCVD [2].

Nevertheless, this study has a limitation as it only included data components of the Framingham risk scoring risk factors such as gender, age, HDL cholesterol levels, total cholesterol levels, blood pressure, type 2 diabetes mellitus status, and smoking status. Data such as body weight, family history, low-density lipoprotein cholesterol levels, triglyceride levels, and physical activity were not included in this study. In addition, the method used was FRS, in which the score is only based on data from the white population. These considerations are expected to become an evaluation for further research.

6. Conclusion

Based on the results of the patients' risk score, the highest percentage of atherosclerotic cardiovascular disease risk using FRS is patients with a moderate-risk level, followed by a high-risk level and a low-risk level. The analysis shows that T2DM participants are at risk for ASCVD that is influenced by gender, age, current smoking, diabetes, blood pressure, HDL cholesterol, and TC.

Ethical approval

We have conducted an ethical approval based on the Declaration of Helsinki with registration research at the Health Research Ethics Committee in Dr. Soetomo General Academic Hospital, Surabaya, Indonesia.

Funding

None.

Author contributor

All authors contributed toward data analysis, drafting and revising the paper, gave final approval of the version to be published and agree to be accountable for all aspects of the work.

Downloaded from http://journals.lww.com/annals-of-medicine-and-surgery by BnDMf5ePPhKav1zEoum1tQIN44+KjLhEz9shH04XMh0CwC1AWnVOp/IIQHhD3i3D00dRv/TVSF14C13VCA/OAVpDda8k2+Y6h5t15kE= on 10/05/2023

Registration of research studies

Name of the registry: Health Research Ethics Committee in the Dr. Soetomo General Academic Hospital, Surabaya, Indonesia.

Unique identifying number or registration ID: 0181/LOE/301.4.2/XI/2020.

Hyperlink to your specific registration (must be publicly accessible and will be checked): .

Guarantor

Soebagijo Adi Soelistijo.

Consent

Written informed consent was obtained from the patient.

Provenance and peer review

Not commissioned, externally peer-reviewed.

Declaration of competing interest

The authors declare no conflict of interest.

Acknowledgment

We thank our editor "Fis Citra Ariyanto".

Abbreviations

ASCVD	atherosclerosis vascular disease
ABCT A1	ATP binding cassette transporter A1
CAD	coronary artery disease
CVD	Cardiovascular disease
FRS	Framingham Risk Score
HDL	high-density lipoprotein
IHD	ischemic heart disease
LDL	low-density lipoprotein
STROCSS	Strengthening the Reporting of Cohort Studies in Surgery
T2DM	type 2 diabetes mellitus
TC	total cholesterol

References

- [1] W. Hinton, A. McGovern, R. Coyle, T.S. Han, P. Sharma, A. Correa, et al., Incidence and prevalence of cardiovascular disease in English primary care: a cross-sectional and follow-up study of the Royal College of General Practitioners (RCGP) Research and Surveillance Centre (RSC), *BMJ Open* 8 (8) (2018), e020282, <https://doi.org/10.1136/bmjopen-2017-020282>.
- [2] A. Crisafulli, P. Pagliaro, S. Roberto, L. Cugusi, G. Mercurio, A. Lazou, et al., Diabetic cardiomyopathy and ischemic heart disease: prevention and therapy by exercise and conditioning, *Int. J. Mol. Sci.* 21 (8) (2020), <https://doi.org/10.3390/ijms21082896>.
- [3] A. Maharani, Sujarwoto, D. Praveen, D. Oceandy, G. Tampubolon, A. Patel, Cardiovascular disease risk factor prevalence and estimated 10-year cardiovascular risk scores in Indonesia: the SMARTHealth Extend study, *PLoS One* 14 (4) (2019), e0215219, <https://doi.org/10.1371/journal.pone.0215219>.
- [4] R. Naito, K. Miyauchi, Coronary artery disease and type 2 diabetes mellitus, *Int. Heart J.* 58 (4) (2017) 475–480, <https://doi.org/10.1536/ihj.17-191>.
- [5] S. De Rosa, B. Arcidiacono, E. Chiefari, A. Brunetti, C. Indolfi, D.P. Foti, Type 2 diabetes mellitus and cardiovascular disease: genetic and epigenetic links, *Front. Endocrinol.* 9 (2018) 2, <https://doi.org/10.3389/fendo.2018.00002>.
- [6] M.A. Abdul-Ghani, A. Jayyousi, R.A. DeFronzo, N. Asaad, J. Al-Suwaidi, Insulin resistance the link between T2DM and CVD: basic mechanisms and clinical implications, *Curr. Vasc. Pharmacol.* 17 (2) (2019) 153–163, <https://doi.org/10.2174/15701611156661710101151119>.
- [7] U. Galicia-Garcia, S. Jebari, A. Larrea-Sebal, K.B. Uribe, H. Siddiqi, H. Ostolaza, et al., Statin treatment-induced development of type 2 diabetes: from clinical evidence to mechanistic insights, *Int. J. Mol. Sci.* 21 (13) (2020), <https://doi.org/10.3390/ijms21134725>.
- [8] K.C. Maki, V. Diwadkar-Navsariwala, M.W. Kramer, Statin use and risk for type 2 diabetes: what clinicians should know, *PGM (Postgrad. Med.)* 130 (2) (2018) 166–172, <https://doi.org/10.1080/00325481.2018.1402658>.
- [9] A.B. Pulungan, I.T. Afifa, D. Annisa, Type 2 diabetes mellitus in children and adolescent: an Indonesian perspective, *Ann Pediatr Endocrinol Metabol* 23 (3) (2018) 119–125, <https://doi.org/10.6065/apem.2018.23.3.119>.
- [10] D. Ardiany, A. Pranoto, S.A. Soelistijo, Widjaja SA. Libriansyah, Association between neutrophil-lymphocyte ratio on arterial stiffness in type-2 diabetes mellitus patients: a part of DiORS Study, *Int. J. Diabetes Dev. Ctries.* (2021), <https://doi.org/10.1007/s13410-021-00965-1>.
- [11] K. Pippitt, M. Li, H.E. Gurgle, Diabetes mellitus: screening and diagnosis, *Am. Fam. Physician* 93 (2) (2016) 103–109.
- [12] A. Kumar, T. Kumar, M. Bhargava, R. Raj, V. Vaibhav, J. Kishore, Salivary and serum glucose levels in diabetes mellitus patients versus control - a randomised control trial, *J Med life* 13 (2) (2020) 235–240, <https://doi.org/10.25122/jml-2020-0062>.
- [13] G. Mathew, R. Agha, Stross 2021: Strengthening the reporting of cohort, cross-sectional and case-control studies in surgery, *Int. J. Surg.* 96 (2021) 106165, <https://doi.org/10.1016/j.jisu.2021.106165>.
- [14] C.A. Schulz, L. Mavarani, N. Reinsch, S. Albayrak-Rena, A. Potthoff, N. Brockmeyer, et al., Prediction of future cardiovascular events by Framingham, SCORE and aSCVD risk scores is less accurate in HIV-positive individuals from the HIV-HEART Study compared with the general population, *HIV Med.* 22 (8) (2021) 732–741, <https://doi.org/10.1111/hiv.13124>.
- [15] W. Weng, Y. Tian, S.X. Kong, R. Ganguly, M. Hersloev, J. Brett, et al., Impact of atherosclerotic cardiovascular disease on healthcare resource utilization and costs in patients with type 2 diabetes mellitus in a real-world setting, *Clin Diabetes Endocrinol* 6 (2020) 5, <https://doi.org/10.1186/s40842-019-0090-y>.
- [16] A. Herbert, J.K. Cruickshank, S. Laurent, P. Boutouyrie, Establishing reference values for central blood pressure and its amplification in a general healthy population and according to cardiovascular risk factors, *Eur. Heart J.* 35 (44) (2014) 3122–3133, <https://doi.org/10.1093/eurheartj/ehu293>.
- [17] A. Ozder, Lipid profile abnormalities seen in T2DM patients in primary healthcare in Turkey: a cross-sectional study, *Lipids Health Dis.* 13 (2014) 183, <https://doi.org/10.1186/1476-511x-13-183>.
- [18] D.M. Maahs, A.K. Maniatis, K. Nadeau, R.P. Wadwa, K. McFann, G.J. Klingensmith, Total cholesterol and high-density lipoprotein levels in pediatric subjects with type 1 diabetes mellitus, *J. Pediatr.* 147 (4) (2005) 544–546, <https://doi.org/10.1016/j.jpeds.2005.04.068>.
- [19] T. Head, S. Daunert, P.J. Goldschmidt-Clermont, The aging risk and atherosclerosis: a fresh look at arterial homeostasis, *Front. Genet.* 8 (2017) 216, <https://doi.org/10.3389/fgene.2017.00216>.
- [20] Y. Yang, J. Tian, C. Zeng, J. Wei, L.J. Li, X. Xie, et al., Relationship between hyperuricemia and risk of coronary heart disease in a middle-aged and elderly Chinese population, *J. Int. Med. Res.* 45 (1) (2017) 254–260, <https://doi.org/10.1177/0300060516673923>.
- [21] M. Rafieian-Kopaei, M. Setorki, M. Doudi, A. Baradaran, H. Nasri, Atherosclerosis: process, indicators, risk factors and new hopes, *Int. J. Prev. Med.* 5 (8) (2014) 927–946.
- [22] L. Badimon, G. Vilahur, LDL-cholesterol versus HDL-cholesterol in the atherosclerotic plaque: inflammatory resolution versus thrombotic chaos, *Ann. N. Y. Acad. Sci.* 1254 (2012) 18–32, <https://doi.org/10.1111/j.1749-6632.2012.06480.x>.
- [23] H.H. Wang, G. Garruti, M. Liu, P. Portincasa, D.Q. Wang, Cholesterol and lipoprotein metabolism and atherosclerosis: recent advances in reverse cholesterol transport, *Ann. Hepatol.* 16 (Suppl. 1) (2017) s27–s42, <https://doi.org/10.5604/01.3001.0010.5495>, s3-105.
- [24] R. Quispe, M.B. Elshazly, D. Zhao, P.P. Toth, R. Puri, S.S. Virani, et al., Total cholesterol/HDL-cholesterol ratio discordance with LDL-cholesterol and non-HDL-cholesterol and incidence of atherosclerotic cardiovascular disease in primary prevention: the ARIC study, *Eur J Prev Cardiol* 27 (15) (2020) 1597–1605, <https://doi.org/10.1177/2047487319862401>.
- [25] R. Nakanishi, L. Baskaran, H. Gransar, M.J. Budoff, S. Achenbach, M. Al-Mallah, et al., Relationship of hypertension to coronary atherosclerosis and cardiac events in patients with coronary computed tomographic angiography, *Hypertension (Dallas, Tex : 1979)* 70 (2) (2017) 293–299, <https://doi.org/10.1161/hypertensionaha.117.09402>.
- [26] Y. Zhou, R. Zhang, G. Wang, A. Wang, C. Zhong, M. Zhang, et al., Coexistence effect of hypertension and angiotensin II on the risk of coronary heart disease: a population-based prospective cohort study among Inner Mongolians in China, *Curr. Med. Res. Opin.* 35 (8) (2019) 1473–1478, <https://doi.org/10.1080/03007995.2019.1601430>.
- [27] G. Siasos, V. Tsigkou, E. Kokkou, E. Oikonomou, M. Vavuranakis, C. Vlachopoulos, et al., Smoking and atherosclerosis: mechanisms of disease and new therapeutic approaches, *Curr. Med. Chem.* 21 (34) (2014) 3936–3948, <https://doi.org/10.2174/092986732134141015161539>.