

paper ficus

by turnitin turnitin

Submission date: 16-Dec-2023 06:08PM (UTC+0700)

Submission ID: 2260655235

File name: 10.1515_jbcpp-2020-0413.pdf (307.36K)

Word count: 4436

Character count: 24612

Siti Qamariyah Khairunisa*, Dwi Wahyu Indriati, Lidya Tumewu, Aty Widyawaruyanti and Nasronudin Nasronudin

Screening of anti-HIV activities in ethanol extract and fractions from *Ficus fistulosa* leaves

<https://doi.org/10.1515/jbcpp-2020-0413>

Received November 27, 2020; accepted March 8, 2021

Abstract

Objectives: Human immunodeficiency virus (HIV) infection is considered as a major immunosuppressive disease linked to malignancies and other opportunistic infections. Recently, the high prevalence of HIV drug-resistant strains required a high demand for novel antiviral drug development, especially in herbal medicine approaches. The objective of this study was to evaluate the possibility of *Ficus fistulosa* leaves can inhibit HIV replication in ethanol extract form as well as its fractions using chloroform, ethyl acetate, and butanol solvents.

Methods: *F. fistulosa* leaves were extracted using ethanol as a solvent and further gradually fractionated in chloroform, ethyl acetate, and butanol solvents. The targeted persistently infected virus (MT4/HIV) cell lines were cocultured with ethanol extract and fractions at different time points. The syncytium formation and cytotoxicity assays were performed to evaluate the potential antiviral activity of *F. fistulosa* leaves.

Results: One of the four tested extract/fractions showed antiviral activity against HIV. The ethanol extract showed weak inhibition with a high level of

toxicity ($IC_{50} = 8.96 \mu\text{g/mL}$, $CC_{50} \geq 50 \mu\text{g/mL}$, and $SI = 5.58$). Meanwhile, chloroform fraction effectively inhibited the MT4/HIV cell proliferation while keeping the toxicity to a minimal level ($IC_{50} = 3.27 \mu\text{g/mL}$, $CC_{50} = 29.30 \mu\text{g/mL}$, and $SI = 8.96$). In contrast of ethyl acetate fraction and butanol fraction showed no anti HIV activity with a high level of toxicity ($CC_{50} \geq 50 \mu\text{g/mL}$) and low SI value ($>2.17 \mu\text{g/mL}$ and $>0.97 \mu\text{g/mL}$).

Conclusions: Chloroform fraction of *F. fistulosa* leaves showed effectively as anti-viral activity against MT4/HIV cells.

Keywords: anti HIV; *Ficus fistulosa*; *in vitro*; medicinal plant.

Introduction

Since being discovered in 1983, human immunodeficiency virus (HIV-1) has been infecting over 38 million people worldwide based on WHO global data tracking with over 25 million (>65%) infected individuals reside in Africa [1]. The distinctive characteristic of HIV-1 is the substantial genetic diversity that builds up within and between hosts [2]. The high mutation rate in HIV-1 is due to the absence of DNA repair enzymes, creating approximately one nucleotide mutation per cycle during viral replication [3]. This unique characteristic of HIV-1 has become a big obstacle to researchers in studying the new approach of HIV-1 antiviral drug discovery.

Diverse novel approaches to anti HIV treatments are currently being developed worldwide and mostly focusing on the initial step of viral entry. One approach targeted the gp120 receptor on the HIV-1 envelope to create a novel antiviral drug by using boronic acid materials as an anti-retroviral agent (ART) [4, 5]. Another therapeutic method, frequently named “shock and kill” used latency-reversing agents (LRAs) to stimulate viral expression (“shock”), and afterward the latent HIV-infected cells could be exterminated by viral cytopathic effects or host immune responses (“kill”) [6, 7]. A recent study showed LRAs extracted from untested marine natural products was efficient to induce antiviral activity *in vitro* [8]. One alternative method is polypharmacology in which two or more multitarget or hybrid drugs can be used simultaneously to increase the antiviral activity against HIV-1 [9, 10]. To date, currently available anti HIV drugs remain nonoptimal

*Corresponding author: Siti Qamariyah Khairunisa, Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia, Phone: +6281331843627, E-mail: skhairunisa@staf.unair.ac.id
Dwi Wahyu Indriati, HIV Study Group, Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia; and Departement of Health, Faculty of Vocational Studies, Universitas Airlangga, Surabaya, Indonesia
Lidya Tumewu, Natural Product Medicine Research and Development, Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia
Aty Widyawaruyanti, Natural Product Medicine Research and Development, Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia; and Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia
Nasronudin Nasronudin, HIV Study Group, Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia; Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia; and Airlangga University Hospital, Universitas Airlangga, Surabaya, Indonesia

due to complicated procedures, uncontrolled cytotoxicity, and unpredicted side effects.

The failure in HIV-1 treatment especially in low-middle income countries was due to a shortage of drugs and the high cost of treatment [11]. However, the main factor of treatment failure is the lack of patient's discipline to take their prescribed drugs daily in the correct dose or not showing up at doctor's appointments for follow-up checks. The patient's social behavior could impact the development of drug-resistant therapy as well as increasing the fatality rate. The urge for *de novo* anti HIV drug development that is cost-efficient and has fewer side effects has become the priority in AIDS pharmacological research, preferably compounds extracted from local natural sources.

The research about natural products in drug development has been used since centuries ago, pioneered by the Egyptians in the discovery of Penicillin from *Penicillium chrysogenum* recorded in 1928 and was clinically accessible in the 1940s [12, 13]. The medicinal properties of active compounds from plants were derived from the anti-inflammatory and protective defense mechanisms of these products [14, 15]. Calanolides from *Calophyllum lanigerum* was one of the first plant-derived compounds discovered to have antiviral activity against HIV-1 [16, 17]. Betulinic acid extracted from the Chinese herb *Syzygium claviflorum* was succeeded to be synthesized as drug, Bevirimat, and continue in clinical trials since 2007 [18]. Other plant-based antiviral candidates have been studied intensively over the years and the results are promising. Additionally, the utilization of locally sourced natural products could be beneficial economically in the development of more affordable medicines.

The genus *Ficus* has approximately 725 species worldwide and three species namely *Ficus hispida*, *Ficus septica*, and *F. fistulosa* have been studied, and they exhibited anticancer, anti-inflammatory, and antiviral activities [19–21]. However, only Indriati et al. reported the potential anti HIV activity, and this recent study will investigate further the hidden anti-viral activity of *Ficus fistulosa* against HIV-1. In the present study, four samples derived from *F. fistulosa*, ethanol extract, ethyl acetate, chloroform, and butanol fractions were used in *in vitro* assay to assess their antiviral activity.

Materials and methods

Cells and viruses

The human acute T-lymphoblastic leukemia cells (MOLT-4/MT-4) were cultured in RPMI-1640 medium and DMEM medium (GIBCO,

USA) with the addition of 10% fetal bovine serum (Sigma, USA), 100 U/mL penicillin G, and 100 µg/mL streptomycin. The cells were incubated at 37 °C and 5% CO₂ humidity for three days prior to further usage. In addition, persistently infected cells were previously made from coculture of peripheral blood mononuclear cells (PBMC) from HIV-1 patients and healthy blood donors. Initially, the PBMC were pre activated with 10 µg/mL phytohemagglutinin (PHA), mitogen, and later on, induced with interleukin-2 (IL-2) [22]. The mix of MT-4 and the in-house persistently infected cells HIV-1 (MT-4/HIV-1) were cocultured according to the protocol from Gyuris et al. [23]. All HIV-1 isolates derived from the stock were collected from Surabaya, Indonesia.

Plant materials

The leaves from *F. fistulosa* were obtained from Cangar Conservation Forest, Malang, West Java, Indonesia. The verification and identification of the plant were supervised by Purwodadi Botanical Garden – Indonesia Institute of Science, East Java, Indonesia. All plant specimens were stored in Natural Product Medicine Research and Development Laboratorium (NPMRD), Institute of Tropical Disease (ITD), Universitas Airlangga, Surabaya, Indonesia.

Plant extraction and fractionations

F. fistulosa leaves as much as 2.5 kg were drying at room temperature and were powdered into approximately 250 g dry stocks. Later on the 250 g of dry stocks were soaked in 1,250 mL of 80% ethanol and sonicated for 2 min (three times) to extract the constituents. The extract was filtered and residue was extracted again using 1,250 mL of 80% ethanol by the same procedure. The extraction of residue was repeated once again. The extraction was conducted by ultrasonic assisted extraction method for three times and using 3,750 mL of 80% ethanol (3 × 1,250 mL) as a solvent in total. The ethanol extract was evaporated to obtain 13 g of dried ethanol extract (E). The 10 g of dried ethanol extract was suspended in 100 mL of distilled water and liquid-liquid partitioned successively with chloroform, ethyl acetate, and butanol solvent as much as 3 × 100 mL each solvent to obtain 4.03 g of chloroform fraction (C), 0.19 g of ethyl acetate fraction (EA) and 0.80 g of butanol fraction (B). The residue was dried to obtain 4.71 g of aqueous fraction (A). All samples were tested for the antiviral activity against HIV-1 except for aqueous fraction was not tested. The stock solutions were stored in –20 °C freezer until being used further.

Phytochemical analysis

The phytochemical analysis of ethanol extract (E), chloroform fraction (C), ethyl acetate fraction (EA), butanol fraction (B), and aqueous fraction (A) was conducted by thin layer chromatography (TLC) method. Extract and fractions as much as 10 mg were dissolved in 1 mL of methanol. Extract and fractions as much as 3 µL then spotted on silica gel F254 plate (Merck) as stationary phase and chloroform: methanol (Merck) (9:1 v/v) as mobile phase. The spots were identified using H₂SO₄ 10% as a spray reagent and observed under UV 254 and 365 nm.

Analysis of anti HIV potentials of plant extract and fractions

Anti-viral activity (syncytia formation assay): After coculturing the MT-4 cells and persistently infected HIV-1 cells derived from PBMC (MT-4/HIV-1), the syncytia formation occurred on the cultured cells. The next step was to apply the coculture cells into syncytia inhibition assay. Two-fold serial dilution of extract/fractions was prepared (50; 25; 12.5; 6.25; 3.13 µg/L). Upon 30 min incubation at 37 °C, MOLT-4 cells (acute lymphoblastic human leukemia cell line) were added into the culture (2.105 cells/mL for a multiplicity of infection (MOI) of 1/20), and the cocultured cells were incubated at 37 °C for one week (seven days). As a negative control (NC), the MT-4/HIV-1 cells were mixed on cultured MOLT-4 cells and incubated at 37 °C for one week (seven days). After incubation, the number of produced syncytia was microscopically counted. The data were furnished in the percent of inhibition compared with negative control by the below equation.

$$\% \text{ Inhibition} = \left[100 - \left(\frac{\text{NC} - \text{Syncytia on sample}}{\text{NC}} \right) \right] \times 100\%$$

Inhibition activity was also evaluated on half concentration of inhibition (IC₅₀) value. This assay was also done in duplicate replicates to ensure the valid result of the toxicity effect of these anti-viral substances.

Cytotoxicity assay: Follow up the Syncytia assay; the cytotoxicity assay was conducted to measure how potent the tested compound was in terms of antiviral activity against HIV-1. This test only targeted the viable infected cells, not the healthy cells. WST-1 Cell Proliferation Reagent (Roche Applied Science, Switzerland) assay was used in this test by converting tetrazolium salt into a formazan product by viable cells. Serial dilution containing *F. fistulosa* extract/fractions were added into MOLT4 cell culture, using only cell culture without extract/fraction as a positive control and plain RPMI medium as a negative control. The cell culture containing extract/fractions were incubated at 37 °C for seven days. Following the incubation period, the MTT reagent was added into the cell culture and the absorbance was read at 450 nm, measured at two-time points of pre incubation and postincubation at 37 °C for 2 h. The cytotoxicity concentration (CC₅₀) score was measured by assessing in which virus dilution that the 50% viable cells detected. This assay was also done in duplicate replicates to ensure the valid result of the toxicity effect of these antiviral substances.

Statistical analysis: In this study, we were used randomized complete block design (RCBD), a design which each experiment unit was divided into randomized block. This design were applied to analyzed the effect of extract and its fractions that were divided into experimental groups (ethanol extract, butanol fraction, ethyl acetate fraction, and chloroform fraction) and the control group on anti HIV effectiveness. Analysis that used for the above design were using two way ANOVA method, which was a type of parametric statistical test that aims to determine whether there were effect differences/effect on two factors (concentration and compound) that cause variations. If the test result showed different effect on each extract and its fraction, further test was needed for the next test step. One of the test methods is using Tuckey test. This test was used when

the extract and its fraction were different. IC₅₀ and CC₅₀ were calculated using probit regression analysis.

Results

Phytochemical properties from *F. fistulosa* are known to have anti-viral activity, and those subfractions contained flavonoids, terpenoids, and chlorophyll compounds. Genus *Ficus* have several phytochemical compounds such as triterpenoid; flavonoids; sterols; coumarin; and anthocyanins in every part of the plant [24]. Previous study reported that ethanol extract of *F. fistulosa* leaves was exhibited anti hepatitis C virus activity against JFH1a. The phytochemical identification revealed that the ethanol extract was contained flavonoids, terpenoids, and chlorophyll compounds [25]. Meanwhile, alkaloids compounds as antifungal against *Aspergillus fumigatus* and *Candida albicans* were isolated from *F. fistulosa* stem bark [26]. Phytochemical analysis was conducted in this study by TLC analysis. The result showed that *F. fistulosa* leaves was contained flavonoids, terpenoids and chlorophyll compounds as previous reported [25]. The ethanol extract and chloroform fraction showed a similar TLC profile. Chlorophyll was detected in ethanol extract and chloroform fraction under UV light 365 nm which was indicated by red spots. Terpenoids and flavonoids were detected as well after sprayed with H₂SO₄ 10% which was indicated by violet spots and yellow-dark brown spots. Meanwhile, ethyl acetate fraction was dominated with flavonoids and contains minor terpenoids. Butanol fraction and aqueous fraction were not eluted well on silica gel due to polar compounds as a dominant content (Figure 1).

In the present study, one extract and three fractions; ethanol extract, chloroform, ethyl acetate, and butanol fractions extracted from *F. fistulosa* leaves were examined for their anti HIV activities *in vitro* (Table 1). Of four samples, only one sample namely chloroform fraction exhibited the antiviral activities against HIV-1. The ethanol extract showed a low IC₅₀ value calculated as 8.96 µg/mL and a high CC₅₀ of >50 µg/mL while these extract exhibited weak anti HIV activity. Additionally, the chloroform fraction displayed interesting values analyzed as 3.27 and 29.30 µg/mL for IC₅₀ and CC₅₀, respectively. These high values of SI (8.96) detected in these fraction suggested that the chloroform fraction could be utilized safely under the parameters applied in this study. In contrast of ethyl acetate fraction and butanol fraction showed no anti HIV activity, while we detected the high value of CC₅₀ in both fractions (>50).




Figure 1: Thin layer chromatography (TLC) profile of ethanol extract (E), chloroform fraction (C), ethyl acetate fraction (EA), butanol fraction (B), and aqueous fraction (A) of *Ficus fistulosa* leaves using silica gel as a stationary phase and chloroform-methanol (9/1 v/v) as a mobile phase. The TLC was observed under UV light 254 nm (a), 365 nm (b), white light (c), and UV light 365 nm after sprayed with H_2SO_4 10% and heated at 105 °C for 5 min (d).

Table 1: The results of syncytium formation and cytotoxicity assays depicted in calculated IC_{50} , CC_{50} and SI values from *Ficus fistulosa* leaves.

Samples	IC_{50} , $\mu g/ml$	CC_{50} , $\mu g/ml$	SI*
Ethanol extract	8.96	>50	>5.58
Chloroform fraction	3.27	29.30	8.96
Ethyl acetate fraction	23.04	>50	>2.17
Butanol fraction	53.98	>50	>0.97

*SI, selectivity index; CC_{50} value was divide by the IC_{50} value.

Discussion

To date, over 40 currently accessible antiretroviral therapy (ART) drugs have been circulating worldwide converting HIV/AIDS from a lethal infection into a manageable chronic disease [27]. However, due to the high rate of mutations detected in the HIV-1 viral replication cycle, drug resistance mutants have become a big obstacle in HIV-1 treatment accomplishment [28, 29]. The other challenge was to offer more affordable HIV-1 drugs, especially in low-income countries. Therefore, new approaches of using local resources plant-based medicines have been introduced in several Asian countries like Thailand and Indonesia [21, 30].

Indonesia has known for its tropical climate and home of the second-largest biodiversity in the world spreading in over 17,000 islands. Due to its geographical location, the flora of Indonesia displays the Asian, Australian, and native collections. Plant-based medicines using indigenous Indonesian flora have been developed rapidly in recent years. Different studies focusing on antiviral, antibacterial, antimalarial, and antifungal potentials of Indonesian medicinal plants reported promising results to be used further for drugs development. Different parts of the plants like stem, leaves, and roots of the following species of native ginger rhizome, *F. fistulosa*, *Garcinia mangostana*, *Melanolepis multiglandulosa*, and *Melicope latifolia* have been studied intensively in the past years [21, 31, 32].

F. fistulosa is a part of the genus *Ficus* in the family of Moraceae commonly found in Asia and New Guinea. In ancient medicine history, *Ficus* genus species have been used for traditional remedies to cure headaches, breathing problems, diarrhea, cough, toothache, eye infection, and scabies [33]. The extensive pharmacological properties of *Ficus* have been reported including antimicrobial, antioxidant, antiviral, antiparasitic, anti-inflammatory, and anticancer [34].

Wahyuni et al. reported that *F. fistulosa* was efficient in inhibiting the growth of two hepatitis C virus (HCV) strains *in vitro*. The study showed that the ethanol extract from *F. fistulosa* leaves successfully inhibits the virus growth during the initial step of inoculation. The study conducted by Indriati et al. observed that the n-hexane fraction of *F. fistulosa* showed a potent activity as an anti HIV drug candidate. This study result was in accordance with a previously reported study in which n-hexane fraction was exhibited anti HIV activity. Furthermore, crude extract (ethanol extract) showed the highest activity among other samples. The phytochemical content of ethanol extract and chloroform fraction was probably the same due to their similar TLC profile. Both ethanol extract and chloroform fraction contained chlorophyll, terpenoids, and flavonoids compounds which were possible to take a role in their anti HIV activity.

Flavonoids inhibit HIV replication in PBMC in dose dependent manner as it is shown in other herbal medicine known as *Scutellaria baicalensis* [35]. Other type of flavonoids, gallate ester and quercetin 3-O-(2-galloyl) a-L arbinopyranose inhibit the activity of integrase enzyme. Flavonoids also showed inhibition to RT activity [36]. While triterpenoid acts as anti HIV in several step of HIV cycle such as entry inhibitor which block membrane fusion, inhibit HIV enzymes (protease, reverse transcriptase (RT), and integrase), and viral maturation [37]. Further study needs to be conducted for isolation and identification of active compounds from ethanol extract and chloroform fraction.

Conclusions

In conclusion, our recent study reported the novel potential of chloroform fraction from *F. fistulosa* as antiviral against HIV-1 in terms of HIV-1 growth inhibition *in vitro* and proven nontoxic to healthy cells since these fractions contain chlorophyll, terpenoid, and flavonoid which can act as anti HIV.

Acknowledgments: The authors would like to send gratitude towards the staffs in NPMRD group at Institute of Tropical Disease for providing the *Ficus fistulosa* extracts and Airlangga University Hospital for their generous support of PBMC collection in this study.

Research funding: The present study was supported by a Grant-in-Aid from Universitas Airlangga (Penelitian Unggulan Fakultas).

Author contributions: SQK performed HIV *in vitro* analysis and drafted the manuscript, as well as involved in the design and coordination of the study; DWI performed HIV *in vitro* analysis and helped in drafting the manuscript; LT performed plant extraction and phytochemical analysis; AW and N were involved in the design and coordination of the study. All authors read and approved the final version of the manuscript.

Competing interests: The authors declare no potential conflict of interests.

Informed consent: Informed consent was obtained from all individuals included in this study.

Ethical approval: The local Institutional Review Board deemed the study exempt from review.

References

1. WHO. WHO global HIV statistical data. Available from: <https://apps.who.int/gho/data/view.main.22100WHO?lang=en> [Accessed 12 Oct 2020].
2. Bale MJ, Kearney MF. Review: HIV-1 phylogeny during suppressive antiretroviral therapy. *Curr Opin HIV AIDS* 2019;14:188–93.
3. Mansky LM. HIV mutagenesis and the evolution of antiretroviral drug resistance. *Drug Resist Updates* 2002;5:219–23.
4. Fahmi MZ, Sukmayani W, Khairunisa SQ, Witaningrum AM, Indriati DW, Matondang MQY, et al. *RSC Adv* 2016;6:92996–3002.
5. Aung YY, Kristanti A, Khairunisa SQ, Nasronudin N, Fahmi MZ. Inactivation of HIV-1 infection through integrative blocking with amino phenylboronic acid attributed carbon dots. *ACS Biomater Sci Eng* 2020;6:4490–501.
6. Deeks SG. HIV: shock and kill. *Nature* 2012;487:439–40.
7. Kim Y, Anderson JL, Lewin SR. Getting the “kill” into “shock and kill”: strategies to eliminate latent HIV. *Cell Host Microbe* 2018;23:14–26.
8. Richard K, Williams DE, de Silva ED, Brockman MA, Brumme ZL, Andersen RJ, et al. Identification of novel HIV-1 latency-reversing agents from a library of marine natural products. *Viruses* 2018;10:348.
9. Teiten MH, Dicato M, Diederich M. Hybrid curcumin compounds: a new strategy for cancer treatment. *Molecules* 2014;19:20839–63.
10. De Castro S, Camarasa MJ. Polypharmacology in HIV inhibition: can a drug with simultaneous action against two relevant targets be an alternative to combination therapy? *Eur J Med Chem* 2018;150:206–27.
11. Capetti A, Rizzardini G. Choosing appropriate pharmacotherapy for drug-resistant HIV. *Expert Opin Pharmacother* 2019;20:667–78.
12. Society AC. Discovery and development of penicillin. Available from: www.acs.org/content/acs/en/education/whatischemistry/landmarks/flemingpenicillin.html [Accessed 23 Oct 2017].
13. Hare R. New light on the history of penicillin. *Med Hist* 1982;26:1–24.
14. Dias DA, Urban S, Roessner U. A historical overview of natural products in drug discovery. *Metabolites* 2012;2:303–36.
15. Cary DC, Peterlin BM. Natural products and HIV/AIDS. *AIDS Res Hum Retrovir* 2018;34:31–8.
16. Kashman Y, Gustafson KR, Fuller RW, Cardellina JH, McMahon JB, Currens MJ, et al. The calanolides, a novel HIV-inhibitory class of coumarin derivatives from the tropical rainforest tree, *Calophyllum lanigerum*. *J Med Chem* 1992;35:2735–43.
17. McKee TC, Covington CD, Fuller RW, Bokesch HR, Young S, Cardellina JH, et al. Pyranocoumarins from tropical species of the genus *Calophyllum*: a chemotaxonomic study of extract in the National Cancer Institute collection. *J Nat Prod* 1998;61:1252–6.
18. Smith PF, Ogundele A, Forrest A, Wilton J, Salzwedel K, Doto J, et al. Phase I and II study of the safety, virologic effect, and pharmacokinetics/pharmacodynamics of single-dose 3-*o*-(3',3'-dimethylsuccinyl)betulinic acid (bevirimat) against human immunodeficiency virus infection. *Antimicrob Agents Chemother* 2007;51:3574–81.
19. Lansky EP, Paaivilainen HM, Pawlus AD, Newman RA. *Ficus* spp. (fig): ethnobotany and potential as anticancer and anti-inflammatory agents. *J Ethnopharmacol* 2008;119:195–213.
20. Al-Khdhairawi AAQ, Krishnan P, Mai CW, Chung FFL, Leong CO, Yong KT, et al. A bis-benzopyrroloisoquinoline alkaloid incorporating a cyclobutane core and a chlorophenanthroindolizidine alkaloid with cytotoxic activity from *Ficus fistulosa* var. *tengerensis*. *J Nat Prod* 2017;80:2734–40.
21. Indriati DW, Tumewu L, Widyawaruyanti E, Khairunisa SQ. The activities of methanol extract, hexane and ethyl acetate fractions from *Ficus fistulosa* in HIV inhibition *in vitro*. *Res J Pharm Technol* 2020;13:187–90.
22. Vicenzi E, Poli G. Infection of CD4⁺ primary T cells and cell lines, generation of chronically infected cell lines, and induction of HIV expression. *Curr Protoc Im* 2005;69:12.3.1–18.
23. Gyuris A, Vajda G, Földes I. Establishment of an MT4 cell line persistently producing infective HIV-1 particles. *Acta Microbiol Hung* 1992;39:271–9.
24. Ahmad S, Bhatti FR, Khaliq FH, Irshad S, Madni A, Medicine A. A review on the prosperous phytochemical and pharmacological effects of *Ficus carica*. *Int J Bioassays* 2013;2:843–9.
25. Hafid AF, Permanasari AA, Tumewu L, Adianti M, Aoki C, Widyawaruyanti A, et al. Activities of *Ficus fistulosa* leave extract and fractions against Hepatitis C virus. *Procedia Chem* 2016;18:179–84.

26. Subramaniam G, Ang KKH, Ng SB, Buss AD. A benzopyrroloisoquinoline alkaloid from *Ficus fistulosa*. *Phytochem Lett* 2009;2:88–90.
27. AIDSinfo. FDA-approved HIV medicines; 2020. Available from: <https://aidsinfo.nih.gov/understandinghiv-aids/fact-sheets/21/58/fda-approved-hiv-medicines> [Accessed 15 Oct 2020].
28. Menéndez-Arias L. Targeting HIV: antiretroviral therapy and development of drug resistance. *Trends Pharmacol Sci* 2002;23:381–8.
29. Ji H, Sandstrom P, Paredes R, Harrigan PR, Brumme CJ, Rios SA, et al. Are we ready for NGS HIV drug resistance testing? The second “winnipeg consensus” symposium. *Viruses* 2020;12:586.
30. Bunluepuech K, Tewtrakul S. Anti-HIV-1 integrase activity of Thai medicinal plants in longevity preparations. *Songklanakarin J Sci Technol* 2011;33:693–7.
31. Nugraha AS, Keller PA. Revealing indigenous Indonesian traditional medicine: anti-infective agents. *Nat Prod Commun* 2011;6:1953–66.
32. Wahyuni TS, Tumewu L, Permanasari AA, Apriani E, Adianti M, Rahman A, et al. Anti-viral activities of Indonesian medicinal plants in the East Java region against hepatitis C virus. *Virology* 2013;10:259.
33. Chee-Yan C, Sulong SY. A review on the phytochemicals, ethnomedicine uses and pharmacology of *Ficus* species. *Curr Tradit Med* 2016;2:3–17.
34. Sánchez-Valdeolivar CA, Alvarez-Fitz P, Zacapala-Gómez AE, Acevedo-Quiroz M, Cayetano-Salazar L, Olea-Flores M, et al. Phytochemical profile and antiproliferative effect of *Ficus crocata* extract on triple-negative breast cancer cells. *BMC Compl Med Ther* 2020;20:191.
35. Ohtake N, Nakai Y, Yamamoto M, Sakakibara I, Takeda S, Amagaya S, et al. Separation and isolation methods for analysis of the active principles of Sho-saiko-to (SST) oriental medicine. *J Chromatogr* 2004;15:D84–90.
36. Kim HJ, Woo E-R, Shin C-G, Park H. A new flavonol glycoside gallate ester from *Acer okamotoanum* and its inhibitory activity against human immunodeficiency virus-1 (HIV-1) integrase. *J Nat Prod* 1998;61:145–8.
37. Han B, Peng Z. Anti-HIV triterpenoid components. *J Chem Pharmaceut Res* 2014;6:438–43.

paper focus

ORIGINALITY REPORT

12%

SIMILARITY INDEX

8%

INTERNET SOURCES

9%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	biomedres.us Internet Source	1%
2	Wang, Y.. "Studies on bioactivities of tea (Camellia sinensis L.) fruit peel extracts: Antioxidant activity and inhibitory potential against @a-glucosidase and @a-amylase in vitro", Industrial Crops & Products, 201205 Publication	1%
3	respiratory-research.biomedcentral.com Internet Source	1%
4	www.nature.com Internet Source	1%
5	www.frontiersin.org Internet Source	1%
6	ar.iiarjournals.org Internet Source	1%
7	"Anti-Viral Metabolites from Medicinal Plants", Springer Science and Business Media LLC, 2024 Publication	1%

8

Submitted to Higher Education Commission
Pakistan

Student Paper

<1 %

9

Nishino, Y.. "Major core proteins, p24s, of human, simian, and feline immunodeficiency viruses are partly expressed on the surface of the virus-infected cells", Vaccine, 1992

Publication

<1 %

10

Brenda De la Cruz-Concepción, Mónica Espinoza-Rojo, Patricia Álvarez-Fitz, Berenice Illades-Aguiar et al. "Cytotoxicity of Ficus Crocata Extract on Cervical Cancer Cells and Protective Effect against Hydrogen Peroxide-Induced Oxidative Stress in HaCaT Non-Tumor Cells", Plants, 2021

Publication

<1 %

11

taeppo.at.webry.info

Internet Source

<1 %

12

mdpi.com

Internet Source

<1 %

13

Jinfeng Yang, Yong Soo Kwon, Myong Jo Kim. "Antimicrobial activity and active compounds of a Rhus verniciflua Stokes extract", Zeitschrift für Naturforschung C, 2018

Publication

<1 %

14

e-journal.unair.ac.id

Internet Source

<1 %

15	f1000research.com Internet Source	<1 %
16	virologyj.biomedcentral.com Internet Source	<1 %
17	HIROTO HARA. "Tetrahydronaphthalene Lignan Compounds as Potent Anti-HIV Type 1 Agents", AIDS Research and Human Retroviruses, 05/20/1997 Publication	<1 %
18	M. Yunoki. "Production of infectious particles from defective human immunodeficiency virus type 1 (HIV-1)-producing cell clones by superinfection with infectious HIV-1", Archives of Virology, 03/1991 Publication	<1 %
19	Masfria , Urip Harahap, Maratua Pandapotan Nasution, Syafruddin Ilyas. "The activity of <i>Rhaphidophora pinnta</i> Lf. Schott leaf on MCF-7 cell line", Advances in Biological Chemistry, 2013 Publication	<1 %
20	retrovirology.biomedcentral.com Internet Source	<1 %
21	www.freeradicalscience.com Internet Source	<1 %

22	Odilon Djakpo. "Rhus chinensis and Galla Chinensis - folklore to modern evidence: review", <i>Phytotherapy Research</i> , 06/17/2010 Publication	<1 %
23	ijpsr.com Internet Source	<1 %
24	mdpi-res.com Internet Source	<1 %
25	www.liebertpub.com Internet Source	<1 %
26	Yang, Zhi-Gang, Liu-Nan Jia, Yan Shen, Atsuko Ohmura, and Susumu Kitanaka. "Inhibitory Effects of Constituents from Euphorbia lunulata on Differentiation of 3T3-L1 Cells and Nitric Oxide Production in RAW264.7 Cells", <i>Molecules</i> , 2011. Publication	<1 %
27	scholar.sun.ac.za Internet Source	<1 %
28	www.mdpi.com Internet Source	<1 %
29	www.smujo.id Internet Source	<1 %
30	Ikuta, K.. "Expression of human immunodeficiency virus type 1 (HIV-1) gag antigens on the surface of a cell line	<1 %

persistently infected with HIV-1 that highly expresses HIV-1 antigens", *Virology*, 198906

Publication

31

Sathit Panvilai, Chanita Napaswad, Jitra Limthongkul, Radeekorn Akkarawongsapat. "Aqueous Extracts of Thai Medicinal Plants Possess anti-HIV-1 Activity", *Journal of Herbs, Spices & Medicinal Plants*, 2020

Publication

32

Subramoniam, Appian. "Anti-Diabetes Mellitus Plants", *Plants with Anti-Diabetes Mellitus Properties*, 2016.

Publication

33

Khumoekae Richard, David Williams, E. de Silva, Mark Brockman, Zabrina Brumme, Raymond Andersen, Ian Tietjen. "Identification of Novel HIV-1 Latency-Reversing Agents from a Library of Marine Natural Products", *Viruses*, 2018

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

<1 %

<1 %

<1 %