

The Role of transdermal carbon dioxide

by Aditiawarman Aditiawarman

Submission date: 06-May-2024 11:10AM (UTC+0800)

Submission ID: 2371815920

File name: The_Role_of_transdermal_carbon_dioxide.pdf (513.11K)

Word count: 4991

Character count: 27060

The role of transdermal carbon dioxide on changes in malondialdehyde levels as a marker of ischemia-reperfusion injury in patients with placenta accreta spectrum underwent temporary abdominal aortic cross-clamping as an adjunct procedure during cesarean hysterectomy

Hari Daswin Pagehgiri^{1*}, Ito Puruhito¹, Aditiawarman², Pudji Lestari³, Yan Efrata Sembiring¹, Dhihintia Jiwangga¹, Arief Rakhman Hakim¹, Rozi Aditya Aryananda²

¹Department of Thoracic, Cardiac, and Vascular Surgery, Dr. Soetomo General Academic Hospital, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia;

²Department of Obstetrics & Gynecology, Dr. Soetomo General Academic Hospital, Universitas Airlangga, Surabaya, East Java, Indonesia;

³Department of Public Health and Preventive Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia;

*Corresponding author:

Hari Daswin Pagehgiri;
Department of Thoracic, Cardiac and Vascular Surgery, Dr. Soetomo General Academic Hospital, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia;
hari.daswin.pagehgiri-2016@fk.unair.ac.id

Received: 2022-06-12

Accepted: 2022-07-29

Published: 2022-08-31

ABSTRACT

Background: Temporary abdominal aortic cross-clamping is often applied as an adjunct procedure to control bleeding in patients with placenta accreta spectrum during cesarean hysterectomy. It is claimed to reduce the blood loss need for transfusion and improve visualization of the operating field. After the cross-clamp is removed, the tissue distal to the occlusion, which was initially in an ischemic state, gets a sudden blood flow causing ischemia-reperfusion injury due to the release of ROS. Transdermal administration of carbon dioxide is expected to reduce the release of ROS through the Bohr Effect to protect against ischemia-reperfusion injury, which can be seen from the level of malondialdehyde.

Method: This experimental study recruited all patients with placenta accreta spectrum who underwent temporary abdominal aortic cross-clamping during cesarean hysterectomy from January to June 2022. Subjects were divided into control groups and treatment groups. The treatment group was given transdermal CO₂ immediately after the aortic cross-clamp was removed. The plasma MDA levels were examined before and after aortic cross-clamping.

Results: The number of subjects in each group was 7 subjects. There was an increase in MDA levels from 19.779±0.870nmol/ml to 21.104±1.053nmol/ml after cross-clamp in all groups, with an average increase of 1.325±0.801nmol/ml ($p=0.00$). The treatment group that received transdermal CO₂ had a lower tendency to increase MDA levels, 1.063±0.803nmol/ml, compared to the control group at 1.586±0.766nmol/ml.

Conclusion: There was an increase in MDA levels as a predictor of ischemia-reperfusion injury in patients undergoing temporary abdominal aortic cross-clamping. The administration of transdermal CO₂ tends to suppress ischemia-reperfusion injury.

Keywords: malondialdehyde, placenta accreta spectrum, temporary abdominal aortic cross-clamping, transdermal carbon dioxide.

Cite This Article: Pagehgiri, H.D., Puruhito, I., Aditiawarman., Lestari, P., Sembiring, Y.E., Jiwangga, D., Hakim, A.R., Aryananda, R.A. 2022. The role of transdermal carbon dioxide on changes in malondialdehyde levels as a marker of ischemia-reperfusion injury in patients with placenta accreta spectrum underwent temporary abdominal aortic cross-clamping as an adjunct procedure during cesarean hysterectomy. *Bali Medical Journal* 11(2): 1023-1029. DOI: 10.15562/bmj.v11i2.3646

INTRODUCTION

The placenta accreta spectrum is defined as the abnormal invasion of trophoblast on one part or all parts of the placenta into the myometrial layer of the uterine wall, which results in a placenta that is morbidly adherent to the uterine wall. As a consequence of the morbidly adherent placenta, the risk of hemorrhage, which prompts blood transfusion and peripartum

hysterectomy, increases.¹ According to a systematic review by Jauniaux et al, the prevalence of placenta accreta spectrum is 0,17% (95% CI, 0,14 – 0,19), with 0,5 (95% CI, 0,3 – 0,36) per 1000 birth and 0,3 (95% CI, 0,2 – 0,4) per 1000 birth as the prevalence for adherent grade and invasive grade, respectively. The incidence of hemorrhage requiring blood transfusion is 46,9% (95% CI, 34 – 66,4), whereas the incidence of peripartum hysterectomy is

52,2% (95% CI, 38,3 – 66,4). Estimated maternal mortality on placenta accreta spectrum is 0,05% (95% CI, 0,06 – 0,69).² In our center at Dr. Soetomo General Academic Hospital, Surabaya, Indonesia, placenta accreta spectrum disorder was increased in the last 6 years. The prevalence was 1% in 2015, 4% in 2017, and increased to 6% in 2019.^{3,4} Extensive bleeding in the placenta accreta spectrum is hypothesized to be caused by the development of

extensive intrapelvic vascular anastomosis by arteries to the gravid uterus.

A clear and dry surgical field is important to properly perform surgery and minimize operative blood loss. This can be achieved by reducing uteroplacental blood flow. Cross-clamping of the abdominal aorta is one of the techniques to reduce uteroplacental blood flow.⁵ Temporary aortic cross-clamp is often applied as an adjunct procedure to control bleeding in patients with placenta accreta during cesarean hysterectomy. It is claimed to reduce the blood loss need for transfusion and improve visualization of the operating field. After the cross-clamp is removed, the tissue distal to the occlusion, which was initially in an ischemic state, gets a sudden blood flow causing ischemia-reperfusion injury due to the release of ROS.

Visceral ischemia and reperfusion injury are significant side effects of the aortic cross-clamp.⁶ In a state of prolonged ischemia, tissue reperfusion can be accompanied by inflammation that results in tissue cell damage and death.⁷ During aortic cross-clamping, an ischemic condition occurs. After the aortic cross-clamp is opened, the distal tissue from the occlusion, which was initially in an ischemic state, gets sudden blood flow (reperfusion), causing ischemia-reperfusion injury.

The most widely accepted molecular mechanism underlying the occurrence of ischemia-reperfusion injury is an increase in Reactive Oxygen Species (ROS).⁸ Tissue damage that occurs due to reperfusion begins with the release of ROS. The imbalance between increased ROS and the ability to neutralize free radicals by antioxidant defense causes oxidative stress.⁸ ROS will then affect lipid degradation, and this can be seen in the final product, such as the detection of Malondialdehyde.⁹

Malondialdehyde (MDA), a three-carbon chain with a low molecular weight aldehyde, is one of the end products of the peroxidation of polyunsaturated fatty acids in cells.⁹⁻¹¹ Malondialdehyde levels are generally recognized as markers of oxidative stress.⁹ ROS causes cell damage by negatively affecting antioxidant defense mechanisms by reducing superoxide dismutase (SOD) levels and increasing

MDA levels by inflammatory processes mediated by tumor necrosis factor-alpha (TNF- α).¹² The increase in free radicals leads to an overproduction of MDA.

Several studies have demonstrated a significant effect of using transdermal CO₂ on microcirculation.^{13,14} Carbon dioxide therapy refers to the transdermal application of CO₂ for therapeutic purposes with the effect of increasing O₂ pressure in tissues, known as the 'Bohr effect'.^{14,15} Following the 'Bohr Effect', there is an increase in oxygen delivery to the tissues due to CO₂ and Hydrogen, which shifts the Oxygen-Hemoglobin dissociation curve, thus forcing O₂ to be released from hemoglobin and consequently increasing the amount of O₂ in the tissues.¹⁶ CO₂ is believed to be a strong inhibitor of ROS through inhibition of the activity of NADPH-oxidase.¹⁷ Transdermal administration of CO₂ is expected to reduce ROS production through the Bohr Effect so that there is protection against the incidence of ischemia-reperfusion injury, which can be seen from MDA levels.

The purpose of this study was to determine changes in MDA levels as a predictor of ischemia-reperfusion injury in patients undergoing temporary abdominal aortic cross-clamping and to determine the effect of transdermal administration of CO₂ as a protective factor for ischemia-reperfusion injury.

METHOD

Study design

This is a clinical trial study of 40 patients with placenta accreta spectrum who underwent temporary abdominal aortic cross-clamping. Subjects were divided into control groups and treatment groups. The treatment group was given transdermal CO₂ immediately after the aortic cross-clamp was removed. Randomization is done by order. Blinding is performed on the patient and the outcome assessor. The patient did not know the treatment received. MDA levels were checked by laboratory staff who did not know the treatment group. The ethical clearance was approved by Ethics Committee in Health Research Dr. Soetomo Academic General Hospital number 0319/KEPK/XII/2021 on December 1st, 2021.

Subject

This study recruited all patients with placenta accreta spectrum who underwent temporary abdominal aortic cross-clamp during cesarean hysterectomy from January 2022 to June 2022 at Dr. Soetomo General Academic Hospital Surabaya, Indonesia. The inclusion criteria were patients with placenta accreta who were diagnosed with placental accreta index (PAI) based on the results of ultrasonography performed by obstetrics and gynecology specialists at Dr. Hospital. Soetomo Surabaya underwent temporary abdominal aortic cross-clamping as an adjunct procedure during a cesarean hysterectomy. Patients with severe comorbid disorders such as chronic kidney disease, heart disease, patients with septic shock and sepsis, patients with peripheral artery disease, and patients with pneumonia covid-19 were excluded.

Treatment Protocol

The patients were put under general anesthesia. The blood samples for MDA I (basal) were collected after anesthesia. The surgical approach was obtained by median laparotomy. A midline or transversal fundal incision is performed to deliver the baby. The umbilical cord was clamped and ligated at the placenta end, and the hysterotomy wound was clamped with forceps. The obstetrician then evaluated the segment involved by the abnormality accreta to determine whether to perform a hysterectomy or a uterine conservative resection surgery. The temporary abdominal aortic cross-clamping was performed by a cardiovascular surgeon if a hysterectomy was planned.⁴⁹

The retroperitoneum between the inferior mesenteric artery and aortic bifurcation was opened. The uterus was pushed downward for a better surgical field. The intestine was retracted by a Deaver retractor and big gauze to expose retroperitoneal fascia. After dissecting the tissue surrounding the aorta, a right-angled was used to place a vessel loop under the aorta avoiding injury to lumbar vessels. Unfractionated heparin 2500 I.U. was administered through the peripheral intravenous line. The vessel loop was used to lift the aorta. The aorta was clamped with a professional atraumatic flexible cardiovascular clamp.

44

After completion of the hysterectomy procedure, the aortic cross-clamp was removed, and bleeding was evaluated. Transdermal CO₂ is immediately given to the thumb with a D'Oxyva device at a dose of 16g for 2-5 minutes. Then a second MDA II blood sample (reperfusion) was collected after 60 minutes after reperfusion.

Laboratory test

The blood was collected preoperative (basal) and 60 minutes after abdominal aortic cross-clamp off, and the plasma MDA levels were measured. Blood samples were centrifugated to separate the serum from the blood. The separated serum was then stored at -70oC for later examination of MDA using the thiobarbituric acid method. The MDA concentration was measured by the spectrophotometric method for the reaction with thiobarbituric acid reactive substances (TBARS).

Data collection and analysis

The data were presented in mean and standard deviation (S.D.) and then tested using the Kolmogorov-Smirnov normality test to assess the normality of the data. Paired t-test was used to determine the differences in MDA level before and after the temporary aortic cross-clamp. Independent t-test (2-tailed) was used for data with normal distribution, and Mann-Whitney (2-tailed) was used for data with non-normal distribution to compare the difference between the control and treatment groups. All data analysis was carried out in SPSS v23.

RESULT

From January to June 2022, there were 38 patients diagnosed with placenta accreta based on preoperative ultrasonography, and 15 patients (39,4%) underwent temporary abdominal aortic cross-clamping as an adjunct procedure for bleeding control during cesarean hysterectomy. One patient was excluded because of comorbid pneumonia covid-19. The number of subjects who were successfully recruited was 14 patients. The number of subjects who were successfully recruited was 14 patients. The demographic characteristics of the patients are presented in Table 1. It can be seen that

Table 1. Patient Characteristic.

Demographic	Treatment group (n=7) Mean (S.D.)	Control group (n=7) Mean (S.D.)
Age	31.57 (3.55) years	36.14 (3.34) years
Weight	69.00 (5.77) kg	66.14 (7.22) kg
Height	154.29 (3.15) cm	155.71 (2.56) cm
Number of Pregnancy	4.0 (1.73) times	4.14 (1.46) times
History of cesarean delivery	1.71 (0.76) times	1.71 (0.49) times

Table 2. Operative Variable.

Operative Variable	Treatment group (n=7)	Control group (n=7)
Placenta invasion type*		
Accreta	0	0
Increta	2 (28.57%)	2 (28.57%)
Percreta	5 (71.43%)	5 (71.43%)
FIGO grading*		
Grade 1	1 (14.29%)	0
Grade 2	1 (14.29%)	0
Grade 3A	2 (28.57%)	2 (28.57%)
Grade 3B	2 (28.57%)	4 (57.14%)
Grade 3C	1 (14.29%)	1 (14.29%)
PAS staging*		
PAS 1	2 (28.57%)	1 (14.29%)
PAS 2	1 (14.29%)	2 (28.57%)
PAS 3	4 (57.14%)	4 (57.14%)
Segment involved*		
Segment 1	2 (28.57%)	2 (28.57%)
Segment 2	5 (71.43%)	5 (71.43%)
Aortic Cross-Clamp duration**	61.43 (15.74) minutes	70.71 (28.79) minutes
Estimated Blood Loss**	2,435.71 (1,367.70) ml	2,807.14 (1,241.78) ml

*n(%)

**mean(SD)

Table 3. Malondialdehyde level in all group.

n=14	Mean	SD	P
MDA 1 (basal), nmol/ml	19.779	0.870	0.00*
MDA 2 (reperfusion), nmol/ml	21.104	1.053	

*Paired-T test (2-tailed)

Table 4. Malondialdehyde level between group.

	Treatment group (n=7) Mean(S.D.)	Control group (n=7) Mean(S.D.)	P
MDA 1 (basal), nmol/ml	20.220 (0.543)	19.377 (0.943)	0.053*
MDA 2 (reperfusion), nmol/ml	21.283 (1.034)	20.923 (1.121)	0.544*
ΔMDA, nmol/ml	1.063 (0.803)	1.586 (0.766)	0.402**

*Independent-T test (2-tailed)

**Mann-Whitney (2-tailed)

both control and treatment groups have comparable characteristics, so it can be concluded that both groups have the same prognostic and confounding factors.

Based on the surgical variables in Table 2, it appears that both control and

treatment groups also have comparable characteristics, so it can be concluded that both groups have the same prognostic and confounding factors. The vascular control with temporary abdominal aortic cross-clamping was mostly used in percreta

Table 5. Secondary outcome.

	Treatment group (n=7) Mean(S.D.)	Control group (n=7) Mean(S.D.)	P
Intubation time, hours	14.82 (8.42)	17.89 (19.65)	0.803**
Hospital length of stay, days	7.86 (3.13)	6.71 (2.14)	0.531**
Mortality	0	0	-
Hemoglobin, gram/dL	9.46 (1.13)	8.04 (1.83)	0.685**
Haematocrit, %	29.04 (3.61)	25.30 (6.42)	0.779**
White Blood Cell count, 10 ³ /uL	17,695 (6,138)	14,641 (7,588)	0.424*
Platelet count, 10 ³ /uL	173,000 (81,973)	145,428 (108,669)	0.602*
Albumin, gram/dL	2.86 (0.28)	2.64 (0.37)	0.224*

*Independent-T test (2-tailed)



**Mann - Whitney (2-tailed)

invasion type, high FIGO grade, and PAS 3 stage, with most in segment 2 involved. The mean estimated blood loss (EBL) was 2,435.71±1,367.70 mL in treatment group and 2,807.14±1,241.78) mL in control group. The aortic cross-clamp duration was 61.43±15.74 minutes in the treatment group and 70.71±28.79 minutes in the control group.

In all groups, it is shown that there was an increase in MDA levels from 19.779±0.870nmol/ml to 21.104±1.053nmol/ml after cross-clamp with an average increase of 1.325±0.801nmol/ml ($p=0.00$) (Table 3). It was concluded that there was an increase in MDA levels as a predictor of ischemia-reperfusion injury in patients who underwent temporary abdominal aortic cross-clamping.

Based on the results shown in Table 4, the treatment group that received transdermal CO₂ had a lower tendency to increase MDA levels, 1.063±0.803nmol/ml, compared to the control group at 1.586±0.766nmol/ml. However, based on statistical tests, the results were not significant. Changes in MDA levels between groups can be seen in Figure 1 until Figure 3.

Table 5 shows the mean intubation time and length of stay in the hospital. There were no statistical differences in the length of intubation and hospitalization in the two groups. There were no patients who died during the study period. Based on the results of routine postoperative laboratory examinations, there were no differences in hemoglobin levels, white blood cell counts, platelets, and albumin levels.

**Figure 1.** MDA level in the control group.**Figure 2.** MDA level in the treatment group.




Figure 3. Boxplot of MDA level between the control and the treatment groups on basal and reperfusion examination (top), and boxplot of change in MDA level in control and treatment groups (bottom).

DISCUSSION

Abdominal aortic occlusion has been performed to control severe hemorrhage in patients with the placenta accreta spectrum. This can be achieved via a direct external approach and endovascular approach.^{5,18,19} Temporary cross-clamping of the abdominal aorta is one of the methods to control severe hemorrhage.²⁰ Implementation of abdominal aortic cross-clamping to control operative blood loss was first described in a case report by Chou et al. The technique starts with the separation of the inferior abdominal aorta (IAA) from the inferior vena cava between the fourth lumbar and aortic bifurcation. It was also emphasized that the procedures were performed by a team consisting of one cardiovascular surgeon and two obstetricians.⁵ Yoshida et al. emphasized that temporary clamping of the abdominal aorta should be done as close to the renal artery as possible.²¹

Based on the results of this study, it was found that there was an increase in MDA levels before and after the abdominal aortic temporary cross-clamping was performed. It can be concluded that MDA can be used as a predictor of ischemia-reperfusion injury after temporary abdominal aortic cross-clamping. The ischemia-reperfusion injury occurs when the blood supply to an organ is disrupted (ischemia) and then restored (reperfusion), which causes the release of Reactive Oxygen Species (ROS) from the mitochondria.²² Reperfusion after ischemia produces ROS, mitochondrial failure, endothelial dysfunction, and sterile inflammation. Under physiological conditions, mitochondria produce

small amounts of ROS through electron leakage.²³ Cells that are ischemic have lower antioxidants.⁷ When exposed to ischemia, the mitochondrial complex is damaged, resulting in excessive ROS generation beyond the antioxidant restoring capacity after reperfusion. ROS, in turn, causes damage to membrane lipids, proteins, and nucleic acids that can lead to cell death.²³

ROS is not very stable, so the measurement is carried out indirectly by the release of MDA as a result of ROS reactions to lipids degradation that is found in cell membranes. Malondialdehyde levels are generally recognized as a marker of oxidative stress.⁹ Plasma MDA levels increased significantly within 1 hour after the onset of reperfusion in all patients who had successful revascularization. MDA levels then slowly decreased to near basal values after 4 hours of revascularization.^{8,24}

In the condition of ischemia-reperfusion injury, there is an increase in NADPH-oxidase activity and a decrease in antioxidants.^{7,22} NADPH-oxidase will cause the conversion of O_2 to O_2^- . O_2^- will be converted into H_2O_2 by Superoxide dismutase (SOD) and then into H_2O by Catalase or Glutathione peroxidase (GPX). The most important radicals or pro-oxidant molecules involved in the disease process are superoxide (O_2^-), hydroxyl radicals ($H.O.$), and 1O_2 .^{7,10} The imbalance between increased ROS and the ability to neutralize free radicals by antioxidant defenses causes oxidative stress with the result of increased ROS.⁸ ROS is not very stable at the time of examination. So that the examination is carried out indirectly, namely the formation of MDA as a result of ROS reactions to cell lipids

that are abundant in cell membranes. Malondialdehyde levels are generally recognized as markers of oxidative stress.⁹

Efforts are currently being made to reduce the incidence of ischemia-reperfusion injury. one of them is by giving transdermal CO_2 . Several studies have demonstrated a significant effect of using transdermal CO_2 on microcirculation.¹⁵ Finzgar *et al.* found that transdermal CO_2 had a significant effect on skin microcirculation, as seen from the increase in laser Doppler flux.¹⁴ The use of transdermal CO_2 in patients with intermittent claudication significantly increased walking distance, peripheral systolic pressure, and pO_2 .¹³ In patients with Diabetic Foot Ulcers, the use of transdermal CO_2 showed an increase in dermal microcirculation based on perfusion index and tissue oxygenation, therefore helping in the wound healing process.²⁵ Damanik & Puruhito found that administration of transdermal CO_2 significantly increased $TcPCO_2$ and increased tissue perfusion index 5.6 times in patient with placenta accreta spectrum underwent temporary abdominal aortic cross-clamping without the presence of adverse events or complications.²⁶ Following the 'Bohr Effect', there is an increase in oxygen delivery to tissues due to CO_2 and Hydrogen which shifts the Oxygen-Hemoglobin dissociation curve so that O_2 is released from hemoglobin and consequently increases the amount of O_2 in the tissues.^{15,16} CO_2 is believed to be a strong inhibitor of ROS through inhibition of the activity of NADPH-oxidase.¹⁷ Based on the results of this study, transdermal CO_2 administration tends to reduce the

incidence of ischemia-reperfusion injury after temporary abdominal aortic cross-clamping seen from the less change of malondialdehyde in the treatment group than in the control group. However, it is not statistically significant.

The population of this study was patients with placenta accreta spectrum who underwent aorta abdominal temporary cross-clamping during cesarean hysterectomy. The non-significant MDA concentration in the two groups may also be due to lipid peroxidation during abdominal aortic temporary clamping surgery in addition to the distal ischemia of the clamp, also due to high oxidative stress conditions in pregnant patients. Research by Khan *et al.* showed that pregnant women experienced an increase in oxidative stress and a decrease in antioxidants, with the mean MDA levels found to be significantly higher in pregnant women compared to non-pregnant women.²⁷ Antioxidant activities such as SOD and Catalase was found to be significantly lower in pregnant women compared to non-pregnant women.^{27,28} Oxidative stress in pregnancy may be a result of increased oxygen consumption and metabolism due to energy and oxygen requirements for fetal growth and development.^{27,28}

The limitations obtained during this study were that the research subjects were limited and carried out during the pandemic era. Further research needs to be done with a larger sample size to obtain greater statistical significance.

CONCLUSION

This study concluded that there was an increase in MDA levels as a predictor of ischemia-reperfusion injury in patients undergoing temporary abdominal aortic cross-clamping. The administration of transdermal CO₂ tends to suppress ischemia-reperfusion injury. Suggestions for further research are: further research needs to be done with a larger sample size to get greater statistical significance. It is necessary to conduct another research to determine the effect of transdermal CO₂ administration on other subjects who are not pregnant or under conditions of oxidative stress.

ETHICAL CLEARANCE

The ethical clearance of this study was approved by Ethics Committee in Health Research Dr. Soetomo Academic General Hospital number 0319/KEPK/XII/2021 on December 1st, 2021.

CONFLICT OF INTEREST

There is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

FUNDING

This research was not funded by a grant from any organization.

AUTHOR CONTRIBUTION

All authors contributed equally to the manuscript. All authors read and approved the final version of the manuscript.

REFERENCE

1. Society of Gynecologic Oncology; American College of Obstetricians and Gynecologists and the Society for Maternal-Fetal Medicine, Cahill AG, et al. Placenta Accreta Spectrum. *Am J Obstet Gynecol.* 2018;219(6):B2-B16. doi:10.1016/j.ajog.2018.09.042.
2. Jauniaux E, Bunce C, Grønbeck L, Langhoff-Roos J. Prevalence and main outcomes of placenta accreta spectrum: a systematic review and meta-analysis. *Am J Obstet Gynecol.* 2019;221(3):208-218. doi:10.1016/j.ajog.2019.01.233.
3. Aditiawarman AR, Akbar MA, Darmawan E, and Sulistyono A. Epidemiology and the Emerging of PASD in Indonesia. 2020. p284-287.
4. Aryananda RA. Resurgence of placenta accreta in Indonesia. *Maj. Obstet. Ginekol.* 2019;(26)98. DOI:10.20473/mog.V26I32018.98-99.
5. Chou MM, Chen MJ, Su HW, et al. Vascular control by infrarenal aortic cross-clamping in placenta accreta spectrum disorders: description of technique. *BJOG.* 2021;128(6):1030-1034. doi:10.1111/1471-0528.16605.
6. Zammert M, Gelman S. The pathophysiology of aortic cross-clamping. *Best Pract Res Clin Anaesthesiol.* 2016;30(3):257-269. doi:10.1016/j.bpa.2016.07.006.
7. Wu MY, Yang GT, Liao WT, et al. Current Mechanistic Concepts in Ischemia and Reperfusion Injury. *Cell Physiol Biochem.* 2018;46(4):1650-1667. doi:10.1159/000489241.
8. Ismael A, Lavado R, Smith RS, et al. Effects of Limb Revascularization Procedures on Oxidative Stress. *J Surg Res.* 2018;232:503-509. doi:10.1016/j.jss.2018.07.024.
9. Gawel S, Wardas M, Niedworok E, Wardas P. Dialdehyd malonowy (MDA) jako wskaźnik

procesów peroksydacji lipidów w organizmie [Malondialdehyde (MDA) as a lipid peroxidation marker]. *Wiad Lek.* 2004;57(9-10):453-455.

10. Grotto D. et al. Importance of the lipid peroxidation biomarkers and methodological aspects for malondialdehyde quantification. *Quim Nova.* 2009;32. p169-174. DOI:10.1590/S0100-40422009000100032.
11. Satria, D., Sembiring, Y. E., Kurniasari, N. & Lestari, P. Combination Therapy of N-Acetylcysteine and Simvastatin Reduce Malondialdehyde (MDA) Levels in Animal Models with Ischemic/Reperfusion Injury (IRI). *Annals of the Romanian Society for Cell Biology.* 2021;25(6). p18712-18719.
12. Pergel A, Tümkaya L, Demiral G, et al. The protective effects of adalimumab on intestinal injury induced with infrarenal aortic occlusion. Infrarenal aort oklüzyonu ile oluşan intestinal hasarında adalimumab'ın koruyucu etkisi. *Ulus Travma Acil Cerrahi Derg.* 2020;26(3):366-372. doi:10.14744/tjtes.2019.59607.
13. Fabry R, Monnet P, Schmidt J, et al. Clinical and microcirculatory effects of transcuteaneous CO₂ therapy in intermittent claudication. Randomized double-blind clinical trial with a parallel design. *Vasa.* 2009;38(3):213-224. doi:10.1024/0301-1526.38.3.213.
14. Finzgar M, Melik Z, Cankar K. Effect of transcuteaneous application of gaseous carbon dioxide on cutaneous microcirculation. *Clin Hemorheol Microcirc.* 2015;60(4):423-435. doi:10.3233/CH-141898.
15. Sakai Y, Miwa M, Oe K, et al. A novel system for transcuteaneous application of carbon dioxide causing an "artificial Bohr effect" in the human body. *PLoS One.* 2011;6(9):e24137. doi:10.1371/journal.pone.0024137.
16. Hall JE and Hall ME. Transport of Oxygen and Carbon Dioxide in Blood and Tissue Fluids. *Guyt Hall Textb Med Physiol.* 2016;(14). p524.
17. Bolevich S, Kogan AH, Zivkovic V, et al. Protective role of carbon dioxide (CO₂) in generation of reactive oxygen species. *Mol Cell Biochem.* 2016;411(1-2):317-330. doi:10.1007/s11010-015-2594-9.
18. Chen L, Wang X, Wang H, Li Q, Shan N, Qi H. Clinical evaluation of prophylactic abdominal aortic balloon occlusion in patients with placenta accreta: a systematic review and meta-analysis. *BMC Pregnancy Childbirth.* 2019;19(1):30. Published 2019 Jan 15. doi:10.1186/s12884-019-2175-0.
19. Riley DP, Burgess RW. External abdominal aortic compression: a study of a resuscitation manoeuvre for postpartum haemorrhage. *Anaesth Intensive Care.* 1994;22(5):571-575. doi:10.1177/0310057X9402200512.
20. Chou MM, Ke YM, Wu HC, et al. Temporary cross-clamping of the infrarenal abdominal aorta during cesarean hysterectomy to control operative blood loss in placenta previa increta/percreta. *Taiwan J Obstet Gynecol.* 2010;49(1):72-76. doi:10.1016/S1028-4559(10)60013-7.
21. Yoshida A, Miura K, Hasegawa Y, et al. A case of placenta previa accreta successfully treated by

- open infrarenal aortic clamping. *Acta Medica Nagasakiensia*. 2013;58, p93-95.
22. Chouchani ET, Pell VR, James AM, et al. A Unifying Mechanism for Mitochondrial Superoxide Production during Ischemia-Reperfusion Injury. *Cell Metab*. 2016;23(2):254-263. doi:10.1016/j.cmet.2015.12.009.
 23. Slegtenhorst BR, Dor FJ, Rodriguez H, Voskuil FJ, Tullius SG. Ischemia/reperfusion Injury and its Consequences on Immunity and Inflammation. *Curr Transplant Rep*. 2014;1(3):147-154. doi:10.1007/s40472-014-0017-6.
 24. Rabl H, Khoschorur G, Colombo T, Tatzber F, and Esterbauer H. Human plasma lipid peroxide levels show a strong transient increase after successful revascularization operations. *Free Radic. Biol. Med*. 1992;13. p281-288. [https://doi.org/10.1016/0891-5849\(92\)90175-G](https://doi.org/10.1016/0891-5849(92)90175-G).
 25. Rahmi C dan Puruhito I. Pengaruh Pemberian Transdermal CO₂ Terhadap Hasil Perawatan Luka Kaki Diabetik Tipe I & II. *IR - Perpust. Univ. AIRLANGGA* 3. 2018.
 26. Damanik, G., Puruhito, I., Aryananda, R. A., Lestari, P. & Soebroto, H. Transdermal CO₂ Increases Perfusion Index in Patients with Placenta Accreta Following Temporary Abdominal Aortic Cross-Clamping. *Ann. Rom. Soc. Cell. Biol*. 2021;25. p19918-19923.
 27. Khan S, Khan S, Khan MM, Alam R, and Khan A. Estimation of The Status of MDA Levels and SOD Activity in Pregnant Women. *Biochem. Cell. Arch*. 2019;19, p169-173.
 28. Sharma P, Prabha SS, Kumar P, and Sharma R. Estimation of malondialdehyde and catalase in pregnant & non-prenant women. *Santosh Univ J Heal Sci*. 2020;6. p21-25. DOI:10.18231/j.sujhs.2020.006.



This work is licensed under a Creative Commons Attribution

The Role of transdermal carbon dioxide

ORIGINALITY REPORT

16%

SIMILARITY INDEX

9%

INTERNET SOURCES

12%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Ahmed Ismaeel, Ramon Lavado, Robert S. Smith, Jack L. Eidson et al. "Effects of Limb Revascularization Procedures on Oxidative Stress", Journal of Surgical Research, 2018
Publication 1%
- 2 www.minervamedica.it
Internet Source 1%
- 3 Putnam, K.M.. "Depressive symptoms and baseline prefrontal EEG alpha activity: A study utilizing Ecological Momentary Assessment", Biological Psychology, 200802
Publication 1%
- 4 aunilo.uum.edu.my
Internet Source 1%
- 5 doi.org
Internet Source 1%
- 6 brieflands.com
Internet Source <1%
- 7 "Kidney transplantation: the recipient", Elsevier BV, 2024 <1%

8

Chou, M.M.. "Temporary Cross-clamping of the Infrarenal Abdominal Aorta During Cesarean Hysterectomy to Control Operative Blood Loss in Placenta Previa Increta/Percreta", Taiwanese Journal of Obstetrics & Gynecology, 201003

Publication

<1 %

9

www.ncbi.nlm.nih.gov

Internet Source

<1 %

10

www.cell.com

Internet Source

<1 %

11

orca.cf.ac.uk

Internet Source

<1 %

12

www.omicsonline.org

Internet Source

<1 %

13

Denise Grotto, Lucas Santa Maria, Juliana Valentini, Clóvis Paniz et al. "Importance of the lipid peroxidation biomarkers and methodological aspects FOR malondialdehyde quantification", Química Nova, 2009

Publication

<1 %

14

Miha Finžgar, Helena Ban Frangež, Ksenija Cankar, Igor Frangež. "Transcutaneous application of the gaseous CO2 for improvement of the microvascular function in

<1 %

patients with diabetic foot ulcers",
Microvascular Research, 2021

Publication

15

Subrata Ray, Anuradha Mitra, Ramapati Sanyal, Tabish Hussain, Anjum Naz. "Analgesic efficacy of suprascapular nerve block versus combined suprascapular and axillary nerve block in patients undergoing shoulder arthroscopy: A prospective, randomized, comparative study", Research Square Platform LLC, 2022

<1 %

Publication

16

Swati V. Elchuri, Briana C. Patterson, Milton R. Brown, Iris Buchanan, Ann C. Mertens, Lillian R. Meacham. "Anti-Mullerian hormone levels in American girls by age and race/ethnicity", Journal of Pediatric Endocrinology and Metabolism, 2015

<1 %

Publication

17

Xiaoling Feng, Xun Mao, Jianlin Zhao. "Clinical Characteristics, Prenatal Diagnosis and Outcomes of Placenta Accreta Spectrum in Different Placental Locations: A Retrospective Cohort Study", International Journal of Women's Health, 2024

<1 %

Publication

18

Yoshitada Sakai, Masahiko Miwa, Keisuke Oe, Takeshi Ueha, Akihiro Koh, Takahiro Niikura,

<1 %

Takashi Iwakura, Sang Yang Lee, Masaya Tanaka, Masahiro Kurosaka. "A Novel System for Transcutaneous Application of Carbon Dioxide Causing an "Artificial Bohr Effect" in the Human Body", PLoS ONE, 2011

Publication

19

hygeiadesenv.fsp.usp.br

Internet Source

<1 %

20

rcastoragev2.blob.core.windows.net

Internet Source

<1 %

21

www.abap.co.in

Internet Source

<1 %

22

www.degruyter.com

Internet Source

<1 %

23

Vamsi, Kalepu. "Comparison of Antiplatelet Efficacy of Xylitol/Sorbitol and Sorbitol Containing Chewing Gums on Plaque Regrowth at Smooth and Occlusal Surfaces", Rajiv Gandhi University of Health Sciences (India), 2023

Publication

<1 %

24

Wahyuni Hariyanto, Laksmi Wulandari, Isnu Pradjoko, Gatot Soegiarto. "Comparison of Vascular Endothelial Growth Factor-A (VEGF-A) Level in Pleural Fluid of Patients with Malignant and Nonmalignant Pleural Effusion", Indian Journal of Surgery, 2020

<1 %

25	9lib.net Internet Source	<1 %
26	clinicaltrials.gov Internet Source	<1 %
27	e-journal.unair.ac.id Internet Source	<1 %
28	jogh.org Internet Source	<1 %
29	moscow.sci-hub.se Internet Source	<1 %
30	vetdergikafkas.org Internet Source	<1 %
31	Kun-Long Huang, Leo Leung-Chit Tsang, Yu-Fan Cheng, Fu-Jen Huang et al. "Planned conservative management of placenta increta and percreta with prophylactic transcatheter arterial embolization and leaving placenta in situ for women who desire fertility preservation", Placenta, 2020 Publication	<1 %
32	N. Lalak. "THE EFFECT OF INFRARENAL AORTIC CROSS CLAMPING AND DECLAMPING ON RENAL FUNCTION IN THE PIG", ANZ Journal of Surgery, 3/1995 Publication	<1 %

33 Paul J. Garvin. "RENOPROTECTIVE EFFECTS OF THE 21-AMINOSTEROID U74389G IN ISCHEMIA-REPERFUSION INJURY AND COLD STORAGE PRESERVATION^{1,2}", Transplantation, 01/1997
Publication

34 Riyan Hidayat, Wan Nurazlina Wan Mohd Noor, Nurihan Nasir, Ahmad Fauzi Mohd Ayub. "The role of GeoGebra software in conceptual understanding and engagement among secondary school student", Infinity Journal, 2024
Publication

35 kclpure.kcl.ac.uk
Internet Source

36 s3-eu-west-1.amazonaws.com
Internet Source

37 scindeks.ceon.rs
Internet Source

38 www.nsveterina.edu.rs
Internet Source

39 "Peer-Reviewed Abstracts", Research Quarterly for Exercise and Sport, 2014
Publication

40 Andreia Fonseca, Diogo Ayres de Campos. "Maternal Morbidity and Mortality Due to

Placenta Accereta Spectrum Disorders", Best Practice & Research Clinical Obstetrics & Gynaecology, 2020

Publication

41

Atsushi Iwata. "Limitations of internal iliac artery ligation for the reduction of intraoperative hemorrhage during cesarean hysterectomy in cases of placenta previa accreta", Journal of Obstetrics and Gynaecology Research, 04/2010

Publication

<1 %

42

Hiroyuki Tokue, Azusa Tokue, Yoshito Tsushima, Takashi Kameda. "Safety and Efficacy of Aortic Vs Internal Iliac Balloon Occlusion for Cesarean Delivery in Coexisting Placenta Accreta and Placenta Previa", CardioVascular and Interventional Radiology, 2020

Publication

<1 %

43

Simon Gelman. "The Pathophysiology of Aortic Cross-clamping and Unclamping", Anesthesiology, 04/1995

Publication

<1 %

44

Vivek Madhusudan Joshi, Suhas Raghuvir Odiv, Yogesh Bhairav Sovani, Prasad Kisanrao Kulat. "Aortic clamping for limiting blood loss at cesarean hysterectomy for placenta

<1 %

percreta: A case series", International Journal of Gynecology & Obstetrics, 2021

Publication

45

doaj.org
Internet Source

<1 %

46

hdl.handle.net
Internet Source

<1 %

47

www.mdpi.com
Internet Source

<1 %

48

www.rusmedreview.com
Internet Source

<1 %

49

www.thieme-connect.com
Internet Source

<1 %

50

Angelo D'Alessandro, Travis Nemkov, Lori K. Bogren, Sandra L. Martin, Kirk C. Hansen. "Comfortably Numb and Back: Plasma Metabolomics Reveals Biochemical Adaptations in the Hibernating 13-Lined Ground Squirrel", Journal of Proteome Research, 2016

Publication

<1 %

51

F Nisi, L Carengo, N Ruggieri, A Reda, MG Pascucci, A Pignataro, E Civilini, F Piccioni, E Giustiniano. "The anesthesiologist's perspective on emergency aortic surgery: preoperative optimization, intra-operative

<1 %

management, and post-operative surveillance", Seminars in Vascular Surgery, 2023

Publication

52

tessera.spandidos-publications.com

Internet Source

<1 %

53

Peng Li, Xia Liu, Xiangkui Li, Xinchuan Wei, Juan Liao. "Clinical outcomes and anesthetic management of pregnancies with placenta previa and suspicion for placenta accreta undergoing intraoperative abdominal aortic balloon occlusion during cesarean section", BMC Anesthesiology, 2020

Publication

<1 %

54

Vito Chiantera, Evrim Erdemoglu, Giuseppe Vercellino, Magdalena Straube, Achim Schneider. "Laparoscopic Management of External Iliac Artery Injury Using Yasargil Clamps and Intracorporeal Suture", Journal of Minimally Invasive Gynecology, 2011

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On