

Risk factors associated with the Case

by Aditiawarman Aditiawarman

Submission date: 06-May-2024 10:43AM (UTC+0800)

Submission ID: 2371780379

File name: Risk_factors_associated_with_the_Case.pdf (193.35K)

Word count: 3945

Character count: 22628

Risk Factors Associated With the Case Fatality Rate of Pulmonary Tuberculosis in Pregnancy: A Five-Year Retrospective Study From a Developing Country

Review began 10/29/2023
Review ended 11/05/2023
Published 11/07/2023

© Copyright 2023
Ernawati et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Ernawati Ernawati¹, Rizka E. Prasetya¹, Aditiawarman Aditiawarman¹, Agus Sulistyono¹, Muhammad Ilham A. Akbar¹

1. Obstetrics and Gynaecology, Universitas Airlangga, Surabaya, IDN

Corresponding author: Ernawati Ernawati, ernawati@fk.unair.ac.id

Abstract

Background

Tuberculosis is a leading cause of maternal and fetal mortality in women of reproductive age. Tuberculosis is frequently misdiagnosed and treated inadequately during pregnancy. Although the global case fatality rate of tuberculosis is decreasing annually, the trend of tuberculosis mortality in Indonesia remains relatively high. Most tuberculosis reports do not include pregnancy status because most countries do not routinely screen for tuberculosis in pregnant women and do not report pregnancy status in female cases. In Southeast Asia, there is currently insufficient data regarding the risk factors associated with maternal mortality due to tuberculosis. This study aimed to identify the risk factors associated with tuberculosis-related mortality during pregnancy.

Methodology

This retrospective study was conducted at Dr. Soetomo General Hospital, Surabaya. Data were collected from patients' medical records. The samples were all pulmonary tuberculosis cases in pregnancy (suspected, bacteriological, and radiologically confirmed cases) from 2014 to 2018. Data on maternal characteristics, underlying risk factors, and maternal outcomes in pregnant women with tuberculosis were collected from medical records. A total of 77 cases of pulmonary tuberculosis in pregnancy were obtained and analyzed using the chi-square test for differences between pregnant women with tuberculosis who survived and those who did not.

Results

In total, 77 cases of pulmonary tuberculosis out of 7,242 deliveries were found during the past five years (incidence per year was 1.07), of whom 20.8% (16/77) died. Eight patients died before the gestational age reached 28 weeks. Most of the non-surviving women were aged <35 years (93.8%; 15/16). More than 30% (5/16) of the patients had human immunodeficiency virus co-infection, and the highest risk factors were pneumonia and miliary tuberculosis. Miliary tuberculosis was significantly associated with maternal mortality in pulmonary tuberculosis ($p = 0.004$) with a relative risk of 3.43.

Conclusions

According to the findings of this study, miliary tuberculosis is a significant risk factor for maternal mortality during pregnancy.

Categories: Epidemiology/Public Health, Obstetrics/Gynecology, Pulmonology

Keywords: human immunodeficiency virus, infection, maternal mortality, miliary tuberculosis, pregnancy, pulmonary tuberculosis

Introduction

Tuberculosis is a leading cause of maternal and fetal mortality in women of reproductive age. According to the WHO Tuberculosis Report 2014, there were 5.5 million female tuberculosis cases in 2015, with 510,000 deaths, and one-third of these women had human immunodeficiency virus (HIV) co-infection. This report did not include pregnancy status because the majority of countries did not routinely screen for tuberculosis in pregnant women and did not report pregnancy status in female cases [1]. In high-burden countries, the prevalence of active tuberculosis in pregnancy and postpartum was greater than 60 cases per 100,000 people per year, whereas it was less than 20 cases per 100,000 people per year in low-burden countries [2]. The highest number of tuberculosis cases in 2016 (45%) occurred in Southeast Asia, where Indonesia was included and classified as a high-burden nation for tuberculosis. Although the global case fatality rate of tuberculosis decreases by 3% per year, the trend of tuberculosis mortality in Indonesia remains relatively

How to cite this article

Ernawati E, Prasetya R E, Aditiawarman A, et al. (November 07, 2023) Risk Factors Associated With the Case Fatality Rate of Pulmonary Tuberculosis in Pregnancy: A Five-Year Retrospective Study From a Developing Country. *Cureus* 15(11): e48446. DOI 10.7759/cureus.48446

high [3]. The incidence of tuberculosis in Indonesia was greater than the global average in 2016, with 1.02 million cases (391 cases per 100,000 population) and a mortality rate of 37 cases per 100,000 individuals. Among all the cases reported in Indonesia, 41.7% were female [4].

Although pregnancy does not affect tuberculosis pathogenesis, reactivation of latent infection, or therapy response, the symptoms of tuberculosis in pregnancy are similar to those in non-pregnant patients, including fever, cough, weight loss, night sweats, and malaise. As a result, tuberculosis is frequently misdiagnosed and inadequately treated during pregnancy [5]. Most mortality associated with tuberculosis is due to non-tuberculosis factors (kidney failure, malignancy, and cirrhosis), with miliary tuberculosis and pneumonia being the leading causes of mortality associated with tuberculosis [6]. The co-occurrence of tuberculosis and human immunodeficiency virus/acquired immunodeficiency syndrome (HIV/AIDS) has significantly amplified mortality rates among women during their reproductive years. Both disorders are considered risk factors that contribute to higher maternal mortality in Sub-Saharan Africa [7]. In Southeast Asia, there is currently insufficient data regarding the risk factors associated with maternal mortality due to tuberculosis. Therefore, this study aimed to identify risk factors associated with tuberculosis-related maternal mortality during pregnancy.

17 Materials And Methods

This retrospective study was conducted at Dr. Soetomo General Academic Hospital, the largest referral hospital in East Indonesia. The data were obtained from the patient's medical records. The study sample consisted of pregnant women admitted to the hospital's delivery room between 2014 and 2018. These women were suspected cases, bacteriologically proven cases, or radiologically confirmed cases of pulmonary tuberculosis. Their medical records were thoroughly documented and included in the analysis. This study employed a comprehensive sampling approach, wherein individuals with incomplete medical record data were eliminated from the analysis. Data on several maternal variables, including maternal age, gestational age, parity, HIV status, presence of intrauterine growth restriction (IUGR), and mode of delivery, were collected.

Additionally, information on underlying risk factors and maternal outcomes, including survival or non-survival, was gathered. SPSS version 23 (IBM Corp., Armonk, NY, USA) was employed for data analysis. The chi-square test was utilized to examine the distinct characteristics among the cohorts of pregnant women diagnosed with tuberculosis who survived and those who did not. Before starting the investigation, the research protocol underwent a comprehensive evaluation and received approval from the Ethics Committee of Dr. Soetomo General Academic Hospital under reference number 0119/LOE/301.4.2/IX/2020.

Results

There were 77 cases of pulmonary tuberculosis out of 7,242 births over the course of five years, of whom 16. Figure 1 displays the data trends, case fatality rates, and case years. The annual incidence was 1.07%, and the average case fatality rate over the past five years was 20.82%.



FIGURE 1: The number of cases, incidence, and case fatality rate of tuberculosis in pregnancy at Dr. Soetomo General Academic Hospital, Surabaya (2014-2018)

24 The characteristics of pregnant women with pulmonary tuberculosis are shown in Table 1. Most births were

performed via cesarean section. The most prevalent indications were obstetric reasons (abnormal fetal presentation and dystocia) and fetal distress, and there were 12 cases with indications of severe lung damage and deteriorating maternal conditions. In contrast, only 13 patients underwent vaginal delivery (three in the non-survival group). Eight patients in the survivor group were not delivered because of early gestation.

Clinical Characteristics	N (%)
Age (year)	
<35	64 (83.1)
≥35	13 (16.9)
Gestational age (weeks)	
<28	27 (35.1)
28–34	22 (28.6)
>34	28 (36.4)
Parity	
Primipara	21 (27.3)
Multipara	56 (72.7)
Maternal outcome	
Survived	61 (79.2)
Did not survive	16 (20.8)
HIV status	
Positive	14 (18.1)
Negative	50 (64.9)
Unknown	13 (16.8)
IUGR	6 (7.7)
Mode of delivery	
Vaginal birth	13 (19.5)
Cesarean section	40 (51.9)
Not delivered	24 (31.2)

20

TABLE 1: Clinical characteristics of pregnant women with tuberculosis in Dr. Soetomo General Academic Hospital.

32

IUGR = intrauterine growth restriction; HIV = human immunodeficiency virus

Table 2 presents the factors associated with maternal mortality in pulmonary patients. Eight of the 16 pregnant women who died of pulmonary tuberculosis died before 28 weeks of gestation, and the majority of the non-survivors were under 35 years of age (15/16). There was a higher mortality rate among pregnant women aged 35 years owing to a variety of complications. Two patients had HIV co-infection with miliary tuberculosis, one patient had heart defects, one patient had HIV co-infection and pneumonia, six patients were co-infected with bacterial pneumonia, and three patients had miliary tuberculosis. More than 30% of individuals were HIV positive (5/16).

Maternal characteristics	Survivor (N = 61)	Non-survivor (N = 16)	P-value*
	N (%)	N (%)	
Maternal age (years)			
<35	49 (80.3)	15 (93.8)	0.202
≥35	12 (19.7)	1 (6.3)	
Gestational age (weeks)			
<28	19 (31.1)	8 (50)	0.082
28–34	16 (26.2)	6 (37.5)	
>34	26 (42.6)	2 (12.5)	
Parity			
Primipara	17 (27.9)	4 (25)	0.819
Multipara	44 (72.1)	12 (75)	
Mode of delivery			
Vaginal birth	10 (16.4)	3 (18.8)	0.122
Cesarean section	35 (57.4)	5 (31.2)	
Not delivered	16 (26.2)	8 (50)	
HIV status			
Positive	9 (14.8)	5 (31.3)	0.268
Negative	42 (68.9)	8 (50)	
Unknown	10 (16.4)	3 (18.8)	
Risk factor			
Hypertensive disorder**	11 (18)	0 (0)	0.067
Heart disease	4 (6.6)	1 (6.3)	0.965
Pneumonia	15 (24.6)	7 (43.8)	0.131
PROM	3 (4.9)	2 (12.5)	0.273
Miliary tuberculosis	4 (6.6)	5 (31.3)	0.004

TABLE 2: Maternal factors associated with maternal mortality with pulmonary tuberculosis at Dr. Soetomo General Academic Hospital.

*: chi-square test; **: including preeclampsia, eclampsia, and chronic hypertension.

HIV = human immunodeficiency virus; PROM = premature rupture of membrane

Pneumonia and miliary tuberculosis were the most significant risk factors in this study. Only miliary tuberculosis was significantly associated with maternal mortality in pulmonary tuberculosis ($p = 0.004$, relative risk = 5.45) of 12 risk factors predicted to be associated with maternal mortality.

Discussion

This study aimed to identify the risk factors associated with tuberculosis-related maternal mortality during pregnancy. The study found that patients with miliary tuberculosis had a 5.45-fold increased risk of death compared to patients without miliary tuberculosis. However, this study found that maternal age, gestational age, parity, and mode of delivery were not associated with pulmonary tuberculosis mortality in pregnant

women. This study revealed that tuberculosis cases with HIV co-infection had a higher mortality rate than tuberculosis cases without HIV. Additionally, the HIV-positive group had more severe complications than the HIV-negative group. This study also found no significant difference between the survival and non-survival groups regarding the risk factors for heart disease during pregnancy. Moreover, the non-survivor group had a higher proportion of tuberculosis cases with pneumonia co-infection, although the difference was not significant.

This study found that age was not associated with pulmonary tuberculosis mortality in pregnant women and that there were no significant differences in gestational age and parity between the tuberculosis group that survived and the tuberculosis group that did not. Demographically, multipara dominated both groups. Nonetheless, 50% of non-surviving cases were reported at a younger gestational age (less than 28 weeks). Pregnancy may inhibit the pro-inflammatory T-helper1 cell response, which may mask tuberculosis symptoms while increasing susceptibility to new infections and tuberculosis reactivation [8]. Several alterations in cellular immunity occur during the final stage of pregnancy, including an increase in phagocyte number and activity, plasmoid dendritic cells, and a decrease in natural killer cell cytotoxicity and interferon-gamma production. This demonstrates the suppression of innate cellular immunity [9].

There was no significant difference between the survival and non-survival groups in terms of mode of delivery. Previous research found no significant difference between pregnant women with tuberculosis and pregnant women without tuberculosis in terms of mode of delivery. There were no significant postpartum complications, except for one case of postnatal maternal death due to shock and end-organ failure as a consequence of military tuberculosis [10]. According to a different study, tuberculosis was associated with the mode of delivery (vaginal delivery, lower segment cesarean section, and instrumental vaginal delivery), with a p-value of 0.009 [11].

Multiple variables influence the impact of tuberculosis on pregnancy, including disease severity, site of infection, human HIV co-infection, gestational age, initial therapy, and patient compliance [12]. This study revealed that tuberculosis cases with HIV co-infection had a higher mortality rate than tuberculosis cases without HIV. Additionally, the HIV-positive group had more severe complications than the HIV-negative group. According to a previous study conducted in Sudan, 11.9% of pregnant women with tuberculosis were also HIV positive. The rate of maternal mortality due to tuberculosis was 4.8%, and all of these deaths were due to co-infection with HIV [13]. Women with HIV and latent *Mycobacterium tuberculosis* infections are more likely to develop active tuberculosis during pregnancy [14]. The difference between our study and previous studies may be that HIV status information was not available for all samples. Thirteen pregnant women had unknown HIV status.

In this study, one of the 16 deceased patients had a history of mitral stenosis, pulmonary hypertension, and pneumonia. Pregnancy with heart disease has unfavorable outcomes, particularly in cases in which heart disease and tuberculosis infection worsen the mother's condition. Pregnancy heart disease is a significant factor in maternal mortality after preeclampsia [15]. According to previous research, the prevalence of heart disease in pregnant women at Dr. Soetomo General Hospital is approximately 0.5%, with heart disease accounting for 14% of all maternal deaths [16].

In women with heart disease, pregnancy-related physiological changes may increase cardiac load and cause perinatal complications [17]. In the third trimester, increased cardiac load causes chronic maternal hypoxia, which is further exacerbated by tuberculosis. Compared with complications from tuberculosis alone, maternal mortality due to underlying heart disease may be a significant cause of death. However, this study found no significant difference between the survival and non-survival groups regarding the risk factors for heart disease during pregnancy. The limited number of heart disease cases may account for this finding.

The non-survivor group had a higher proportion of tuberculosis cases with pneumonia co-infection, although the difference was not statistically significant. Other factors may also contribute to pulmonary tuberculosis mortality in pregnant women. A 2016 retrospective study found that pneumonia was the primary predictor of mortality in patients with tuberculosis [6]. Patients with tuberculosis are more susceptible to bacterial infections, such as *Streptococcus pneumoniae*, which can activate latent tuberculosis, and vice versa. Consequently, diagnosis and treatment are challenging [18]. Pneumonia, pulmonary edema, pleural effusion, tuberculosis, and asthma are frequent lung diseases that occur during pregnancy and are associated with poor maternal and fetal outcomes [19].

This study found that patients with miliary tuberculosis had a 3.45-fold increased risk of death compared to patients without miliary tuberculosis. According to previous studies, cavities, miliary tuberculosis, and pneumonia are predictors of tuberculosis-related mortality. Extrapulmonary involvement and cirrhosis of the liver also contribute to deaths caused by tuberculosis [6]. Another study reported a case of miliary tuberculosis with a negative outcome during pregnancy. In this case, ascites, epistaxis, melena, convulsions, and abnormal laboratory findings (hemoglobin 7% and thrombocyte $60,000/\text{mm}^3$) were reported, and maternal death occurred after delivery of a stillborn infant [20].

Age, immunodeficiency, diabetes, mental disorders, elevated liver enzyme levels, kidney dysfunction,

malnutrition, thrombocytopenia, and radiological findings of ground-glass opacity are poor prognostic factors for miliary tuberculosis [21,22]. Multiorgan failure, septic shock, and acute respiratory distress syndrome are examples of fulminant acute conditions. With proper treatment, the mortality rate of miliary tuberculosis decreases from close to 100 percent to 7.1-30% [23].

Strengths and limitations

Indonesia is a high-burden country for tuberculosis and ranks among the countries with the highest incidence of the disease. Since the past decade, only a handful of studies have examined tuberculosis during pregnancy in Southeast Asia, and the majority of these studies are case reports or case series. This study had a larger sample of pregnant women with tuberculosis than the previous study on the same subject conducted in Indonesia or Southeast Asia. This cohort retrospective study found that miliary tuberculosis was significantly associated with maternal mortality in pulmonary tuberculosis ($p = 0.004$), indicating that patients with miliary tuberculosis have a 3.43-fold increased risk of death compared with patients without miliary tuberculosis. This study has some limitations. As a retrospective study, the data collected were limited. Additional prospective studies are required to further identify the risk factors associated with maternal mortality due to tuberculosis during pregnancy.

Research with larger samples should be conducted in the future to investigate why the incidence of miliary tuberculosis is higher in some regions of the world, such as Southeast Asia. Besides poor access to healthcare or inadequate treatment, there may be other factors that play a role in the development of miliary tuberculosis, such as genetics. Identifying the risk factors associated with tuberculosis-related maternal mortality during pregnancy can help improve the outcomes of pregnant women with tuberculosis.

Conclusions

In summary, compared to patients without miliary tuberculosis, those with miliary tuberculosis have a 3.43-fold increased risk of death. The risk of developing severe complications from tuberculosis is especially high in developing countries; therefore, early detection of either the onset of the disease or the onset of complications is required to reduce the mortality rate. Unfortunately, pregnancy can further mask tuberculosis symptoms, making diagnosis difficult. Consequently, tuberculosis is frequently misdiagnosed and inadequately treated during pregnancy. If the patient receives appropriate treatment, it can reduce the mortality rate due to tuberculosis and its complications, such as miliary tuberculosis.

Additional Information

Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

Concept and design: Ernawati Ernawati

Acquisition, analysis, or interpretation of data: Ernawati Ernawati, Rizka E. Prasetya, Aditiawarman Aditiawarman, Agus Sulistyono, Muhammad Ilham A. Akbar

Drafting of the manuscript: Ernawati Ernawati, Rizka E. Prasetya, Aditiawarman Aditiawarman, Muhammad Ilham A. Akbar

Critical review of the manuscript for important intellectual content: Ernawati Ernawati, Agus Sulistyono

Supervision: Ernawati Ernawati, Agus Sulistyono

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. Ethics committee of the Dr. Soetomo General Academic Hospital issued approval 0119/LOE/301.4.2/IX/2020. **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

Acknowledgements

We would like to thank Annisa and Esmond who helped compile some data.

References

1. WHO. Global tuberculosis report. (2014). <https://www.who.int/publications-detail-redirect/9789241564809>.
2. Mathad JS, Gupta A: Tuberculosis in pregnant and postpartum women: epidemiology, management, and research gaps. *Clin Infect Dis*. 2012, 55:1532-49. [10.1093/cid/cis732](https://doi.org/10.1093/cid/cis732)
3. Kemenkes RI. Infodatin Tuberkulosis. Jakarta: Kementerian Kesehatan RI - Pusat Data dan Informasi . (2018). <https://www.depkes.go.id/article/view/18030500005/waspadai-peningkatanpenyakit-menular.html%0A>.
4. World Health Organization. Global tuberculosis report. (2021). <https://apps.who.int/iris/bitstream/handle/10665/259366/9789241565516-eng.pdf?sequence=1>.
5. Friedman LN, Tanoue LT: Tuberculosis in pregnancy. UptoDate. Baron EL (ed): Wolters Kluwer, Philadelphia, PA; 2022.
6. Lin CH, Lin CJ, Kuo YW, et al.: Tuberculosis mortality: patient characteristics and causes. *BMC Infect Dis*. 2014, 14:5. [10.1186/1471-2354-14-5](https://doi.org/10.1186/1471-2354-14-5)
7. Adhikari M: Tuberculosis and tuberculosis/HIV co-infection in pregnancy. *Semin Fetal Neonatal Med*. 2009, 14:254-40. [10.1016/j.siny.2009.02.001](https://doi.org/10.1016/j.siny.2009.02.001)
8. Piccinni MP: T cell tolerance towards the fetal allograft . *J Reprod Immunol*. 2010, 85:71-5. [10.1016/j.jri.2010.01.006](https://doi.org/10.1016/j.jri.2010.01.006)
9. Kraus TA, Engel SM, Sperling RS, et al.: Characterizing the pregnancy immune phenotype: results of the viral immunity and pregnancy (VIP) study. *J Clin Immunol*. 2012, 32:300-11. [10.1007/s10875-011-9627-2](https://doi.org/10.1007/s10875-011-9627-2)
10. Regitz-Zagrosek V, Blomstrom Lundqvist C, Borghi C, et al.: ESC Guidelines on the management of cardiovascular diseases during pregnancy: the Task Force on the Management of Cardiovascular Diseases during Pregnancy of the European Society of Cardiology (ESC). *Eur Heart J*. 2011, 32:3147-97. [10.1093/eurheartj/ehr218](https://doi.org/10.1093/eurheartj/ehr218)
11. Wakamatsu K, Nagata N, Kumazoe H, et al.: Prognostic factors in patients with miliary tuberculosis . *J Clin Tuberc Other Mycobact Dis*. 2018, 12:66-72. [10.1016/j.jctube.2018.07.001](https://doi.org/10.1016/j.jctube.2018.07.001)
12. Loto OM, Awowole I: Tuberculosis in pregnancy: a review . *J Pregnancy*. 2012, 2012:379271. [10.1155/2012/379271](https://doi.org/10.1155/2012/379271)
13. Ali AA, Abdallah TM, Rayis DA, Adam I: Maternal and perinatal outcomes of pregnancies associated with tuberculosis in eastern Sudan. *Int J Gynaecol Obstet*. 2011, 114:286-7. [10.1016/j.ijgo.2011.02.023](https://doi.org/10.1016/j.ijgo.2011.02.023)
14. Bates M, Ahmed Y, Kapata N, Maeurer M, Mwaba P, Zumla A: Perspectives on tuberculosis in pregnancy . *Int J Infect Dis*. 2015, 32:124-7. [10.1016/j.ijid.2014.12.014](https://doi.org/10.1016/j.ijid.2014.12.014)
15. Stangl V, Schad J, Gossing G, Borges A, Baumann G, Stangl K: Maternal heart disease and pregnancy outcome: a single-centre experience. *Eur J Heart Fail*. 2008, 10:855-60. [10.1016/j.ejheart.2008.07.017](https://doi.org/10.1016/j.ejheart.2008.07.017)
16. Muninggar L, Yusuf M, Prasetyo B: Maternal mortality risk factor in pregnancy with heart disease at Dr. Soetono General Hospital, Surabaya, Indonesia. *Maj Obs Gin*. 2019, 27:17-23.
17. Puspa Pitaloka C, Secka A, Ernawati E, Sulistyono A, Juwono HT, Gumilar Dachlan E, Aditiawarman A: Characteristics shifting of heart disease in pregnancy: a report from low middle-income country . *J Public Health Res*. 2021, 10:2137. [10.4081/jphr.2021.2137](https://doi.org/10.4081/jphr.2021.2137)
18. Garcia R: Community-acquired pneumonia due to Streptococcus pneumoniae: when to consider coinfection with active pulmonary tuberculosis. *Case Rep Infect Dis*. 2019, 2019:4618413. [10.1155/2019/4618413](https://doi.org/10.1155/2019/4618413)
19. Han Y, Lee JH, Chang JH, Shim SS, Kim Y, Ryu YJ: Clinical features and outcomes in patients with pulmonary complications during pregnancy and peripartum. *J Matern Fetal Neonatal Med*. 2020, 33:1191-6. [10.1080/14767058.2018.1517313](https://doi.org/10.1080/14767058.2018.1517313)
20. Yadav V, Sharma JB, Kachhawa G, Kulshrestha V, Mahey R, Kumari R, Kriplani A: Obstetrical and perinatal outcome in pregnant women with extrapulmonary tuberculosis. *Indian J Tuberc*. 2019, 66:158-62. [10.1016/j.ijtb.2018.10.010](https://doi.org/10.1016/j.ijtb.2018.10.010)
21. Lee K, Kim JH, Lee JH, et al.: Acute respiratory distress syndrome caused by miliary tuberculosis: a multicentre survey in South Korea. *Int J Tuberc Lung Dis*. 2011, 15:1099-103. [10.5588/ijtld.10.0557](https://doi.org/10.5588/ijtld.10.0557)
22. Lee J, Lim JK, Seo H, et al.: Clinical relevance of ground glass opacity in 105 patients with miliary tuberculosis. *Respir Med*. 2014, 108:924-30. [10.1016/j.rmed.2014.03.016](https://doi.org/10.1016/j.rmed.2014.03.016)
23. Touré NO, Cissé MF, Dia Kane Y, et al.: [Miliary tuberculosis: a report of 49 cases]. *Rev Mal Respir*. 2011, 28:512-6. [10.1016/j.rmr.2011.02.005](https://doi.org/10.1016/j.rmr.2011.02.005)

Risk factors associated with the Case

ORIGINALITY REPORT

16%

SIMILARITY INDEX

11%

INTERNET SOURCES

13%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

cris.unica.edu.pe

Internet Source

1%

2

M. E. Roerink, J. W. M. Lenders, I. C. Schmits, A. M. A. Pistorius, J. W. Smit, H. Knoop, J. W. M. van der Meer. "Postural orthostatic tachycardia is not a useful diagnostic marker for chronic fatigue syndrome", *Journal of Internal Medicine*, 2017

Publication

1%

3

sakthi s, durairaj d, james antony bhagat m, nathiya b, thennarasu ar, nalin kumar. "The Impact of 4% Articaine on ECG: Insights from a Randomised Control Trial", *Research Square Platform LLC*, 2024

Publication

1%

4

www.smj.org.sg

Internet Source

1%

5

Yassine Eddair, Elmehdi Belouad, Elmostafa Benaissa, Tilila Abassor, Fatna Bsaibiss, Adil Maleb, Mostafa Elouennass. "Phenotypic and genotypic characterization of ESBL

1%

Enterobacteriaceae clinical isolates in a Moroccan hospital", Microbiology Society, 2024

Publication

6	repository.unair.ac.id Internet Source	1 %
7	cdr.lib.unc.edu Internet Source	1 %
8	Vasakova, Martina. "Challenges of Antituberculosis Treatment in Patients with Difficult Clinical Conditions", The Clinical Respiratory Journal, 2014. Publication	1 %
9	Rahul Kaul, H. S. Chhabra, Vijayanth Kanagaraju, Rajat Mahajan, Vikas Tandon, Ankur Nanda, Gururaj Sangondimath, Nishit Patel. "Antepartum surgical management of Pott's paraplegia along with maintenance of pregnancy during second trimester", European Spine Journal, 2015 Publication	1 %
10	link.springer.com Internet Source	1 %
11	"Nutrition and Health in a Developing World", Springer Science and Business Media LLC, 2017 Publication	<1 %

12	www.mdpi.com Internet Source	<1 %
13	scholar.unair.ac.id Internet Source	<1 %
14	Matthew Bates, Yusuf Ahmed, Nathan Kapata, Markus Maeurer, Peter Mwaba, Alimuddin Zumla. "Perspectives on tuberculosis in pregnancy", International Journal of Infectious Diseases, 2015 Publication	<1 %
15	f1000research.com Internet Source	<1 %
16	jtd.amegroups.com Internet Source	<1 %
17	www.researchgate.net Internet Source	<1 %
18	apps.who.int Internet Source	<1 %
19	Vikas Yadav, J.B. Sharma, Garima Kachhawa, Vidushi Kulshrestha, Reeta Mahey, Rajesh Kumari, Alka Kriplani. "Obstetrical and perinatal outcome in pregnant women with extrapulmonary tuberculosis", Indian Journal of Tuberculosis, 2018 Publication	<1 %

20

Frederick Berro Rivera, John Vincent Magalong, Ourlad Alzeus Tantengco, Gerard Francis Mangubat et al. "Maternal and neonatal outcomes among pregnant women with cardiovascular disease in the Philippines: a retrospective cross-sectional study from 2015–2019", *The Journal of Maternal-Fetal & Neonatal Medicine*, 2022

Publication

<1 %

21

es.scribd.com

Internet Source

<1 %

22

journals.lww.com

Internet Source

<1 %

23

www.frontiersin.org

Internet Source

<1 %

24

AbdelAzim A. Ali, Tajeldin M. Abdallah. "Clinical presentation and epidemiology of female genital tuberculosis in eastern Sudan", *International Journal of Gynecology & Obstetrics*, 2012

Publication

<1 %

25

Hanula, Ryan Justin. "Influenza Therapeutics: Evaluating the Benefits and Harms of Oseltamivir and Corticosteroids.", *McGill University (Canada)*, 2023

Publication

<1 %

26 Wallace, A. E., R. Fraser, and J. E. Cartwright. "Extravillous trophoblast and decidual natural killer cells: a remodelling partnership", Human Reproduction Update, 2012.

Publication

<1 %

27 downloads.hindawi.com

Internet Source

<1 %

28 Verena Stangl, Johanna Schad, Gabriele Gossing, Adrian Borges, Gert Baumann, Karl Stangl. "Maternal heart disease and pregnancy outcome: A single-centre experience", European Journal of Heart Failure, 2008

Publication

<1 %

29 repository.publisso.de

Internet Source

<1 %

30 repub.eur.nl

Internet Source

<1 %

31 www.jcgo.org

Internet Source

<1 %

32 www.researchsquare.com

Internet Source

<1 %

33 www.scielo.br

Internet Source

<1 %

34 www.thelancet.com

Internet Source

<1 %

35

"S290 topic 4 autoimmunity 1 WEB077021 ",
Open University

Publication

<1 %

36

Claudia L. Royo-Pabon, Carlos M. Perez-Velez.
"Tuberculosis exposure, infection and disease
in children: a systematic diagnostic
approach", Pneumonia, 2016

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On