

Enhanced recovery after vascular surgery

by Yan Efrata Sembiring

Submission date: 11-May-2024 05:46AM (UTC+0800)

Submission ID: 2374084999

File name: Enhanced_recovery_after_vascular_surgery.pdf (1.67M)

Word count: 8359

Character count: 47573

REVIEW

Enhanced recovery after vascular surgery

Yeri H. SARAGI ¹, I. PUTRA ¹, Yan E. SEMBIRING ^{2*}, Heroe SOEBROTO ², Ketut P. YASA ³, Cheong LIM ⁴

⁴⁰ ⁷
¹Department of Cardiac and Vascular Surgery, Dr. Soetomo Academic General Hospital, Airlangga University, Surabaya, Indonesia; ²Department of Thoracic, Cardiac and Vascular Surgery, Faculty of Medicine, Dr. Soetomo General Hospital, Airlangga University, Surabaya, Indonesia; ³Department of Thoracic, Cardiac and Vascular Surgery, Faculty of Medicine, Prof. IGNG Ngoerah General Hospital, Udayana University, Bali, Indonesia; ⁴Department of Thoracic and Cardiovascular Surgery, Bundang Hospital, Seoul National University, Seongnam, South Korea

⁶
^{*}Corresponding author: Yan E. Sembiring, Department of Thoracic, Cardiac and Vascular Surgery, Faculty of Medicine, Dr. Soetomo Academic General Hospital, Universitas Airlangga, Surabaya, East Java, Indonesia. E-mail: yan-e-s@fk.unair.ac.id

ABSTRACT

INTRODUCTION: A multimodal perioperative care measure, the enhanced recovery after surgery (ERAS) method, is intended to accomplish early recovery following surgical procedures. It aims to preserve preoperative organ function and mitigate the significant stress response that typically occurs during recovery.

EVIDENCE ACQUISITION: The goal of this systematic review and meta-analysis was to evaluate the advantages of enhanced recovery in the scope of vascular surgery. Following PRISMA Guidelines, a systematic search was conducted on various electronic reference databases (Web of Sciences, PubMed and Cochrane library). The keywords employed were ("Enhanced Recovery After Surgery" OR "ERAS") AND (vascular) AND ("surgery" OR "operation" OR "procedure"). Inclusion criteria are articles published in English, and full-text was available, published between 2013-2023. Data was obtained on hospitalization duration, in hospital mortality, and post-surgical morbidity.

EVIDENCE SYNTHESIS: Five hundred seventeen articles were identified and seven papers involving 1954 patients included for systematic review. The incidence of postoperative morbidity demonstrated a significant reduction when utilizing the ERAS approach in comparison to non-ERAS protocols (OR=0.21 [95%CI, 0.08 to 0.54], P=0.001). Additionally, the implementation of an ERAS protocol resulted in a notable reduction in hospitalization duration (MD=-0.59 [95%CI, -1.13 to -0.04], P=0.04). Furthermore, no significant difference was identified in hospital mortality rates.

CONCLUSIONS: The multimodal perioperative care approach known as Enhanced Recovery After Surgery (ERAS) method is intended to facilitate immediate postoperative recovery by safeguarding preoperative organ function and reducing the visceral stress response post-surgery. The utilization of ERAS, coupled with enhanced perioperative care, brings substantial advantages to patients undergoing vascular surgery as well as those undergoing a range of other surgical specialties.

(Cite this article as: Saragi YH, Putra I, Sembiring YE, Soebroto H, Yasa KP, Lim C. Enhanced recovery after vascular surgery. Ital J Vasc Endovasc Surg 2023;30:147-57. DOI: 10.23736/S1824-4777.23.01603-0)

KEY WORDS: Enhanced recovery after surgery; Vascular surgical procedures.

Introduction

¹³
A multimodal perioperative care measure, the enhanced recovery after surgery (ERAS) method, is intended to accomplish early recovery following surgical procedures. It aims to preserve preoperative organ function and mitigate the significant stress response that typically occurs during recovery.¹

The initial establishment and successful execution of ERAS method occurred in colorectal surgery. Subsequent-

ly, there have been reports of its implementation in orthopedic, cardiac, and vascular surgery as well.²

Due to the diverse range of vascular surgery types, such as endovascular, open, and hybrid procedures, patients may encounter various complications during their post-procedural recovery. As a result, each type of vascular surgery is associated with its own specific set of complications. Unlike other surgical specialties that typically concentrate on a specific body region, vascular surgeons undertake a wide array of surgeries, including cervical,

retroperitoneal, transabdominal, and upper and lower extremity procedures.³

The ideal approach to enforcing ERAS involves skilled and adaptable vascular team comprising surgeons, anesthesiologists, nurses, and therapists, who can follow a clinical pathway or protocol. Patient participation in smoking cessation, preoperative nutrition, and exercise programs would also be beneficial to reduce known risks. However, in the absence of clear guidance and assistance from the vascular team, geriatric patients often encounter comorbidities, decreased mobility, and access issues. The utilization of ERAS, coupled with enhanced perioperative care, brings substantial advantages to patients undergoing vascular surgery as well as those undergoing a range of other surgical specialties.³

Evidence acquisition

Research methodology

¹¹ The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 guidelines were adhered to in this systematic review. Relevant studies published until April 23, 2023 were identified through electronic data searches in Web of Science, Cochrane Library, and PubMed. The search terms utilized were as follows: (“Enhanced Recovery After Surgery” OR “ERAS”) AND (vascular) AND (“surgery” OR “operation” OR “procedure”). In addition, manual searches were conducted to identify potentially pertinent articles from the source list of collected studies. The search was restricted to English-language articles, and all searches were restricted to publications between the years 2013 and 2023.

Selection of studies and eligibility criteria

The process of selecting studies involved multiple stages and specific criteria for eligibility. Initially, the remaining articles underwent filtering based on their titles, followed by screening the abstracts of potentially relevant articles. Subsequently, articles that satisfied the eligibility criteria and had complete full texts were retrieved and evaluated. The entire study selection procedure was carried out separately by two researchers, and any disputes that arose were settled through discussions with the other investigators until a consensus was reached.

We included any single arm or comparative observational studies or randomized clinical trials. The study participants were exclusively adults (over 18 years old) going through either endovascular or open procedures on the venous or arterial system. The focus of interest was on

the implementation of a standard ERAS-like pathway or specific components of an ERAS pathway (including early mobility, multimodal analgesia, coordinated perioperative care, and patient education) during the prompt perioperative period. Exclusion criteria encompassed studies with inaccessible full texts, impertinent titles or abstracts, case series, review articles, letters to the editors, conference abstract, or case reports. Additionally, non-English studies, research involving children (<18 years old), postoperative or outpatient recovery pathway, and studies lacking adequate data to compute effect sizes for the outcomes were likewise excluded. The outcomes of interest were post-operative morbidity (involving any adverse events), duration of hospitalization, and in-hospital mortality.

Data retrieval

Two writers independently conducted the data extraction process, and the extracted data were subsequently cross-checked for accuracy. In the event of any discrepancies, a discussion was held to reach a consensus. Pertinent information from each included study was collected, including study-related details (such as first author, study design, study location, study period, and year of publication), demographic data (including age and sample size), as well as outcome data.

Statistical analysis

¹⁴ All analyses were performed using Review Manager version 5.4 (The Cochrane Collaboration, The Nordic Cochrane Centre, Copenhagen, Denmark). Odds Ratio (OR) and mean difference (MD) were utilized as the outcome measures for dichotomous data and continuous, respectively. The P value and 95% confidence interval (CI) were reported for the measures. Meta-analysis was performed for each outcome if at least two studies reported the same type of data. Heterogeneity among the studies was assessed using Cochran's Q and I² statistics. In cases where there was no significant heterogeneity (P value >0.1, I²<50%), a fixed-effects model was utilized for meta-analysis. Alternatively, a random-effects model was employed when heterogeneity was present. Potential publication bias was examined visually using Begg's funnel plots. Significance in all analyses was indicated by a P value <0.05.

Evidence synthesis

Study selection

We retrieved 511 records from the initial database query and six more records via manual searching. After screen-




Figure 1.—Flow diagram illustrating the study selection process based in PRISMA guidelines (Adapted from Page *et al.*).⁴

ing the titles and abstracts, 19 articles were identified as potentially eligible for review. Following a thorough assessment of the full texts, seven studies met the criteria and were included. The PRISMA flow chart (Figure 1) provides a summary of the process of selecting studies.⁴

Study characteristics

The characteristics of the seven included studies, which are relevant to this analysis, are presented in Table I.^{2, 5-10} It comprises one randomized controlled trial (RCT), five observational studies, and one descriptive study. These studies collectively involved a total of 1954 patients who underwent vascular surgery. Among the included studies, the majority (N.=6) were conducted in Europe, while one study was conducted in Asia. For detailed information on the outcomes of each individual study, please refer to Table II.^{2, 5-10}

Patient's characteristics

In this systematic review, we only included adult patients who underwent ERAS elements included in vascular surgery. In terms of types of vascular surgery, five studies^{2, 7-10} investigating the impact of ERAS on abdominal aortic an-

eurysm repair, one study⁶ on varicose vein surgery and one study on coronary artery surgery.⁵

Postoperative morbidity

In the current study, post-operative morbidity was defined as any complications that occurred following vascular surgery. Out of the six studies available on post-operative morbidity, the meta-analysis included data from five studies, which involved 873 patients. A pooled estimate of five studies showed significant differences in the incidence of postoperative morbidity (OR=0.21 [95%CI, 0.08 to 0.54], P=0.001), in which individuals who underwent ERAS elements included in vascular procedure had lower odds in developing any complications following vascular surgery; thus, suggesting the benefit of ERAS protocol implementation (Figure 2).^{2, 5, 6, 9, 10} In terms of potential publication bias, Begg's funnel plots analysis was performed qualitatively by visual inspections as documented in Figure 3.

In hospital mortality

Information regarding hospital mortality was obtained from a total of five studies involving 1511 patients.⁵⁻¹⁰ A total of 23 mortality was reported among patients who underwent abdominal aortic aneurysm repair utilizing ERAS protocol, whereas no mortality reported among patients who underwent coronary artery surgery implementing ERAS protocol. Nevertheless, since the majority of the studies reported no mortality and lack of comparison group; thus, meta-analysis cannot be generated.

Duration of hospitalization

Information on the duration of hospitalization following vascular surgery was provided by a total of seven studies. However, the meta-analysis included only three studies involving 246 patients.⁵⁻⁸ A pooled estimate of three studies showed significant differences in the mean hospital stay (MD=-0.59 [95%CI, -1.13 to -0.04], P=0.04), in which individuals who underwent ERAS elements included in vascular surgery had shorter period of hospitalization compared to control group; thus, suggesting the benefit of ERAS protocol implementation (Figure 4).^{5, 8, 10}

Discussion

Counseling, risk assessment, and optimization in preoperative

The ERAS protocol guides postoperative mobilization, discharge planning, and nutrition in the perioperative pe-

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The production of derivative works from the Article is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

TABLE I.—Characteristics of the included studies.^{2, 5-10}

Authors	Country	Study Design	Study Period	Population	Age (mean±SD)/ median (IQR)	Intervention	Comparison	Outcome
Renghi <i>et al.</i> ¹⁰	Italy	RCT	2005-2006	59 patients underwent mini-invasive abdominal aortic surgery in a fast-track	I=72.2±9.6 C=68.9±10	Thoracic epidural infusion	Local analgesia	Postoperative recovery, day of discharge, postoperative complication
Martelli <i>et al.</i> ⁹	Italy	Retrospective cohort	2006-2010	97 patients with abdominal aortic aneurysms underwent EVAR vs. fast-track open protocol	I=83±2 C=82±1.5	Left subcostal incision, no opioids, epidural analgesia, early mobilization and feeding	Endovascular aneurysm repair (EVAR) group	Intraoperative outcome (length of surgery, infusion, blood and liquid loss), in hospital mortality, postoperative complication
Hayashi <i>et al.</i> ⁸	Japan	Descriptive study	2012-2014	67 patients underwent open abdominal aortic aneurysm repair	I=68.7±6.1 C=69.0±7.6	Preoperative 6-minute walk test	Patients who achieved ambulation independence on/ after postoperative day 3	Pre, intra, and postoperative recovery outcome
Feo <i>et al.</i> ²	Italy	Retrospective cohort	2005-2013	221 patients underwent abdominal aortic aneurysm repair	I=69.9±8.6 C=71.1±8.2	Preoperative consultation, Two-hour preoperative fasting, NG tube removed in PACU, epidural analgesia, early mobilization and feeding	Patients who had undergone conventional perioperative protocols	Postoperative morbidity and mortality, time to functional recovery, hospital length of stay, readmission rate
Brustia <i>et al.</i> ⁷	Italy	Retrospective study	2000-2014	1,014 patients underwent abdominal aortic aneurysm repair	70.49±8.2	Left subcostal incision, epidural analgesia, early mobilization and feeding	N/A	The incidence and characteristics of postoperative complications, the mortality rate, and the time of discharge to home for patients
Licker <i>et al.</i> ⁶	Switzerland	Prospective study	2009-2011	376 patients underwent varicose vein surgery	I=47±14 C=47±14	Local analgesia plus sedation vs. general anesthesia	Patients receiving general anesthesia or spinal anesthesia	Post-operative recovery time, length of stay, hospital costs, adverse events
Anastasiadis <i>et al.</i> ⁵	Greece	Prospective study	2010-2011	120 patients underwent elective coronary artery surgery	I=64.4±9.9 C=65.1±7.8	Minimal extracorporeal circulation (MECC)	Patients who underwent conventional cardiopulmonary bypass	Rate of fast-track recovery, overall morbidity and mortality, need for perioperative blood transfusion, chest tube drainage, and duration of hospitalization

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

TABLE II.—Outcomes of individual studies.^{2, 5-10}

Authors	Post-operative morbidity, n	In hospital mortality, n	Hospital Length of Stay, days, mean±SD / median (range)
Renghi <i>et al.</i> ¹⁰	I=15/29 C=24/30	I=0 C=0	I=3.2±1.4 C=3.4±1.7
Martelli <i>et al.</i> ⁹	I=11/55 C=9/42	I=0 C=0	4 (mean of both groups); on the third day: I=64% of patients C=42.9% of patients
Hayashi <i>et al.</i> ⁵	NR	N/A	I=9.8±2.1 C=11.1±2.8
Feo <i>et al.</i> ²	I=24/130 C=62/91	I=3/130 C=3/91	I=5 (4-7) C=7 (6-9)
Brustia <i>et al.</i> ⁷	104/1,014	17/1,014	3.97±2.7
Licker <i>et al.</i> ⁶	I=8/176 C=82/200	NR	I=patients were eligible for discharge on the day of surgery; C=14 patients required overnight hospital admission
Anastasiadis <i>et al.</i> ⁵	I=3/60 C=8/60	I=0 C=0	I=10.8±2.6 C=11.5±2.8




Figure 2.—Forest plot.^{2, 5, 6, 9, 10}




Figure 3.—Funnel plot.




Figure 4.—Forest plot.^{5, 8, 10}

riod. ERAS can also be used in vascular surgery to implement strategies to optimize preoperative chronic medical conditions and lifestyle factors as presented in Table III.³

The preoperative evaluation is of utmost importance as it allows the physician to assess risks, evaluate the patient's stability in relation to preexisting conditions, and maximize pre-operative organ function. This is particularly significant as postoperative dysfunction often correlates with preexisting comorbidities. When ERAS anxiety is present, it is projected that the duration of hospitalization and requirement for analgesics can be decreased. Table IV illustrates the suggested guidelines for ERAS implementation in vascular surgery (including preoperative, intraoperative, and postoperative periods).²

Patients should prioritize premedication to minimize

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The production of derivative works from the Article is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

TABLE III.—ERAS in vascular surgery:²

Respiratory
Preoperative antibiotics use
Early extubation
Protective modes of ventilation
Prevention of ventilator-associated pneumonia
High-flow oxygen therapy or intermittent nasal continuous positive airway pressure after extubation
1 Breathing exercises
Cardiovascular
Monitoring for signs of myocardial ischemia
Goal directed fluid therapy
1 Maintaining the MAP above 80–90 mmHg
Renal
Monitoring of the amount of urine and creatinine clearance
Maintaining the normovolemia and electrolyte balance
Use diuretics with caution
Avoid fluid overload and dopamine
Nutrition
Without preoperative mechanical bowel preparation
Maintenance the glucose level <215 mg/dL
Oral nutrition within the first 24–48 h after operation
Use prokinetics (metoclopramid and erythromycin)
Avoidance of nasogastric drainage or early removal
Monitoring of IAP
Pain management
PCEA 48 h before and 48 h after intervention
Avoid systemic opioid use
Other
Stabilize already existing disease and optimize organ dysfunction before surgery
Oral carbohydrate drinks before surgery
Minimize the time for surgical intervention
Patient education
Consider thromboprophylaxis
Avoid hypothermia
Early mobilization

TABLE IV.—ERAS in vascular surgery:¹

Preoperative	Perioperative	Postoperative
Patient counseling/education	Abdominal drains (none)	Regular diet
Fasting (reduced)	Nasogastric tube (none)	9 Early feeding
9 Use of non-steroidal anti-inflammatory drugs (NSAIDs)	Fluid restriction	Pain management
Carbohydrate enriched drinks	Analgesia – NSAIDS, narcotics, use of epidural analgesia	Chewing gum/ileus
Stress management	Fluid restriction	Early ambulation
Patient expectations (managing)		Exercise therapy
		Patient discharge planning
Accelerated rehabilitation	Urinary catheter (time of removal)	Removal of epidural catheter

the surgical stress response. Beta-blockers are effective in reducing catecholamine levels and cardiovascular complications both during and after surgery. Given that surgical procedures trigger sympathetic system activation and stress-induced catabolism, the use of anti-catabolic medications can positively impact the postoperative recovery. Alpha-2 adrenergic agonists like clonidine and dexmedetomidine are particularly beneficial, as they help reduce postoperative nausea and vomiting, intraoperative blood loss, and myocardial ischemia.²

Fasting overnight before surgery serves multiple purposes, including preventing dehydration and minimizing the likelihood of postoperative complications such as pain and nausea. Clear carbohydrate fluids may reduce postoperative anxiety and endocrine response if administered during this period.²

Cardiovascular risk evaluation and optimization

Optimal management of cardiovascular risk is crucial in vascular patients due to the high prevalence of cardiac comorbidities, which can significantly impact surgical outcomes. In alignment with the guidelines set forth by the American Heart Association (AHA), the implementation of ERAS method should adhere to the AHA's recommendations for preoperative cardiovascular evaluation and optimization. This may involve conducting diagnostic tests such as electrocardiograms, echocardiograms, and stress testing to assess and manage the cardiovascular risk in patients with preexisting heart conditions. Continuation of medications such as statins, angiotensin-converting enzyme inhibitors/angiotensin receptor blockers, beta-blockers, and antiplatelet agents during the surgical period is crucial as part of the comprehensive medical management of vascular disease. When appropriate, higher-risk inpatients should be referred to a cardiologist for overall preoperative assistance and optimization and postoperative care.³

Anticoagulants

Patients undertaking vascular rehabilitation may be limited in their selection of regional anesthetic techniques due to the routine use of systemic anticoagulants. Anticoagulants pose a risk of bleeding complications in vascular patients. ERAS facilitates the implementation of protocols for managing anticoagulation, including discontinuation and reintroduction, to ensure patient safety and optimize outcomes. The operation of the ERAS system will include instructions on preoperative anticoagulant use, the scheduling of anticoagulant discontinuation, and postoperative anticoag-

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The production of derivative works from the Article is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

ulant resumption. Anticoagulant reversal agents are typically reserved for emergent or emergency situations.³

Tobacco cessation

Tobacco cessation is crucial in vascular patients due to its detrimental effects on vascular health, perioperative complications, and wound healing. With a high prevalence of tobacco use among vascular patients, incorporating tobacco cessation as part of the ERAS pathway would greatly benefit their outcomes. Hospitalization presents an opportune moment to motivate patients to quit smoking and promote tobacco cessation. Patients are often more receptive to quitting during this time of treatment, making it an ideal opportunity to provide support and encourage tobacco cessation efforts. For patients who are not ready to quit smoking, ERAS combines additional nicotine treatment regimens (*i.e.*, gum, patches) with counseling efforts.^{3, 11}

Physical exertion

According to Hayashi *et al.*, regular physical activity before surgery can improve the condition of AAA patients. Regular physical activity refers to exercising for at least 30 minutes, twice a week, over a duration of at least one year. The specific amount and duration of activity required for visible results remain uncertain. Nevertheless, patients should be notified that initiating or maintaining an exercise program before surgery may lead to a reduce duration of hospitalization and accelerated healing. The patient's preoperative activity level should also be considered when discussing expected postoperative recovery time.⁸

Perioperative pain control

Aortic surgery is one of the most important interventions in vascular surgery. Today, endovascular management and open surgery are the most common surgical techniques. According to Hertzner *et al.*, elective open infrarenal aortic surgery is associated with an average ICU stay of approximately three days and a mortality rate ranging from 1.2% to 10.5%.¹² Both endovascular and open surgery are prone to postoperative complications such as graft migration, thrombosis, and aneurysm rupture. However, endovascular management offers advantages in terms of shorter procedure duration, reduced blood loss risk, decreased risk of malnutrition, and shortened hospitalization.^{11, 13, 14}

Regional anesthesia

Epidural anesthesia is found to be more commonly employed compared to regional anesthesia among the 12

studies investigating ERAS-like pathways in open aortic surgery. Studies that incorporate epidural anesthesia as part of the ERAS pathway have shown positive results, such as reduced complication rates, faster extubation after surgery, shorter ICU stays, and shorter length of stays. Multiple of these studies report effective pain management without the use of analgesics. Epidural anesthesia has been found to decrease the release of epinephrine in patients undergoing coronary artery bypass graft surgery, potentially lowering the occurrence of myocardial ischemia and consequently reducing mortality and morbidity rates. The observed decrease in complication rates among patients receiving epidural anesthesia during vascular surgery is thought to be attributed to the attenuated stress response. The combined use of epidural and general anesthesia in aortic surgery has been shown to reduce the requirement for mechanical ventilation compared to general anesthesia alone.³

According to Muehling *et al.*, In patients who received epidural anesthesia, the incidence of postoperative mechanical ventilation was significantly lower (5%) compared to those who received only general anesthesia in the "traditional" group (33%). This is because patients receiving epidural anesthesia use less intraoperative inhalational anesthetics. In different study comparing general anesthesia combined with an epidural to general anesthesia alone in open aortic surgery, no significant differences were observed in ICU stay, hospital stay, time to oral intake, time to ambulation, mortality, or morbidity. However, similar to Muehling's investigation, patients who underwent epidural anesthesia were extubated considerably earlier compared to those who only received general anesthesia.¹³

In all cases, patients were provided with nasogastric tubes until bowel sounds resumed. However, the lack of a feeding protocol and active ambulation might have hindered the potential benefits of epidural anesthesia. The combination of various components in the ERAS pathway demonstrates synergistic and beneficial effects.³

Delirium screening

Delirium is a common coexisting condition in patients with vascular disease and is often overlooked, resulting in a decline in functional ability. To prevent delirium, it is crucial to minimize unnecessary stays in the intensive care unit, minimize opioid usage, promote early ambulation, establish a healthy sleep pattern, optimize the day/night cycle, and provide visual and verbal cues for orientation. Regular and thorough screening for delirium should be implemented as part of routine care.³

Postoperative period

The postoperative period is of utmost importance in the recovery of patients undergoing vascular surgery. Important to a successful recovery is the caliber of postoperative care. The optimization of microcirculation, inflammatory response management, fluid resuscitation, hemodynamic support, blood glucose levels, aerobic metabolism, oxygen delivery, and appropriate use of vasoconstrictors and vasodilators are key aspects in achieving successful outcomes in vascular surgical interventions.³

The occurrence of complications is affected by aspects such as the underlying disease, the specific surgical procedure, and the urgency of the intervention. Several complications may occur in the early postoperative period, while others may occur in the late phase.³

The study by Crimi and Hill demonstrated the significance of early management of hemodynamic, pulmonary, renal, neurological, hematologic, and gastrointestinal complications immediately after surgery. Ischemic-reperfusion injury results in dysfunctions of these organ systems. Treatment on the first postoperative day is primarily determined by the kind of surgical procedure, the operative time, and the patient's condition. Throughout this duration, it is crucial to be aware of complications, particularly hemodynamic ones.¹⁵

Intravenous fluid management

In aortic surgery, effective fluid management is crucial in the postsurgical period. It includes maintaining a balanced fluid status to prevent complications like acute kidney injury and issues associated with fluid overload, such as heart failure, impaired oxygenation, and delayed bowel function. Generally, patients are already in a euvoletic state after surgery, eliminating the need for additional fluid boluses or diuretics. With careful clinical assessment, the fluid balance of postoperative patients can be optimized.¹⁶

In cases where a euvoletic patient is capable of oral fluid intake, intravenous fluids may be unnecessary to maintain adequate fluid levels. Nonetheless, when fluid replacement is necessary, a restrictive fluid therapy approach tends to yield better outcomes compared to liberal fluid administration. Research has shown that patients receiving 1.5 L/day of fluids experienced lower risk of cardiopulmonary complications, fewer complications, and shorter lengths of stay in the ICU and overall duration of hospitalization compared to those in the free fluid therapy group. In abdominal aortic surgery, a Cochrane review found no evidence supporting the use of specific fluids, crystalloids, or colloids.¹⁶

Pain management

The utilization of analgesics for managing chronic pain in vascular patients has witnessed a notable rise. A considerable number of patients undergoing lower extremity vascular surgery experience chronic pain, which was traditionally addressed with analgesics. The regular administration of analgesics can significantly impact the perioperative pain management of these patients.¹⁷

The ERAS pathway for lower extremity revascularization patients needs to consider both opioid-naive individuals and those dependent on opioids. Managing chronic opioid users can pose a challenge. Since surgery can alleviate or remove the underlying cause of chronic pain, the ERAS pathway should take into account the patient's opioid usage level and reasons for opioid use. For patients whose chronic pain improves following revascularization, the prescribed regimen should involve dose reduction. Patients with chronic pain unimpacted by surgery may require additional analgesia for adequate postoperative pain control. Continuous peripheral nerve blocks (CPNB) provide significant benefits for lower extremity vascular surgery patients.¹⁸

Evidence supports the use of standard, multimodal, and opioid-sparing analgesia to reduce perioperative opioid consumption and expedite recovery in opioid-naive patients. Continuous peripheral nerve blocks (CPNB) hold the capacity to reduce additional opioid usage in opioid-tolerant patients during the perioperative period. The use of continuous peripheral nerve blocks in lower limb vascular surgery has not been reported, despite its mention in the context of the ERAS pathway for other lower limb procedures like knee replacement. However, research suggests that incorporating local analgesia as a key component of pain control in lower extremity vascular surgery is both feasible and beneficial. For instance, Licker et al. successfully implemented a local analgesia and sedation protocol in 176 patients undergoing saphenous vein ligation and phlebectomy, as an alternative to previous general anesthesia used in 200 patients.¹⁹

The occurrence of postoperative nausea, dizziness, and migraines was significantly lower at 4% compared to 41% ($P=0.001$). The average duration of discharge from the ambulatory surgical center was reduced by 364 minutes, enabling the scheduling of an additional case per day without a rise in complications. In another study, the utilization of local analgesia and sedation showed excellent outcomes in performing popliteal bypass sheath. All ten patients in the case series successfully underwent the procedure and were able to walk within eight hours without any complications (Figure 5, 6).²⁰⁻²²




Figure 5.—ERAS in aorta vascular surgery.




Figure 6.—ERAS principle in vascular surgery at lower extremity.

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The production of derivative works from the Article is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

Mobilization and implants

Postoperative mobilization is a vital aspect of the ERAS pathway, particularly for patients undergoing lower extremity vascular surgery. These patients often have preexisting mobility limitations and reduced functional capacity, making preoperative conditioning challenging. Therefore, specialized personnel, such as physical therapists, may be required to facilitate postoperative mobilization due to the specific challenges involved. Mobilization is the initial (key) ERAS pathway used for other surgeries, including open aortic surgery, but in actuality, postoperative ambulation requires the assistance of a nurse or even a family member. Patient education and psychological support regarding recovery, physical therapy, prosthetics, discharge planning, and rehabilitation are essential for lower extremity amputees. The ERAS pathway for amputees should include preoperative education, physiotherapy consultations, and ideally medical rehabilitation peer visits and limb care education.¹⁹ Marzen-Groller *et al.* establish protocols for hospitalization for ambulation after amputation. Patients undergoing below-knee amputation, above-knee amputation, or transtatarsal amputation received preoperative physical therapy evaluations and continued with postoperative care. The study showed improved mobility scores and a potential decrease in DVT rates, although not statistically significant.¹⁹

Conclusions

The ERAS protocol aims to promote early recovery after surgical procedures by preserving preoperative organ function and minimizing the physiological stress response to surgery. It employs a multimodal perioperative care approach. Given the diverse range of vascular surgeries, such as open, endovascular, and hybrid procedures, there are specific complications associated with each. To address these challenges, the implementation of Enhanced Recovery After Surgery (ERAS) and comprehensive perioperative care can greatly benefit patients undergoing vascular surgery, as well as those undergoing various other surgical specialties.

References

1. Gotlib Conn L, Rotstein OD, Greco E, Tricco AC, Perrier L, Soobiah C, *et al.* Enhanced recovery after vascular surgery: protocol for a systematic review. *Syst Rev* 2012;1:52.
2. Feo CV, Portinari M, Tsolaki E, Romagnoni G, Verri M, Camerani S, *et al.* The effect of an Enhanced Recovery Program in elective retroperitoneal abdominal aortic aneurysm repair. *J Vasc Surg* 2016;63:888–94.
3. Ljungqvist O, Francis NK, Urman RD, editors. *Enhanced Recovery After Surgery: A Complete Guide to Optimizing Outcomes*. Cham: Springer International Publishing; 2020.
4. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, *et al.* The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
5. Anastasiadis K, Asteriou C, Antonitsis P, Argiriadou H, Grosomanidis V, Kyparissa M, *et al.* Enhanced recovery after elective coronary revascularization surgery with minimal versus conventional extracorporeal circulation: a prospective randomized study. *J Cardiothorac Vasc Anesth* 2013;27:859–64.
6. Licker M, Brandao-Farinelli E, Cartier V, Gemayel G, Christenson JT. Implementation of a fast-track-pathway including analgo-sedation with local anaesthesia for outpatient varicose vein surgery: a cohort study. *Phlebology* 2013;28:418–25.
7. Brustia P, Renghi A, Aronici M, Gramaglia L, Porta C, Musiani A, *et al.* Fast-track in abdominal aortic surgery: experience in over 1,000 patients. *Ann Vasc Surg* 2015;29:1151–9.
8. Hayashi K, Hirashiki A, Kodama A, Kobayashi K, Yasukawa Y, Shimizu M, *et al.* Impact of preoperative regular physical activity on postoperative course after open abdominal aortic aneurysm surgery. *Heart Vessels* 2016;31:578–83.
9. Martelli M, Renghi A, Gramaglia L, Casella F, Brustia P. Abdominal aortic aneurysm treatment: minimally invasive fast-track surgery and endovascular technique in octogenarians. *J Cardiovasc Surg* 2017;58:557–64.
10. Renghi A, Gramaglia L, Casella F, Moniaci D, Gaboli K, Brustia P. Local versus epidural anesthesia in fast-track abdominal aortic surgery. *J Cardiothorac Vasc Anesth* 2013;27:451–8.
11. Chaar CI, Fitzgerald TN, Dewan M, Huddle M, Schlosser FJ, Perkal M, *et al.* Endovascular aneurysm repair is associated with less malnutrition than open abdominal aortic aneurysm repair. *Am J Surg* 2009;198:623–7.
12. Hertzner NR, Mascha EJ, Karafa MT, O'Hara PJ, Krajewski LP, Beven EG. Open infrarenal abdominal aortic aneurysm repair: the Cleveland Clinic experience from 1989 to 1998. *J Vasc Surg* 2002;35:1145–54.
13. Muehling BM, Halter G, Lang G, Schelzig H, Steffen P, Wagner F, *et al.* Prospective randomized controlled trial to evaluate "fast-track" elective open infrarenal aneurysm repair. *Langenbecks Arch Surg* 2008;393:281–7.
14. van Zanten AR. Nutrition barriers in abdominal aortic surgery: a multimodal approach for gastrointestinal dysfunction. *JPEN J Parenter Enteral Nutr* 2013;37:172–7.
15. Crimi E, Hill CC. Postoperative ICU management of vascular surgery patients. *Anesthesiol Clin* 2014;32:735–57.
16. McGinagle KL, Spangler EL, Pichel AC, Ayyash K, Arya S, Settembrini AM, *et al.* Perioperative care in open aortic vascular surgery: A consensus statement by the Enhanced Recovery After Surgery (ERAS) Society and Society for Vascular Surgery. *J Vasc Surg* 2022;75:1796–820.
17. Lindgren H, Gottsäter A, Qvarfordt P, Bergman S. All Cause Chronic Widespread Pain is Common in Patients with Symptomatic Peripheral Arterial Disease and is Associated with Reduced Health Related Quality of Life. *Eur J Vasc Endovasc Surg* 2016;52:205–10.
18. Scarlet S, Isaak RS, McGinagle KL. Design and Implementation of an Enhanced Recovery after Surgery (ERAS) Pathway for Major Limb Amputation in Vascular Surgery. *Am Surg* 2018;84:e147–9.
19. Seretny M, Colvin LA. Pain management in patients with vascular disease. *Br J Anaesth* 2016;117(Suppl 2):ii95–106.
20. Carli F, Kehlet H, Baldini G, Steel A, McRae K, Slinger P, *et al.* Evidence basis for regional anesthesia in multidisciplinary fast-track surgical care pathways. *Reg Anesth Pain Med* 2011;36:63–72.
21. Marzen-Groller KD, Tremblay SM, Kaszuba J, Girodo V, Swavely D, Moyer B, *et al.* Enhanced Recovery After Surgery: A Review. *J Vasc Nurs* 2008;26:74–81.
22. Ljungqvist O, Scott M, Fearon KC. Enhanced recovery after surgery: a review. *JAMA Surg* 2017;152:292–8.

Conflicts of interest

The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

Authors' contributions

Yeri Holo Saragi: conception and design, drafting of the article and collection and assembly of data; Yan E. Sembiring: critical revision of the article for important intellectual content and Final approval of the article; I. Putra: provision of study materials or patients; Heroe Subroto: critical revision of the article for important intellectual content and final approval of the article; Ketut P. Yasa: statistical expertise and critical revision of the article for important intellectual content; Cheong Lim: conception and design. All authors read and approved the final version of the manuscript.

History

Manuscript accepted: July 21, 2023. - Manuscript received: July 18, 2023.

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The production of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

Enhanced recovery after vascular surgery

ORIGINALITY REPORT

11%

SIMILARITY INDEX

9%

INTERNET SOURCES

8%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

mafiadoc.com

Internet Source

1%

2

Katharine L. McGinagle, Jens Eldrup-Jorgensen, Rebecca McCall, Nikki L. Freeman et al. "A systematic review of enhanced recovery after surgery for vascular operations", *Journal of Vascular Surgery*, 2019

Publication

1%

3

www.annalsofafricansurgery.com

Internet Source

1%

4

Tim J. P. Batchelor. "Chapter 53 Thoracic Surgery", Springer Science and Business Media LLC, 2020

Publication

1%

5

Yingping Kuang, Jun Xiong, Zheng Zhang, Kai Liao, Xiaohong Zhou, Jie Xiang, Jun Li, Lingling Xu, Haiping Zhou. "Assessment of the Quality of Randomized Controlled Trials Report of Acupuncture and Moxibustion for allergic rhinitis by the CONSORT Statement、 STRICTA

1%

and STRICTOM Guidelines", Research Square Platform LLC, 2023

Publication

6	Sri P. NEGORO, Yan E. SEMBIRING, Latifah A. ZATI, I G. PUTRA, Jeffrey J. DILLON. "The use of artificial intelligence in the diagnosis of peripheral arterial disease: a systematic review", Italian Journal of Vascular and Endovascular Surgery, 2024	<1 %
<hr/>		
7	repository.unair.ac.id	<1 %
<hr/>		
8	systematicreviewsjournal.biomedcentral.com	<1 %
<hr/>		
9	www.systematicreviewsjournal.com	<1 %
<hr/>		
10	pure.rug.nl	<1 %
<hr/>		
11	mhealth.jmir.org	<1 %
<hr/>		
12	www.scielo.br	<1 %
<hr/>		
13	Feng Chen, Gongchao Wang. "Enhanced Recovery after Surgery for Lung Cancer Patients", Open Medicine, 2020	<1 %

Publication

14	www.wjgnet.com Internet Source	<1 %
15	Katharine L. McGinigle, Emily L. Spangler, Adam C. Pichel, Katie Ayyash et al. "Perioperative care in open aortic vascular surgery: A Consensus Statement by the Enhanced Recovery after Surgery (ERAS®) Society and Society for Vascular Surgery", Journal of Vascular Surgery, 2022 Publication	<1 %
16	medicaljournalssweden.se Internet Source	<1 %
17	www.diva-portal.org Internet Source	<1 %
18	zaguan.unizar.es Internet Source	<1 %
19	stikes-nhm.e-journal.id Internet Source	<1 %
20	www.nature.com Internet Source	<1 %
21	Katharine L. McGinigle, Emily L. Spangler, Adam C. Pichel, Katie Ayyash et al. "Perioperative care in open aortic vascular surgery: A consensus statement by the Enhanced Recovery After Surgery (ERAS)	<1 %

Society and Society for Vascular Surgery",
Journal of Vascular Surgery, 2022

Publication

22

coek.info

Internet Source

<1 %

23

e-sciencecentral.org

Internet Source

<1 %

24

Guillemette Lienhart, Elsa Masson, Pierre Farge, Anne-Marie Schott-Pethelaz, Béatrice Thivichon-Prince, Marc Chanelière. "Factors perceived by health professionals to be barriers or facilitators to caries prevention in children: a systematic review", Research Square Platform LLC, 2023

Publication

<1 %

25

paramed.dums.ac.ir

Internet Source

<1 %

26

repositorio.unesp.br

Internet Source

<1 %

27

www.frontiersin.org

Internet Source

<1 %

28

perioperativemedicinejournal.biomedcentral.com

Internet Source

<1 %

29

www.mdpi.com

Internet Source

<1 %

30

"Essentials of Regional Anesthesia", Springer Science and Business Media LLC, 2018

Publication

<1 %

31

Claire Temple-Oberle, Carmen Webb. "Chapter 45 ERAS for Breast Reconstruction", Springer Science and Business Media LLC, 2020

Publication

<1 %

32

Elena Giacomelli, Walter Dorigo, Marco Campolmi, Andrea Casini, Aaron Fargion, Ruth L. Bush, Gabriele Piffaretti, Carlo Pratesi. "A pilot study of the ERAS (Enhanced Recovery After Surgery) protocol in aortic surgery", Journal of Vascular Surgery, 2020

Publication

<1 %

33

Schraag, Stefan. "Postoperative Management", Best Practice & Research Clinical Anaesthesiology, 2016.

Publication

<1 %

34

Thomas G. Poder, Lucien P. Coulibaly, Myriam Gaudreault, Simon Berthelot, Maude Laberge. "Validated Tools to Measure Costs for Patients: A Systematic Review", The Patient - Patient-Centered Outcomes Research, 2021

Publication

<1 %

35

docero.net

Internet Source

<1 %

36 ss.bjmu.edu.cn Internet Source <1 %

37 www.medrxiv.org Internet Source <1 %

38 www.termedia.pl Internet Source <1 %

39 Neto, Marina Felicidade Dias. "New Insights into Epidemiology and Pathophysiology of Abdominal Aortic Aneurysms", Universidade do Porto (Portugal), 2024
Publication <1 %

40 Yan Efrata Sembiring, Puruhito Puruhito, Heroe Soebroto, Agung Prasmono et al. "Performing cardiac surgery during COVID-19 pandemic in Surabaya, Indonesia: A single-center retrospective observational study", Asian Cardiovascular and Thoracic Annals, 2021
Publication <1 %

41 www.pcori.org Internet Source <1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On