## **ABSTRACT**

Data of the National Commission for Child Protection in 2008, a total of 62.7% high school teens are not virgins and 21.2% adolescents claimed to have an abortion. Primary data in related SMK year 2009 to 2011 obtained the results of 13 students had dropped out because of pregnancy outside marriage. This study aims to analyze the relationship between the psychosocial aspects of adolescent sexual behavior in SMK Swasta Simo Gunung Surabaya.

The method used in this study were analytical observational, cross-sectional approach. Number of samples 76 vocational students. The analysis used is a bivariate analysis with chi square statistical test to analyze the relationship between the psychosocial aspects of adolescent sexual behavior

The results of bivariate analysis illustrates is the more active adolescent sexual behavior (46.7% bad and 53.3% fairly bad) as the psychosocial aspects of adolescent individuals in the negative category. Chi-square test statistic ( $X^2$ ) using SPSS shows the  $X^2$  count (42.754) is greater than  $X^2$  table (5.591) so that it can be seen a significant relationship.

Chi-square test statistic  $(X^2)$  with  $\alpha = 0.05$ , p = 0.000 value obtained, since the p value  $<\alpha$  (0.000 <0.05), then Ho is rejected then it can be seen there is a relationship between the psychosocial aspects of adolescent sexual behavior.

Keywords: psychosocial aspects, sexual behavior, adolescent