

## DAFTAR PUSTAKA

- Awang, K., Chan, G., Litaudon, H., Ismail, N.H., Martin, M., and Gueritte, F., 2010, 4-Phenylcoumarins from *Mesua elegans* with Acetylcholinesterase Inhibitory Activity, *Bioorg. Med. Chem.*, **18**, 7873–7877.
- Bandaranayake, W.M., Selliah, S.S., Sultanbawa, M.U.S., and Games, D.E., 1975, Xanthones and 4-phenylcoumarins of *Mesua thwaitesii*, *Phytochem.*, **14**, 265-269.
- Cao, S.G., Sim, K.Y., Pereira, J., and Goh, S.H., 1998, Coumarins from *Calophyllum teysmanii* (Guttiferae), *Phytochem.*, **47(6)**, 1051–1055.
- Carpenter, I., McGarry, E. J., and Scheinmann, F., 1970, The Neoflavonoids and 4-Alkylcoumarins from *Mammea africana* G Don, *Tet. Lett.*, **46**, 3983–3986.
- Chahar, M.K., Kumar, D.S.S., Lokesh, T., and Manohara, K.P., 2012, *In-vivo* Antioxidant and Immunomodulatory Activity of Mesuol Isolated from *Mesua ferrea* L. Seed Oil, *Inter. Immunopharm.*, **13**, 386-391.
- Chan,G., Kamarudin, M.N.A., Wong, D.Z.H., Ismail, N.H., Latif, F.A., Hasan,A., Awang, K., and Kadir, H.A., 2012, Mitigation of H<sub>2</sub>O<sub>2</sub>-Induced Mitochondrial Mediated Apoptosis in NG108-15 Cells by Novel Mesuagenin C from *Mesua kunstleri* (King) Kosterm, *Evidence-Bas. Comp. and Alter. Med.*, 1-18.
- Crombie, L., Jones, R.C.F., and Palmer, C.J., 1987, Synthesis of the *Mammea* Coumarins. Part 1. The coumarins of the Mammea A, B and C Series, *J. Chem. Soc., Perkin Trans. 1*, **X**, 317-331.
- Diallo, D., Maiga, A., Diakite, C., and Willcox, M., 2004, *Malaria-5 : Development of an Antimalarial Phytomedicine in Mali*, CRC Press LLC, 1-14.
- Ee, G.C.L., Lim, C.K., Ong, G.P., Sukari, M.A. and Lee, H.L., 2006, Daphnifolin, a New Xanthone from *Mesua daphnifolia* (Guttiferae), *J. Asian Nat. Prod. Res.*, **8(6)**, 567-570.

Ee, G.C.L., Teh, S. S., Kwong, H. C, Mah, S. H., Lim, Y.M., and Rahmani, M., 2012, A new benzophenone from *Mesua congestiflora*, an inhibitor against human B lymphocyte cancer cell line, *Phytochem.*, **5(3)**, 545–548.

Elyzaar, I. R. F., Hay, S. I., and Baird, J. K., 2011, Malaria Distribution, Prevalence, Drug Resistance and Control in Indonesia, *Adv. Parasitol.*, **74**, 41–175.

Fidock, D A., Rosenthal, P J., Croft, S L., Brun, R., Nwaka, S., 2004, Antimalarial Drug Discovery: Efficacy Models For Compound Screening, *Nature Reviews Drug Discovery*, **3**, 509-520.

Firmansyah, M., Yusuf, 2014, Isolasi Dan Identifikasi Senyawa Fenolik dari Akar *Erythrina Ovalifolia* Roxb serta Uji Aktivitas Antimalaria Terhadap *Plasmodium falciparum*, Skripsi, Fakultas Sains dan Teknologi, Universitas Airlangga, Surabaya.

Govindachari, T.R., Pai, B.R., Subramaniam, P.S., Rao, U.R. and Muthukumaraswamy, N, 1967, Constituents of *Mesua ferrea* L.- I, Mesuaxanthone A and Mesuaxanthone B, *Tet.*, **23**, 243-248.

Govindachari, T.R., Pai, B.R., Subramaniam, P.S., Rao, U.R. and Muthukumaraswamy, N, 1967, Constituents of *Mesua ferrea* L.- II, Ferruol A, a new 4-alkylcoumarin, *Tet.*, **23**, 4161-4165.

Gunasekera, S.P., Ramachandran, S., Selliah, S., and Sultanbawa, M.U.S., 1975, Chemical investigation of ceylonese plants. Part XVII. Isolation and structures of the xanthones in the extractives of *Mesua ferrea* L. (form *M. salicina* Pl. and Tr.)(Guttiferae), *J. Chem. Soc., Perkin Trans. 1*, 2447-2450.

Harijanto, P.N., Nugroho, A., Gunawan.,Carta A., 2010, Malaria dari Molekuler ke Klinis, Edisi 2. Buku Kedokteran EGC, Jakarta, Indonesia.

Heyne, K, 1987, Tumbuhan Berguna Indonesia, terjemahan Badan Litbang Kehutanan, Jilid III, Cetakan kesatu, Badan Penelitian dan Pengembangan Kehutanan, Departemen Kesehatan.

Iinuma M, Tosa H, Tanaka T, Riswan S., 1996, Two new dimeric xanthones in *Mesua ferrea*, *Hetero.*, **43(9)**, 1999-2004.

- Islam, R., Ahmed, I., Sikder, A.A., Haque, M.R., Al-Mansur, A., Ahmed, M., Rasheed, M., and Rashid, M.A., 2014, Chemical Investigation of *Mesua nagassarium* (Burm. f.) Kosterm, *J. Bas. & App. Sci.*, **10**, 124-128.
- Kitua, A.Y., and Malebo, H.M., 2004, Malaria Control in Africa and the Role of Traditional Medicine, CRC Press LLC, chapter 1.
- Kohler, I and Siems, K., 2002, In Vitro Antiplasmodial Investigation of Medicinal Plants from El Salvador, *Z. Naturforsch.*, **57c**, 277-278.
- Morel, C., Guilet, D., Oger, J.M., Seraphin, D., Sevenet, T., Wiart, C., Hadi, A.H.A., Richomme, P., and Bruneton, J., 1999, 6-Acylcoumarins from *Mesua racemosa*, *Phytochem.*, **50**, 1243-1247.
- Omar, Van, 2007, In Vitro And In Vivo Antiplasmodial Properties of Some Malaysian Plant Used in Traditional Medicine Infectious Disease, *J. of Pakistan*, **16**.
- Peres, V., and Nagem, T.J., 1997, Trioxygenated Naturally Occurring Xanthones, *Phytochem.*, **44(2)**, 191-214
- Peres, V., Nagem, T.J., and Oliveira, F.F., 2000, Tetraoxygenated Naturally Occurring Xanthones, *Phytochem.*, **55**, 683-710.
- Rajesh, K. P., Manjunatha, H., Krishna, V., and Swamy, B.E.K., 2013, Potential *In Vitro* Antioxidant and Protective Effects of *Mesua ferrea* Linn. Bark Extracts on Induced Oxidative Damage, *Indust. Crops & Prod.*, **47**, 186–198.
- Raju, M.S., Srimannarayana, G., and Rao, N.V.S., 1976, Structure of Mesuaferrone-B, a New Biflavanone from The Stamens of *Mesua ferrea* Linn, *Tet. Lett.*, **49**, 4509-4512
- Ravikumar, S., Inbaneson, S. J., and Suganthi, P., 2012, *In Vitro* Antiplasmodial Activity of Ethanolic Extracts of South Indian Medicinal Plants Against *Plasmodium falciparum*, *Asian Pac. J. Trop. Disease*, 180-183.
- Roy, S.K., Kumari, N., Pahwa, S., Agrahari, U.C., Bhutani, K.K., Jachak, S.M., and Nandanwar, H., 2013, NorA Efflux Pump Inhibitory Activity of Coumarins from *Mesua ferrea*, *Fitoterapia*, **90**, 140–150.

- Saputri, R. D., 2011, Isolasi dan Identifikasi Senyawa Alkaloid dari Daun Pule (*Alstonia Scholaris* L.R.Br) serta Uji Aktivitas Antiplasmodial secara *In Vitro* terhadap *Plasmodium falciparum*, Skripsi, Fakultas Sains dan Teknologi, Universitas Airlangga, Surabaya.
- Singh, S., Gray, A.I., and Waterman, P.G., 1993, Mesuabixanthone-A and Mesuabixanthone-B: Novel Bis-Xanthones from The Stem Bark of *Mesua ferrea* (Guttiferea), *Nat. Prod. Lett.*, **3(1)**, 53-58.
- Teh, S.S., Ee, G.C.L., Rahmani, M., Yap, Y.H.T., Go, R., and Mah, S.H., 2011, Pyranoxanthones from *Mesua ferrea*, *Mole.*, **16**, 5647-5654.
- Teh, S.S., Ee, G.C.L., Mah, S. H., Lim, Y.M., and Rahmani, M., 2012, *Mesua beccariana* (Clusiaceae) a Source of Potential Anticancer Lead Compounds in Drug Discovery, *Mole.*, **17**, 10791–10800.
- Tonmunphean S, Parasuk V, and Kokpol S., 2000, QSAR study of antimalaria activities and artemisinin-heme binding properties obtained from docking calculations, *Quant. Struct.-Act. Relat.*, **19**.
- Verotta, L., Lovaglio, E., Vidari, G., Finzi, P.V., Neri, M.G., Raimondi, A., Parapini, S., Taramelli, D., Riva, A., and Bombardelli, E., 2004, 4-Alkyl and 4-phenylcoumarins from *Mesua ferrea* as Promising Multidrug Resistant Antibacterials, *Phytochem.*, **65**, 2867-2879.
- Walia, S., and Mukerjee, S.K., 1984, Ferrxanthone, a 1,3,5,6-tetraoxxygenated xanthone from *Mesua ferrea*, *Phytochem.*, **23(8)**, 1816-1817.
- Whitmore, T.C, 1973, *Tree Flora of Malaya-Volume Two*. Kuala Lumpur: Longman, 162-236.
- Wright, C.W., 2004, *Laboratory Research : Pharmacological Properties of The Active Constituents of Some Traditional Herbal Antimalarials*, CRC Press LLC, Part 4, Chapter 15.