

DAFTAR PUSTAKA

- Abrams, K. & Sanso, B. (1998). Approximate Bayesian Inference For Random Effects Meta-analysis. *STATISTICS IN MEDICINE*, Vol. 17, pp. 201-218.
- Al-Khafaji, A. & Webb, A.R. (2004). Fluid resuscitation. *Continuing Education in Anaesthesia, Critical Care & Pain*, Vol. 4, No. 4, pp. 127-131.
- Annane, D., Siami, S., Jaber, S., Martin, C., Elatrous, S., Declère, A.D., Preiser, J.C., Outin, H., Troché, G., Charpentier, C., Trouillet, J.L., Kimmoun, A., Forceville, X., Darmon, M., Lesur, O., Reignier, J., Abroug, F., Berger, P., Clec'h, C., Cousson, J., Thibault, L. & Chevret, S. (2013). Effects of Fluid Resuscitation With Colloids vs Crystalloids on Mortality in Critically Ill Patients Presenting With Hypovolemic Shock: The CRISTAL Randomized Trial. *JAMA*, Vol. 310(17):1809-1817.
- Barnes, A.D. & Lee, S.H. (2013). *Koda Kimble and Young's Applied Therapeutics: The Clinical Use of Drugs*, Philadelphia: LIPPINCOTT WILLIAMS&WILKINS.
- Borenstein, M., Hedges, L.V., Higgins, J.P.T. & Rothstein, H.R. (2009). *Introduction to Meta-Analysis*, UK: John Wiley & Sons, Ltd.
- Boussekey, N., Darmon, R., Langlois, J., Alfandari, S., Devos, P., Meybeck, A., Chiche, A., Georges, H. & Leroy, O. (2010). Resuscitation with low volume hydroxyethyl starch 130 kDa/0.4 is not associated with acute kidney injury. *Critical Care*, Vol. 14:R40.
- Brunkhorst, F.M., Engel, C., Bloos, F., Meier-Hellmann, A. & Ragaller, M. (2008). Intensive Insulin Therapy and PentastarchResuscitation in Severe Sepsis. *The New England Journal of Medicine*, Vol. 358(2), pp. 125-39.
- Burns, R.P. (2009). *Business Research Methods and Statistics Using SPSS*: SAGE Publications Ltd
- Erstad, B.L. (2011). Hypovolemic Shock. In *Pharmacotherapy a Patophysiology Approach*. Dipiro, J. (Ed.). Philadelphia: McGraw-Hill.
- Guidet, B., Martinet, O., Boulain, T. & Feissel, M. (2012). Assessment of hemodynamic efficacy and safety of 6% hydroxyethylstarch 130/0.4 vs. 0.9% NaCl fluid

- replacement in patients with severe sepsis: The CRYSTMAS study. *Critical Care*, Vol. 16, pp. R94.
- Gurbuz, H.A., Durukan, A.B., Salman, N., Tavlasoglu, M., Durukan, E., Ucar, H.I. & Yorgancioglu, C. (2013). Hydroxyethyl starch 6%, 130/0.4 vs. a balanced crystalloid solution in cardiopulmonary bypass priming: a randomized, prospective study. *Journal of Cardiothoracic Surgery*, Vol. 8:71.
- Hert, S.D. & Baerdemaeker, L.D. (2014). Why hydroxyethyl starch solutions should NOT be banned from the operating room. *Anaesthesiology Intensive Therapy*, Vol. 46(5): 336–341.
- Higgins, J.P.T., Thompson, S.G., Deeks, J.J. & Altman, D.G. (2003). Measuring inconsistency in meta-analyses. *BMJ*, Vol. 327:557-560.
- Hulse, J.D. & Yacobi, A. (1983). Hetastarch:an overview of the colloid and its metabolism. *Drug Intelligence&Clinical Pharmacy*, Vol. 17(5):334-41.
- James, M.F.M. (2011). Pharmacology of Intravenous Fluid. In *Alternatives to Blood Transfusion in Transfusion Medicine*. A, M. & Linden, P.V.D. (Eds.). Singapore: Blackwell Publishing. pp. 111-117.
- Jungheinrich, C. & Neff, T.A. (2005). Pharmacokinetics of Hydroxyethyl Starch. *Clinical Pharmacokinetic*, Vol. 44(7), pp. 681-699.
- Kaplan, D. & Depaoli, S. (2012). Bayesian Statistical Methods. In *Oxford Handbook of Quantitative Methods*. Oxford: Oxford University Press.
- Kaufar, D.S. & Wade, C.E. (2005). The Epidemiology and Modern Management of Traumatic. *Critical Care*, pp. 9(Suppl 5):S1-S9.
- Kumle, B., Boldt, J., Piper, S., Schmidt, C., Suttner, S. & Salopek, S. (1999). The Influence of Different Intravascular Volume Replacement Regimens on Renal Function in the Elderly. *Anesthesia and Analgesia*, Vol. 89, pp. 1124 –30.
- Mitra, S. & Khandelwal, P. (2009). Are All Colloids Same? How to Select the Right Colloid? *Indian Journal of Anaesthesia*, Vol. 53(5), pp. 592-607.
- Myburgh, J.A. & Mythen, M.G. (2013). Resuscitation Fluids. *The New England Journal of Medicine*, Vol. 369(13), pp. 1243-51.

- Nolan, J. (1999). Fluid replacement. *British Medical Bulletin*, Vol. 55(4):821-43.
- Pearson, K. (1904). Report On Certain Enteric Fever Inoculation Statistics. *British Medical Journal*, Vol. 3, pp. 1243-1246.
- Perner, A., Haase, N., Guttermann, A.B., Tenhunen, J., Klemenzson, G., Åneman, A., Madsen, K.R., Møller, M.H., Elkjær, J.M., Poulsen, L.M., Bendtsen, A., Winding, R. & Steensen, M. (2012). Hydroxyethyl Starch 130/0.42 versus Ringer's Acetate in Severe Sepsis. *The New England Journal of Medicine*, Vol. 367, pp. 124-34.
- Raudenbush, S.W. & Bryk, A.S. (1985). Empirical Bayes Meta-analysis. *Journal of Educational Statistics* Vol. 10(2): 75-98
- Rhee, P., Koustova, E. & Alam, H.B. (2003). Searching for the Optimal Resuscitation Method: Recommendations for the Initial Fluid Resuscitation of Combat Casualties. *The Journal Of Trauma*, Vol. 54, pp. S52–S62.
- Sakr, Y., Payen, D., Reinhart, K., Sipmann, F.S., Zavala, E., Bewley, J., Marx, G. & Vincent, J.-L. (2007). Effects of hydroxyethyl starch administration on renal function in critically ill patients. *British Journal of Anaesthesia*, Vol. 98(2):216-24.
- Schabinski, F., Oishi, J., Tuche, F., Luy, A., Sakr, Y., Bredle, D., Hartog, C. & Reinhart, K. (2009). Effects of a predominantly hydroxyethyl starch (HES)-based and a predominantly non HES-based fluid therapy on renal function in surgical ICU patients. *Intensive Care Medicine*, Vol. 35:1539–1547.
- Schortgen, F., Lacherade, J.-C., Bruneel, F., Cattaneo, I., Hemery, F., Lemaire, F. & Brochard, L. (2001). Effects of hydroxyethylstarch and gelatin on renal function in severe sepsis: a multicentre randomised study. *The Lancet*, Vol. 357: 911–16.
- Spaniol, J.R., Knight, A.R., Zebley, J.L., Anderson, D. & Pierce, J.D. (2007). Fluid Resuscitation Therapy for Hemorrhagic Shock. *Journal of Trauma Nursing*, Vol. 14, No. 3, pp. 152-160.
- Stansby, D., MacLennan, S. & Hamilton, P.J. (2000). Management of massive blood loss: a template guideline. *British Journal of Anaesthesia*, Vol. 85, No. 3, pp. 487-91.

- Sunatrio, S. (2000). *Resusitasi Cairan*, Jakarta: Media Aesculapius.
- Sutton, A.J. & Abrams, K.R. (2001). Bayesian methods in meta-analysis and evidence synthesis. *Statistical Methods In Medical Research*, Vol. 10, pp. 277-303.
- Sutton, A.J. & Higgins, J.P.T. (2008). Recent developments in meta-analysis. *STATISTICS IN MEDICINE*, Vol. 27, pp. 625–650.
- Warren, B.B. & Durieux, M.E. (1997). Hydroxyethyl Starch: Safe or Not? *Anesthesia and Analgesia*, Vol. 84:206-12.
- Wiedermann, C.J. (2014). Reporting bias in trials of volume resuscitation with hydroxyethyl starch. *Wien Klin Wochenschr (The Central European Journal of Medicine)*, Vol. 126:189–194.

