

DAFTAR PUSTAKA

1. Welten GM, et al. The Influence of Aging on the Prognostic Value of Revised Cardiac Risk Index for Postoperative Cardiac Complications in Vascular Surgery Patients . *Eur J Vasc Endovasc Surg*; 2007; 34:632-638.
2. Munro J, Booth A & Nicholl J. Routine Preoperative Testing: a Systematic Review of the Evidence. *Health Technol Assesment*; 1997; 1(12).
3. American Society of Anesthesiologist. Practice Advisory for Preanesthesia Evaluation. *Anesthesiology*; 2002; 96:485-96.
4. Marietta, M. et al., 2006. Pathophysiology of Bleeding in Surgery. *Transplantation Proceedings*, 38, pp.812–814.
5. Al-Khajafi, A. & Webb, A., 2004. Fluid resuscitation. *Critical care & Pain Journal*, 4(4), pp.127–131.
6. Krausz, M.M., 2006. Initial resuscitation of hemorrhagic shock. *World journal of emergency surgery : WJES*, 1, p.14. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1488835/> [Accessed June 12, 2014].
7. Rossaint, R. et al., 2010. Management of bleeding following major trauma: an updated European guideline. *Critical care (London, England)*, 14(2), p.R52. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2360176/>
8. Nolan, J., 1999. Fluid Replacement. *British Medical Bulletin*, 55 (No 4), 821-843
9. Bouglé, A., Harrois, A. & Duranteau, J., 2013. Resuscitative strategies in traumatic hemorrhagic shock. *Annals of Intensive Care*, 3, pp.1–9.
10. Myburgh, J.A. & Mythen, M.G., 2013. Resuscitation Fluids. *The New England Journal of Medicine*, 369, pp.1243–1251.
11. Prough DS, Wolf SW, Funston S, Svensen CH. Acid-Base Fluids and Electrolytes. In : Barash PG, Cullen BF, Stoelting RK, eds. *Clinical Anesthesia*. 5th Ed. Philadelphia: Lippincott Williams & Wilkins; 2006. p175-207
12. Stoelting RK, Miller RD, eds. *Basics of Anesthesia*. 4th Ed Philadelphia : Churchill Livingstone; 2000. p233-46
13. Hall BA, Frigas E, Matesic D, Gillet MD, Sprung J. Case Report : Intraoperative Anaphylactoid Reaction and Hydroxyethyl Starch in Balanced Electrolyte Solution. *Can J Anesthesia* 2006 : 53 : 989-93
14. Mills, G.H., 2007. Hydroxyethyl starch: does our choice of colloid prevent or add to renal impairment? *British journal of anaesthesia*, 98(2), pp.157–9. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC17251208/> [Accessed May 11, 2014].
15. Gan, T.J., 2011. Colloid or Crystalloid : Any Differences in Outcomes ? *IARS Review Course Lecture*, pp.27–32.
16. Standl T, Burneister MA, Schroeder F, Currline E. Hydroxy Ethyl Starch (HES) 130/0,4 Provide Larger and Faster Increases in Tissue Oxygen Tension in Comparison with Prehemodilution Values tahn HES 70/0,5 or HES 200/0,5 in Voluunteeers Undergoing Acute Normovolume Hemodilution. *Anesth Analg* 2003; 96 : 936-43
17. Sunatrio S. Tatalaksana Cairan Intraoperatif dan Pilihan Cairan Naskah Lengkap Kongres Nasional Ikatan Dokter Spesialis Anestesi Indonesia, Makasar 2004. 56-73
18. Myburgh, John., et al. 2012. Hydroxyethyl Starch or saline for Fluid Resuscitation in Intensive Care: The CHEST Study. *The New England Journal of Medicine*, 2012., pp.1-11
19. Perter A, Haase N, et al. 2012: 6S Trial Group. Hydroxyethyl Starch 130/0,42 versus Ringer's acetate in Severe Sepsis. *Nengland J Med*, 2012: 367(2): 124-134

20. Hydroxyethyl Starch Solutions: FDA Safety Communication- Boxed Warning on Increased Mortality and Severe Renal Injury and Risk of bleeding . 2013. Available at: <http://www.fda.gov/safety/Medwatch/Safety Information/ Safety Alert for Human Medical Product/ ucm358349.htm>
21. Annane Djillali, Sianni Shidasp, et al. 2013. Effect of Fluid Resuscitation with Colloids vs Crystalloids on Mortality in Critically Ill Patients Presenting with Hypovolemic Shock: The CRISTAL Study. *The Journal of American Medical Association* , 2013: 310(17). Pp. 1811-1817
22. Guidet B, Martinet O, et al. 2012. Assessment of Hemodynamic Efficacy and safety of 6% Hydroxyethyl Starch 130/0,4 vs 0,9% NaCl Fluid Replacement in Patients with Severe Sepsis: The CRYSTMAS Study. *Critical Care* 2012, 310(17), pp: 1809-1817
23. Heinze, H. et al., 2009. Comparison of HES 130/0,42 and HES 200/0,5 for hemodynamic stabilisation in major urological surgery. *Applied Cardiopulmonary Pathophysiology*, 13, pp.11–19.
24. Kozek-Langenecker, S. a et al., 2008. The effects of hydroxyethyl starch 130/0.4 (6%) on blood loss and use of blood products in major surgery: a pooled analysis of randomized clinical trials. *Anesthesia and analgesia*, 107(2), pp.382–90. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18633012> [Accessed May 10, 2014].
25. Neff TA, Doelberg M, Jungheirinch C, Saverland A, Spahn DR, Stocker R. Repetitive Large Dose Infusion of The Novel Hydroxyethyl Starch 130/0,4 in Patients with Severe Head Injury. *Anesth Analg* 2003; 93: 1453-9
26. Kozek-langenecker, S.A. & Scharbert, G., 2007. Effects of hydroxyethyl starches on hemostasis. *Transfusion Alternatives in Transfusion Medicine*, 9, pp.173–181.
27. Navickis, R.J., Haynes, G.R. & Wilkes, M.M., 2012. Effect of hydroxyethyl starch on bleeding after cardiopulmonary bypass: a meta-analysis of randomized trials. *The Journal of thoracic and cardiovascular surgery*, 144(1), pp.223–30. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22578894> [Accessed May 9, 2014].
28. McCahon, R. & Hardman, J., 2006. [Pharmacology of plasma expanders]. *Anesthesia and Intensive Care Medicine*, 8, pp.79–81. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6202449>.
29. Haisch G, Boldt J, Krebs C, Kumle B, Stuttner S, Schulz A. The Influence od Intravascular Volume Therapy with a New Hydroxyethyl Starch Preaparation (6% HES 130/0,4) on Coagulation in Patients Undergoing Major Abdominal Surgery. *Anesth Analg* 2001; 92: 565-71
30. Tigchelaar, I. et al., 1997. Hemostatic Effects of Three Colloid Plasma Substitutes for Priming Solution in Cardiopulmonary Bypass. *European Journal of Cardio-Thoracic Surgery*, 11, 626-632
31. Topcu, I., et al. 2012. Evaluation of Hemostatic Changes using Thromboelastography after Crystalloid or Colloid Fluid Administration during Major Orthopedic Surgery. *Braz J Med Bio Res* 45(9), 869-874
32. Marcucci, C.E & Schoettker, P., 2015. Perioperative Hemostasis, XVIII, Chapter 9 Intravenous Fluids and Coagulation, p138-140
33. Jin & Yu, 2009. Effects of Acute Hypervolemic Fluid Infusion of Hydroxyethyl Starch and Gelatin on Hemostasis and Possible Mechanisms. *Clin Appl Tromb Hemost Vol 00 Numb 00*, p 1-8
34. Sadler JE. Von Willebrand Factor. *J. Biol. Chem* 1991; 266: 22777-80.
35. Mohri H, Fujimura Y, Shima M, Yoshioka A, Houghten RA, Ruggeri ZM, et al. Structure of the von Willebrand factor domain interacting with glycoprotein Ib. *J Biol Chem* 1988; 263: 17901- 4.

36. Pearson JD. The control of production and release of haemostatic factors in the endothelial cel. *Bailliere's Clin Haematol* 1993; 6: 629-51
37. Fischbach F, Dunning MB. A Manual of Laboratory and Diagnostic Tests. 7th ed. Philadelphia: Lippincott Williams & Wilkins, 2004; 134,141
38. Hambleton J. Diagnosis and incidence of inherited von Willebrand disease. *Curr Opin Hematol* 2001; 8: 306–311.
39. Dutton, R.P., 2012. Haemostatic resuscitation. *Brithis Journal of Anaesthesia*, 109, pp.39–46.
40. Gutierrez, G., Reines, H.D. & Wulf-Gutierrez, M.E., 2004. Clinical review: hemorrhagic shock. *Critical care (London, England)*, 8(5), pp.373–81. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1453876/> [Accessed June 3, 2014].
41. Strunden S Mike, Heckel Kai, Goetz Alwin, Reuter AD. Perioperative Fluid and Volume Management: Physiological basis, tools, and strategies. *Annals of Intensive care* 2011;1(2):1-8.
42. Haljamäe H. Rules of Thumb. In R. G. Hahn, ed. Clinical Fluid Therapy in the Perioperative Setting. Cambridge: Cambridge University Press 2011:18–21.
43. Kobayashi, L., Costantini, T.W. & Coimbra, R., 2012. Hypovolemic shock resuscitation. *The Surgical clinics of North America*, 92(6), pp.1403–23. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3453876/> [Accessed May 7, 2014].
44. Sunatrio, S., 2000. *Resusitasi Cairan*, Jakarta: Media Aesculapius.
45. Kozek-Langenecker, S. a, 2005. Effects of hydroxyethyl starch solutions on hemostasis. *Anesthesiology*, 103(3), pp.654–60. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC16129993/>.
46. Warren, B.B. & Durieux, M.E., 1997. Starch: Safe or Not? *Anesthesia and analgesia*, 84, pp.206–212.
47. Jungheinrich, C. & Neff, T. a, 2005. Pharmacokinetics of hydroxyethyl starch. *Clinical pharmacokinetics*, 44(7), pp.681–99. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC15966753/>.
48. De Jonge, E. et al., 2001. Decreased circulating levels of von Willebrand factor after intravenous administration of a rapidly degradable hydroxyethyl starch (HES 200/0.5/6) in healthy human subjects. *Intensive care medicine*, 27(11), pp.1825–9. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC11810130/> [Accessed May 9, 2014].
49. Boldt, J., 2005. Intraoperative Fluid Therapy - Crystalloid / Colloid Debate. Conferencias Magistrales Vol 28. Supl I, pp S23 - S28
50. Evans, P.A et al., 1998. Effects of Gelatin based Resuscitation Fluids on Platelet Aggregation. *British Journal of Anaesthesia* 81: 198 - 202
51. De Jonge et al., 1998. Impaired Hemostasis by Intravenous Administration of a Gelatin based Plasma Expander in Human Subjects. *Thrombosis & Haemostasis*, 79, 286-90
52. Linden , P.V & Ickx, B.E., 2006. The Effects of Colloid Sollutions on Hemostasis. *Can J Anesth* 53-6, PS 30-S39
53. Veeckman, L, 2003. Crystalloids and Colloids, Personal View on Their Perioperative Use. *Acta Anaesth Belg*, 54, 323-331
54. Mitra, S. & Khandelwal, P. 2009. Are Colloid Same ? How to Select the Right Colloid. *Indian Journal of Anaesthesia*. 53 (5): 592
55. Black, L. et al., 2013. Bloody Easy Coagulation Simplified. *Ontario Regional Blood Coordinating Network*, pp.4–10.
56. Mohri H, Fujimura Y, Shima M, Yoshioka A, Houghten RA, Ruggeri ZM, et al. Structure of the von Willebrand factor domain interacting with glycoprotein Ib. *J Biol Chem* 1988; 263: 17901-4

57. Pearson JD. The control of production and release of haemostatic factors in the endothelial cel. *Bailliere's Clin Haematol* 1993; 6: 629-51
58. Stockschlaeder, M., Schneppenheim, R. & Budde, U., 2014. Update on von Willebrand factor multimers: focus on high-molecular-weight multimers and their role in hemostasis. *Blood coagulation & fibrinolysis : an international journal in haemostasis and thrombosis*, 25(3), pp.206–16. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3969155/> [Accessed July 10, 2014].
59. Sadler, J.e., 2009. Low Von Willebrand factor: Somerimes a Risk Factor and Sometimes a Disease. *Hematology*, pp. 106-112
60. Reininger, A.J., 2008. Function of von Willebrand factor in haemostasis and thrombosis. *Haemophilia*, 14, pp.11–26.
61. Sindunata R & Probohoesodo MY. Faktor Patogenesis dan Diagnosis Penyakit von Willebrand. Ind, Vol 13 No1, Nov 2006: 23-30
62. Wilson, D., 2008. *McGraw-Hill's Manual of Laboratory & Diagnostic Test*, McGraw-Hill Companies
63. Kahlon, a et al., 2013. Quantification of perioperative changes in von Willebrand factor and factor VIII during elective orthopaedic surgery in normal individuals. *Haemophilia : the official journal of the World Federation of Hemophilia*, 19(5), pp.758–64. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3711418/> [Accessed September 4, 2014].