

Pakistan
Journal of

NUTRITION



ISSN: 0022-0174
CODEN: PNUTDH
Volume 46 No. 5 2015

Zinaa Oksana Vladimirova, Rebezov Maksim Borisovich and Vlasovskova Evgeniya Sergeevna

Pakistan Journal of Nutrition Volume 15, Number 3, 249-254, 2018

[Abstract](#) [PDF](#) [References](#)

Actual Nutrition of Indigenous Peoples of the North in Today's Socio-Economic Development of the Arctic [On the Example of the Evenk District Olensk the Republic of Sakha (Yakutia)]

Vedim Grigorovich Khrosopkin, Anna Immedentseva Sitsova, Elena Nikolaevna Sitsova, Svetlana Semenovna Madnosov and Marina Vladimirova Khrosopkina

Pakistan Journal of Nutrition Volume 15, Number 3, 255-259, 2018

[Abstract](#) [PDF](#) [References](#)

Correlation Between Body Fat Percentage (BFP) and Basal Energy Expenditure (BEE) in Javanese Women Aged 40-60 Years Old in Sleman, Yogyakarta, Indonesia

Sylvia Salmu, Toto Sudargo and Neni Trihasita Rahmawati

Pakistan Journal of Nutrition Volume 15, Number 3, 260-263, 2018

[Abstract](#) [PDF](#) [References](#)

An Economic Analysis of Kampung Chicken Production Using the Small Water Plant *Azolla microphylla* in Their Feed

Agus Setiadi, Giwanto Imam Santoso, Sumanono, L.D. Hafiduz and A.B. Susanto

Pakistan Journal of Nutrition Volume 15, Number 3, 264-267, 2018

[Abstract](#) [PDF](#) [References](#)

Performance and Protein Efficiency Ratio of Starter Phase Pegagan Ducks Fed Fermented Rations Made from Locally Sourced Materials

Sofia Sandi, Fibri Yosi and Miksuwanti

Pakistan Journal of Nutrition Volume 15, Number 3, 268-273, 2018

[Abstract](#) [PDF](#) [References](#)

LC ESI-MS and FT-IR Analysis of *Dendrothoe pentandra* L. Miq Leaf Methanolic Extracts to Identify Compounds with Progestrone-Like Effects

M. Lusardi and B. Hermanto

Pakistan Journal of Nutrition Volume 15, Number 3, 274-280, 2018

[Abstract](#) [PDF](#) [References](#)

Analyzing Proximate Composition of Macro Nutrients of Sri Lankan Cassava Variety "Kiriakwad"

M.A.D. Somendrika, I. Wickramasinghe, M.A.I. Wasupala and S. Peiris

Pakistan Journal of Nutrition Volume 15, Number 3, 283-287, 2018

[Abstract](#) [PDF](#) [References](#)

Eating Habits, Activity Pattern and Socioeconomic Factors That Affect the Prevalence of Overweight and Obesity among Adolescents in Urban Areas of Enugu State, Nigeria

Ani Peace Nwanneka, Ene-Obong Henrietta Mwach and Uweke Peter Oje

Pakistan Journal of Nutrition Volume 15, Number 3, 288-294, 2018

[Abstract](#) [PDF](#) [References](#)

Relationship of Environmental Condition, Container and Behavior with the Existence of *Aedes aegypti* Mosquito Larvae in an Endemic Area of Dengue Hemorrhagic Fever, Makassar

Ashari Rangid, Ririn Yuchartuti and Hari Ransuki Notobroto

Pakistan Journal of Nutrition Volume 15, Number 3, 295-298, 2018

[Abstract](#) [PDF](#) [References](#)

The Effects of P, S and Mg Supplementation of Oil Palm Fronds Fermented by *Phanerochaete chrysosporium* on Rumens Fluid Characteristics and Microbial Protein Synthesis

Dewi Febriana, Noolman Jamarun, Mardiaty Zain and Khairul

Pakistan Journal of Nutrition Volume 15, Number 3, 299-304, 2018

[Abstract](#) [PDF](#) [References](#)

[Home](#) [Journals](#) [For Authors](#)
[For Subscribers](#) [ASCI](#)

© Science Alert. All Rights Reserved

Editor-in-Chief



Heema Fayez Jayyem
The Hashemite University, Jordan

REGIONAL EDITORS



**Yousef Abd Elwahab Yousef
Mohammed Alfa**
King Abdulaziz University, Saudi Arabia

ASSOCIATE EDITORS



Gubser Ahmad Nayik
Sree Longowal Institute of Engineering
and Technology, India



**Ammar Mohammad Hamood AL-
Faraj**
King Abdulaziz University



Ahsan Akram
University of Agriculture, Faisalabad

TECHNICAL EDITORS



Shreshth Adil Harrod
Shri-Bhadrachalam University of Agricultural
Sciences and Technology of Kadabelli,
India



Anil Parghal
Lovely Professional University, India



Mohammed K. Al-Khuseibi
Sulaiman Caboot University



Wafael Abdel Halimman Ah Sorour
Agricultural Research Center, Cairo
University, Egypt



Nermin Benk
Cankaya Onsekiz Mart University,
Turkey



Motsemmedabach Mohammednasa
Shahid Bahonar University of Kerman



Prithvi Kumar Mandal
Rajiv Gandhi Institute of Veterinary
Education and Research, India



Hawra I. Qudus
North Carolina State University



Moshab Ibrahim-Ahmed Waly
Sulaiman Caboot University, Oman

Article Processing Charge

Subscribe to E-alerts



Indexed In

[Academic Journals Database](#)

[ASCI Database](#)

[Asian Digital Library](#)

[Cambridge Scientific Abstract](#)

[Chemical Abstract Service](#)

[EBSCO](#)

[Genamics](#)

[MLA](#)



UNIVERSITAS AIRLANGGA

FAKULTAS KESEHATAN MASYARAKAT

Kampus C Mulyorejo Surabaya 601115 Telp.031-5920948, 5920949 Fax 031-5924618

Laman : <http://www.fkm.unair.ac.id>; E-mail: info@fkm.unair.ac.id

SURAT KETERANGAN

Nomor : 3148/UN3.1.10/KP/2023

Yang bertandatangan di bawah ini :

Nama : Dr. Santi Martini, dr. M.Kes
NIP : 196609271997022001
Pangkat/Golongan : Pembina / Gol. (IV/a)
Jabatan : Dekan

Dengan ini menerangkan bahwa :

Nama : Dr. Hari Basuki Notobroto, dr., M.Kes
NIP : 196506251992031002
Pangkat/Golongan : Pembina (Gol. IV/a)
Jabatan : Lektor Kepala

Telah melaksanakan penelitian dengan judul sebagai berikut :

No.	Judul Karya Ilmiah	Tahun Pelaksanaan
1	Exploring Midwives' Need and Intention to Adopt Electronic Integrated Antenatal Care (C-14)	2018
2	Risk Factors as an Indicator of Non-Complications Spontaneous Preterm Birth: a Study in Eight Hospitals (C-19)	2020
3	Independency Models of Nursing self-care for Ischemic Stroke Patient (C-23)	2015
4	Indicators of Husband's Role in Pregnancy and Maternity Care (C-27)	2017
5	Container Crane Operator Ergonomics Analysis PT. X Port Of Tanjung Perak, Surabaya (C-40)	2015
6	Influence of the village head leadership-based self-efficacy and personal mastery to total visits of children under 5 years to health (C-42)	2016
7	Relationship of Environmental Condition, Container and Behavior with the Existence of Aedes aegypti Mosquito Larvae in an Endemic Area of Dengue Hemorrhagic Fever, Makassar (C-43)	2016
8	Relationship Early Initiation of Breastfeeding with Exclusive Breastfeeding: A Case-control Study (C-48)	2016
9	Comparison of Results of Measurement Hand Dynamometer with Lactat Acid Blood Plasma for	2016

	Muscle Fatigue Level Indicator Hand Computer Operator (C-52)	
10	Enhancement HIV Health Literacy at Servant of God in Providing Support for Individual at Risk of HIV for Following VCT in Province of NTT (C-55)	2017
11	The Effect of Work Position on Fatigue on the Arm Muscles of Computer Operator (C-59)	2016
12	Warm Pad Reduces Anxiety, Somatic Pain, Strain Pain, Perineal Rupture and Postpartum Blood Volume in Normal Delivery (with Normal Baby Weight Range) (C-65)	2018
13	Hubungan Jumlah Kunjungan ANC dan Stigmatisasi dengan Keikutsertaan Ibu Hamil dalam Tes HIV setelah Konseling oleh Petugas Kesehatan (Di Wilayah Kerja Puskesmas Turi Lamongan) (C-101)	2016
14	Faktor yang Berhubungan dengan Perilaku Ibu Rumah Tangga Melakukan Pemeriksaan Payudara Sendiri (Sadari) (K4) (C-102)	2017
15	Influence of prenatal class to the practice of P4K (Birth Planning and Prevention of Birth Complication) (C-103)	2016
16	Rendahnya Keikutsertaan Pengguna Metode Kontrasepsi Jangka Panjang Pada Pasangan Usia Subur (C-104)	2014
17	Analisis Hubungan Kunjungan Neonatal, Asfiksia dan BBLR dengan Kematian Neonatal (C-106)	2014
18	Pemodelan Regresi Logistik Backward pada Faktor Risiko Kanker Serviks di Yayasan Kanker Wisnuwardhana Surabaya (C-107)	2015
19	Dukungan Informasi tentang Menstruasi kepada Anak Usia Sekolah Dasar (C-108)	2015
20	Pengaruh Aktivitas Seksual Pranikah, Ketaatan Beragama dan Sosial Ekonomi terhadap Kehamilan Remaja di Kecamatan Saptosari Gunungkidul (C-109)	2016
21	Pengaruh Pengetahuan, Kontrol Diri terhadap Perilaku Seksual Pranikah di Kalangan Remaja SMK di Surabaya (C-111)	2016
22	Premarital Sexual Behavior among Papua Women: a Qualitative Research (C-114)	2019
23	The Effect of Organizational Culture and Readiness to Change on Organizational Entrepreneurship: The Role of Competence (C-122)	2021
24	Determinants of Acute Respiratory Infections Among Child Under Five Years in Surabaya (C-131)	2018

Adapun penelitian tersebut layak dilakukan dan menghasilkan output yang sangat baik, meskipun belum ada *Uji Ethical Clearance* karena merupakan penelitian observasional.

Demikian surat keterangan ini kami buat untuk dapat dipergunakan sebagai persyaratan pengusulan Jabatan Fungsional Guru Besar.

Sabaya, 24 April 2023



Dr. Sani Martini, dr. M.Kes
NIP. 96609271997022001

Relationship of Environmental Condition, Container and Behavior with the Existence of *Aedes aegypti* Mosquito Larvae in an Endemic Area of Dengue Hemorrhagic Fever, Makassar

Ashari Rasjid¹, Ririh Yudhastuti² and Hari Basuki Notobroto²

¹Doctoral Program in Health Sciences, Faculty of Public Health, Airlangga University, Surabaya, Indonesia

²Faculty of Public Health, Airlangga University, Campus C Airlangga University, Surabaya-60115, Mulyorejo, Indonesia

Abstract: In Rappocini District, Makassar City, dengue hemorrhagic fever (DHF) is an endemic area. The high incidence of dengue is closely related to the population of *Aedes aegypti* mosquitoes. This study aimed to analyze the relationship between environmental conditions, container and behavior on the existence of the mosquito of *Aedes aegypti*. This was an observational study with a cross-sectional study design. The sample size was 100 houses and they were selected via simple random sampling. In depth interviews, measurements and observations were conducted to achieve the research objective. Data were analyzed by Chi-square and Fisher's exact tests to determine the relationship of environmental conditions (temperature, humidity), container (container type) and the characteristics of residents (knowledge, attitude and actions) with the presence of *Aedes aegypti* larvae. This study found a significant relationship between humidity ($p = 0.000$), type of container ($p = 0.004$), knowledge ($p = 0.001$) and practice ($p = 0.001$) in the presence of *Aedes aegypti* mosquito larvae. Meanwhile, the air temperature and the attitude of the respondents were not significantly related with the results of the study of *Aedes aegypti* larvae. In conclusion, the environmental conditions, container and community characteristics in Rappocini District, Makassar City are not conducive to mosquito nest cleaning (PSN); thus, the density of *Aedes aegypti* is high in this region. The density of larvae showed high dispersion, i.e., HI = 68%, CI = 30.6%, BI = 82% and DF = 8. Public participation in the eradication of dengue disease via eradication of the mosquito *Aedes aegypti* is recommended and would involve cleaning the environment, especially by draining water stores, which are nests for mosquito larvae.

Key words: *Aedes aegypti*, behavior, environmental, mosquito larvae

INTRODUCTION

Dengue hemorrhagic fever (DHF) is an environmental health problem and the number of individuals with DHF tends to increase in increasingly widespread areas, in line with increased population mobility and population density. DHF is caused by dengue virus, which is transmitted by *Aedes aegypti* and *Aedes albopictus*. *Aedes aegypti* has a role in the transmission of this disease, because it lives in and around human homes. Dengue disease is endemic in Makassar because every year definitive cases of dengue fever are recorded somewhere in Makassar. From 2011 to 2013, the number of reported dengue fever cases increased from 1741 (CFR = 0.5) to 2143 (CFR = 0.2) (Dinas Kota Makassar, 2013). The results of periodic monitoring of larvae (CHD) in 2013 showed that the average of larva-free rate in the Rappocini Sub District, Makassar was 87% that indicating that the density of *Aedes aegypti* was still high at that time (Dirjen PPPL, 2009). The existence of *Aedes aegypti* in a geographic area is an indicator of the existence of an *Aedes aegypti* population in the area. Implementing counter measures for DHF is difficult, because the disease is currently incurable. However, the

best way to prevent this disease is to eradicate vector mosquito larvae through eradication of *Aedes spp* nests. This study examined the impact of the eradication of *Aedes aegypti* mosquito larvae on the incidence of dengue in the Village Rappocini Makassar. In order for this intervention to be successful, the relationship of physical environmental conditions, containers and characteristics of residents with the presence of *Aedes aegypti* mosquito larvae must be determined.

MATERIALS AND METHODS

This experiment was conducted in 100 lived-in houses with containers (water reservoirs) in the Village of Rappocini, District Rappocini, Makassar City. This was an observational study with a cross-sectional study design. The study was designed to analyze the state of the object being observed while at the same time attempting to analyze the existing problems. The independent variables were environmental conditions, including air temperature and humidity; type of container and individuals' characteristics, such as knowledge, attitudes and actions. The dependent variable was the

presence of *Aedes aegypti* larvae in Rappocini Village, Rappocini District, Makassar.

The primary data were obtained through interviews and observations. The existence of *Aedes Aegypti* larvae at the homes and the type of container were determined by observation according to the Technical Instructions manual for the eradication of mosquito-borne dengue fever (Dirjen, 2009). Interviews were conducted with the head of the household or its representative at the homes being sampled to determine the characteristics and behavior of members of the society in response to the presence of *Aedes aegypti* larvae. Air temperature data were obtained using a thermometer and air humidity data were obtained using a hygrometer. Secondary data were obtained from the Provincial Health Office of South Sulawesi, Makassar City Health Department, Rappocini Health Center, District Office and Rappocini village and included geographic data, demographics, government and the incidence of dengue hemorrhagic fever (DHF).

RESULTS AND DISCUSSION

Characteristics of respondents: Most respondents were female. Most respondents were between 46 and 55 years of age (38.38%) and 11 respondents (4%) were 15-25 years of age. The education level of the respondents varied; most respondents finished junior level school (35.35%), while only 3 respondents (3%) graduated from a university. Most respondents were house wives who did not work and, thus, had an important role in the mosquito nest eradication program for the prevention of dengue hemorrhagic fever (PSN-DBD) and dengue disease.

Density of *Aedes aegypti* larva: Examination of *Aedes aegypti* larvae in Rappocini sub district, Makassar was conducted by visually examining containers in and outside the home.

Table 1 shows that 100 houses were inspected: 58 (58%) were positive for *Aedes aegypti* mosquito larvae and 42 (42%) were not, resulting in a house index (HI) of 58. Regarding container inspection, of the 268 containers inspected, 82 (30.6%) contained *Aedes aegypti*, resulting in a container index (CI) of 30.6 and a Breteau index (BI)

Relationship between air temperature and *Aedes* mosquito larva: Of 58 homes that contained larvae, 25 (43.1%) had an air temperature conducive to the development of *Aedes aegypti* larvae and 33 (56.9%) did not. Forty percent of homes had an air temperature conducive to the development of *Aedes aegypti* larvae (20-30°C), while 60% had an air temperature that was poor for the development of *Aedes aegypti* larvae. This is because the measurement was only conducted once at the time of the survey, which was between approximately 08:00 and 11:00. The Chi-Square test

Table 1: Distribution of total larvae based on existence of *Aedes aegypti* mosquito larvae in Rappocini Village Makassar in 2014

Checked	Total	---- LQ ----		HI	CI	BI	DF
		Yes	No				
House	100	58	32	58	30.6	82	7
Container	268	82	186				

LQ: Larvae quantity

resulted in a p value = 0.591 ($p > .05$), indicating that there was no significant relationship between air temperature and the presence of *Aedes aegypti* larvae in the Village of Rappocini, Makassar.

Air humidity: In addition to air temperature, air humidity is another environmental condition that can affect the development of *Aedes aegypti* larvae. According to (Glen, 2008; Ramesh *et al.*, 2010; Brisbois and Ali, 2010; Barrera *et al.*, 2006) humidity in the range of 81.5 to 89.5% is optimal for the embriosasi process and mosquito embryo survival.

Container type: The container types observed in the respondents' homes were divided into 3 categories: shelter water for everyday purposes and natural shelter water not for everyday purposes (WHO SEARO, 2004). Of those numbers, about 252 containers (94.0%) were filled for daily use whereas 16 (6%) were not.

Relationship between container type and presence of *Aedes* spp: Two hundred fifty-two containers were filled for daily use and 82 of them were positive for mosquito larvae. This is because the Village of Rapocini, Makassar is a dense residential area with slums and poor environmental sanitation, such that many breeding sites for *Aedes aegypti* mosquito larvae that are present, such as bathtubs, toilets, reservoirs and jars, are rarely cleaned. The mentioned sixteen containers that contain some water were not used for daily drinking purpose, but they were used for vases and for pets drinking and none of them were positive for mosquito larvae. The lack of mosquito larvae in those containers was because the water in them was replaced every day. This is supported by research conducted by Glen (2008) in Manila, Philippines. This study involved several surveys conducted in several cities in Indonesia and showed that the containers with the highest potential for mosquito breeding were those that were used day-to-day, such as drums, jars, bathtubs, toilets, buckets and the like.

Knowledge: Fifty-three respondents (91.4%) had a poor level of knowledge and had larvae in their homes. In contrast, only 5 respondents (8.6%) had a good level of knowledge. This finding was supported by the results of the Chi-Square test, which obtained $p = 0.001$ ($p < .05$), indicating a significant relationship between the level of

knowledge and the presence of *Aedes aegypti* in homes in the Village of Rappocini Makassar. A study by Notoatmodjo (2005); EHP (2008) showed that knowledge of the cognitive domain is very important for the formation of one's actions (overt behavior) and that behavior that is based on knowledge is more lasting than behavior that is not based on knowledge. If a person does not fully understand how mosquito eradication is achieved and is not aware of the factors that affect the existence of larvae, then he/she cannot take appropriate actions; thus, the homes of such unknowledgeable respondents contained *Aedes aegypti*.

Attitude: A good attitude with regard to efforts towards mosquito nest eradication, more than abatisation, was observed in 89% of the respondents. This is because the respondents always answered that their efforts towards PSN are acceptable. The respondents answered "no" when asked whether they approved of the draining of the water reservoirs, despite being conscious of the purpose, namely, to remove *Aedes aegypti* larvae. However, this response is related to the poor condition of the water. The attitude of the respondents was that the question of draining the reservoir remains closed and that it was an unrealistic option; consequently, they agreed with the implementation of PSN efforts and abatisasi despite these efforts not necessarily being in accordance with the respondent's attitude. The results of the Fisher's exact test ($p = 0.113$) ($p > \alpha$) supported this finding and showed no significant relationship between respondent attitudes and the presence of *Aedes aegypti* larvae in the Village of Rappocini, Makassar.

Action: Among respondents with unfavorable actions, 65.5% had *Aedes aegypti* larvae in their homes, while for all respondents 34.5% had *Aedes aegypti* larvae in their homes. This shows that the action of the respondents are very closely linked to the presence of larvae in their homes. The Chi-Square test statistic obtained a $p = 0.001$ ($p < \alpha$), indicating a significant correlation between the actions of the respondents with the presence of *Aedes aegypti* mosquito larvae. According to WHO-SEARO (2004); Gubler *et al.* (2001) people's behavior has an effect on the environment because the environment is an area for the development of such behavior. If people are willing to perform mosquito nest eradication regularly and continuously, they can prevent the growth *Aedes aegypti* mosquito larvae and prevent the illness dengue hemorrhagic fever.

Conclusions: The density of *Aedes aegypti* larvae in the Village of Rappocini, Makassar measured by the parameters HI = 58%, CI = 30.6%, BI = 82% and DF = 7

indicates an increasingly high transmission of *Aedes aegypti* mosquitoes, that is sufficiently fast to enable the spread of the disease DHF. No association was found between temperature and humidity and the presence of larva. The type of container used by the people in the village of Rappocini Makassar was related to the presence of *Aedes aegypti* mosquito larvae. The knowledge and action to reduce or suppress the density of *Aedes aegypti* mosquito larvae were related to the presence of *Aedes aegypti* mosquito larvae. However, the attitudes of the respondents were not related to the presence of *Aedes aegypti* larvae in the Village of Rappocini, Makassar.

Suggestion: The members of the urban Village Community of Rappocini Makassar need to take a more active role in the eradication of dengue disease through eradication of *Aedes aegypti* mosquito breeding by 3 M, especially in draining water reservoirs with regular base and wall brushing and sprinkling abate powder into containers that cannot be drained. These methods of eradication should be disseminated throughout the Village of Rappocini, Makassar via the mass media, schools, places of worship, "PKK/Dasa Wisma/Jumantik" cadres and other community groups.

ACKNOWLEDGEMENTS

The authors are grateful to the Rappocini District, Makassar Municipality for providing permission for the performance of this research and are grateful to the local community who were very kind in providing the necessary information and allowing us to collect some samples from their homes.

REFERENCES

- Barrera, R., M. Amador and G.G. Clark, 2006. Ecological Factors Influencing *Aedes aegypti* (Diptera: Culicidae) Productivity in Artificial Containers in Salinas. Puerto Rico. J. Med. Entomol.
- Brisbois, B.W. and S.H. Ali, 2010. Climate Change, Vector-Borne Disease and Interdisciplinary Research: Social Science Perspectives on an Environment and Health Controversy. Econ. Health, Heidelberg: Springer, 2010.
- Dinas Kesehatan Kota Makassar, 2013. Laporan Tahunan Kegiatan Surveilans DBD Makassar.
- Direktorat Jenderal Pengendalian Penyakit dan Penyehatan Lingkungan, 2009. Profil Pengendalian Penyakit dan Penyehatan Lingkungan Tahun 2008. Jakarta. Dirjen PP and PL.
- EHP, 2008. Dengue Reborn Widespread Resurgence of A Resilient Vector. Environmental Health Perspectives, 9: 116.
- Glen, L. Sia Su, 2008. Correlation of Climatic Factors and Dengue Incidence in Metro Manila, Philippines. Ambio Vol. 37, No.4, June 2008.

- Gubler, J. Duane, Paul Reiter, Kristie L. Ebi, Wendy Yap and Roger Nasci and Jonathan A. Parz, 2001. Climate Variability and Change in the United States: Potential Impacts on Vector and Rodent-Borne Diseases. *Environmental Health Perspectives* Volume 109 May 2001.
- Notoatmodjo, S., 2005. *Metodologi Penelitian*. Jakarta: Rineka Cipta Noor,
- Ramesh, C.D., P. Sharmila, G.P.S. Dhillon and P.D. Aditya, 2010. Climate change and threat of vector-borne diseases in India: Are we prepared? New York/Heidelberg: Springer-Verlag, 106: 763-773.
- WHO-Searo, 2004. *Pencegahan dan Pengendalian Penyakit Demam Dengue dan Demam Berdarah Dengue*. Depkes RI. Jakarta.