

THESIS

**THE EFFECT ORAL ADMINISTRATION OF L-  
ARGININE ON SPERMATOGENESIS AND SERTOLI  
CELLS OF RABBITS TESTES (*Oryctolagus cuniculus*)**



By:

**HENING TYAS PITALOKA**

**SIN 061111122**

**FACULTY OF VETERINARY MEDICINE  
AIRLANGGA UNIVERSITY  
SURABAYA  
2016**

**ENDORSEMENT FORM**


**THE EFFECT ORAL ADMINISTRATION OF L-  
ARGININE ON SPERMATOGENESIS AND SERTOLI  
CELLS OF RABBITS TESTES (*Oryctolagus cuniculus*)**

Research Result  
Submitted in partial fulfillment of the requirement for the degree of  
Bachelor of Veterinary Medicine  
at  
Faculty of Veterinary Medicine, Airlangga University, Surabaya

By:

HENING TYAS PITALOKA  
SIN 061111122

Supervising Committee



Prof. Mas'ud Hariadi, M.Phil, Ph.D., drh  
Supervisor



Dr. Nove Hidajati, drh., M.Kes.  
Co-Supervisor

## DECLARATION

Hereby, I declare that in this thesis entitled:

### **THE EFFECT ORAL ADMINISTRATION OF L- ARGININE ON SPERMATOGENESIS AND SERTOLI CELLS OF RABBITS TESTES (*Oryctolagus cuniculus*)**

There is no work ever published to obtain a college degree in a certain college and to my knowledge there is also no work or opinion ever written or published by others, except those in writing referred to in this paper and mentioned in the references.

Surabaya, 10 August 2016



**Hening Tyas Pitaloka**  
SIN 061111122

Has been assessed in Result Seminar

Date : 1<sup>st</sup> August 2016

**RESULT SEMINAR ASSESSEMENT COMMITTEE**


Head : Suzanita Utama, drh.,M.Phil., Ph.D.

Secretary : Dr. Kadek Rachmawati,drh., M.Kes.

Member : Djoko Legowo, drh., M.Si.

Supervisor : Prof. Mas'ud Hariadi, M.Phil, Ph.D., drh

Co-Supervisor : Dr. Nove Hidajati, drh., M.Kes.





Has been examined

Date : 10<sup>th</sup> August 2016

**THESIS DEFENSE ASSESSMENT COMMITTEE**


Head : Suzanita Utama, drh., M.Phil., Ph.D.

Members : Dr. Kadek Rachmawati, drh., M.Kes.

Djoko Legowo, drh., M.Si.

Prof. Mas'ud Hariadi, M.Phil, Ph.D., drh

Dr. Nove Hidajati, drh., M.Kes.



Surabaya, 10<sup>th</sup> August 2016  
Faculty of Veterinary Medicine  
Universitas Airlangga  
Dean,

A handwritten signature in black ink, appearing to read 'Pudji Srianto', is written over a horizontal line.

Prof. Dr. Pudji Srianto, drh., M.Kes.  
NIP. 195601051986011001

# **THE EFFECT ORAL ADMINISTRATION OF L-ARGININE ON SPERMATOGENESIS AND SERTOLI CELLS OF RABBITS TESTES (*Oryctolagus cuniculus*)**

Hening Tyas Pitaloka

## **ABSTRACT**

L-Arginine is the precursor of nitric oxide (NO). The aim of this research was to know the effect oral administration of L-arginine on spermatogenesis and the number of Sertoli cell of seminiferous tubules of rabbit (*Oryctolagus cuniculus*). This research used sixteen rabbits (6 months old and  $\pm 3$  kg of body weight) were divided into two groups, P0 and P1. P0 is a control group and P1 was given L-arginine 6g/300ml/day for 30 consecutive days. After 30 days treatment, their testicles were collected and processed for histological examination. The data were analyzed by Kruskal Wallis Test for Johnsen's Score and ANOVA for Sertoli cells counting at the significancy level of 5%. The result from statistical analysis showed that treatment with L-arginine did not increase spermatogenesis compared with control group, while the Sertoli cells of rabbit testicular showed that L-arginine increased the number of Sertoli cells.

**Keywords** : spermatogenesis, Sertoli cells, L-Arginine, nitric oxide, testis

## ACKNOWLEDGMENT

First and most of all, praise and thank to ALLAH SWT because of His mercy and guidance this thesis entitled **The Effect Oral Administration of L-Arginine on Spermatogenesis and Sertoli Cells of The Testes of Rabbits (*Oryctolagus cuniculus*)**, have finished.

On this occasion the author would like to give thanks to :

Prof. Dr. Pudji Srianto, drh.m M.Kes, the dean of Faculty of Veterinary Medicine, Universitas Airlangga, for the occasion to study in Faculty of Veterinary Medicine, Universitas Airlangga.

The supervisor committee, Prof. Mas'ud Hariadi, M.Phil, Ph.D., drh, and Dr. Nove Hidajati, drh., M.Kes., for all the time, patience, advice and support that given to the author.

The examiner committee, Suzanita Utama, drh.,M.Phil., Ph.D., Dr. Kadek Rachmawati, drh., M.Kes., Djoko Legowo, drh., M.Si., for all the advices, knowledge and guidance in the completion of my study.

Dr. Agnes Theresia Soelih Estoepangestie, Drh., as my trustee lecturer for her kindness, help, support and had being my second parent during my study in Faculty of Veterinary Medicine, Universitas Airlangga.

All lecturers and staff Faculty of Veterinary Medicine, Universitas Airlangga for the knowledge during my study in Faculty of Veterinary Medicine, Universitas Airlangga.

My research and thesis partners, Arif Prasetyo, dr., Syahroni Ibnu, dr., Fachruddin Aziz, Firdausy Kurnia Maulana, Gavrila Amadea, for great teamwork and motivation.

My father Asmuin, S.Pd., my mother Zuliana Surani for your endless affection, education, and motivation for the success of my study all this time. And my beloved twin sister Wening Tyas Pitaloka thank for always beside me, accompany me to finish this thesis.

My dearest best friends Wulan Arfina Ryani, Putri Agnia Kuswandi dan Eka Ardianta for support, prayed, love, advices, patience and motivation all this time.

And all of my vacationer friends, for the experiences and stories that we have created. All the member of IC 32, for being my second family along this years, and the joyful day we spent together.

The author acknowledges that this writing is still lacking and far from perfect. Therefore, the author expects critics and recommendations that will help in the improvement of this thesis. With a humble heart, the author wishes that this research will be useful for the advancement of science and may give a contribution to the veterinary medicine field and all the people who needs it.

Surabaya, 10 August 2016

Author



## LIST OF CONTENTS

|   | Page |
|---|------|
| ENDORSEMENT FORM .....                                  | ii   |
| DECLARATION .....                                       | iii  |
| IDENTITY .....  | iv   |
| ABSTRACT .....  | vi   |
| ACKNOWLEDGEMENT .....                                   | vii  |
| LIST OF CONTENTS .....                                  | ix   |
| LIST OF TABLES .....                                    | xi   |
| LIST OF FIGURES .....                                   | xii  |
| APPENDICES .....  | xiii |
| ABBREVIATIONS AND SYMBOLICS MEANING .....               | xiv  |
| CHAPTER 1 INTRODUCTION .....                            | 1    |
| 1.1. Background of Research .....                       | 1    |
| 1.2. Problem Statement .....                            | 3    |
| 1.3. Theoretical Base .....                             | 3    |
| 1.4. Aim of Research .....                              | 5    |
| 1.5. Benefit of Research .....                          | 5    |
| 1.6. Hypothesis .....                                   | 5    |
| CHAPTER 2 LITERATURE REVIEW .....                       | 6    |
| 2.1. Reproduction of Male .....                         | 6    |
| 2.1.1. Testes .....                                     | 6    |
| 2.1.2. Seminiferous Tubules.....                        | 7    |
| 2.1.3. Spermatogenesis.....                             | 8    |
| 2.1.4. Sertoli Cells.....                               | 10   |
| 2.2. Nitric Oxide .....                                 | 12   |
| 2.3. L-Arginine .....                                   | 14   |
| 2.3.1. L-Arginine in Male Reproductive System .....     | 17   |
| 2.4. Rabbit ( <i>Oryctolagus cuniculus</i> ) .....      | 18   |
| 2.4.1. Characteristic of Male Rabbit Reproduction ..... | 20   |
| CHAPTER 3 MATERIALS AND METHODS .....                   | 21   |

|   |           |
|---|-----------|
| 3.1. Time and Location of Research .....                      | 21        |
| 3.2. Materials of Research .....                              | 21        |
| 3.2.1. Experimental Animals .....                             | 21        |
| 3.2.2. Research Materials .....                               | 21        |
| 3.2.3. Research Equipments .....                              | 22        |
| 3.3. Research Methods .....                                   | 22        |
| 3.3.1. Animal Preparation .....                               | 22        |
| 3.3.2. Treatment .....  | 22        |
| 3.3.3. Organ Collecting .....                                 | 23        |
| 3.3.4. Histological Preparation of Testes .....               | 24        |
| 3.3.5. Seminiferous Tubules Histological Examination .....    | 24        |
| 3.4. Research Design .....                                    | 25        |
| 3.5. Research Variables .....                                 | 25        |
| 3.5.1. Dependent Variable .....                               | 25        |
| 3.5.2. Independent Variable .....                             | 25        |
| 3.5.3. Control Variable .....                                 | 25        |
| 3.6. Operational Definition of Variables .....                | 26        |
| 3.7. Data Analysis .....                                      | 26        |
| 3.7. Research Framework .....                                 | 27        |
| <b>CHAPTER 4 RESULT .....</b>                                 | <b>28</b> |
| 4.1. Spermatogenesis .....                                    | 28        |
| 4.2. Sertoli Cells .....                                      | 29        |
| <b>CHAPTER 5 DISCUSSION .....</b>                             | <b>31</b> |
| 5.1. Spermatogenesis .....                                    | 31        |
| 5.2. Sertoli Cells .....                                      | 33        |
| 5.3. Relation Between Spermatogenesis and Sertoli Cells ..... | 35        |
| <b>CHAPTER 6 CONCLUSION AND SUGGESTIONS .....</b>             | <b>36</b> |
| 6.1. Conclusion .....   | 36        |
| 6.2. Suggestions .....  | 36        |
| <b>SUMMARY .....</b>  | <b>37</b> |
| <b>REFERENCES .....</b>                                       | <b>39</b> |
| <b>APPENDICES .....</b>                                       | <b>47</b> |

## LIST OF TABLES

|  | Page |
|--|------|
| Table 4.1. The mean and deviation standard of spermatogenesis and Sertoli cell number..... | 28   |




**LIST OF FIGURES**

|   | Page |
|---|------|
| Figure 2.1. Scheme of testis (Trainer, 2007).....   | 7    |
| Figure 2.2. Diagram of spermatogenesis (O'Day, 2010) .....  | 10   |
| Figure 2.3. Histology of testis stains H&E with 1000x magnification<br>(Junqueira, 2007). .....   | 12   |
| Figure 2.4. Schematic presentation of the presumed NO systems in<br>Leydig cells, testicular blood vessels, and seminiferous<br>tubules (Middendorff et al. 1997) ..... | 14   |
| Figure 2.5. NO formation in endothelial cells of blood vessel (Koolman<br>and Roehm, 2005).....   | 15   |
| Figure 2.6. New Zealand White rabbit.....   | 19   |
| Figure 4.1. Histological features of spermatogenesis of rabbit testis.....  | 29   |
| Figure 4.2 Histological features of Sertoli cells.....  | 30   |



## APPENDICES

|  | <b>Page</b> |
|--|-------------|
| Appendix 1. Figures of Seminiferous Tubules .....                      | 47          |
| Appendix 2. Testes Histology Preparation .....                         | 49          |
| Appendix 3. Histology Observation Result of Seminiferous Tubules ..... | 50          |
| Appendix 4. Results of Data Analysis of Spermatogenesis .....          | 54          |
| Appendix 5. Results of Data Analysis of Sertoli Cells Number .....     | 56          |
| Appendix 6. Documentation of Research .....                            | 58          |



**ABBREVIATIONS AND SYMBOLICS MEANING**

|                |   |
|----------------|---|
| <i>et al</i>   | : <i>et alii</i>                              |
| NO             | : Nitric Oxide                                |
| e.g.           | : <i>exempli gratia</i>                       |
| %              | : percent                                     |
| g              | : gram  |
| NOS            | : Nitric Oxide Synthase                       |
| nNOS           | : Neural Nitric Oxide Synthase                |
| eNOS           | : Endothelial Nitric Oxide Synthase           |
| iNOS           | : Inducible Nitric Oxide Synthase             |
| cGMP           | : cyclic Guanosine Monophosphate              |
| H&E            | : Hematoxylin and Eosin                       |
| FSH            | : Follicle-stimulating hormone                |
| cNOS           | : Constitutive Nitric Oxide Synthase          |
| NADPH          | : Nicotinamide Adenine Dinucleotide Phosphate |
| O <sub>2</sub> | : Oxygen                                      |
| ml             | : mililiter                                   |
| mg             | : miligrams                                   |
| kg             | : kilograms                                   |
| h              | : hours                                       |
| ®              | : Registered                                  |
| IM             | : Intra Muscular                              |

## CHAPTER 1 INTRODUCTION

### 1.1 Background of Research

Humans at this time almost all use supplements to improve performance. The management of unusual stress therefore has acquired enormous significance in day-to-day life. It is possible to support the body's adaptation by using food supplements, dietary elements, herbs and minerals for increasing physical and mental performance (Gupta *et al.*, 2004).

Dietary supplements are intended to provide nutrients that may otherwise not be consumed in sufficient quantities. The most supplement common consumed are purposed to improve the energy, to increase the immunity, to maintain the fitness of body condition or even for sexual necessity.

Sometimes male sexual health eventually turned to dysfunction of performance. In male experience it as impotence, known technically as erectile dysfunction. Many environmental, physiological and genetic factors have been implicated in poor sperm function and infertility (Garg *et al.*, 2011). To help boost male sexual performance, they consume L-Arginine as dietary supplement. L-Arginine has become known as a safe and effective sexual nutrient for men. Therefore, oral dosage of L-Arginine as a supplement readily available and with prices which are relatively cheap. L-Arginine treatment was well tolerated, there were neither overt signs of toxicity and stress, nor gross behavioral abnormalities. In the treated rats, the food and water intake, the body weight and the rectal

temperature were not significantly different from those of the controls (Ratnasooriya and Dharmasiri, 2001).

L-Arginine known as semi-essential amino acid, because even though the body normally makes enough of it, supplementation is sometimes needed. L-arginine plays a key role in modulating host defences and cellular immunity. Environmental factors, such as pesticides, exogenous estrogens, and heavy metals may negatively impact spermatogenesis. Arginine therapies have been shown to improve sperm counts and sperm motility (Husien *et al.*, 2011).

L-Arginine is mainly as the precursor of Nitric Oxide (NO). NO is now known to be produced by various cells in different organs, including smooth muscle cells, mesangial cells, neurons, platelets, hepatocytes, macrophages, fibroblasts and epithelial cells. NO regulates smooth muscle cell tone, platelet aggregation and adhesion, cell growth, apoptosis, neurotransmission and injury as well as infection-induced immune reactions. Because these neurons, blood vessels and cells of the immune system are integral part of reproductive organs, and in view of the important functional role that NO plays in those system, it is likely that NO is an important regulator of the biology and physiology of the reproductive system (Rosselli *et al.*, 1998).

Arginine supplementation significantly improved sperm motility and abnormality without any side effect (Scibona *et al.*, 1994). Researcher found out that feeding arginine-deficient diet to adult men for 9 days decreased sperm counts by ~90% and increased the presentage of non-motile sperm approximately 10 fold (Gad, 2010). A study by Chen *et al* (1999) revealed a significant



subjective improvement in sexual function in men with organic erectile dysfunction after oral intake of 5 g L-Arginine for 6 weeks, but only if they had decreased NO excretion or production. Long term oral administration of Arginine to diabetic rabbits increased endothelium-dependent relaxation of rabbit corpus cavernosum (Hupertan *et al.*, 2012)

Supplementation of L-Arginine is expected to affect the regulation of the entire part in the seminiferous tubules. The seminiferous tubules are the site of the germination, maturation and transportation of the sperm cells within the male testes. Spermatogenesis through the process of meiosis takes place within seminiferous tubules. Along the way, the maturing sperm cells receive nutrients and raw materials from Sertoli cells which are located within seminiferous tubules. These component in seminiferous tubules are important roles in reproductive system.

Based on the background above the research untitled the effect of oral L-Arginine administration on spermatogenesis and Sertoli cells of the testes rabbits (*Oryctolagus cuniculus*) has been conducted.

## 1.2 Problem Statement

1. Does oral administration of L-arginine increase spermatogenesis of rabbit?
2. Does oral administration of L-arginine increase Sertoli cells of rabbit?

## 1.3 Theoretical Base

L-Arginine is a physiological nitrogen donor for NOS-catalysed reactions; therefore, availability of this essential substrate could determine rates of NO generation. The present report shows that NO-producing enzymes (NOS) and

NO receptors present in peritubular lamina propria, Sertoli, and blood vessel cells, suggesting production and activity of NO in these structures (Middendorff *et al.*, 1997).

The endothelium (inner lining) of blood vessels uses NO to signal the surrounding smooth muscle to relax, thus resulting in vasodilation and increasing blood flow. Nitric oxide is highly reactive, yet diffuses freely across membranes (Stryer, 1995). Nitric Oxide has been shown to influence permeability of vascular endothelial cells (Noel *et al.*, 1995; Rumbaut *et al.*, 1995).

In seminiferous tubules, NO-induced cGMP production may mediate relaxation of myofibroblasts. Whereas endothelin, for example, has been shown to be involved in peritubular cell contraction in the rat (Tripiciano *et al.*, 1996). In the peritubular lamina propria, NO may participate in the regulation of the peristaltic activity of the tubules, which in turn, is necessary for sperm transport (Setchell *et al.*, 1994). Furthermore, NO may influence the permeability of the lamina propria and, by this, the transport of nutrients into the tubular lumen for spermatogenesis process (Holstein *et al.*, 1996).

NOS is present and functionally active in testicular blood vessels and seminiferous tubules refers to a local production of NO in these structures, the NO-induced increases in cGMP production observed in isolated tubules may, in part, be caused by NO produced by Sertoli cells. Hence, testicular vasculature and seminiferous tubules are sites of NO production and activity. It is an attractive idea that NO acts locally to regulate the distribution of oxygen, nutrients, and

hormones by testicular vessels, as well as the peristaltic activity of tubules in context of sperm transport (Middendorff *et al.*, 1997).

Because NOS is found in testicular endothelial cells, Sertoli cells, and regulation of eNOS by germ cells, this seemingly suggests that NO is involved in regulating spermatogenesis.

#### **1.4 Aim of Research**

1. To know the effect of oral administration of L-arginine on increasing the spermatogenesis in rabbit.
2. To know the effect of oral administration of L-arginine on increasing Sertoli cell number in rabbits.

#### **1.5 Benefit of Research**

The benefits of this research are:

1. Providing knowledge and information for researchers and public about the effect of L-Arginine as supplementation on male fertility.
2. Leading the therapy development as therapies to address some types of infertility in male.

#### **1.6 Hypothesis**

The hypothesis of this research are:

1. Oral administration of L-arginine increases spermatogenesis in rabbit.
2. Oral administration of L-arginine increases Sertoli cells in rabbit.

## CHAPTER 2 LITERATURE REVIEW

### 2.1 Reproduction of Male

#### 2.1.1 Testes

The testes are a pair of ovoid glands organs are essential for the function of male reproductive system. Testicular cells responsible for the production of sperm and male sex hormone testosterone


Embryological testes develop in the dorsal part of the abdominal cavity in a retroperitoneal position. As the fetus develops, the testicles migrate the abdomen and descend through the inguinal canal into the scrotum where they are found in adults. Because the testicles develop in the lumbar position, they took the blood and nerve supply from this area. The blood supply and the inervation follow their testicles descend. (Breazile *et al.*, 1971).

Testis consists of two networks that include, vaginal tunic viceral and tunica albuginea. Internal, seminiferous tubules, interstitial cells, nerves, blood vessels, rete testis and efferent ductules found. Each testis is covered with glossy hard connective tissue called visceral vaginal tunic (tunica vaginalis propria). It comes from the peritoneum as the testes descend. It serves to support the testis. Directly below the tunica albuginea testes, which also supports the testes and is the connective tissue capsule that surrounds the blood vessels that meander near the surface of the testicle. Septula testis are strands of connective tissue which is a branch of this layer and connected to the mediastinum testis, which is the core of connective tissue of the testes. This network holds the connective tissue of the



seminiferous tubules and interstitial cells in place and give shape and to support the testicles (Sorensen, 1979).

The mediastinum testis contains blood vessels, nerves, lymphatics, and the rete testis which is a portion of duct system of the testis. Rete testis is the mouth of production per lobule before exit from testis through efferen ducts to the epididymis (Figure 2.1). A vascular layer in the tunica albuginea assists in regulating the temperature of the testis. The bulk of the testis consists of seminiferous tubules and interstitial cells. The seminiferous tubules produce the sperm (Breazile *et al.*, 1971).



**Figure 2.1.** Scheme of testis (Trainer, 2007)

### 2.1.2 Seminiferous Tubules

Seminiferous tubule is the site of germination, maturation and transport of sperm in the testes of men. Seminiferous tubules lined by stratified epithelium complexes containing two different populations of cells, spermatogenic cells,

which develop into spermatozoa and Sertoli cells that have the function to support and nutrition.

The epithelium which lining the seminiferous tubules contains two cell types, sustentacular, or Sertoli cells and spermatozoa and their developing germ cells precursor. Sustentacular large cells extend from the base of the epithelium into the lumen of the seminiferous tubules. The shape is irregular because they surround the developing germ cells. Sustentacula cells secrete fluids that bathe the developing germ cells and help with transport spermatozoa out of the tubules to the rete testis and they are required for maturation of spermatozoa (Frandsen *et al.*, 2003). Along the way, the maturing sperm cells receive nutrients and raw materials from the vascular Sertoli cells located in the tubule walls until they become mature primary sperm cells ( spermatozoa).

Line of tubular basement membrane, and the rest in this is the beginning of spermatogenic cells, the spermatogonia. Supporting the basement membrane are collagen fibers (Constantinides, 1974). There is a developmental progression of cells from the periphery of the tubule to the internal lumen called spermatogenesis (Sorensen, 1979).

### **2.1.3 Spermatogenesis**


Spermatogenesis is the term for the processes involved in the formation of a mature male gametes mostly undifferentiated germ cells. Spermatogenesis takes place in the seminiferous tubules. Rounded immature sperm cells undergo mitosis in a row and meiotic division (spermatocytogenesis) and metamorphic changes (spermiogenesis) to produce spermatozoa.

During spermatocytogenesis, spermatogonia proliferate by mitosis. This is the division of cells from the beginning of sperm formation until a change in shape occurs. The original cells in the process of spermatogenesis are the Type A spermatogonia, which may require testosterone for development from embryonic gonocyte (Steinberger and Duckett, 1967). The A type spermatogonia are containing two or more nucleoli and lie dormant on the basement membrane until they divide mitotically, to be other A type cell in dormant form A cells and active B cells that contain only one nucleolus (Sorensen, 1979). Mitosis ends when a B spermatogonium yields two primary spermatocytes.

The diploid number of primary spermatocytes is halved during meiosis. A primary spermatocyte is transformed into two secondary spermatocytes during meiosis I, these cells then in turn are converted into (1N) spermatids during meiosis II. The second meiotic division is rapid. Spermatocytes and spermatids tend to be larger than their ancestral spermatogonia.

In spermiogenesis, stage of development characterized by spermatids transformation into spermatozoa. Spermatids undergo changes in shape during spermiogenesis, into spermatozoa streamline adjusted for fertilization. Each spermatid is transformed into a functional unit of the mature spermatozoon (Sorensen, 1979). Spermiogenesis involving nuclear condensation, acrosome cap formation, and development of the tail. head formed at one end, and the Golgi apparatus creates enzymes that would be the acrosome. The tail of one of centrioles elongated cells into sperm tails (Figure 2.2).

The mature spermatozoa are released from the protective Sertoli cells into the lumen of the seminiferous tubule. The resulting spermatozoa are now mature but lack motility. The non-motile spermatozoa are transported to the epididymis in testicular fluid secreted by Sertoli cells with the aid of peristaltic contraction.



**Figure 2.2.** Diagram of spermatogenesis (O'Day, 2010)

#### 2.1.4 Sertoli Cell

Testicular function is under the control of expression and repression of several gene and gene products, and many of these work through Sertoli cells (Johnson *et al.*, 2008). Sertoli cells are tall simple columnar cells, which span from the basement membrane to the lumen. They surround the proliferating and differentiating germ cells forming pockets around these (Figure 2.3). Sertoli cell functions include: support and nutrition of germ cells; release of mature germ



cells into the lumen; translocation of developing germ cells in an adluminal direction; secretion of androgen binding protein, transferrin, inhibin; cell-cell communication via gap junctions to coordinate spermatogenesis; blood-testis barrier.


In the process of spermatogenesis, Sertoli cells provide structural support and cytokines for sperm cells, regulate nutrition development, and play a crucial role in maintaining a stable micro-environment for spermatogenesis (Lee and Cheng, 2004). The microtubules of Sertoli cells are involved in spermatogenesis through the regulation of cell morphology, intracellular transport, organelle positioning, cell motility, cell division and other physiological processes (Li *et al.*, 2009)

Sertoli cells are the somatic cells of the testis that are essential for testis formation and spermatogenesis. Sertoli cells facilitate the progression of germ cells to spermatozoa via direct contact and by controlling the environment within seminiferous tubules. The regulation of spermatogenesis by FSH and testosterone occurs by the action of these hormones on the Sertoli cells. While the action of testosterone is necessary for spermatogenesis, the action of FSH minimally serves to promote spermatogenic output by increasing the number of Sertoli cells (Griswold, 1998).

Like all epithelial cells, the Sertoli cells are avascular. Sertoli cells support the germ cell progenitors and help to transfer nutrients from the nearby capillaries. The developing spermatogonia rely on the Sertoli cells for all of their nourishment. The blood-testis barrier formed by the Sertoli cells effectively

isolates the developing spermatogonia, spermatocytes, spermatids and mature spermatozoa from blood. Sertoli cells also produce testicular fluid, including a protein that binds to and concentrates testosterone, which is essential for the development of the spermatozoa

The number of germ cells supported by a single Sertoli cells is the best reflection of the functional efficiency of the cells and are usually highly correlated with spermatogenic efficiency (daily sperm production per gram testis) (Russell and Peterson, 1984; França and Russell, 1998).



**Figure 2.3.** Histology of testis stains H&E with 1000x magnification (Junqueira, 2007).

## 2.2 Nitric Oxide

Nitric oxide (NO) is synthesized from L-arginine by the action of NO synthase (NOS), an enzyme existing in three isoforms. Brain NOS (bNOS) or neuronal NOS (nNOS or NOS1) and endothelial NOS (eNOS or NOS3), also referred to as constitutive NOS (cNOS), are responsible for the continuous basal

release of NO and both require calcium/calmodulin for activation (Griffith and Stuehr, 1995; Snyder, 1995). A third isoform is an inducible calcium-independent form (iNOS or NOS2) that is expressed only in response to inflammatory cytokines and lipopolysaccharides (Nussler and Billiar, 1993; Morris and Biliar, 1994).


The varied expression and activity patterns of the NOS enzymes uniquely suit the different functions of the NO generated from each isoform in normal physiologic functions and in disease states (Lei *et al.*, 2013). The neuronal NOS (nNOS) and endothelial NOS (eNOS) are calcium-dependent and produce low level of NO. The inducible NOS (iNOS) is calcium independent and produce large amount of NO. Under normal condition, the activity of iNOS is very low, but it is stimulated during inflammation by bacterial endotoxins (Braga, 2012).

Nitric oxide functions as a neurotransmitter in the nervous system and as a mediator of endothelium dependent relaxation of blood vessels and mediate macrophage tumoricidal and bactericidal action. Nitric oxide is one of the reactive oxygen species has been involved in various physiological mechanisms of cell signaling in many tissues. Nitric oxide is produced by various cells in different organs, including smooth muscle cells, mesengeal cells, neurons, plateletes, hepatosites, macrophages, fibroblasts and epithelial cells. Nitric oxide set the tone of smooth muscle cells, platelete aggregation and adhesion, cell growth, apoptosis, neurotransmission and injuries and infections caused by immune reactions. NO possibility of production sites and activities are shown. physiological effects may include the impact on the production of testosterone by



the Leydig cells, peritubular myofibroblasts relaxation and dilation of blood vessels testis. Because NO can freely diffuse across the membrane, three different systems can affect one another (Middendorff *et al.* 1997) (Figure 2.4).

The processes are known to be associated with biology, physiology and pathophysiology of various reproductive processes and NO has been recognized as a molecule that importantly regulates the biology and physiology of reproductive function (VidyaGarg and Garg, 2011). The nitrogen derivate free radical nitric oxide also appear to play a significant role in reproduction and fertilisation (Rosselli *et al.*, 1998).



**Figure 2.4.** Schematic presentation of the presumed NO systems in Leydig cells, testicular blood vessels, and seminiferous tubules (Middendorff *et al.* 1997).


### 2.3 L-Arginine

L-Arginine (2-amino-5-guanidino-pentanoic acid) is a conditionally essential amino acid that is a natural constituent of dietary proteins. Arginine is a



stable nutrient in an aqueous solution, and is not destroyed by sterilization conditions (e.g. high temperature and high pressure). Arginine is not toxic and its administration is generally safe for human and animals (Flynn *et al.*, 2002).

L-arginine is the biological precursor of nitric oxide, an endogenous gaseous messenger molecule involved in a variety of endothelium-dependent physiological functions (Wu and Meininger, 2000), including its critical role in cardiovascular protection and immune support (Duru *et al.*, 2011; Sunita *et al.*, 2000).



**Figure 2.5.** NO formation in the endothelial cells of blood vessel (Koolman and Roehm, 2005).

L-arginine is the only physiological nitrogen donor for NOS-catalysed reaction; hence availability of this essential substrate could determine rates of NO generation. Arginine synthesis as well as its transport into the cells can also influence NO synthesis. Arginine is synthesized from citrulline by the actions of

arginosuccinate synthetase and arginosuccinatelyase (Morris, 1992) and catabolized by arginase (Albina *et al.*, 1988). NO arises from arginine in the endothelial cells of blood vessels triggered by  $\text{Ca}^{2+}$ -calmodulin. NO diffuses from endothelial cells into vascular muscle cells where it leads to the formation of the cGMP, which is the trigger of relaxation of the smooth muscle and vessels dilatation (Koolman and Roehm, 2005) (Figure 2.5).

Arginine rapidly enters the circulation reaches a plasma peak 40-60 minutes after intake (pharmacokinetics are similar in the rat, the pig, the rabbit and human) and slowly decreases during the next 24 hour (Boger *et al.*, 1998). It is thus clear that upon administration Arginine has prolonged contact with the vascular endothelium prior to any other tissue. Arginine is also very well tolerated and can be administrated at high dosages (Barbul, 1986).

It has been shown that endogenous arginine synthesis is couples to NO synthesis (Morris *et al.*, 1994; Nussler *et al.*, 1994). Low concentrations of arginine caused by release of arginase in wound are responsible for reduced NO synthesis (Albina *et al.*, 1988). Apart from synthesis, transport of arginine into the cells can also regulate NOS activity. In this regard, it has been shown that hepatic L-arginine transport, which is normally low, is enhanced during sepsis (Sax *et al.*, 1988; Inoue *et al.*, 1993)

Dietary L-arginine supplementation attenuates the oxidative stress induced by burn injury with a better macrophage response (Tsai *et al.*, 2002). These beneficial effect of L-arginine have been attributed to its dependent formation of NO within the endothelial lining ( Brandes *et al.*, 2000).

### 2.3.1 L-Arginine in Male Reproductive System

L-arginine is a dietary supplement often used to improve the performance of sexual health, for example, erectile dysfunction. L-arginine is also known as sexual nutrients that safe and effective. L-arginine plays an important role in regulating the body's defenses, as well as play an active role in the formation of sperm. Arginine amino acid deficiency can cause a metabolic disorder that causes a decrease in sperm motility and disruption of spermatogenesis process.

Effect of amino acid arginine improves the quality of sperm occurs through multiple mechanisms. Arginine amino acids can protect the sperm plasma membrane from damage caused by lipid peroxidation that is by increasing the production of nitric oxide. The mechanism is similar to the mechanism of antioxidant to protect cells from free radicals (Adams *et al.*, 1994). A study of nitric oxide (NO) have been carried out both in vitro and in vivo and the results showed that nitric oxide (NO) has multiple biological activities and pharmacological among others are increasing fertility through increased testosterone levels and activates the release of luteinizing hormone - releasing hormone (LHRH), which is a hormone stimulator for secretion of FSH and LH hormones that affect the improvement of the process of spermatogenesis ( *Rosselli et al.*, 1998; Elgohary *et al.*, 1999).

A deficiency in L-arginine causes derangement of sperm metabolism leading to decrease in motility loss of spermatogenesis and increase morphology abnormality (Holt and Albanesi, 1994). Furthermore L-arginine plays an

important role in stimulating sperm motility in humans, rabbits, and goats under in vitro conditions (Radany *et al.*, 1981).

L-Arginine is capable of regulating penile erection. Because NO acts as a mediator of penile erection and expressed in the cavernosal smooth muscle cells of the penis (Garban *et al.*, 1997). Within the testis, NO has also been shown to regulate blood flow, cell permeability and contractile function of myofibroblasts, which in turn regulate steroid synthesis and transport (Rosselli *et al.*, 1998).

#### **2.4 Rabbit (*Oryctolagus cuniculus*)**

Rabbit race classification according to binominal system is as follows:

(Linnaeus, 1758)

Kingdom : Animalia  
Phylum : Chordata  
Sub phylum : Vertebrata  
Class : Mamalia  
Ordo : Lagomorpha  
Family : Leporidae  
Sub family : Leporinae  
Genus : *Oryctolagus*  
Species : *Oryctolagus cuniculus*





**Figure 2.6.** New Zealand White rabbit.

Rabbits have a thin skin and dense fur that consists of a soft undercoat and stiff guard hairs. They do not have footpad; instead the feet are covered with thick fur, the skin on the neck is loose and pendulous and forms a pronounced dewlap in females of some breed. The nostrils are sensitive with large number of tactile vibrissae. Scent glands are situated in the deep inguinal spaces that are found on either side of the anus immediately dorsal to urogenital opening. In the male rabbit, the testicles are found in hairless scrotal sacs on either side of the penis. The inguinal canal remains open and the testicles can be retracted into the abdomen. Retraction occurs during periods of sexual inactivity or during periods of insufficient food. Male rabbits have rudimentary nipples (Harcourt-Brown, 2002)

New Zealand white rabbits are the albino offspring of coloured rabbits. The snowy coat of a New Zealand white rabbit is a normal length like other rabbit breed. The body is well haired with both underfur and guard hairs being present. Naked areas of skin are located on the tip of the nose, a small portion of scrotum

of the male and the inguinal spaces in both sexes (Weisbroth *et al.*, 1974). New Zealand white rabbits have a life span of over 6-13 years. The urine volume is usually about 130ml/kg BW/24 h and water intake in rabbits is 50-100 ml/kgBW/24 h (Harcourt-Brown, 2002).

Rabbit has served as a good and practical model in studies of basic spermatogenesis as well as for investigating effects of potential toxins or other agents on that process (Castro *et al.*, 2002).

#### **2.4.1 Characteristic of Male Rabbits Reproduction.**

In the male, the oval-shaped testes within the scrotum remain in communication with the abdominal cavity, where they were at birth. The testicles descend at about two months and can move freely from the scrotum to the abdomen through an opening in the inguinal canal (Campos *et al.*, 2014). The short, back-slanting penis points forward when erect. The male rabbit tends to mature slower than the female. Spermatogenesis begins between days 40 and 50. The testicular tubes become active at about 84 days. The first spermatozoa are present in the ejaculate at about 110 days. Sexual maturity, defined as the moment when daily sperm production ceases to increase, is reached at 32 weeks by New Zealand White rabbits in temperate climates. However, a young buck in these same conditions can be used for reproduction from the age of 20 weeks. Coitus may occur for the first time at about 100 days, but the viability of the sperm cells is very weak or nil in the first ejaculates. So first mating should be timed for age 135 to 140 days (Lebas *et al.*, 1997).

## CHAPTER 3 MATERIALS AND METHODS

### 3.1 Time and Location of Research

This research was held at non-infectious animal laboratory, the Institute of Tropical Disease, Universitas Airlangga, Surabaya, as place for treatment, termination and collecting testes. Preparation and staining of seminiferous tubules histological slides were done at The Laboratory of Pathology, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya. This research was held on February-June 2015.

### 3.2 Material of Research

#### 3.2.1 Expeerimental Animals

This research used 16 healthy male New Zealand White rabbit (*Oryctolagus cuniculus*) with average 6-12 months of ages, 3 kg of body weight collected from rabbit farmers located in Malang, East Java.

#### 3.2.2 Research Materials

Materials used in this research were mineral water, Susu PAP® feed concentrate (PT. Japfa Comfeed Indonesia, Tbk), GNC L-Arginine 500® (Nutra Manufacturing USA) for maintenance and treatment.

Alcohol 70%, ketamine HCL, xylazine, formalin 10% used for organ collecting and preservation.



Absolute alcohol and 70%, 80%, 90%, 95% alcohol concentration, xylol, paraffin, *Hematoxylin-eosin* staining used for staining and preparation of histological materials.

### **3.2.3 Research Equipments**

Equipments used for maintenance in this research were rabbit cages and bottles for drink.

Mask, gloves, syringe 1ml, small plastic containers, forceps, scalpel, blades, surgical scissors were used for preparation of the organ samples.

Object glass, cover glass, pipette, tissue processor automatic, water bath, hot plate, microtome, light microscope (Olympus® CX-21). and camera were used for histological observation.

## **3.3 Research Methods**

### **3.3.1 Animal Preparation**

Rabbits were divided into two groups, then adapted to their new environment for one week. The rabbits were given feed and drunk *ad libitum* and placed in cages made of steel with size 90x60x40 cm, in the room with normal temperature, humidity, and good air ventilation.

### **3.3.2 Treatment**

Rabbits were divided into two groups. First group was used as control, while rabbits in the other group received L-arginine in their drinking water.



L-arginine dosage is determined using dose 2g/kg of body weight/day. L-arginine dissolved in drinking water, 6 gram of L-arginine (12 capsules of L-arginine). The treatment was given for 4 weeks (Okazaki *et al.*, 1997).

Treatment type in this study:

P0(control group) : The rabbits were given drink water only for four weeks.

P1(treatment group) : The rabbits were given treatment with L-arginine dissolved in drinking water for four weeks.

### 3.3.3 Organ Collection

At the end of the period of treatment, rabbits were sacrificed instead of castration, since it was a research groups. All groups of rabbits were injected with a combination of ketamine-xylazine with dose of 25 mg/kg/IM for ketamine and 3mg/kg/IM for xylazine.

After the rabbit has reached a stable level of anaesthesia at about 10 minutes after the injection, and positioned in dorsal recumbence, the fur around the scrotum was shaved. The incision was made through the skin, at the cranial end of the scrotum. The fibrous tunic was incised then the vaginal tunic was exposed. The testis, epididymis and the deferent duct were pushed and advanced through the incision. Removed the testis by cutting through the spermatic cord and cut the ligament of the epididymis and the testis.

The testes from each rabbit were exposed and collected, then fixed in 10% formalin to be used for histology slide.

### 3.3.4 Histological Preparation of Testes

Preparation process of testis and *Hematoxylin- Eosin* staining can be seen in (Appendix 2).

### 3.3.5 Seminiferous Tubules Histological Examination

Data is determined by calculating the average score of spermatogenesis in 20 visual field of seminiferous tubules of each slide preparation in a light microscope with 400 times magnification. For standardization, it used scoring system for testicular biopsies (Johnsen score) (Yama *et al.*, 2013). For the figures of Johnsen's score can be seen in the Appendix 1.

- 1 = no seminiferous epithelium
- 2 = no germinal cells, Sertoli cells only
- 3 = spermatogonia only
- 4 = no spermatozoa or spermatids, few spermatocytes
- 5 = no spermatozoa or spermatids, many spermatocytes
- 6 = no spermatozoa, no late spermatids, few early spermatids
- 7 = no spermatozoa, no late spermatids, many early spermatids
- 8 = less than five spermatozoa per tubule, few late spermatids
- 9 = slightly impaired spermatogenesis, many late spermatids, disorganized epithelium
- 10 = full spermatogenesis

Counting the amount of Sertoli cells can be done by observing 20 of view on 20 seminiferous tubules on each slide preparation. Calculation results of each field

of view in preparation are summed and then calculate the average. Observation are done using a light microscope with magnification of 400 times.

### 3.4 Research Design

This research is an experimental study with Completely Randomized Design (CRD), with two treatments. The number of repetitions in each treatment is determined by formulation  $t(n-1) \geq 15$ , whereas  $n$  for quantity of sample for each treatment and  $t$  is number of experiment (Kusriningrum, 2008).

$$t(n-1) \geq 15$$

$$2(n-1) \geq 15$$

$$2n - 2 \geq 15$$

$$n \geq 8.5 (8)$$

Based on the calculation above, this research used eight rabbits repetition of each group.

### 3.5 Reasearch Variables

#### 3.5.1 Dependent Variable

The histology score of spermatogenesis and the amount of Sertoli cells.

#### 3.5.2 Independent Variable

Oral administration of L-Arginine

#### 3.5.3 Controlled Variables

Breed, sex, age and weight of rabbits.

### 3.6 Operational Definition of Variables

3.6.1 L-Arginine is an amino acid supplement that formed into capsule.

3.6.2 Spermatogenesis is the process in male gametes are produced from male primordial germ cells and transform into spermatozoa. With hematoxylin and eosin staining, nucleus of spermoatogonium, primary and secondary spermatocytes are stained in blue or purple, while the cytoplasm are pink.


3.6.3. Sertoli cells extend from the basement membrane to the luminal surface of the seminiferous epithelium. With hematoxylin and eosin staining, Sertoli cells are recognized in the seminifereous tubules by their pale invaginated irregular nuclei. Nuclei should have a slightly wrinkled nuclear membrane and have prominent nucleoli.

### 3.7 Data Analysis

This research is an experimental study with Completely Randomized Design (CRD). The data were obtained in the form of semi-quantitative data of spermatogenesis and quantitative of Sertoli cells of rabbits testes. Data were presented as mean  $\pm$  standard deviation.. To determine treatment differences was analyzed by Kruskal Wallis Test for Johnsen's Score and ANOVA for Sertoli cells counting at the significancy level of 5% (Kusriningrum, 2008). The data was processed using the program Statistical Package for the Social Sciences (SPSS) 21.0 for Windows.



### 3.8 Research Framework





## CHAPTER 4 RESULT

Microscopic observation was conducted using histological preparations with H.E staining of the rabbit testicles to observed the spermatogenesis and the number of Sertoli cells after given treatment for 4 weeks. There were two groups, in control group (P0) and group with L-arginine administration orally (P1). Observation was done by using a microscope magnification of 400 times on spermatogenesis of 20 seminiferous tubules of each slide preparation then averaged and Sertoli cells were observed in 20 different field of view of each slide preparation then averaged.

### 4.1. Spermatogenesis

The data result of observation and assessment was obtained by using Johnsen's Score method. Data of Johnsen's score for spermatogenesis in each treatment are shown in Table 4.1 (Appendix 3).


**Table 4.1** The mean and deviation standard of spermatogenesis and Sertoli cells number. P0 = for spermatogenesis and number of Sertoli cells without L-Arginine administration and P1 = with L-Arginine administration.

| Group                              | Johnsen's Score for Spermatogenesis (Mean $\pm$ SD) | Amount of Sertoli Cells per Seminiferous Tubules (Mean $\pm$ SD) |
|------------------------------------|---|--|
| P0 ( Control )                     | (8.69 $\pm$ 0.25) <sup>a</sup>                      | (29.11 $\pm$ 1.39) <sup>a</sup>                                  |
| P1 ( L-Arginine 2g/kg of BW/ day ) | (8.59 $\pm$ 0.23) <sup>a</sup>                      | (31.19 $\pm$ 0.92) <sup>b</sup>                                  |

Description: superscript (<sup>a,b</sup>) in the same column showed significant differences between treatments (p <0.05).

Kruskall-Wallis test result (Appendix 4), shows the score of spermatogenesis in the control treatment without the administration of L-arginine not significant different ( $p>0.05$ ) from the treatment of l-arginine administration of a dose of 6g / 300ml / day.

Spermatogenesis of rabbit testis both in control group (P0) and treatment group (P1) can be seen in Figure 4.1.



**Figure 4.1.** Histological features of spermatogenesis of rabbit testis. P0=without L-Arginine administration and P1=with L-Arginine administration of rabbit testis showed spermatogonia (white arrow), Sertoli cells (blue arrow), primary spermatocyte (orange arrow), secondary spermatocyte (red arrow), spermatid (green arrow), spermatozoa (black arrow) (*Hematoxylin-Eosin* staining; 400x magnification).

#### 4.2. Sertoli Cells


Histological observation of Sertoli cells in rabbit testis control and treatment group data are showed in Appendix 3. The means and deviation standards of Sertoli cells in rabbit testis both in the control group and the treatment group are presented in Table 4.1.

ANOVA on the number of Sertoli cells (Appendix 5), shows the number of Sertoli cells in the control without the administration of L-arginine different



significantly ( $p < 0.05$ ) from the treatment of L-arginine administration of a dose of 6g / 300ml / day.

The number of Sertoli cells of rabbit testis both in control group (P0) and treatment group (P1) can be seen in Figure 4.2.



**Figure 4.2.** Histological features of Sertoli cells. P0=without L-Arginine administration P1=with L-Arginine administration . Seminiferous tubule with L-Arginine administration shown by higher amount of Sertoli cells (white arrow) (Hematoxylin-Eosin staining; 400x magnification).



## CHAPTER 5 DISCUSSION

In recent time, the attention is to the use of L-Arginine supplementation used widely to enhance tissue growth and general performance, in treatment of men sterility and prevention of male impotence. Nitric oxide (NO) is synthesized from L-arginine by the action of NO synthase (NOS). Nitric oxide serves as a neurotransmitter in the nervous system and as a mediator of endothelium-dependent relaxation of blood vessels (Moncada *et al.*, 1991; Schmidt and Walter, 1994). Research on nitric oxide (NO) have been carried both in vitro and in vivo and the results showed that nitric oxide (NO) has multiple biological and pharmacological activity, among others, are increasing fertility.

In the testis, NOS has been shown to regulate an array of functions, including sperm motility and maturation, as well as germ cell apoptosis in the testis. NO production and NOS expression was detected in testes (Lee and Cheng, 2004). Peritubular cells, Sertoli cells and testicular blood vessels appear to be sites of NO production and activity (Fujisawa *et al.*, 2001).

This study obtained results that L-arginine may affect the amount of the Sertoli cells and increase spermatogenesis of the testes of rabbits (*Oryzctolagus cuniculus*).

### 5.1 Spermatogenesis

The administration of L-arginine in rabbits for 30 days with a dose 6gram / 300ml / day showed no significant increase in spermatogenesis (Table 4.1).



The lack of statistical significance might be due to that spermatogenesis of experimental animals could have been normal in this study. This result similar to recent studies, the absence of L-arginine effect in healthy young men on vessel size or endothelium-dependent or smooth muscle-dependent vasodilatation given oral L-arginine 21 g daily for three days (Adams *et al.*, 1995) and in experimental conducted by Duru *et al* (2011) in adult Sprague-Dawley rats administrated 15 mg/100 mg body weight L-arginine orally for 8 weeks that showed the normal histological architecture of the seminiferous epithelium and interstitial spaces were essentially normal. Husein *et al* (2011) investigated the effects of L-arginine orally administrated with 0,1mg/100ml water on sperm morphology in albino male mice for 30 days and considered that not showed histological changes compared with control. Same results of another research conducted by Jarad *et al* (2011) in healthy male rats were administrated L-Arginine intraperitonically 200mg/kg body weight for 60 days did not showed histological changes in spermatogenesis. Another similar result obtained from healthy crossbred albino rats were orally administrated 100 mg/kg/day and 200 mg/kg/day for 7 days and had no significant effect on sexual competence (in term of sexual arousal, libido, sexual vigour and sexual performance) (Ratnasooriya and Dharmasiri, 2001).

The most likely explanation is that normal endothelium, which already secretes nitric oxide in the basal state, does not normally produce significantly more nitric oxide even if extra substrate is available (Adams *et al.*, 1995). Another possibly explanation is that supplementation of L-arginine directed for treatment of men sterility, prevention of male impotence (Kocic *et al.*,



2012) and supporting abnormal condition like in treatment for diabetic rat (Jarad *et al.*, 2011), the treatment of L-arginine deficiency and spermatogenic arrest (Jungling and Bunge, 1976). In this study L-Arginine suggested as supplementation that can contribute to the maintenance of normal the spermatogenesis.

## 5.2 Sertoli Cells

The administration of L-arginine in rabbits for 30 days with a dose 6gram / 300cc / day showed that it increased the number of the Sertoli cells (Table 4.2).

The theoretical basis of this study is the NO-induced increases in cGMP production observed in isolated tubules may, in part, be caused by NO produced by Sertoli cells. NO is also produced from the synthesis of endothelial cells formed by the oxidation of L-arginine into citrulline with the help of NOS through biochemical reactions. on vascular endothelial cells are the role of calcium-calmodulin complex required to activate eNOS. NO diffuses from endothelial cells into vascular muscle cells where it leads to the formation of the cGMP, which is the trigger of relaxation of the smooth muscle and vessels dilatation and by this, NO may influence the transport of nutrients (Koolman and Roehm, 2005).

The increase of Sertoli cells number coresponds with previous study that reveals the positive correlation between changes in NO release and FSH secretion. Synthesis of NO is necessary for the manifestation as well as maintenance on FSH secretion. This might be due to the fact that nitric oxide (NO) can activate

guanylate cyclase and thus increase the synthesis of cGMP, which is responsible for the release of FSH (Kosior-Korzecka and Bobowiec, 2006).

FSH increases the rate of proliferation of Sertoli cells (Sharpe *et al.*, 2003). FSH minimally serves to promote spermatogenic output by increasing the number of Sertoli cells (Griswold, 1998). FSH has been implicated as a major factor regulating Sertoli cell proliferation. The increase in secretion of Follicle Stimulating Hormone (FSH) increase the number of Sertoli cells through proliferation (Maechem, *et al.*, 1996).

The number of Sertoli cells determines testicular size. Change in diameter of seminiferous tubules is a major determinant of testicular weight. Increases in seminiferous tubule length and in the absolute volume of seminiferous epithelium and lumen. This increase in epithelial volume was associated with proportionate increases in numbers of Sertoli cells (Orth, *et al.*, 1988).

It was concluded that testes containing a larger total number of Sertoli cells were heavier than testes with fewer Sertoli cells. Larger testes would contain a greater volume of seminiferous tubules, characterized by either greater total tubular length and (or) diameter (Berndtson *et al.*, 1987). These statement relates to a previous study conducted by Aziz (2015) that the administration of the same dose of L-arginine in rabbits for four weeks increased the diameter of the seminiferous tubules and thickness of epithelium of seminiferous tubules.

### 5.3 Relationship between spermatogenesis and Sertoli cells

The function and efficiency of Sertoli cells appear to be limiting to germ cell numbers that the number of germ cells appears to be directly related to the number of Sertoli cells and probably to their synthetic capability (Griswold, 1995).

Because of their intimate physical relationship with the germ cells and the variety of supportive roles they perform, it seemed reasonable to speculate that the absolute numerical size of the Sertoli cell population might establish the upper limit for spermatozoal production in testis. The adult population of Sertoli cells to be stable throughout life and therefore complicated equipped to fully support spermatogenesis (Russell et al., 1990).

Based on this study from the obtained result that there were no significant changes in spermatogenesis but did significant changes in number of Sertoli cells. This condition is contraditive with the basic theory in male reproductive system. This can occur due to the formation of spermatogenesis in experimental animals has been in a state of maximum level, so that the increase of Sertoli cells number does not affect the increase in spermatogenesis. Hence, the L-arginine administered increased the number of Sertoli cells did not associated by spermatogenesis changes, but no study has yet shown any explanation for this condition.

## CHAPTER 6 CONCLUSION AND SUGGESTIONS

### 6.1 Conclusion

The conclusion of this study is

1. Oral administration of L-arginine can not increase spermatogenesis of rabbit.
2. Oral administration of L-arginine can increase number of Sertoli cells of rabbit.

### 6.2 Suggestions

Based on the result of research, there are some suggestions:

1. While the explanation about the increase number of Sertoli cells do not associated with spermatogenesis changes in rabbit testes after supplementation of L-arginine, further research is needed.
2. Need some continuation researches about varying doses, periods of use, the way of administration and with different species of experimental animals to know about potential of L-arginine against testicular tissue.



## SUMMARY

**HENING TYAS PITALOKA.** The effect oral administration of L-Arginine to spermatogenesis and sertoli cells of rabbits testes (*Oryctolagus cuniculus*). This research was conducted under the supervision of Prof. Mas'ud Hariadi, M.Phil, Ph.D., drh, as supervisor and Dr. Nove Hidajati, drh., M.Kes., as co-supervisor.

L-Arginine known as semi-essential amino acid, because even though the body normally makes enough of it, supplementation is sometimes needed. L-Arginine mainly as the precursor of Nitric Oxide (NO). NO regulates smooth muscle cell tone, platelet aggregation and adhesion, cell growth, apoptosis, neurotransmission and injury as well as infection-induced immune reactions. Nitric oxide (NO) has multiple biological and pharmacological activity, among others, are increasing fertility carried both *in vitro* and *in vivo*. L-Arginine plays an important role in improvement of erectile dysfunction and stimulating sperm motility in humans, rabbits, and goats under *in vivo* conditions. L-Arginine supplementation significantly improved sperm motility and abnormality without any side effects.

This research used 16 healthy male rabbits. It used complete randomized design with two groups and eight repetitions each. The treatment given by the criteria of male rabbits, 6 months old, with an average weight of 3 kg type of New Zealand and the treatments: P0: standard ration, P1: standard ration with L-Arginine (2g/ kg of BW/ day) administration of l-arginine held for 30 days

preceded by a period of adaptation for seven days. L-arginine doses used in this study is a series of research studies main groups entitle "Pengaruh pemberian l-arginine secara oral pada penggunaan allograft arteri yang dipreservasi dengan tehnik beku kering (freeze drying) pada arteri carotis kelinci (*Oryctolagus cuniculus*) untuk mengurangi stenosis akibat intimal hiperplasia". Observation was done on the spermatogenesis and Sertoli cell microscopically using histological preparations. Data were analyzed by Kruskall Wallis Test for Johnsen's Score and ANOVA for Sertoli cells counting at the significancy level of 5%. Noted that spermatogenesis after treatment of administration of oral L-arginine showed not significant different ( $p>0.05$ ) between group. While on the observation of the Sertoli cells showed significant differences ( $P <0.05$ ).

The results of this study provide information that L-Arginine does not improve spermatogenesis but may increase the number of Sertoli cells of rabbit testes.

## REFERENCES

- Adams, M.R., Forsyth, C.J., Jessup, W., Robinson, J., Celermajer, D.S. 1995. Oral L-Arginine Inhibits Platelet Aggregation but Does Not Enhance Endothelium-Dependent Dilatation in Healthy Young Men. *Journal of The American College of Cardiology*. 26: 1054-1061.
- Albina, J.E., Mills, C.D., Barbul, A. 1988. Arginine Metabolism in Wounds. *American Journal of Physiology*. 254: E459-E467.
- Aziz, F. 2015. Pengaruh Pemberian L-Arginine Secara Oral terhadap Gambaran Histologi Tubulus Seminiferus Testis Kelinci (*Oryctolagus cuniculus*) [Skripsi]. Fakultas Kedokteran Hewan. Universitas Airlangga.
- Barbul, A. 1986. Arginine: Biochemistry, Physiology, and Therapeutic Implications. *Journal of Parenteral and Enteral Nutrition* 10 (2):227-238.
- Berndtson, W.E., Igboeli, G., Pickett, B.W. 1987. Relationship of Absolute Numbers of Sertoli Cells to Testicular Size and Spermatogenesis in Young Beef Bulls. *J. Anita. Sci.* 64: 241-246.
- Braga, M. 2012. Perioperative Immunonutrition and Gut Function. *Curr. Opin. Clin. Nutr. Metab. Care*. 15 (5): 485-488
- Brandes, R.P., Brandes, S., Boger, R.H., Bode-Boger, S.M. and Mugge, A. 2000. L-Arginine Supplementation in Hypercholesterolemic Rabbits Normalizes Leukocyte Adhesion to Non-Endothelial Matrix. *Life Sciences*. 66: 1519-1524.
- Breazile, J.E., Beames, G., Cardielhac, P.T., Newcomer, W.S. 1971. *Textbook of Veterinary of Physiology*. Lea & Febiger. Philadelphia.
- Bode-Boger, S.M., Boger, R.H., Galland, A., Tsikas, D., & Frolich, J.C. 1998. L-Arginine Induced Vasodilation in Healthy Humans, Pharmacokinetic-Pharmacodynamic Relationship. *British Journal of Clinical Pharmacology*. 46: 489-497.
- Campos, A.C.N., Gadelha, C.R.F., Guerreiro, M.E.F., Pereira, E.S., Lima, I.C.S., Linard, M.A.B., Meneses, H.M., Castelo-Branco K.F. and Estevam, F.N.L. 2014. Male Rabbit Reproductive Physiology. *Standard Research Journal of Agricultural Sciences*. 2(8): 120-128.
- Castro, A.C.S., Berndtson, W.E., Cardoso, F.M. 2002. Plasma and Testicular Testosterone Levels, Volume Density and Number of Leydig Cells and



- Spermatogenic Efficiency of Rabbits. *Brazilian Journal of Medical and Biological Research*. 35 (4): 493-498.
- Chen, J., Wollman, Y., Chernichovsky, T., Iaina, A., Sofer, M., Matzkin, H. 1999. Effect of Oral Administration of High-Dose Nitric Oxide Donor L-Arginine in Men with Organic Erectile Dysfunction: Results of A Double-Blind, Randomized, Placebo-Controlled Study. *BJU International*. 83(3): 269-73.
- Constantinides, P. 1974. *Functional Electronic Histology*. Elsevier, New York.
- Duru, F.I.O., Olalekan, O.O., Azu, O.O., Okoko, I.I. 2011. L-Arginine Augments Oxidative Stress in Cryptorchid Testes of Adult Spague-Dawley Rats. *Journal of Medicine and Medical Science*. 2 (4): 777-782.
- Elgohary, M., W.M. Awara., S. Nassar., S. Hawas. 1999. Deltamethrin-Induced Testicular Apoptosis in Rats: The Protective Effect of Nitric Oxide Synthase Inhibitor. *Toxicol* 132: 1-8.
- Franca, L.R., Russell, L.D. 1998. *The Testis of Domestic Animals*. Churchill Livingstone, Madrid.
- Frandsen, R.D., Wilke, W.L., Fails, A.D. 2003. *Anatomy and Physiology of Farm Animals*, 6th ed. Lippincott Williams & Wilkins, Baltimore.
- Flynn, N.E., Meininger, C.J., Haynes, T.E., Wu, G. 2002. The Metabolic Basis of Arginine Nutrition and Pharmacotherapy. *Biomedicine and Pharmacotherapy*. 56: 427-438.
- Fujisawa, M., Yamanaka, K., Tanaka, H., Tanaka, H., Okada, H., Arakawa, S., Kamidono, S. 2001. Expression of Endothelial Nitric Oxide Synthase in Sertoli Cells of Men with Infertility of Various Causes. *British Journal of Urology International*. 87: 85-88.
- Gad, M.Z. 2010. Anti-aging Effects of L-Arginine. *Journal of Advanced Research*. 1: 169-177.
- Garban, H., Marquez, D., Magee, T., *et al.* 1997. Cloning of Rat and Human Inducible Penile Nitric Oxide Synthase: Application for Gene Therapy of Erectile Dysfunction. *Biology of Reproduction*. 56: 954-963.
- Griffith, O.W., Stuehr, D.J. 1995. Nitric Oxide Synthase Properties and Catalytic Mechanism. *Annual Review of Physiology*. 57: 707-736
- Griswold, M.D. 1995. Interactions Between Germ Cells and Sertoli Cells in The Testis. *Biology of Reproduction*. 52: 211-216.



- Griswold, M.D. 1998. The Central Role of Sertoli Cells in Spermatogenesis. *Seminars in Cell and Developmental Biology*. 9 (4): 411-416.
- Yao, G.D., Shu, Y.M., Shi, S.L., Peng, Z.F., Song, W.Y., Jin, H.X., Sun, Y.P. 2014. Expression and Potential Roles of HLA-G in Human Spermatogenesis and Early Embryonic Development. *Plos One*. 9(3): 1-9.
- Gupta, V., Gupta, A., Saggu, S., Divekar, H.M., Grover, S.K. and Kumar, R. 2004. Anti-stress and Adaptogenic Activity of L-Arginine Supplementation. *Oxford University Press*. 2 (1): 93-97.
- Harcourt-Brown, F. 2002. *Textbook of Rabbit Medicine*. Butterworth Heinemann, Oxford.
- Henwood, A. 2010. *Connection Special Stains and H&E*. Volume 14. Dako. USA. 115-120.
- Hikim, A. P. S., Vera, Y., Vernet, D., Castanares, M., Diaz-Romero, M., Ferrini, M., Swerdloff, R. S., Gonzalez-Cadavid, N. F. and Wang, C. 2005. Involvement of Nitric Oxide-Mediated Intrinsic Pathway Signaling in Age-Related Increase in Germ Cell Apoptosis in Male Brown-Norway Rats. *Journal of Gerontology: Biological Sciences*. 60A (6): 702–708.
- Holstein, A.F., Maekawa, M., Nagano, T., Davidoff, M.S. 1996. Myofibroblasts in The Lamina Propria of Human Seminiferous Tubules are Dynamic Structures of Heterogenous Phenotype. *Archives of Histology and Cytology*. 59: 109-125.
- Holt, L.E., Albanesi, A.A. 1994. Observation of Amino Acids Deficiencies in Man. *Transactions of the Association of Americanphysicians Journal*. 58: 143-156.
- Hupertan, V., Neuzillet, Y., Stücker, O., Pons, C., Leammel, E., Leuret, T. 2012. Effects of Nucleotides Adenosine Monophosphate and Adenosine Triphosphate in Combination with L-Arginine on Male Rabbit Corpus Cavernosum Tissue. *International Journal of Andrology*. 35 (6): 860-866.
- Husein, R.H., Ahmed, M.O., Muhammed, S.M. 2011. Effect of L-Arginine, Vitamin E and Their Combinations on Sperms Morphology in Albino Male Mice. *Journla of Al-Nahrain University*. 14 (2): 137-143.
- Inoue, Y., Bode, B.P., Beck, D.J. 1993. Arginine Transport in Human Liver. Characterization and Effects of Nitric Oxide Synthase Inhibitors. *Annals of Surgery*. 218: 350-363.

- Jarad, A.S., Al-Samawy, E.R.M., Al-Badran, A.S.H. 2011. Effect of L-Arginine on Spermatogenesis of The Diabetic Rat. *Basrah Journal of Veterinary Research*. 10 (2): 19-32.
- Johnson, L., Thompson, D.L. Jr., Varner, D.D. 2008. Role of Sertoli Cell Number And Function on Regulation of Spermatogenesis. *Animal Reproduction Science*. 105 (1): 23-51.
- Jungling, M.L. and Bunge, R.G. 1976. The Treatment of Spermatogenic Arrest with Arginine. *Fertility and Sterility*. 27: 282-283.
- Junqueira, LC., 2007. *Persiapan Jaringan untuk Pemeriksaan Mikroskopik. Histology Dasar: teks dan atlas. Edisi 10. Jakarta : EGC.* 3-5.
- Jusuf, A. A. 2009. *Histoteknik Dasar: Rangkaian Proses Histoteknik Pembuatan Sediaan Histologi. Fakultas Kedokteran Universitas Indonesia. Depok.*
- Kocic, G., Nikolic, J., Jevtovic-Stolmenov, T., Sokolovic, D., Kocic, H., Cvetkovic, T., Pavlovic, D., Cencic, A. and Stojanovic, D. 2012. L-Arginine Intake Effect on Adenine Nucleotide Metabolism in Rat Parenchymal and Reproductive Tissues. *The Scientific World Journal*. 2012: 1-4.
- Koolman, J., Roehm, K.H. 2005. *Color Atlas of Biochemistry, 2nd edition. Thieme. Stuttgart.*
- Kosior-Korzecka, U., Bobowiec, R. 2006. Leptin Effect on Nitric Oxide and GnRH-Induced FSH Secretion from Ovine Pituitary Cells in Vitro. *Journal of Physiology and Pharmacology*. 57 (4): 637-647.
- Kusriningrum, R.S. 2008. *Perancangan Percobaan. Edisi Ke-2. Airlangga University Press. Surabaya.* 15.
- Lebas, F., Coudert, P., de Rochambeau, H., Thébault, R.G. 1997. *The Rabbit-Husbandry, Health and Production. Food and Agriculture Organization of The United Nations. Rome.*
- Lee, N.P., Cheng, C.Y. 2004. Adaptors, Junction Dynamics And Spermatogenesis. *Biology of Reproduction*. 71: 392-404.
- Lei, J., Y. Vodovotz, E. Tzeng and T.R. Billiar. 2013. Nitric oxide, A Protective Molecule in The Cardiovascular System. A review. *Nitric Oxide*. 35: 175–185.

- Li, Y., Xian-zhong, W., Meng-bo, Y., Jia-hua, Z. 2009. Effect of Nitric Oxide on Sertoli Cell Microtubule of Piglets. *Chinese Journal of Agricultural Biotechnology*. 6 (3): 257-263.
- Linnaeus, C. 1758. Tomus I. *Systema Naturae Per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, Cum Characteribus, Differentiis, Synonymis, Locis*. Editio Decima, Reformata. Stockholm. Laurentii Salvii: (1-4), 1-824.
- Maechem, S.J., McLachlan, R.I., de Kretser.D.M., Robertson, D.M., Wreford, N.G. 1996. Neonatal Exposure of Rats to Recombinant Follicle Stimulating Hormone Increases Adult Sertoli and Spermatogenic Cell Numbers. *Biology of Reproduction*. 54: 36-44.
- Middendorff, R., Muller, D., Wichers, S., Holstein, A.F. and Davidoff, M.S. 1997. Evidence for Production and Functional Activity of Nitric Oxide in Seminiferous Tubules and Blood Vessels of the Human Testis. *Journal of Clinical Endocrinology and Metabolism*. 82 (12): 4154-4161.
- Moncada, S., Palmer, R.M.J., Higgs, E.A. 1991. Nitric Oxide Physiology, Pathophysiology and Pharmacology. *Pharmacological Reviews*. 43: 109-142.
- Morris, S.M, Jr. 1992. Regulation of Enzymes of Urea and Arginine Synthesis. *Annual Review of Nutrition*. 12: 81-101.
- Morris, S.M., Billiar, T.R. 1994. New Insights into The Regulation of Inducible Nitric Oxide Synthesis. *American Journal of Physiology*. 266: E829-E839.
- Muntiha, M. 2001. Teknik Pembuatan Preparat Histopatologi dari Jaringan Hewan Dengan Pewarnaan Hematoksilin dan Eosin (H&E). Balai penelitian Veteriner. Bogor. 156-163.
- Noel, A.A., Fallek, S.R., Hobson, R.W., Duran, W.N. 1995. Inhibition of Nitric Oxide Synthase Attenuates Primed Microvascular Permeability in The *in vivo* Microcirculation. *Journal of Vascular Surgery*. 22: 661-670.
- Nussler, A.K., Billiar, T.R. 1993. Inflammation, Immunoregulation and Inducible Nitric Oxide Synthase. *Journal of Leukocyte Biology*. 54: 171-178.
- Nussler, A.K., Billiar, T.R., Liu, Z.Z., Morris, S.M. 1994. Coinduction of Nitric Oxide Synthase and Arginosuccinate Synthetase in A Murine Macrophage Cell Line Implications For Regulation of Nitric Oxide Production. *The Journal of Biological Chemistry*. 269: 1257-1261.




- O'Day, D.H. 2010. Formation of the Male Sex Cells: Spermiogenesis. Human Development. University of Toronto, Mississauga.
- Okazaki, J., Komori, K., Kawasaki, K., Eguchi, D., Ishida, M., Sugimachi, K. 1997. L-Arginine Inhibits Smooth Muscle Cell Proliferation of Vein Graft Intimal Thickness in Hypercholesterolemic Rabbits. Cardiovascular Research. 36: 429-436.
- Orth, J.M., Gunsalus, G.L. & Lamperti, A.A. 1988. Evidence from Sertoli Cell-Depleted Rats Indicates That Spermatid Number in Adults Depends on Numbers of Sertoli Cells Produced During Perinatal Development. Endocrinology. 122: 787-794.
- Radany, E.W., Atherton, R.W., Forrester, I.T. 1981. Arginine Uptake in Rabbit Spermatozoa. Archives of Biochemistry and Biophysics. 210: 770-774.
- Ratnasooriya, W. D., Dharmasiri, M.G. 2001. L-Arginine, The Substrate of Nitric Oxide Synthase, Inhibit Fertility of Male Rats. Asian Journal of Andrology. 3 (2): 97-103.
- Rodriguez, I., Ody, C., Araki, K., Garcia, I., Vassalli, P. 1997. An Early and Massive Wave of Germinal Cell Apoptosis is Required for The Development of Functional Spermatogenesis. EMBO Journal. 16 :2262-2270.
- Rosselli, M., Keller, P.J., Dubey, R.K. 1998. Role of Nitric Oxide in The Biology, Physiology and Pathophysiology of Reproduction. Human Reproduction. 4 (1): 3-24.
- Rumbaut, R.E., McKay, M.K., Huxley, V.H. 1995. Capillary Hydraulic Conductivity Is Decreased by Nitric Oxide Synthase Inhibition. American Journal of Physiology. 268: H1856-H1861.
- Russell, L.D., Ettlin, R.A., Sinha-Hikim, A.P. and Clegg, E.D. 1990. Histological and Histopathological Evaluation of the Testis. Cache River Press, Vienna, IL, USA.
- Russell, L.D., Peterson, R.N. 1984. Determination of The Elongate Spermatid-Sertoli Cell Ratio in Various Mammals. Journal of Reproduction and Fertility. 70: 635-664.
- Saeed, K., Tahir, M., Lone, K.P. 2015. Effect of Phoenix Dactylifera (Date Palm) Pit Powder on Nicotine Induced Spermatotoxicity in Adult Albino Mice. Journal of Pakistan Medical Association. 65 (1): 43-48.
- Sandford, J.C. 1996. The Domestic Rabbit, 5th edn. Blackwell Science.




- Sax, H.C., Hasselgren, P.O., Talamini, M.A. 1988. Amino Acid Uptake in Isolated, Perfused Liver: Effect of Trauma and Sepsis. *Journal of Surgical Research*. 45: 50-55.
- Schmidt, H.H.W., Walter, U. 1994. NO at Work. *Cell*. 78: 919-925.
- Scibona, M., Meschini, P., Capparelli, S. 1994. L-Arginine and Male Infertility. *Minerva Urologica e Nefrologica*. 46: 251-253.
- Setchell, B.P., Maddocks, S., Brooks, D.E. 1994. *Anatomy, Vasculature, Innervation, and Fluids of The Male Reproductive Tract*. Raven Press, New York.
- Sharma, R.K., Bhat, R.A., Goyal, A.K., Bhardwaj, J.K. 2015. Germ Cells Apoptosis during Spermatogenesis in Mammals. *Journal of Entomology and Zoology Studies*. 3(3): 506-515.
- Sharpe, R.M., McKinnell, C., Kivlin, C., Fisher, J.S. 2003. Proliferation and Functional Maturation of Sertoli Cells, And Their Relevance to Disorders of Testis Function in Adulthood. *Reproduction*. 125: 769-784.
- Snyder, S.H. 1995. Nitric Oxide: NO Endothelial NO. *Nature*. 377: 196-197.
- Sorensen., A.M. 1979. *Animal Reproduction*. McGraw-Hill. New York.
- Steinberger, E., Duckett, G.E. 1967. Hormonal Control of Spermatogenesis. *Journal of Reproduction and Fertility Supplement*. 2:75-87.
- Sunita, R., Goutam, R., Mishra, S.C., Raviprakash, V. 2000. Role of Nitric Oxide in Central Regulation of Humoral Response in Rats. *Indian Journal of Pharmacology*. 32: 318-320.
- Tripiciano, A., Filippini, A., Giustiniani, Q., Palombi, F. 1996. Direct Visualization of Rat Peritubular Myoid Cell Contraction in Response to Endothelin. *Biology of Reproduction*. 55: 25-31.
- Tsai, H.J., Shang, H.F., Yeh, C.L., Yeh, S.L. 2002. Effects of Arginine Supplementation on Antioxidant Enzyme Activity and Macrophage Response in Burned Mice. *Burns*. 28: 258-263.
- VidyaGarg, Garg, S.P. 2011. Role of Nitric Oxide in Male Infertility. *Journal of Indian Academy of Forensic Medicine*. 33 (1): 65-68.
- Weisbroth, S.H., Flatt, R.E., Kraus, A.L. 1974. *The Biology of The Laboratory Rabbit*. Academic Press, Inc. California.

- Trainer TD. 2007. Testis and Excretory Duct System. In: Mills SE, (ed.). Histology for Pathologists. 3rd edition ed. Virginia: Lippincott Williams & Wilkins, p. 944-60.
- Wu, G., Meininger, C.J. 2000. Arginine Nutrition and Cardiovascular Function. Journal of Nutrition. 130: 2626-2629.
- Yama, O.E., Amah, C.I., Okoko, I.E., Kusemiju, T.O., Ekundayo, O.T. 2013. Fertility Ratio and The Johnsen Histological Scoring of Testicular Tissues in *Momordica charantia*-Treated Male Sprague-Dawley Rats. Journal of Basic Medical Sciences. 1 (1): 22-26.






### Appendix 1. Figures of Seminiferous Tubules



(A) Johnsen Score 10. Spermatogonia close to the basement membrane (dark red arrow) primary spermatocytes (dark blue arrow), round spermatid (brown arrow), elongated spermatid (green arrows) and mature spermatozoa (black arrow) seen. Interstitial cells of Leydig (light blue arrow) conspicuous between

intertubular area. Sertoli cells seen (pink arrow). (B) Johnsen score of 9. All stages of spermatogenic cell line are seen. Round spermatids (green arrow), elongated spermatid (yellow arrow) and mature spermatozoa in large number (dark blue arrow) seen. (C) Johnsen score of 8. Partial restoration of spermatogenic cell line. Elongated spermatids seen (dark blue arrow). (D) Johnsen score of 6. Mature spermatozoa are not discernible. Few elongated spermatids with many rounded spermatids seen. There is sequestration of epithelium towards the center of lumen (dark blue arrow). (E) Johnsen score of 5. Tubule contains spermatogonia (blue arrow), primary spermatocytes (yellow arrow). Spermatids and spermatozoa are not observed. Myoid cells (green arrow) seen outside the basement membrane and disruption of normal arrangement of seminiferous epithelium. HE staining of testicular tissues 400x (Saeed *et al.*, 2015).



(F) Johnsen Score 4. Sertoli cells, spermatogonia and few spermatocytes but not spermatids or mature spermatozoa were found in seminiferous tubules. (G) Johnsen Score 3. Sertoli cells and few spermatogonia were seen in seminiferous tubules. (H) Johnsen Score 2. Only sertoli cells were observed in seminiferous tubules. HE staining of testicular tissues 400x. Source: Yao., *et al.* 2014.



## Appendix 2. Testes Histology Preparation (Jusuf, 2009)

Fixation, collect tissue immediately after euthanasia to prevent post mortem autolysis and decomposition. Place the organ in the container filled with 10% formalin with 1:10 ratio of organ to formalin least 24 hours for preservation. The fixed tissue will dehydration by passing the tissue blocks through ascending grades of alcohol 70%, 80%, 96% and absolute alcohol I, II and III respectively 2 hours each. Clearing by put them into xylol I and II respectively 2 hours each. Embedding the tissues are immersed in melted paraffin I and II then put them inside an oven for 2 hours. Some iron molds are smeared with glycerin in order to prevent adhesion of paraffin to the mold and then tissues that have been sliced into pieces are put in with forceps until paraffin solidified. Paraffin block is cut by microtome about 5  $\mu\text{m}$  thickness, then dip into 40°C of warm water until tissue expanded well. Put down the cutting tissue on an object glass that have been smeared with albumin then dry on a hot plate. Sliced dried tissue is immerses in xylol I and II for 2 minutes each. Next, slice dried tissue rehydrated successively in absolute alcohol, 96%, 80%, 70% alcohol and water for 2 minutes. Then it is immersed in *hematoxylin* for 1 minute, washed by tap water for one minute, immersed into eosin for 5 minutes, put into distilled water for 1-2 minutes, and immersed into 70%, 80%, 90% and absolute alcohol for 5-10 times. Immersed into xylol I and II for 3 minutes. Staining tissues are mounted with liquid adhesive then covered with cover glass.

**Appendix 3. Histology Observation Result of Seminiferous Tubules**

| No | Spermatogenesis |       |      |       |       |      | Sertoli cell Counting |       |       |       |       |       |
|----|-----------------|-------|------|-------|-------|------|-----------------------|-------|-------|-------|-------|-------|
|    | Group           | Score | Mean | Group | Score | Mean | Group                 | Score | Mean  | Group | Score | Mean  |
| 1  | P0.1            | 7     | 8,8  | P1.1  | 10    | 8,55 | P0.1                  | 22    | 27,95 | P1.1  | 34    | 31,3  |
| 2  |                 | 10    |      |       | 9     |      |                       | 32    |       |       | 35    |       |
| 3  |                 | 7     |      |       | 9     |      |                       | 20    |       |       | 39    |       |
| 4  |                 | 10    |      |       | 9     |      |                       | 31    |       |       | 35    |       |
| 5  |                 | 7     |      |       | 10    |      |                       | 23    |       |       | 44    |       |
| 6  |                 | 10    |      |       | 10    |      |                       | 30    |       |       | 43    |       |
| 7  |                 | 10    |      |       | 8     |      |                       | 34    |       |       | 23    |       |
| 8  |                 | 7     |      |       | 9     |      |                       | 24    |       |       | 33    |       |
| 9  |                 | 9     |      |       | 8     |      |                       | 28    |       |       | 24    |       |
| 10 |                 | 7     |      |       | 9     |      |                       | 23    |       |       | 38    |       |
| 11 |                 | 9     |      |       | 9     |      |                       | 31    |       |       | 30    |       |
| 12 |                 | 10    |      |       | 8     |      |                       | 29    |       |       | 28    |       |
| 13 |                 | 9     |      |       | 7     |      |                       | 27    |       |       | 24    |       |
| 14 |                 | 8     |      |       | 8     |      |                       | 27    |       |       | 26    |       |
| 15 |                 | 8     |      |       | 8     |      |                       | 25    |       |       | 25    |       |
| 16 |                 | 9     |      |       | 9     |      |                       | 28    |       |       | 32    |       |
| 17 |                 | 9     |      |       | 7     |      |                       | 31    |       |       | 28    |       |
| 18 |                 | 10    |      |       | 7     |      |                       | 34    |       |       | 23    |       |
| 19 |                 | 10    |      |       | 7     |      |                       | 29    |       |       | 25    |       |
| 20 |                 | 10    |      |       | 10    |      |                       | 31    |       |       | 37    |       |
| 1  | P0.2            | 10    | 8,2  | P1.2  | 8     | 8,55 | P0.2                  | 33    | 26,75 | P1.2  | 25    | 32,45 |
| 2  |                 | 7     |      |       | 8     |      |                       | 19    |       |       | 31    |       |
| 3  |                 | 7     |      |       | 7     |      |                       | 22    |       |       | 25    |       |
| 4  |                 | 8     |      |       | 8     |      |                       | 28    |       |       | 24    |       |
| 5  |                 | 9     |      |       | 7     |      |                       | 29    |       |       | 28    |       |
| 6  |                 | 10    |      |       | 8     |      |                       | 35    |       |       | 38    |       |
| 7  |                 | 8     |      |       | 9     |      |                       | 27    |       |       | 34    |       |
| 8  |                 | 7     |      |       | 10    |      |                       | 23    |       |       | 28    |       |
| 9  |                 | 8     |      |       | 9     |      |                       | 28    |       |       | 38    |       |
| 10 |                 | 9     |      |       | 7     |      |                       | 28    |       |       | 42    |       |
| 11 |                 | 9     |      |       | 10    |      |                       | 32    |       |       | 30    |       |
| 12 |                 | 7     |      |       | 10    |      |                       | 21    |       |       | 26    |       |
| 13 |                 | 7     |      |       | 10    |      |                       | 20    |       |       | 36    |       |
| 14 |                 | 8     |      |       | 8     |      |                       | 28    |       |       | 35    |       |
| 15 |                 | 7     |      |       | 8     |      |                       | 20    |       |       | 38    |       |
| 16 |                 | 9     |      |       | 9     |      |                       | 34    |       |       | 33    |       |
| 17 |                 | 7     |      |       | 8     |      |                       | 19    |       |       | 37    |       |
| 18 |                 | 9     |      |       | 9     |      |                       | 30    |       |       | 33    |       |
| 19 |                 | 10    |      |       | 9     |      |                       | 33    |       |       | 39    |       |

|    |      |    |      |      |    |     |      |    |      |      |    |       |
|----|------|----|------|------|----|-----|------|----|------|------|----|-------|
| 20 |      | 8  |      |      | 9  |     |      | 26 |      |      | 29 |       |
| 1  | P0.3 | 9  | 9,1  | P1.3 | 8  | 8,4 | P0.3 | 28 | 31   | P1.3 | 35 | 31,55 |
| 2  |      | 8  |      |      | 7  |     |      | 29 |      |      | 26 |       |
| 3  |      | 7  |      |      | 8  |     |      | 27 |      |      | 25 |       |
| 4  |      | 8  |      |      | 9  |     |      | 34 |      |      | 32 |       |
| 5  |      | 9  |      |      | 8  |     |      | 27 |      |      | 28 |       |
| 6  |      | 10 |      |      | 8  |     |      | 34 |      |      | 34 |       |
| 7  |      | 10 |      |      | 7  |     |      | 28 |      |      | 24 |       |
| 8  |      | 10 |      |      | 10 |     |      | 30 |      |      | 34 |       |
| 9  |      | 10 |      |      | 8  |     |      | 37 |      |      | 31 |       |
| 10 |      | 10 |      |      | 9  |     |      | 36 |      |      | 38 |       |
| 11 |      | 8  |      |      | 9  |     |      | 30 |      |      | 33 |       |
| 12 |      | 10 |      |      | 10 |     |      | 31 |      |      | 45 |       |
| 13 |      | 9  |      |      | 9  |     |      | 29 |      |      | 32 |       |
| 14 |      | 8  |      |      | 10 |     |      | 29 |      |      | 43 |       |
| 15 |      | 10 |      |      | 7  |     |      | 36 |      |      | 29 |       |
| 16 |      | 10 |      |      | 9  |     |      | 31 |      |      | 32 |       |
| 17 |      | 9  |      |      | 7  |     |      | 34 |      |      | 22 |       |
| 18 |      | 10 |      |      | 9  |     |      | 38 |      |      | 37 |       |
| 19 |      | 9  |      |      | 8  |     |      | 29 |      |      | 23 |       |
| 20 |      | 8  |      |      | 8  |     |      | 23 |      |      | 28 |       |
| 1  | P0.4 | 10 | 8,75 | P1.4 | 8  | 8,6 | P0.4 | 36 | 30,5 | P1.4 | 30 | 30,85 |
| 2  |      | 8  |      |      | 8  |     |      | 29 |      |      | 34 |       |
| 3  |      | 7  |      |      | 9  |     |      | 25 |      |      | 29 |       |
| 4  |      | 8  |      |      | 9  |     |      | 33 |      |      | 31 |       |
| 5  |      | 9  |      |      | 8  |     |      | 30 |      |      | 28 |       |
| 6  |      | 10 |      |      | 9  |     |      | 40 |      |      | 30 |       |
| 7  |      | 10 |      |      | 8  |     |      | 39 |      |      | 31 |       |
| 8  |      | 8  |      |      | 10 |     |      | 28 |      |      | 35 |       |
| 9  |      | 9  |      |      | 10 |     |      | 37 |      |      | 39 |       |
| 10 |      | 10 |      |      | 8  |     |      | 28 |      |      | 25 |       |
| 11 |      | 10 |      |      | 7  |     |      | 29 |      |      | 21 |       |
| 12 |      | 10 |      |      | 7  |     |      | 41 |      |      | 23 |       |
| 13 |      | 7  |      |      | 8  |     |      | 20 |      |      | 41 |       |
| 14 |      | 10 |      |      | 10 |     |      | 29 |      |      | 35 |       |
| 15 |      | 10 |      |      | 10 |     |      | 34 |      |      | 29 |       |
| 16 |      | 8  |      |      | 8  |     |      | 25 |      |      | 25 |       |
| 17 |      | 7  |      |      | 8  |     |      | 28 |      |      | 43 |       |
| 18 |      | 7  |      |      | 10 |     |      | 25 |      |      | 28 |       |
| 19 |      | 7  |      |      | 8  |     |      | 20 |      |      | 31 |       |
| 20 |      | 10 |      |      | 9  |     |      | 34 |      |      | 29 |       |
| 1  | P0.5 | 10 | 8,6  | P1.5 | 9  | 8,3 | P0.5 | 28 | 29,1 | P1.5 | 37 | 30,9  |
| 2  |      | 8  |      |      | 9  |     |      | 29 |      |      | 33 |       |

|    |      |    |     |      |    |      |      |    |       |      |    |      |
|----|------|----|-----|------|----|------|------|----|-------|------|----|------|
| 3  |      | 7  |     |      | 9  |      |      | 25 |       |      | 24 |      |
| 4  |      | 8  |     |      | 10 |      |      | 31 |       |      | 36 |      |
| 5  |      | 9  |     |      | 8  |      |      | 30 |       |      | 38 |      |
| 6  |      | 10 |     |      | 10 |      |      | 29 |       |      | 42 |      |
| 7  |      | 10 |     |      | 9  |      |      | 33 |       |      | 31 |      |
| 8  |      | 10 |     |      | 10 |      |      | 31 |       |      | 40 |      |
| 9  |      | 10 |     |      | 7  |      |      | 36 |       |      | 21 |      |
| 10 |      | 8  |     |      | 9  |      |      | 29 |       |      | 35 |      |
| 11 |      | 7  |     |      | 7  |      |      | 27 |       |      | 28 |      |
| 12 |      | 8  |     |      | 7  |      |      | 28 |       |      | 20 |      |
| 13 |      | 7  |     |      | 8  |      |      | 30 |       |      | 29 |      |
| 14 |      | 10 |     |      | 9  |      |      | 29 |       |      | 28 |      |
| 15 |      | 9  |     |      | 7  |      |      | 26 |       |      | 25 |      |
| 16 |      | 9  |     |      | 7  |      |      | 30 |       |      | 31 |      |
| 17 |      | 7  |     |      | 8  |      |      | 21 |       |      | 34 |      |
| 18 |      | 8  |     |      | 8  |      |      | 30 |       |      | 28 |      |
| 19 |      | 8  |     |      | 7  |      |      | 32 |       |      | 25 |      |
| 20 |      | 9  |     |      | 8  |      |      | 28 |       |      | 33 |      |
| 1  | P0.6 | 8  | 8,6 | P1.6 | 10 | 8,6  | P0.6 | 31 | 28,3  | P1.6 | 39 | 29,5 |
| 2  |      | 8  |     |      | 8  |      |      | 27 |       |      | 24 |      |
| 3  |      | 7  |     |      | 10 |      |      | 29 |       |      | 42 |      |
| 4  |      | 9  |     |      | 10 |      |      | 18 |       |      | 38 |      |
| 5  |      | 10 |     |      | 9  |      |      | 32 |       |      | 30 |      |
| 6  |      | 10 |     |      | 7  |      |      | 29 |       |      | 24 |      |
| 7  |      | 10 |     |      | 10 |      |      | 35 |       |      | 32 |      |
| 8  |      | 8  |     |      | 7  |      |      | 31 |       |      | 30 |      |
| 9  |      | 10 |     |      | 9  |      |      | 34 |       |      | 21 |      |
| 10 |      | 7  |     |      | 7  |      |      | 33 |       |      | 18 |      |
| 11 |      | 7  |     |      | 9  |      |      | 19 |       |      | 28 |      |
| 12 |      | 10 |     |      | 9  |      |      | 20 |       |      | 29 |      |
| 13 |      | 7  |     |      | 8  |      |      | 38 |       |      | 21 |      |
| 14 |      | 8  |     |      | 9  |      |      | 21 |       |      | 29 |      |
| 15 |      | 9  |     |      | 9  |      |      | 24 |       |      | 27 |      |
| 16 |      | 9  |     |      | 8  |      |      | 30 |       |      | 31 |      |
| 17 |      | 9  |     |      | 8  |      |      | 29 |       |      | 34 |      |
| 18 |      | 9  |     |      | 10 |      |      | 28 |       |      | 41 |      |
| 19 |      | 9  |     |      | 8  |      |      | 25 |       |      | 33 |      |
| 20 |      | 8  |     |      | 7  |      |      | 33 |       |      | 19 |      |
| 1  | P0.7 | 10 | 8,7 | P1.7 | 10 | 8,65 | P0.7 | 34 | 29,65 | P1.7 | 43 | 30,8 |
| 2  |      | 7  |     |      | 8  |      |      | 25 |       |      | 29 |      |
| 3  |      | 7  |     |      | 9  |      |      | 22 |       |      | 31 |      |
| 4  |      | 10 |     |      | 9  |      |      | 39 |       |      | 30 |      |
| 5  |      | 9  |     |      | 7  |      |      | 29 |       |      | 29 |      |



|    |      |    |     |      |    |     |      |    |      |      |    |      |
|----|------|----|-----|------|----|-----|------|----|------|------|----|------|
| 6  |      | 10 |     |      | 9  |     |      | 37 |      |      | 24 |      |
| 7  |      | 10 |     |      | 8  |     |      | 39 |      |      | 28 |      |
| 8  |      | 10 |     |      | 9  |     |      | 43 |      |      | 26 |      |
| 9  |      | 7  |     |      | 9  |     |      | 22 |      |      | 28 |      |
| 10 |      | 8  |     |      | 10 |     |      | 25 |      |      | 40 |      |
| 11 |      | 9  |     |      | 8  |     |      | 30 |      |      | 41 |      |
| 12 |      | 8  |     |      | 9  |     |      | 24 |      |      | 36 |      |
| 13 |      | 8  |     |      | 9  |     |      | 28 |      |      | 28 |      |
| 14 |      | 7  |     |      | 7  |     |      | 24 |      |      | 19 |      |
| 15 |      | 9  |     |      | 9  |     |      | 28 |      |      | 22 |      |
| 16 |      | 8  |     |      | 8  |     |      | 28 |      |      | 29 |      |
| 17 |      | 9  |     |      | 9  |     |      | 23 |      |      | 30 |      |
| 18 |      | 9  |     |      | 10 |     |      | 28 |      |      | 43 |      |
| 19 |      | 9  |     |      | 8  |     |      | 34 |      |      | 31 |      |
| 20 |      | 10 |     |      | 8  |     |      | 31 |      |      | 29 |      |
| 1  | P0.8 | 10 | 8,8 | P1.8 | 9  | 9.1 | P0.8 | 34 | 29,6 | P1.8 | 23 | 32,2 |
| 2  |      | 8  |     |      | 8  |     |      | 30 |      |      | 33 |      |
| 3  |      | 7  |     |      | 10 |     |      | 24 |      |      | 45 |      |
| 4  |      | 8  |     |      | 8  |     |      | 33 |      |      | 38 |      |
| 5  |      | 9  |     |      | 8  |     |      | 24 |      |      | 34 |      |
| 6  |      | 10 |     |      | 9  |     |      | 32 |      |      | 30 |      |
| 7  |      | 8  |     |      | 10 |     |      | 35 |      |      | 33 |      |
| 8  |      | 9  |     |      | 8  |     |      | 38 |      |      | 28 |      |
| 9  |      | 10 |     |      | 8  |     |      | 28 |      |      | 23 |      |
| 10 |      | 10 |     |      | 10 |     |      | 28 |      |      | 41 |      |
| 11 |      | 7  |     |      | 10 |     |      | 31 |      |      | 43 |      |
| 12 |      | 8  |     |      | 9  |     |      | 25 |      |      | 24 |      |
| 13 |      | 7  |     |      | 9  |     |      | 27 |      |      | 26 |      |
| 14 |      | 9  |     |      | 10 |     |      | 29 |      |      | 44 |      |
| 15 |      | 9  |     |      | 9  |     |      | 21 |      |      | 27 |      |
| 16 |      | 8  |     |      | 10 |     |      | 25 |      |      | 31 |      |
| 17 |      | 10 |     |      | 9  |     |      | 28 |      |      | 27 |      |
| 18 |      | 10 |     |      | 10 |     |      | 31 |      |      | 31 |      |
| 19 |      | 10 |     |      | 8  |     |      | 41 |      |      | 29 |      |
| 20 |      | 9  |     |      | 10 |     |      | 28 |      |      | 34 |      |

### Appendix 4. Results of Data Analysis of Spermatogenesis

#### Case Processing Summary

| Group              | Cases |         |         |         |       |         |
|--------------------|-------|---------|---------|---------|-------|---------|
|                    | Valid |         | Missing |         | Total |         |
|                    | N     | Percent | N       | Percent | N     | Percent |
| P0=Control Group   | 8     | 100.0%  | 0       | 0.0%    | 8     | 100.0%  |
| P1=Treatment Group | 8     | 100.0%  | 0       | 0.0%    | 8     | 100.0%  |

#### Descriptives

| Group              | Statistic                        | Std. Error  |        |
|--------------------|----------------------------------|-------------|--------|
| P0=Control Group   | Mean                             | 8.6938      | 08986  |
|                    | 95% Confidence Interval for Mean | Lower Bound | 8.4813 |
|                    |                                  | Upper Bound | 8.9062 |
|                    | 5% Trimmed Mean                  | 8.6986      |        |
|                    | Median                           | 8.7250      |        |
|                    | Variance                         | .065        |        |
|                    | Std. Deviation                   | .25416      |        |
|                    | Minimum                          | 8.20        |        |
|                    | Maximum                          | 9.10        |        |
|                    | Range                            | .90         |        |
|                    | Interquartile Range              | .20         |        |
|                    | Skewness                         | -.608       | .752   |
|                    | Kurtosis                         | 2.256       | 1.481  |
| P1=Treatment Group | Mean                             | 8.5938      | .08315 |
|                    | 95% Confidence Interval for Mean | Lower Bound | 8.3971 |
|                    |                                  | Upper Bound | 8.7904 |
|                    | 5% Trimmed Mean                  | 8.5819      |        |
|                    | Median                           | 8.5750      |        |
|                    | Variance                         | .055        |        |
|                    | Std. Deviation                   | .23519      |        |
|                    | Minimum                          | 8.30        |        |
|                    | Maximum                          | 9.10        |        |
|                    | Range                            | .80         |        |
|                    | Interquartile Range              | .20         |        |
|                    | Skewness                         | 1.422       | .752   |
|                    | Kurtosis                         | 3.455       | 1.481  |

#### Tests of Normality

| Group              | Kolmogorov-Smirnov <sup>a</sup> |    |      | Shapiro-Wilk |    |      |
|--------------------|---------------------------------|----|------|--------------|----|------|
|                    | Statistic                       | df | Sig. | Statistic    | df | Sig. |
| P0=Control Group   | .231                            | 8  | .200 | .922         | 8  | .446 |
| P1=Treatment Group | .280                            | 8  | .064 | .853         | 8  | .102 |

\*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

## Kruskal-Wallis Test

| Group                              | N  | Mean Rank |
|------------------------------------|----|-----------|
| P0=Control Group                   | 8  | 10.19     |
| Spermatogenesis P1=Treatment Group | 8  | 6.81      |
| Total                              | 16 |           |

|             | Spermatogenesis |
|-------------|-----------------|
| Chi-Square  | 2.049           |
| df          | 1               |
| Asymp. Sig. | .152            |

a. Kruskal Wallis Test  
b. Grouping Variable: Group

### Appendix 5. Results of Data Analysis of Sertoli Cells Number

#### Case Processing Summary

| Group              | Cases |         |         |         |       |         |
|--------------------|-------|---------|---------|---------|-------|---------|
|                    | Valid |         | Missing |         | Total |         |
|                    | N     | Percent | N       | Percent | N     | Percent |
| P0=Control Group   | 8     | 100.0%  | 0       | 0.0%    | 8     | 100.0%  |
| P1=Treatment Group | 8     | 100.0%  | 0       | 0.0%    | 8     | 100.0%  |

#### Descriptives

|                    |                                  | Statistic   | Std. Error |  |
|--------------------|----------------------------------|-------------|------------|--|
| P0=Control Group   | Mean                             | 29.1063     | .49294     |  |
|                    | 95% Confidence Interval for Mean | Lower Bound | 27.9406    |  |
|                    |                                  | Upper Bound | 30.2719    |  |
|                    | 5% Trimmed Mean                  | 29.1319     |            |  |
|                    | Median                           | 29.3500     |            |  |
|                    | Variance                         | 1.944       |            |  |
|                    | Std. Deviation                   | 1.39423     |            |  |
|                    | Minimum                          | 26.75       |            |  |
|                    | Maximum                          | 31.00       |            |  |
|                    | Range                            | 4.25        |            |  |
|                    | Interquartile Range              | 2.25        |            |  |
|                    | Skewness                         | -.378       | .752       |  |
|                    | Kurtosis                         | -.380       | 1.481      |  |
|                    | Mean                             | 31.1938     | .32616     |  |
| P1=Treatment Group | 95% Confidence Interval for Mean | Lower Bound | 30.4225    |  |
|                    |                                  | Upper Bound | 31.9650    |  |
|                    | 5% Trimmed Mean                  | 31.2181     |            |  |
|                    | Median                           | 31.1000     |            |  |
|                    | Variance                         | .851        |            |  |
|                    | Std. Deviation                   | .92251      |            |  |
|                    | Minimum                          | 29.50       |            |  |
|                    | Maximum                          | 32.45       |            |  |
|                    | Range                            | 2.95        |            |  |
|                    | Interquartile Range              | 1.23        |            |  |
|                    | Skewness                         | -.470       | .752       |  |
|                    | Kurtosis                         | .690        | 1.481      |  |



**Tests of Normality**

| Group              | Kolmogorov-Smirnov <sup>a</sup> |    |      | Shapiro-Wilk |    |      |
|--------------------|---------------------------------|----|------|--------------|----|------|
|                    | Statistic                       | df | Sig. | Statistic    | df | Sig. |
| P0=Control Group   | .138                            | 8  | .200 | .975         | 8  | .935 |
| P1=Treatment Group | .210                            | 8  | .200 | .945         | 8  | .663 |

\*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

**Test of Homogeneity of Variances**

Sertoli\_cells

| Levene Statistic | df1 | df2 | Sig. |
|------------------|-----|-----|------|
| 1.379            | 1   | 14  | .260 |

**Oneway****ANOVA**

Sertoli\_cells

|                | Sum of Squares | df | Mean Square | F      | Sig. |
|----------------|----------------|----|-------------|--------|------|
| Between Groups | 17.431         | 1  | 17.431      | 12.473 | .003 |
| Within Groups  | 19.564         | 14 | 1.397       |        |      |
| Total          | 36.995         | 15 |             |        |      |


### Appendix 6. Documentation of Research



Rabbit were placed in each cage



L-Arginine 500®



Combination of anesthetic drugs



The process of collecting testis



Testes stored in 10% formalin solution