


Indonesian Journal of Tropical and Infectious Disease



Medical Audit of the Management of Patients with Sepsis in The Intermediate Care Unit of Department Internal Medicine School of Medicine Airlangga University/Dr. Soetomo Hospital

Comparison of Neutralizing Antibody Response to Type 1 Polio Virus on Healthy Infants Receiving either Oral Monovalen Polio Vaccine Type 1 or Oral Trivalen Polio Vaccine Given with Basic DTP/Hb Immunization

Cognitive Behavior Change for the Improvement of Health Care, Cognitive Function and School Achievement in Helminth Infected Children

3'UTR Polymorphism of NRAMPI Gene and Susceptibility to Lung Tuberculosis among Patients and Nurses in Surabaya, Indonesia

Risk Factors of Neonatal Sepsis: A Preliminary Study in Dr. Soetomo Hospital

Histoid Leprosy

Correlation between soluble urokinase plasminogen activator receptor with CD4 T lymphocyte and WHO clinical staging of HIV infection

Intestinal Parasites Infection in AIDS Patients with Chronic Diarrhea at Dr. Soetomo General Hospital Surabaya

TTC Repeats Variation of Mycobacterium leprae Isolates for Analysis of Leprosy Transmission in Leprosy Endemic Area in East Java, INDONESIA

The Role Activity of Complement, TNF α & IL12 in Pathogenesis Dengue Virus

New Biotype of *Vibrio cholerae* O1 from Clinical Isolates in Surabaya

Indonesian Journal of Tropical and Infectious Disease

EDITORIAL BOARD OF INDONESIAN JOURNAL OF TROPICAL AND INFECTIOUS DISEASE

Editor in Chief

Nasronuddin, MD., Ph.D. (Indonesia)

Co Editor

Prof. Henri A. Verbrugh, M.D., Ph.D. (The Netherlands)

Prof. Hak. Hotta, M.D., Ph.D. (Japan)

Prof. Ismoedianto, MD, Ph.D. (Indonesia)

Maria Inge Lusida, MD., Ph.D (Indonesia)

Editorial Board

Prof. Yoshitake Hayashi (Japan)

Prof. Fumihiko Kawamoto, Ph.D. (Japan)

Prof. Hartmut Kuehn, M.D (Germany)

Prof. Dr.Med. Puruhito (Indonesia)

Prof. Soemarno, Ph.D. (Indonesia)

Prof. Guntur Hermawan, Ph.D. (Indonesia)

Prof. Abdul Rahman Omar, PhD (Malaysia)

Prof. Shinzo Izumi (Japan)

Prof. Retno Handayani, MD., Ph.D. (Indonesia)

Prof. Fedik A. Rantam, DVM, Ph.D. (Indonesia)

Prof. Soetjipto, MD., Ph.D. (Indonesia)

Prof. Soegeng Soegianto, MD., Ph.D. (Indonesia)

Prof. Indropo Agusni, MD., Ph.D. (Indonesia)

Prof. Kuntaman, MD., Ph.D. (Indonesia)

Prof. Ni Made Mertaniasih, MD., Ph.D. (Indonesia)

Prof. Yoes P. Dachlan, MD., M.Sc., Ph.D. (Indonesia)

Prof. Suharto, MD., MSc, Ph.D. (Indonesia)

Prof. Dr.Med. Soekry Erfan Kusuma (Indonesia)

Prof. Kuntoro, MD., Ph.D. (Indonesia)

Prof. Soehartono Taat Putra, MD., Ph.D. (Indonesia)

Ni Nyoman Tri Puspaningsih, Ph.D. (Indonesia)

Prof. Bambang Prajogo, Ph.D. (Indonesia)

Eddy Bagoes Wasito, MD., Ph.D. (Indonesia)

Chairul A. Nidom, DVM., Ph.D. (Indonesia)

Editorial Assistant

Prihartini Widiyanti, DDM, Ph.D.

Garry Cores de Vries, DVM, M.Sc., Ph.D

Retno Pudji Rahayu, DDM, Ph.D.

E. Bimo Aksono, DVM., Ph.D

Rahayu Anggraini, Ph.D.

Indah S. Tantular, MD., Ph.D.

Agung Sosiawan, DDM, Ph.D.

Dadik Rahardjo, DVM

Vitanata Arfijanto, MD.

Secretariat

Pudjiono, Drs., M.Si.

Titi Savitri, S.Pd.

Kris Cahyo Mulyatno, S.KM.

Ismulawati Kusumaningrum, Dra.

Wartono, S.H.

Levi Prahastuti, S.S.

Zakaria Pamungkas

Secretariat Office

Publishing Unit of Tropical and Infectious Disease, Institute of Tropical Disease Airlangga University
Kampus C, Jalan Mulyorejo Surabaya 60115, Jawa Timur - Indonesia. Phone 62-31-5992445-46 Faximile 62-31-5992445
E-mail: itd@unair.ac.id. Homepage: www.itd.unair.ac.id

Indonesian Journal of Tropical and Infectious Disease

CONTENTS

	<i>Page</i>
1. Medical Audit of the Management of Patients with Sepsis in the Intermediate Care Unit of Department Internal Medicine School of Medicine Airlangga University/Dr. Soetomo Hospital Usman Hadi and Erwin Astha Triyono	1-4
2. Comparison of Neutralizing Antibody Response to Type 1 Polio Virus on Healthy Infants Receiving either Oral Monovalent Polio Vaccine Type 1 or Oral Trivalent Polio Vaccine Given with Basic DTP/Hb Immunization Edim Hartati, Ismoedijanto, and Soegeng Soegijanto	5-9
3. Cognitive Behavior Change for the Improvement of Health Care, Cognitive Function and School Achievement in <i>Helminth</i> Infected Children Hastaning Sakti	10-16
4. 3'UTR Polymorphism of <i>NRAMP1</i> Gene and Susceptibility to Lung Tuberculosis among Patients and Nurses in Surabaya, Indonesia Rahayu Anggraini	17-22
5. Risk Factors of Neonatal Sepsis: A Preliminary Study in Dr. Soetomo Hospital Martono Tri Utomo	23-26
6. Histoid Leprosy Umi Rinasari, Sawitri, M. Yulianto Listiawan, Cita Rosita S Prakoeswa, Indropo Agusni, Rachmat Santoso, and Shinzo Izumi	27-31
7. Correlation between soluble urokinase plasminogen activator receptor with CD4 T lymphocyte and WHO clinical staging of HIV infection Shinta Oktya Wardhani, Niniek Burhan, Gatoet Ismanoe, and Tri Yudani MR	32-35
8. Intestinal Parasites Infection in AIDS Patients with Chronic Diarrhea at Dr. Soetomo General Hospital Surabaya R. Heru Prasetyo	36-37 ✓
9. TTC Repeats Variation of <i>Mycobacterium leprae</i> Isolates for Analysis of Leprosy Transmission in Leprosy Endemic Area in East Java, INDONESIA Dinar Adriaty, Ratna Wahyuni, Iswahyudi, Indropo Agusni, and Shinzo Izumi	38-43
10. The Role Activity of Complement, TNF α & IL12 in Pathogenesis Dengue Virus Soegeng Soegijanto, Dian Dwi Sary, Budi Setiawan, and Atsushi Yamanaka	44-47
11. New Biotype of <i>Vibrio cholerae</i> O1 from Clinical Isolates in Surabaya Garry Cores de Vries, Emy Koestanti Sabdoningrum, and Dadik Rahardjo	48-53

Research Report

Intestinal Parasites Infection in AIDS Patients with Chronic Diarrhea at Dr. Soetomo General Hospital Surabaya

R. Heru Prasetyo

Departement of Parasitology, School of Medicine, Airlangga University

ABSTRACT

One of the mayor health problems among HIV-infected patients is opportunistic infection due to the defect of immunity. A cross sectional study was showed current the prevalence of intestinal parasites infection in AIDS patients with chronic diarrhea were being treated at Dr. Soetomo General Hospital Surabaya during the period June 2007 to May 2008. Fecal samples were collected from 122 AIDS patients with chronic diarrhea. Samples were examined for intestinal parasites using modified acid fast stain. Ninety-seven (79.5%) of the 122 were found to have intestinal parasitic infections, *Entamoeba histolytica* (61.5%; 75/122), *Cryptosporidium* (52.5%; 64/122), *Giardia lamblia* (5.73%; 7/122), and *Trichuris trichiura* (0,82%; 1/122). This study shows that parasitic infections are important clinical complications in AIDS patients in Surabaya Indonesia.

Keywords: HIV-infected patients, AIDS, chronic diarrhea, intestinal parasites

INTRODUCTION

Human immunodeficiency virus (HIV) infection is an extremely serious worldwide problem to day. As of 2006, about 169–216 thousands Indonesian were estimated to be infected with HIV by Departement of Health, Republic of Indonesia. East Java holds the rank of provinces with high prevalence of HIV/AIDS cases after Papua and DKI Jakarta, followed by West Java, Bali and Riau. Total number of HIV/AIDS patients in Surabaya is the highest among other 37 districts in East Java, comprising 50% of total HIV/AIDS patients in East Java.¹

Since the beginning of HIV epidemic in the world, opportunistic infection have been recoqnized as an important clinical complication. In particular, opportunistic intestinal parasites plays a mayor role as etiologic agent of chronic diarrhea in HIV patient.² Chronic diarrhea in such patients can profoundly compromise the absorptive function of the small intestine, and cause significant mortality. Several species of protozoan parasites, *Cryptosporidium parvum*, *Isoospora belli*, *Cyclospora cayetanensis*, *Entamoeba histolytica*, and *Giardia lamblia* are the most common opportunistic infection associated with acquired immunodeficiency syndrome (AIDS) in developing countries.³

Diarrhea is a common complication of HIV patients, it occurs in almost 90% of AIDS patients in developing countries.⁴ However, no studies have focused specifically on opportunistic intestinal parasitic infections among HIV infected patients in the Republic of Indonesia. Thus, the present study was undertaken to determine the prevalence of intestinal parasitic infections among AIDS patients in Surabaya.

MATERIALS AND METHODS

Patients

AIDS patients with chronic diarrhea who were treated the Dr. Soetomo General Hospital Surabaya, during the periode from June 2007 to May 2008, were enrolled in this study. Fecal samples were obtained from 122 patients. Chronic diarrhea was defined as three or more loose stool passed or watery bowel movements daily, for more than two weeks.⁵

Examination of parasitic infections

Fecal samples were immediately fixed in 10% formalin, and processed using sedimentation technique. The sediment examined as iodine wet mounts, and observed under a light microscope to detect ova, larva and cyst. In addition,

air dried smears from fresh fecal samples were stained by modified acid fast (Ziehl-Neelsen) procedure to detect *Cryptosporidium*, *Isospora*, and other coccidian protozoa under a light microscope ($\times 600$).^{5,6}

RESULT

Ninety seven of 122 fecal samples (79.5%) were found to have intestinal parasitic infection. Opportunistic protozoans predominant, *Entamoeba histolytica* (61.5%; 75/122), *Cryptosporidium parvum* (52.5%; 64/122), *Giardia lamblia* (5.73%; 7/122), and *Trichuris trichiura* (0.8%; 1/122).

DISCUSSION

HIV is retrovirus infecting CD4-positive cells causing profound immunosuppression, and clinically manifest as AIDS. The cells principally infected by HIV are T4 lymphocytes (helper) and macrophages. The eventual loss of helper cell function is the main reason for immunodeficiency in the individual susceptible to opportunistic infections.⁷

Virtually every organ system in the body can be affected clinically during the course of HIV infection. The gastrointestinal tract is a major target, and the physiological sequelae are an important cause of morbidity and mortality. Two main mechanisms have been postulated. The first is reduced intestinal immunity resulting in chronic opportunistic infections, which themselves cause altered intestinal function. The second is that HIV itself affects the intestinal mucosa, causing malfunction.⁷

The clinical stage of HIV infection at which intestinal mucosal immunity fails is by definition when opportunistic infection occur (that is, clinical progression to stage IV disease), namely AIDS. Diarrhea is a significant problem in HIV infected patients in developing countries.⁷ It occurs in almost 90% of AIDS patients in developing countries.⁴

In this cross sectional study of AIDS patients, the overall prevalence of intestinal parasites in fecal samples was 79.5%. This prevalence is higher than those previously reported in other studies conducted in Africa. Various intestinal parasites were found in the fecals of the patients. Pathogenic helminth was found in one patient or 0.8% of the patients, a prevalence lower to that reported in other studies of HIV-infected patients. Seventy five (61.5%) patients had *Entamoeba histolytica*, a prevalence that was higher than the prevalence rate of reported in patients with diarrhea. The finding of this study shows that prevalence of *Cryptosporidium parvum* pathogenic and opportunistic intestinal parasite is high among Dr. Soetomo

General Hospital AIDS patients with chronic diarrhea. The prevalence is higher than those previously reported in other studies conducted in Africa,^{5,8} India,^{4,9} and in United States and Western Europe.¹⁰ In patients with AIDS cryptosporidiosis, commonly, a permanent diarrhea illness that leads to chronic malabsorption of fluids, nutrients, vitamins, and electrolytes-with resulting wasting. In patients who fail to respond to therapy, death usually occurs in three to six months.¹⁰ For this reason, cryptosporidiosis is a secondary infection should be suspected in any AIDS patients with chronic diarrhea.

The result of the present study demonstrate that opportunistic protozoan infections are prevalent among Indonesian AIDS patients and they represent an important source of clinical complications among these patients with chronic diarrhea, and the results of our study should therefore prompt physicians caring for HIV infected patients to request fecal examination and specific test for *Cryptosporidium*.

ACKNOWLEDGMENT

We are thankful to Dr. Nasronudin, dr., Sp.PD., K-PTI, Dr. Soetomo General Hospital of Internal Medicine for providing fecal specimens from AIDS patients.

REFERENCES

1. Information and Communication Bureau, East Java, 2005. Jatim Urutan ketiga Prevalensi Tinggi HIV/AIDS, March 3.
2. Wiwanitkit V, 2001. Intestinal Parasitic Infections in Thai HIV-infected Patients with Different Immunity Status, BMC Gastroenterology. Downloaded from www.biomedcentral.com. by on November 8, 2007.
3. Guk SM, Seo M, Park YK, Oh MD, Choe KW, Kim JL, et al., 2005. Parasitic infection in HIV-infected patients who visited Seoul National University Hospital during the period 1995-2003, The Korean Journal of Parasitology, Vol. 43, No. 1, 1-5, March.
4. Sathesh KS, Ananthan S, Lakshmi P., 2002. Intestinal parasitic infection in HIV infected patients with diarrhea in Chennai, Indian Journal of Medical Microbiology, Volume 20, Issue 2, 88-91.
5. Awole M, Selassie SG, Kassa T, Kibru G., 2003. Prevalence of Intestinal Parasites in HIV-infected adult Patients in Southwestern Ethiopia, Ethiop. J. Health Dev; 17(1): 71-78.
6. WHO, 2003. Manual of Basic Techniques for Health Laboratory, Geneva, 123-124.
7. Griffin GE. Human immunodeficiency virus infection and the intestine. Downloaded from www.sciencedirect.com. by on January 12, 2008.
8. Sarfiati C, Bourgeois A, Menotti J, Liegeois F, Somo RM, Delaporte E, et al., 2006. Prevalence of Intestinal Parasites including Microsporidia in Human Immunodeficiency Virus-infected Adult in Cameroon: A cross-sectional Study.
9. Mohandas K, Sehgal R, Sud A, Malla N, 2002. Prevalence of Intestinal Parasitic Pathogens in HIV-Seropositive Individual in Northern India, Jpn. J. Infect. Dis., 2002; 55, 83-84.
10. Gilson I, and Buggy BP., 1996. HIV News line, Oktober.