

**INTERNATIONAL JOURNAL
OF PHARMACY AND
PHARMACEUTICAL SCIENCES**



Editorial Board

Editor-in-Chief

Prof. M. S. Bhatia, India
(Bharati Vidyapeeth College of Pharmacy, Kolhapur, India)
Email: manish.bhatia@bharatividyaapeeth.edu
Email: editor@ijppsjournal.com

Associate Editors

Dr. Avijeet Jain, India
(Shri Sathya Sai Institute of Pharmaceutical Sciences, Bhopal, India)
Email: avijeet_9826275340@rediffmail.com

Diah Ayu Maharani
Department of Preventive and Public Health, Dentistry Faculty of Dentistry, University of Indonesia
Email: raniabdillah@gmail.com

Dr. Subhash C Mandal, India
(Department of Pharmaceutical Technology, Jadavpur University, Kolkata, India)
Email: subhashmandal@yahoo.com

Dr. Lokesh Deb, India
(Medicinal Plants and Horticultural Resources Division, Institute of Bioresources and Sustainable Development (IBSD), Department of Biotechnology, Government of India, Takyelpat Institutional Area, Imphal, Manipur, India)
Email: lokesh_deb@rediffmail.com

Dr. Wong Tin Wui, Malaysia
(Non-Destructive Biomedical and Pharmaceutical Research Center, University of Technology MARA, Malasiya)
Email: wongtinwui@yahoo.com

Assistant Editors

Dr. Idress Hamad Attittala, Libya
(Omar El-Mukhtar University, Faculty of Science, Botany Department, El-Beida, Libya)
Email: idressattitalla2004@yahoo.com

Dr. Alok Nahata, Malaysia
(Alor Star, Malaysia)
Email: aloknahata@gmail.com

Executive Editor

Mr. Niranjan Pathak, India

(Dept. of Pharmaceutical Sciences, Dr H S Gour Central University, Saugor, India)

Email: niranjanpathaklib@gmail.com

Editorial Board Members

Dr. Furhan Iqbal
Bahauddin Zakariya University Multan, Pakistan

Dr. Ebtessam Ahmed Mohammed Essa
Department of Pharmaceutical Technology,
Faculty of Pharmacy, Tanta University, Tanta,
Egypt

Dr. Syed Muhammad Farid Hasan
Faculty of Pharmacy, University of Karachi,
Karachi, Pakistan

Dr. Abdel Raheim Mohammed Ahmed Donia
College of Pharmacy – Salman Bin Abdul Aziz
University, Egypt

Dr. Pranay Jain
Institute of Engineering & Technology,
Kurukshetra University, Kurukshetra, Haryana,
India

Dr. Dr. Niaz Ali
Institute of Basic Medical Sciences, Khyber
Medical University, Peshawar, Pakistan

Dr. M. Saeed Arayne
Chairman, Department of Chemistry, University of
Karachi, Pakistan

Dr. Wanzala Wycliffe
School of Pure and Applied Sciences, South
Eastern Kenya University, Kenya

Dr. Mayuree Tangkiatkumjai
Faculty of Pharmacy, Srinakharinwirot University,
Ongkharak, Nakhonayok, Thailand

Dr. Anup Naha
Dept. of Pharmaceutic, MCOPS, Manipal,
Karnataka, India

Dr. Jagdish Labhubhai Kakadiya
Indubhai Patel College of Pharmacy and Research
Centre, Petlad-Khambhat Road, Dharmaj, Anand,
Gujarat, India

Dr. Maha Ali Eissa Ahmed
Department of Pharmacology, Faculty of
Pharmacy, MISR University for Science and
Technology (MUST), Giza Governorate, Egypt.

Dr. Rajesh Mukthavaram
Health Science Drive, University of California, San

Dr. Javed Intekhab
G. F. College (Rohilkhand University),
Shahjahanpur, U.P., India

Dr. Manish P. Patel
Dept. of Pharmaceutics and Pharmaceutical
Technology, Nootan Pharmacy College, Visnagar,
Gujarat, India

Dr. Narendra Babu Shivanagere Nagojappa
J.N. Medical College, KLE University, Belgaum,
Karnataka, India

Dr. C. Chellaram
Vel Tech Multi Tech Engg. College, Chennai, India

Dr. Mehdi Shafiee Ardestani
Department of Medicinal Chemistry and
Radiopharmacy, Tehran University of Medical
Sciences, Tehran, Iran

Dr. Amal Amin Mohamed
Plant Biochemistry Department; Agriculture
Division -National Research Center, Dokki, Cairo,
Egypt

Dr. Rabab Kamel Mahmoud
National Research Center, Cairo, Egypt

Dr. Syed Adnan Ali Shah
Universiti Teknologi MARA (UiTM), Puncak Alam
Campus, Bandar Puncak Alam, Selangor D. E.,
Malaysia

Dr. Sat Pal Singh Bisht
Roland Institute of Pharmaceutical Sciences
Berhampur, Orissa, India.

Dr. Shahu Ingole
Smt. Kashibai Navale Medical College & Hospital,
Pune India

Dr. Gina Samy El-Feky
Pharmaceutics Department, Faculty of Pharmacy,
Modern Science and Arts University, Egypt

Dr. Abdalla Ahmed Elbashir Ahmed
Khartoum University, Sudan

Dr. Yesudass Dominic Ravichandran
School of Advanced Sciences, VIT University,
Vellore, Tamil Nadu, India

Diego, Lajolla, CA, California, USA

Dr. Saifullah Khan
International Islamic University, Malaysia

Norhaniza Aminudin
Institute of Biological Sciences, Faculty of Science,
University of Malaya, Kuala Lumpur, Malaysia.

Dr. Shazia Jamshed
Kulliyyah of Pharmacy, International Islamic
University Malaysia (IIUM), Kuantan, Pahang,
Malaysia

Dr. Mayuree Tangkiatkumjai
Department of Clinical Pharmacy and Social
Pharmacy Faculty of Pharmacy, Srinakharinwirot
University, Nakhonnayok, Thailand

Dr. İsmail Murat Palabiyik
Faculty of Pharmacy, University of Ankara,
Tandoğan, Ankara, Turkey

Nadeem A. Kizilbash
Faculty of Medicine, Northern Border University
Arar, Saudi Arabia

Dr. Debajit Kalita
Department of Botany, Morigaon College, Assam,
India

Dr. Seyed Mohammad
Department of Biology, Faculty of Basic Sciences,
University of Mazandaran, Babolsar, Iran

Dr. Shalini Sivadasan
Faculty of pharmacy, AIMST University, Semeling,
Kedah, Malaysia

Dr. Sujimon Tanvichien
Srinakharinwirot University Nakornayok Rd. A.
Ongkarak T.Ongkarak, Nakornayok, Thailand

Prof. Dr.-Ing. habil. Dr. h. c. Lothar Mörl
Institute für Apparate- und Umwelttechnik Otto-
von-Guericke-Universität Magdeburg, Germany

Dr. V. Ravichandran
Faculty of Pharmacy, AIMST University, Semeling,
Kedah, Malaysia

Dr. Zahid Hussain
Department of Pharmaceutics, Faculty of
Pharmacy, Universiti Teknologi MARA, Puncak
Alam Campus, Malaysia

Dr. Ajay Kumar Meena
Department of AYUSH, Ministry of Health &
Family Welfare, Government of India, India

Dr. Ashish C. Suthar
Herbal R & D, Piramal Life Sciences Ltd., Mumbai,
India

Dr. Seema Akbar
Research Institute of Unani Medicine, The
University of Kashmir Campus, Srinagar, J. & K.,
India

Dr. P. Thillai Arasu
Department of Chemistry, Wollega University,
Nekemta, Ethiopia

Dr. Sooraj S. Nath
Safi Institute of Advanced Study, Kozhikode, India

Dr. Erum Shireen
Dept. of Biochemistry, University of Karachi,
Pakistan

Dr. M. M. Gupta
School of Pharmacy, Faculty of Medical Sciences,
The University of The West India, India

Dr. Najma Sultana
United Biotechnologies, Karachi, Pakistan

Dr. Sivakumar P
Department of Petroleum Engineering, JCT
College of Engineering and Technology, Pichanur,
Tamil Nadu, India

Dr. Evren Algin Yapar
Department in Ministry of Health, Turkish
Medicines and Medical Devices Agency, Ankara,
Turkey

Dr. Vishal Vijay Pandey
Jayawantrao Sawant College of Pharmacy &
Research, Hadapsar, Pune, India

Dr. Shamkuwar Prashant Babarao
Government College of Pharmacy, Thiba Palace,
Ratnagiri, India

Dr. S. K. Starling
Department of Chemistry, Mewar University
Chittorgarh, Rajasthan, India

Dr. Syed Sajjad Hussien
Manipal College of Pharmaceutical Sciences,
Manipal University, India

Dr. Ahmed Osman
Department of Psychology, Faculty of Education,
Assiut University, Malaysia

Dr. Abdel-Tawab Halim Mossa Abd El-Aziz
Department of Pesticide Chemistry, National
Research Centre (NRC) Dokki, Cairo, Egypt

Dr. V. Vinod Prabhu
Department of Biochemistry, University of
Madras, Guindy campus, Tamil Nadu, India

Dr. Ching Siew Mooi
University Putra, Malaysia

Dr. Manish A. Rachchh
Pharmacological Research and IPR University
road, Rajkot, Gujarat, India

Kiran Kumar Chereddy
Manager at Novartis Pharma AG
Postfach 4002, Basel
Switzerland

Dr. Fahd M. Abd Al Galil
Department of Zoology, Faculty of Applied
Science, Thamar University, Yemen

Dr. Priyanka Bhatt
Department of Pharmaceutical Sciences, College
of Pharmacy, University of South Florida, USA

Dr. Beril Anilanmert
Istanbul University-Cerrahpasa Institute of
Forensic Sciences, Cerrahpasa/Istanbul

Dr. Andleeb Shahzadi
Istanbul University Cerrahpasa-Fatih Istanbul,
Turkey 34098

Dr. Mohd Masih Uzzaman Khan
Unaizah College of Pharmacy Unayzah 56264,
Saudi Arabia

Dr. Asif Husain
Jamia Hamdard University, New Delhi, India

Dr. Muhammad Shahzad Aslam
School of Bioprocess Engineering, Universiti
Malaysia Perlis, Kompleks Pusat Pengajian
Jejawi, Arau, Perlis

Dr. Gláucio Diré Feliciano
State University Center Foundation of the West
Zone / Rio de Janeiro, Brazil

Dr Dilipkumar Pal
Department of Pharmaceutical Sciences, Guru
Ghasidash Vishwavidyalaya, Bilaspur, C.G., 495
009, India

Ali Abdullah Ali Al-yahawi
Assistant Professor of Clinical Pharmacy &
Therapeutics, Yemen

Dr. Maha Zaki Rizk
National Research Center, Therapeutic Chemistry
Dept., National Research Center, Dokki, Cairo,
Egypt



Online ISSN: 0975-1491

Print ISSN: 2656-0097



Journal Metrics 2018

Source Normalized Impact per Paper (SNIP): 2.029

SCImago Journal Rank (SJR): 0.23

ISSN: 0975-1491

**International Journal of
Pharmacy and...**



Visitor No. 107894

[Our Journals](#) || [Open Access Policy](#) || [Publication & Peer Review Policy](#) || [Publication Ethics](#)

The publication is licensed under a [Creative Commons License \(CC BY\)](#). [View Legal Code](#)

Copyright © 2018 All Rights Reserved, **Innovare Academic Sciences** | Powered By **CyberDairy**

[HOME](#) / [ARCHIVES](#) / Vol 7, Issue 12, 2015REVIEW ARTICLE(S)**BRIEF REVIEW ON THE MEDICINAL USES AND ANTIMICROBIAL ACTIVITY OF DIFFERENT PARTS OF SCHINUS TEREBINTHIFOLIUS RADDI**MÃ“NICA REGINA GUIMARÃ“ES DOS SANTOS, JESSICA HELLEN SOUZA DA SILVA, MARINA LETÃ“CIA DO CARMO CAXITO Pages 1-7[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)**CHITOSAN HYDROGELS FOR DRUG DELIVERY AND TISSUE ENGINEERING APPLICATIONS**EDUARDO PEREIRA DE AZEVEDO Pages 8-14[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)ORIGINAL ARTICLE(S)**THEOPHYLLINEÃ“NAÃ“ SACHARRINE SINGLE CRYSTAL ISOLATION FOR ITS STRUCTURE DETERMINATION**ILMA NUGRAHANI, EVALIANTY AGIANA PERTIWI, OKKY DWICHANDRA PUTRA Pages 15-24[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)**ISOLATION AND CHARACTERIZATION OF STIGMASTEROL FROM CHLOROFORM FRACTION OF AERIAL PART OF ARGEMONE MEXICANA L**PRAVEEN S. NAYAK, D. M. KAR, SHWETA P. NAYAK Pages 25-29[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)**ASSAY METHOD DEVELOPMENT AND VALIDATION FOR SIMULTANEOUS QUANTITATIVE ESTIMATION OF DILOXANIDE FUROATE AND TINIDAZOLE IN TABLETS BY RP-HPLC**

PARUPUGALLA SWETHA, D. VIJAY KUMAR, RAKESH B, ASHOK KUMAR AKUTHOTA

Pages 30-35

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

PREPARATION AND EVALUATION OF ANTHRALIN BIODEGRADABLE NANOPARTICLES AS A POTENTIAL DELIVERY SYSTEM FOR THE TREATMENT OF PSORIASIS

GHAREB M. SOLIMAN, SHAABAN K. OSMAN, AHMED M. HAMDAN

Pages 36-40

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

TOXICOLOGICAL AND PHARMACOLOGICAL ASSESSMENT OF GOLD NANORODS IN NORMAL RATS

MOSTAFA A. EL-SAYED, NEMAT A. YASSIN, ALI A. SHABAKA, OSAMA A. EL-SHABRAWY, SAWSAN S. MAHMOUD, SIHAM M. EL-SHENAWY, MARWA A. EL-SHAER, NABILA S. HASSAN, EMAD A. AL-ASHKAR, WAEL H. EISA, REHAB F. ABDEL-RAHMAN

Pages 41-50

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

A STUDY TO PREDICT ANTI-INFLAMMATORY ACTIVITY OF EUGENOL, MYRISTICIN, AND LIMONENE OF CINNAMOMUM SINTOC

SRI ADI SUMIWI, OKTAVIA SARMA SIHOMBING, ANAS SUBARNAS, MARLINE ABDASSAH, JUTTI LEVITA

Pages 51-54

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

PHYTOPHENOLICS COMPOSITION, HYPOLIPIDEMIC, HYPOGLYCEMIC AND ANTIOXIDATIVE EFFECTS OF THE LEAVES OF FORTUNELLA JAPONICA (THUNB.) SWINGLE

NAGAT S. ABU-GABAL, HOWAIDA I. ABD-ALLA, NAIMA Z. MOHAMED, HANAN F. ALY, NAGWA M. M. SHALABY

Pages 55-63

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

2,4-CYCLOADDITION REACTIONS: PREPARATION AND CYTOTOXICITY OF NOVEL QUINOLINE AND PYRROLO [3,4-F] QUINOLINE DERIVATIVES

AHMED S. I. MOHAMEDE, MOHAMED A. A. ELNEAIRY, SANAA M ELDINE

Pages 64-68

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

A STUDY OF CHLOROPHYLLIN OF MEDICINAL PLANTS, ITS CHEMICAL CHARACTERIZATION AND ANTI-PROLIFERATIVE ACTIVITY WITH SPECIAL REFERENCE TO SOLANUM TRILOBATUM L. ON LIVER CELL LINE

N. BANU, PAVITHRA S.

Pages 69-79

[View PDF](#)

[Abstract](#)

[Download PDF](#)

[View HTML](#)

OPTIMIZATION OF PANTOPRAZOLE ENTERIC PELLETS COATING PROCESS BY QBD: EFFECT OF COATING PROCESS VARIABLES ON THE INTERMEDIATE QUALITY OF THE PRODUCT AND SCALE UP

GIRISH SONAR, SWATI RAWAT

Pages 80-87

[View PDF](#)

[Abstract](#)

[Download PDF](#)

[View HTML](#)

EXPRESSION AND ENZYMIC PROPERTIES OF A UNIQUE RECOMBINANT ANTICOAGULANT AND FIBRINOLYTIC ENZYME FROM ACINETOBACTER BAUMANNII TU04

RENUKA KRISHNAN, CHUN SHIONG CHONG, KIAN MAU GOH, FIRDAUS ABDUL WAHAB, HARYATI JAMALUDDIN

Pages 88-94

[View PDF](#)

[Abstract](#)

[Download PDF](#)

[View HTML](#)

ISOLATION OF ENDOPHYTIC FUNGI FROM ARTEMISIA ANNUA, L AND IDENTIFICATION OF THEIR ANTIMICROBIAL COMPOUND USING BIOAUTOGRAPHY METHOD

INDAH PURWANTINI, WAHYONO ., MUSTOFA ., RATNA ASMAH

Pages 95-99

[View PDF](#)

[Abstract](#)

[Download PDF](#)

[View HTML](#)

IN VITRO ANTIOXIDANT AND ANTICANCER ACTIVITY OF MIMOSA PUDICA LINN EXTRACT AND L-MIMOSINE ON LYMPHOMA DAUDI CELLS

FELISA PARMAR, NISHA KUSHAWAHA, HYACINTH HIGHLAND, LINZ-BUOY GEORGE, LINZ-BUOY GEORGE

Pages 100-104

[View PDF](#)

[Abstract](#)

[Download PDF](#)

[View HTML](#)

SIMULTANEOUS DETERMINATION OF ARTESUNATE AND AMODIAQUINE IN HUMAN PLASMA USING LC-MS/MS AND ITS APPLICATION TO A PHARMACOKINETIC STUDY

RAMBABU MADDELA, NAGESWARA RAO PILLI, VASU BABU RAVI, VINAYENDER ADIREDDY, AJITHA MAKULA

Pages 105-112

[View PDF](#)

[Abstract](#)

[Download PDF](#)

[View HTML](#)

DOCKING AND CYTOTOXICITY STUDIES OF 2-VINYLCROMONE DERIVATIVES ON HUMAN BREAST CANCER CELL LINES

SWATI KAUSHIK, MEGHA RIKHI, SEEMA BHATNAGAR

Pages 113-117

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

COMPARATIVE STUDIES OF EFFICACY AND EFFECTS ON OXIDATIVE STRESS OF AMLODIPINE AND RAMIPRIL IN THE HYPERTENSIVE PATIENTS OF NORTH EAST INDIA

SURAJIT TAMULI, SANJEEV KAKATI, SANJIB DAS, KHUMANTHEM DEEPAK SINGH, SURAJIT KUMAR GHOSH Pages 118-121

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)PREVALENCE AND MOLECULAR CHARACTERIZATION OF EXTENDED SPECTRUM β -LACTAMASES IN KLEBSIELLA PNEUMONIAE ISOLATES FROM CANCER PATIENTS AND OTHERS

TAGHRID S. EL-MAHDY, MOHAMMED O. ABDELAZIZ, RAMADAN A. EL-DOMANY Pages 122-127

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

ANTIMICROBIAL POTENTIAL OF FUNGAL ENDOPHYTES ISOLATED FROM LEAVES OF PROSOPIS JULIFLORA (SW.) DC. AN IMPORTANT WEED

APARNA SRIVASTAVA, RAVEESHA KOTESHWAR ANANDRAO Pages 128-136

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

GENETIC SUSCEPTIBILITY OF HCV RNA IN VARIOUS ISONYM GROUPS OF PUNJAB, PAKISTAN

RUBI GHAZALA, SHILU MATHEW, ISHTIAQ QADRI Pages 137-141

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

DESIGN AND DEVELOPMENT OF NOVEL MICROEMULSION BASED TOPICAL FORMULATION OF HESPERIDIN

VAISHALI KILOR, NIDHI SAPKAL, GUNJAN VAIDYA Pages 142-148

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

MOLECULAR ASSESSMENT OF GENETIC DIVERSITY IN INDIAN ACCESSIONS OF ALOE VERA USING SSR MARKER

SANDEEP KUMAR, MANILA YADAV, AMITA YADAV, JAYA PARKASH YADAV Pages 149-154

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

STRUCTURAL AND FUNCTIONAL ANALYSIS OF AF9-MLL ONCOGENIC FUSION PROTEIN USING

HOMOLOGY MODELING AND SIMULATION BASED APPROACH

MEDHA DAVE, ADITI DAGA, RAKESH RAWAL

Pages 155-161

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

ANTIOXIDANT AND ANTI-INFLAMMATORY ACTIVITY OF THE INDONESIAN FERNS, NEPHROLEPIS FALCATA AND PYRROSIA LANCEOLATA

ISMIARNI KOMALA, AZRIFITRIA YARDI, OFA SUZANTI BETHA, FINI MULIATI, MALIYATHUN NIA[™]MAH

Pages 162-165

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

STUDY THE SUSTAIN RELEASE EFFECT OF DIFFERENT POLYMERS USED IN THE FORMULATION OF ASPIRIN-ROSUVASTATIN TABLETS

OMAR S. SALIH

Pages 166-172

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

SOLID-STATE STABILITY AND SOLUBILITY DETERMINATION OF CRYSTALLINE FORMS OF MOXIFLOXACIN HYDROCHLORIDE

TAMĀRIS A. JĀŠLIO, JERUSA S. GARCIA, RUDY BONFILIO, MAGALI B. DE. ARAĀŠJO, MARCELLO G. TREVISAN

Pages 173-177

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

THE EFFECT OF PROPHYLACTIC INHIBITION OF INDUCIBLE NITRIC OXIDE SYNTHASE BY AMINOGUANIDINEON SERUM LEVELS OF SOME ADIPOCYTOKINES IN PRISTANE-INDUCED ARTHRITISIN RATS

SHATHA H. ALI, ALI A. KASIM

Pages 178-183

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

DEVELOPMENT OF STABILITY INDICATING RP-HPLC METHOD FOR THE DETERMINATION OF RELATED SUBSTANCES IN ATORVASTATIN SOLID DOSAGE FORM AND BULK DRUGS

MANOJ KUMAR VADLAMUDI, SANGEETHA DHANARAJ, SANJEEVA YARKALA, JOHN JOSEPH JAYAPAL, PAVAN KUMAR KOMMAVARAPU

Pages 184-194

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

SYNTHESIS OF NOVEL CYCLOHEXANONE DERIVATIVES AS BCR-ABL T1351 INHIBITORS

SULAIMAN ALI MUHAMMAD, SUBASH SUGANYA, SUBBAN RAVI, SENNIAPPAN VENKATACHALAPATHI Pages 195-199

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

MOLECULAR DOCKING, DESIGN, SYNTHESIS, IN VITRO ANTIOXIDANT AND ANTI-INFLAMMATORY EVALUATIONS OF NEW ISOQUINOLINE DERIVATIVES

MANIKANDAN ALAGUMUTHU, KULATHU IYER SATHIYANARAYANAN, SIVAKUMAR ARUMUGAM Pages 200-208

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

FORMULATION AND CHARACTERIZATION OF CARVEDILOL NANOEMULSION ORAL LIQUID DOSAGE FORM

HAYDER KADHIM DRAIS, AHMED ABBAS HUSSEIN Pages 209-216

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

EVALUATION OF CHEMOPREVENTIVE RESPONSE OF PENTOXIPHYLLINE AND SILDENAFIL IN COLORECTAL CARCINOMA EXPERIMENTALLY INDUCED IN RATS: COMPARATIVE STUDY WITH 5-FLUOROURACIL

AHMED HAMEDJWIED ATAIMISH, NADAN. AL-SHAWI, MOHAMMED M. AL-QURTAS Pages 217-222

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

ESTIMATION OF ALKALOIDS AND PHENOLICS OF FIVE EDIBLE CUCURBITACEOUS PLANTS AND THEIR ANTIBACTERIAL ACTIVITY

BIKASH DEBNATH, MD. JASHIM UDDIN, PRASENJIT PATARI, MANIK DAS, DEBASISH MAITI, KUNTAL MANNA Pages 223-227

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

FEASIBILITY AND ACCEPTABILITY OF MY ELECTRONIC PERSONAL HEALTH RECORD MONITOR (MY-EPHRM)

LUA PEI LIN, UMAR IDRIS IBRAHIM Pages 228-232

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

TOTAL PHENOL, FLAVONOID AND ANTIOXIDANT PROPERTIES OF AURICULARIA AURICULA-JUDAE


PACKIALAKSHMI B., SUDHA G., CHARUMATHY M. Pages 233-237

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)


DETERMINATION OF CAPECITABINE-AN ANTICANCER DRUG IN DRIED BLOOD SPOT BY LC-ESI-MS/MS

Quality Supplements at iHerb®
iHerb


Quartiles




SJR




Citations per document




Total Cites Self-Cites



External Cites per Doc Cites per Doc



% International Collaboration



Citable documents Non-citable documents

Cited documents Uncited documents



PURAN SINGHAL, PRIYANKA A. SHAH, JAIVIK V. SHAH, PRIMAL SHARMA, PRANAV S. SHRIVASTAV Pages 238-245

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

CYTOTOXICITY OF CRUDE EXTRACT FROM SPONGE-ASSOCIATED BACTERIA AGAINST MOLT4 LEUKEMIC CELL LINES THROUGH APOPTOSIS

AI KARWATI, JUN NOMURA, NAHROWI RAMLI, ARIS TRI WAHYUDI

Pages 246-249

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

ANTIMALARIAL ACTIVITY AND CYTOTOXICITY STUDY OF ETHANOL EXTRACT AND FRACTION FROM ALECTRYON SERRATUS LEAVES

ATY WIDYAWARUYANTI, USWATUN KHASANAH, LIDYA TUMEWU, HILKATUL ILMI, ACHMAD FUAD HAFID, INDAH S TANTULAR Pages 250-253

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

MICROWAVE-ASSISTED SYNTHESIS OF AGNP USING AQUEOUS LEAVES EXTRACT OF VINCA ROSEA AND ITS THERAPEUTIC APPLICATION

SWETA PATEL, AISHAANI MAHESHWARI, AMRISH CHANDRA

Pages 254-258

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

IMMUNOMODULATORY ROLE OF HONEY AND PROPOLIS ON CARBON TETRACHLORIDE (CCL4) INJECTED RATS

KADRY A. ELBAKRY, CAMELIA A. ABDEL MALAK, MAHMOUD M. HOWAS

Pages 259-262

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

FREE RADICAL LOAD IN LYMPHOID ORGANS (SPLEEN AND THYMUS) OF INDIAN GOAT *CAPRA HIRCUS*: ROLE OF SEX, SEASON AND MELATONIN

SOMENATH GHOSH, KAUSHALENDRA ., CHANDANA HALDAR

Pages 263-269

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

EMBEDMENT TECHNIQUE: AN ALTERNATIVE TO WET GRANULATION FOR BETTER CONTROL OF RELEASE OF HIGHLY WATER SOLUBLE DRUGS—A CASE STUDY WITH DILTIAZEM HCL

GRANDHI SRIKAR, PRAMEELARANI AVULA, SRIHARSHA KOREDDI

Pages 270-276

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

DESIGN, IN-SILICO DOCKING AND PREDICTIVE ADME PROPERTIES OF NOVEL PYRAZOLINE DERIVATIVES WITH SELECTIVE HUMAN MAO INHIBITORY ACTIVITY

VISHNU NAYAK BADAVATH, BARIJ NAYAN SINHA, VENKATESAN JAYAPRAKASH

Pages 277-282

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

MOLECULAR, HISTOLOGICAL, AND ANTI-OXIDANT EVALUATION OF COLITIS INDUCTION IN RATS BY DIFFERENT CONCENTRATION OF DEXTRAN SODIUM SULFATE (5 KDA)

NOPPAWAT PENGKUMSRI, PRASIT SUWANNALERT, BHAGAVATHI SUNDARAM SIVAMARUTHI, RAWIWAN WONGPOOMCHAI, SOPHON SIRISATTHA, ACHIRAYA TAMMASAKCHAI, SIRINYA TAYA, SASITHORN SIRILUN, SARTJIN PEERAJAN, CHAIYAVAT CHAIYASUT

Pages 283-287

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

SHORT COMMUNICATION(S)

QUANTIFICATION OF ASH AND SELECTED PRIMARY METABOLITES FROM NON-EDIBLE PARTS OF SEVERAL FRUITS

URVI GUPTA, HITESH SOLANKI

Pages 288-290

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

FIRST COMPLETE GENOME SEQUENCE OF OCULAR ISOLATE CARRYING BLAVIM-2 MEDIATED MULTI-DRUG-RESISTANT(MDR) PSEUDOMONAS AERUGINOSA VRFPA04 FROM INDIA AND TWO DRAFT GENOME SEQUENCE OF OCULAR ISOLATES PSEUDOMONAS AERUGINOSA VRFPA03 AND VRFPA05 HARBORING NOVEL BLA

N. MURUGAN, J. MALATHI, V. UMASHANKAR, MADHAVAN HN

Pages 291-292

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

ESTIMATION OF BACOSIDE-A IN BACOPA MONNIERI AERIAL PARTS USING TLC DENSITOMETRY

RAJESH KUMAR, TEJPAL SINGH, DEEPAK KUMAR, MANDEEP SINGH, SANDEEP KAUR, RAJEEV GARG

Pages 293-295

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

THE EFFECTS OF COSMOS CAUDATUS AND PIPER SARMENTOSUM AQUEOUS EXTRACTS ON MALE MICE FERTILITY

DZULSUHAIMI DAUD, ADRIANA AZAHAR, SITI SOLEHAH FATINAH ZAINAL ABIDIN, ALENE TAWANG

Pages 296-298

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

GC-MS ANALYSIS OF METHANOLIC EXTRACT OF LEAVES OF RHODODENDRON CAMPANULATUM

SAKSHI PAINULI, NISHANT RAI, NAVIN KUMAR

Pages 299-303

 [View PDF](#)

 [Abstract](#)

 [Download PDF](#)

 [View HTML](#)

CASE STUDY(S)

VALPROIC ACID INDUCED TRANSAMINITIS

VRINDA NAMPOOTHIRI, R LAKSHMI

Pages 304-305

 [View PDF](#)

 [Abstract](#)

 [Download PDF](#)

 [View HTML](#)



Online ISSN: 0975-1491

Print ISSN: 2656-0097



Journal Metrics 2018

Source Normalized Impact per Paper (SNIP): 2.029

SCImago Journal Rank (SJR): 0.23

ISSN: 0975-1491

Journal of Pharmacy and
Pharmaceutical Sciences

← Show this widget in
your own website

Q2

Pharmaceutical
Science

Just copy the code below
and paste within your html
code:

SJR 2019

0.4

powered by scimagor.com



L'il Critters, Complete Multivitamin...

IDR 202k

America's #1 Gummy Vitamin Brand Roarin' Taste Dietary
Supplement Delicious Fruit...

iHerb

Metrics based on Scopus® data as of April 2020

Dr Rashmi Sharma 2 months ago

Excellent.

reply



Melanie Ortiz 1 month ago


Scimagor Team

Dear Dr. Rashmi, thanks for your participation! Best Regards, SCImago Team

Richardson Brown 2 years ago

It's an enormous pleasure and honor to organize "15th World Congress On Bioavailability and Bioequivalence" scheduled during July 29-30, 2019 at Bangkok, Thailand. The conference is mainly focused on the theme of "Essential Innovation in the field of Pharmaceutical Science for Public and Medical health". BABE Conference 2019 is designed with prudent keynote sessions, session lectures, and poster presentations, presentations from the young researchers, panel discussions, and the B2B meetings with world-renowned speakers from the stream of clinical and pharmaceutical sciences. It provides the best platform for the researchers to the researchers all over globe to introduce themselves to the world with their unique researches. It's an open forum to discuss new researches and the challenges faced during the BA/BE studies, manufacturing the

International Journal of
Pharmacy and...



SJR 2018

0.23



powered by scimagojr.com

Visitor No. 107894107894

Our Journals || Open Access Policy || Publication & Peer Review Policy || Publication Ethics

The publication is licensed under a Creative Commons License (CC BY). View Legal Code

Copyright © 2018 All Rights Reserved, **Innovare Academic Sciences** | Powered By **CyberDairy**



International Journal of

Peer Reviewed Online Journal Impact Factor 2019: 7.583

[ijsr.net](#)

[OPEN](#)

Journal of Pharmacy and Pharmaceutical Sciences

Country [Canada](#) -  SCIMAGO INSTITUTIONAL RANKINGS

Subject Area and Category [Medicine](#)
[Medicine \(miscellaneous\)](#)

[Pharmacology, Toxicology and Pharmaceutics](#)
[Pharmaceutical Science](#)
[Pharmacology](#)

Publisher [Canadian Society for Pharmaceutical Sciences](#)

Publication type [Journals](#)

ISSN [14821826](#)

Coverage [1998-2020](#)

Scope The Journal of Pharmacy and Pharmaceutical Sciences (JPPS) is the official journal of the Canadian Society for Pharmaceutical Sciences. JPPS is a broad-spectrum, peer-reviewed, international pharmaceutical journal circulated electronically via the World Wide Web. Subscription to JPPS is free of charge. Articles will appear individually as soon as they are accepted and are ready for circulation.

 [Homepage](#)

[How to publish in this journal](#)

[Contact](#)

 [Join the conversation about this journal](#)

75

H Index

Original Article

ANTIMALARIAL ACTIVITY AND CYTOTOXICITY STUDY OF ETHANOL EXTRACT AND FRACTION FROM *ALECTRYON SERRATUS* LEAVES

ATY WIDYAWARUYANTI^{1,2}, USWATUN KHASANAH⁴, LIDYA TUMEWU², HILKATUL ILMU², ACHMAD FUAD HAFID^{1,2}, INDAH S TANTULAR^{2,3}

¹Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Surabaya 60286, Indonesia, ²Center for Natural Product Medicine Research and Development, Institute of Tropical Disease, Universitas Airlangga, Surabaya 60115, Indonesia, ³Department of Parasitology, Faculty of Medicine, Universitas Airlangga, Surabaya 60286, Indonesia, ⁴Postgraduate student, Faculty of Pharmacy, Universitas Airlangga, Surabaya 60286, Indonesia
Email: aty_ww@yahoo.com

Received: 26 Feb 2015 Revised and Accepted: 02 Nov 2015

ABSTRACT

Objective: The resistance of *Plasmodium falciparum* to standard antimalarial drugs has led the scientist to investigate medicinal plants as new potentials in the treatment or prevention of malaria. Previous study showed that the ethanol extract from *Alectryon serratus* leaves inhibited *P. falciparum* *in vitro*. The purpose of this study was to determine the *in vitro* and *in vivo* antimalarial activity and toxicity of ethanol extract and fractions of *A. serratus* leaves.

Methods: Ethanol extract of *A. serratus* leaves was partitioned with dichloromethane, ethyl acetate and n-butanol successively. Antimalarial activities were determined *in vitro* against *P. falciparum* 3D7 based on HRP2 measurement in a simple enzyme-linked immunosorbent assay (ELISA) and *in vivo* against *Plasmodium berghei* ANKA based on the 4-days suppressive test by Peters. The toxicity study was determined by MTT assay using Huh7it cells.

Results: Ethanol extract and fractions of *A. serratus* exhibited antimalarial activity and was proved to be nontoxic. Ethyl acetate (EA) and butanol (BuOH) fractions performed a higher antimalarial activity (IC₅₀<10 µg/ml) and lower toxicity (SI>10) compared with ethanol extract and dichloromethane (DCM) fraction. EA fraction had IC₅₀ value of 9.45 µg/ml and SI of 10.58, while BuOH fraction had IC₅₀ value of 7.69 µg/ml and SI of 130.04. *In vivo* antimalarial activity was conducted for ethanol extract and EA fraction. The result showed that EA fraction and ethanol extract had ED₅₀ 5.92 mg/kg BW and 13.82 mg/kg BW, respectively.

Conclusion: Ethanol extract and EA fraction of *A. serratus* leaves showed *in vivo* and *in vitro* antimalarial activities and proved to be nontoxic. Both of them are a good candidate of new source in the development of new antimalarial drugs.

Keywords: *Alectryon serratus* leaves, Antimalarial activity, Toxicity.

INTRODUCTION

Malaria is one of the most important infectious diseases in the tropics and sub-tropics region. Each year 300 to 500 million new cases are diagnosed and approximately 1.5 million people, mostly children [1], died because of the disease. *Plasmodium falciparum*, the most widespread etiological agent for human malaria, has become increasingly resistant to standard antimalarial drugs e. g. chloroquine and antifolates [2]. Consequently, new drugs are urgently needed today to treat malaria. These drugs should have novel modes of action or be chemically different from the drugs in current use.

Natural products, particularly plants, might become the strategy to find new antimalarial drugs since nature is a promising source of drugs. Plants have been sources of medicine throughout the history of medicine. For thousands of years natural compounds, mostly derived from plants, have been used for traditional medicine. Plant species have demonstrated their potential to provide effective drugs for the treatment of malaria. Two of the most effective antimalarial drugs available are quinine and artemisinin, which are derived from the plant [3]. Quinine is isolated from *Chinchona* species (Rubiaceae) and artemisinin from *Artemisia annua*. Artemisinin derivatives are now recommended by the World Health Organization (WHO) worldwide in combination with other drugs, such as amodiaquine, mefloquine, lumefantrine as the first-line treatment of malaria [4].

Our preliminary study, the screening of antimalarial activity of some Indonesian plants using HRP2 method, indicated that extracts of *A. serratus* leaves had the highest activity (IC₅₀ value of 12.3µg/ml) and was potential as a target for further study [5]. *A. serratus* is included in the Sapindaceae family, which is rich in substances such as flavonoid, diterpenoid, glycoside, phenol, saponin, kaempferol, quercetin and β-

sitosterol [6]. This information encouraged us to study the plant further. In this study, antimalarial activity and toxicity effect of the extract and fractions *A. serratus* leaves were assessed.

MATERIALS AND METHODS

Plant material

Leaves of *A. serratus* were collected from Alas Purwo National Park, Banyuwangi, East Java, Indonesia. Authentication and identification of plant were carried out at the Purwodadi Botanical Garden, East Java, Indonesia.

Extraction and fractionation

1 kg of the sample was extracted using 80% ethanol by ultrasonic-assisted maceration technique for two minutes to achieve three-time replication. The ethanol extract were filtered, pooled, and dried at 40°C using rotary evaporator and weighed afterward. 100 g of ethanol extract was suspended in distilled water and partitioned with dichloromethane (DCM), ethyl acetate (EA) and n-butanol (BuOH) successively, which were then concentrated to dry in a rotary evaporator. The crude extract and fractions were kept in air tight containers and were stored at 4°C for use in phytochemical screening and antimalarial assay.

Phytochemical screening

Dried ethanol extract and fractions of DCM, EA, BuOH and water (1.0 mg) were diluted in methanol. The phytochemical screening was performed by Thin Layer Chromatography (TLC) method using stationary phase silica gel RP-18 and acetonitrile: methanol: water (2:1:4) mobile phase, as well as 10% sulphuric acid reagent. The spots were observed under UV 254 nm and 366 nm.

***P. falciparum* (3D7 strain) and *in vitro* culture**

P. falciparum 3D7 strain was obtained from Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia and was maintained in our laboratory at 5% hematocrit (human type O-positive red blood cells) in complete RPMI 1640 medium (RPMI 1640 medium supplemented with 5.96 g HEPES, 0.05 g hypoxanthine, 2.1 g NaHCO₃, 50 µg/ml gentamycin and completed with 10% human O+serum) in petri dish by modified candle jar method. Incubations were done at 37 °C. The culture was routinely monitored through Giemsa staining of the thin blood smears. For the experiment, the parasite contain predominantly ring forms. Parasite of stock cultures were further diluted with uninfected type O+human erythrocytes and culture medium to achieve a starting parasitemia of 0.05% and a hematocrit of 1.5%. This final parasite culture was immediately used for antimalarial assay [7, 8].

***In vitro* antimalarial assay**

Antimalarial activity assay used HRP2 (HRP2 Kit Cellabs Pty. Ltd., Brookvale, New South Wales, Australia). The screening assay used single concentration of extract and fractions (average concentration of 100 µg/ml). To each well of a microplate, 100 µl diluted extract solution was added into 100 µl of final parasite culture. The plates were then incubated for 72 h at 37 °C. They were subsequently frozen-thawed twice to obtain complete hemolysis and stored at -30 °C until further processing and 100 µl of each of the hemolysis culture samples was transferred to the ELISA plates. The plates were precoated with monoclonal antibodies against *P. falciparum* HRP2 and incubated at room temperature for 1 hour in humidified chamber. The plates were washed five times with the washing solution (200 µl of each well) and 100 µl of the diluted antibody conjugate were added to each well. After incubation for an additional 1 hour in humidified chamber, the plates were washed with the washing solution (200 µl of each well) and 100 µl of diluted (1:20) chromogen TMB (tetramethylbenzidine) was added to each well. The plates were then incubated for another 15 min in the dark, and 50 µl of the stop solution was added. The optical density values were read with an ELISA plate reader at an absorbance maximum of 450 nm [9]. Inhibition percentage was calculated using the following formula:

$$\frac{(\text{optical density of control well} - \text{optical density of sample well})}{\text{optical density of control well}} \times 100\%$$

Samples which were revealed with inhibition percentage >80% will be further tested to determine the Inhibitory Concentration (IC₅₀) values using serial concentration of 100 µg/ml to 0.01 µg/ml. The IC₅₀ values were determined graphically from dose-response curves (concentration versus percent inhibition curves) with non-linear analysis by SPSS probit.

Toxicity assay and analysis

Toxicity assay used MTT cell proliferation assay for several concentration 1000 µg/ml; 800 µg/ml; 400 µg/ml; 200 µg/ml; 100 µg/ml; 10 µg/ml; 1 µg/ml; 0.1 µg/ml; 0.01 µg/ml. Huh7it cells in 96 well plates were treated with serial dilution of the samples or control. Prepared wells were analyzed using ELISA reader at 720 nm and 650 nm wavelength.

Criteria of antimalarial activity *in vitro* and toxicity

Extract and fraction exhibiting IC₅₀ < 25 µg/ml was considered active. Extract showing IC₅₀ < 100 µg/ml was classified as follows: Marginally active at SI < 4, partially active at SI 4-10 and active SI > 10 [3].

***In vivo* antimalarial assay**

The antimalarial activity of ethanol extract and EA fraction of *A. serratus* leaves were assessed by 4-day suppressive test. On the first day (D0) of the experiment, male BALB/C mice weighing 21-25 g were injected with intraperitoneal (i. p.) inoculation of 5% parasitemia infected by *P. berghei* (ANKA strain) erythrocytes in Alsever's suspension of 0.2 ml.

The mice were randomly divided into 7 groups, control group (no treatment), ethanol extract group (dose of 100 mg/kg BW; 10 mg/kg BW; 1 mg/kg BW) and EA group (dose of 100 mg/kg BW; 10 mg/kg BW; 1 mg/kg BW). Each group consisted of 5 mice.

Ethanol extract and EA fraction were suspended in 0.5% CMC-Na and administered orally for 4 d (day 0-day 3). Thin blood smears were made from the tail blood of the mice every day for 7 d (day-0 until day-6), stained with giemsa and then assessed by microscope. Percentage of parasitemia was counted based on infected erythrocytes calculated per 5000 erythrocytes. Percentage of inhibition growth of *P. berghei* was calculated using the formula:

$$100\% - \left(\frac{XE}{XK} \times 100\% \right)$$

Xe: % parasitaemia growth of experimental group

Xk: % parasitaemia growth of negative control

Criteria of antimalarial activity *in vivo*

Criteria of antimalarial activity *in vivo* was based on Munoz et al. (2000), who suggests that an extract is highly active if it shows a value of ED₅₀ < 100 mg/kg BW, active if ED₅₀ 100-250 mg/kg BW, and moderate if ED₅₀ > 500 mg/kg BW [10].

RESULTS AND DISCUSSION**Phytochemical screening**

A. serratus from the Sapindaceae family has compounds such as flavonoids, diterpenoids, glycoside, phenol, saponin, kaempferol, quersetin and β-sitosterol [6]. Fractionation of *A. serratus* extract yielded 4 fractions, which were DCM fraction, EA fraction, BuOH fraction and an aqueous fraction. These four fractions from *A. serratus* extract were analyzed using TLC with a mobile phase of chloroform: methanol (3:1 v/v) and stationary phase of silica gel F254. The spot were observed under UV 254 nm and 366 nm, and derived by 10% of H₂SO₄. TLC result indicated that DCM fraction containing terpenoid compound, shown by red or brownish red coloration. EA fraction contained flavonoid compound, shown by yellow coloration. TLC analysis was also conducted using mobile phase of acetonitrile: methanol: water (2:1:4 v/v) and stationary phase of RP-18. BuOH fraction also contained flavonoid compound but was derived as glycoside compound. TLC profile from aqueous fraction showed polar compound.

Antimalarial *in vitro* and toxicity

Extract and fractions of *A. serratus* were screened for *in vitro* antimalarial activity against *P. falciparum* (3D7 strain) using HRP2 assay. The result of the antimalarial test indicated that ethanol extract, DCM fraction, EA fraction, and BuOH fraction had high activities because its inhibition percentage was more than 90%. The results are summarized in table 1.

Table 1: *In vitro* antimalarial activity of ethanol extract and fractions from *A. serratus* leaves

Sample	Conc (µg/ml)	% Inhibition*
Ethanol extract	100	95.30±0.203
DCM fraction	100	91.26±0.696
EA fraction	100	96.51±0.406
BuOH fraction	100	92.80±0.319
Aqueous fraction	100	54.18±6.349

*Values are mean±SD of triplicates

The result of table 1 became a basis to determine IC_{50} as a parameter of activity. To estimate the potential of extract to inhibit parasite growth without toxicity, the selectivity index (SI) was introduced. SI is defined as the ratio of the toxicity to antimalarial activity. Low SI indicates that the

antiplasmodial activity is probably due to toxicity rather than activity against the parasite themselves. In contrast, higher selectivity (SI>10) indicates potentially safer therapy [3]. The IC_{50} values, CC_{50} values and SI values of ethanol extract and fraction of *A. serratus* are shown in table 2.

Table 2: IC_{50} values, CC_{50} values and SI of ethanol extract and fractions from *A. serratus*

Sample	CC_{50} *	IC_{50} *	SI*
Ethanol extract	>1000	10.27	97.37
DCM fraction	>100	14.81	6.75
EA fraction	>100	9.45	10.58
BuOH fraction	>1000	7.69	130.04

*Values are mean of duplicates.

Extract and fraction exhibited IC_{50} <100 μ g/ml. Extract, EA fraction and BuOH fraction were classified as active with SI values of 97.37, 10.58, and 130.04 and one as partially active (DCM fraction) exhibiting SI of 6.75. Only two samples were found active: EA fraction and BuOH fraction, showing SI>10 and IC_{50} <10 μ g/ml, respectively. EA and BuOH fractions also had IC_{50} values of 9.45 μ g/ml and 7.69 μ g/ml, respectively. Both of them were potential to be selected for further investigation.

In vivo antimalarial assay

The average of parasitemia percentage resulted from ethanol extract and EA fraction group on first-day observation until fifth day (D0-D4) tended to increase, but the increase was not as high as in the control group. This showed that ethanol extract and EA fraction had an effect on the growth of *P. berghei* (ANKA strain) parasites in mice. The results are summarized in table 3 and 4.

Table 3: Activity of ethanol extract of *A. serratus* leaves on *P. berghei* infected mice

Dose (mg/kg BW)	% Parasitemia					% Inhibition
	D0	D1	D2	D3	D4	
Negative control	1.47±0.07	12.80±0.41	20.53±1.45	27.40±2.62	34.22±3.66	
100	1.04*±0.11	2.93*±0.32	6.30*±1.38	8.82*±1.54	12.55*±2.95	64.87±8.90
10	1.57±0.05	5.20*±0.47	10.24*±0.37	14.15*±0.56	18.03*±0.33	49.74±0.90
1	1.84*±0.22	7.01*±0.36	14.11*±0.42	18.07*±0.73	25.44*±0.35	27.95±0.73

Values are mean±SD (n=5). *Denotes significance at the level of P<0.05 versus the control group.

Table 4: Activity of EA fraction of *A. serratus* leaves on *P. berghei* infected mice

Dose (mg/kg BW)	% Parasitemia					% Inhibition
	D0	D1	D2	D3	D4	
Negative control	1.47±0.07	12.80±0.41	20.53±1.45	27.40±2.62	34.22±3.66	
100	1.02*±0.09	2.27*±0.18	4.66*±0.29	6.66*±0.62	8.91*±0.77	75.91±2.63
10	1.11*±0.09	4.30*±0.43	7.02*±0.71	10.60*±0.64	15.11*±0.77	57.29±2.17
1	1.21*±0.03	5.03*±0.17	12.75*±0.64	18.59*±0.58	23.71*±0.66	31.32±1.96

Values are mean±SD (n=5). *Denotes significance at the level of P<0.05 versus the control group.

Parasitemia percentage data were used to determine inhibition percentage. The highest inhibition of ethanol extract was found at a dose of 100 mg/kg BW with the value of 64.87%. Further percentage inhibition at a dose of 10 mg/kg BW was 49.74% and 27.95% at a dose of 1 mg/kg BW. The highest inhibition percentage on EA fraction was also found at a dose of 100 mg/kg BW with the value of 75.91%. Further inhibition at a dose of 10 mg/kg BW was 57.29% and at a dose of 1 mg/kg BW was 31.32%. Inhibition of *P. berghei* growth followed a dose-dependent manner. Higher doses applied to mice produced higher inhibition.

Inhibition percentage was analyzed using log-probit to determine ED_{50} . The result of the analysis showed that ethanol extract and EA fraction of *A. serratus* was highly active as an antimalarial agent with ED_{50} values of 13.82 mg/kg BW and 5.92 mg/kg BW, respectively. EA fraction had higher antimalarial activity and potential as a new source in the development of the antimalarial drug.

TLC profile showed that EA fraction from *A. serratus* leaves contained flavonoid compounds. The previous study showed that flavonoid compounds were active as antimalarial, e.g. artoindonesianin R, artoindonesianin A-2, and artocarpon A isolated from the bark of *Artocarpus champedon*; garcinia xanthone isolated from *Garcinia dulcis* was active as antimalarial with IC_{50} value of 0.96 μ g/ml; biflavonoids isolated from *Selaginella bryopteris* were

also active as antimalarial. The data of antimalarial activity of flavonoid had been reported, including exigua flavone A and exigua flavone B from the stem of *Artemisia indica*, which had antimalarial activity with IC_{50} value of 50 M and 50 μ g/ml [11-13]. It was possible that flavonoid compounds have a contribution in antimalarial activity of EA fraction from *A. serratus* leaves.

CONCLUSION

Ethanol extract and ethyl acetate (EA) fraction *A. serratus* leaves showed antimalarial activities and proved to be non-toxic. Both of them are a good candidate of new source in the development of new antimalarial drugs. The result of TLC profile indicated that EA fraction contained flavonoid compounds. It was concluded that flavonoid compounds from EA fraction took effect on antimalarial activity.

ACKNOWLEDGEMENT

The authors acknowledge grant support from Indonesian Directorate General of Higher Education DIPA BOPTN 2014, contract no 965/UN3/2014.

CONFLICT OF INTERESTS

Declared None

REFERENCES

1. Greenwood BM, Bojang K, Whitty CJ, Targett GA. Malaria. *Lancet* 2005;365:1487-98.
2. Ramalheite C, Lopes D, Mulhovo S, Rosario VE, Jose M, Ferreira U. Antimalarial activity of some plants traditionally used in Mozambique. Workshop Plants Medicinails and Fitoterapeuticas. IICT/CCCM; 2008.
3. Valdes AF, Martinez JM, Lizama RS, Gaiten YG, Rodriguez DA, Payrol JA. *In vitro* antimalarial activity and cytotoxicity of some selected cuban medicinal plants. *Rev Inst Med Trop Sao Paulo* 2010;52:197-201.
4. Mutabingwa TK. Artemisinin-based combination therapies (ACTs) best hope for malaria treatment but inaccessible to the needy. *Acta Trop* 2005;95:305-15.
5. Widyawaruyanti A, Devi A, Fatria N, Tumewu L, Tantular IS, Hafid AF. *In vitro* antimalarial activity screening of several indonesian plants using HRP2 assay. *Int J Pharm Pharm Sci* 2014;6:6-9.
6. Adema F, Leenbouts PW, Van Weizen PC. *Flora Malesiana*. Leiden University; 1994.
7. Trager W, Jensen JB. Human malaria parasites in continuous culture. *Science* 1976;193:673-75.
8. Srivastava K, Singh S, Singh P, Puri SK. *In vitro* cultivation of *Plasmodium falciparum*: Studies with modified medium supplemented with Albumax II and various animal sera. *Exp Parasitol* 2007;116:171-4.
9. Noedl H, Wernsdorfer WH, Miller RS, Wongsrichanalai C. Histidine-rich protein II: a novel approach to malaria drug sensitivity testing. *Antimicrob Agents Chemother* 2002;46:1658-64.
10. Munoz V, Sauvain M, Bourdy G, Callapa J, Bergeron S, Rojas I, et al. A search for natural bioactive compounds in Bolivia through a multidisciplinary approach: Part I. Evaluation of the antimalarial activity of plants used by the Chacobo Indians. *J Ethnopharmacol* 2000;69:127-37.
11. Widyawaruyanti A, Syafrudin, Zaini NC. Antimalarial activity and mechanism of action of flavonoid compound isolated from *Artocarpus chempeden* Spreng stembark. *Biosains* 2011;13:67-77.
12. Saxena S, Pant N, Jain DC, Bhakuni RS. Antimalarial agent from plant source. *Curr Sci* 2003;85:1314-29.
13. Batista R, Junior AJS, Olievera AB. Plant-derived antimalarial agents: new leads and efficient phytomedicines. Part II. Non-alkaloidal natural products. *Molecules* 2009;14:3037-72.