

Antiviral Activity of Marine Sponges *Homaxinella tanitai* and *Microxina Subtilis* against Hepatitis C Virus

by Suciati Suciati

Submission date: 26-Sep-2018 02:32PM (UTC+0800)

Submission ID: 1008646702

File name: RJPBCS_2017.pdf (956.48K)

Word count: 2899

Character count: 14946



ISSN: 0975-8585

Research Journal of Pharmaceutical, Biological and Chemical Sciences

Antiviral Activity of Marine Sponges *Homaxinella tanitai* and *Microxina Subtilis* against Hepatitis C Virus.

Suciati^{1*}, Irsyad Abdillah¹, Muhammad Ihsan Uddin¹, Achmad Fauzi¹, Myrna Adianti², and Achmad Fuad^{1,2}.

¹Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy Universitas Airlangga, Jl. Dharmawangsa Dalam Surabaya 60286, East Java, Indonesia;

²Institute of Tropical Diseases, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java Indonesia

ABSTRACT

The aim of the study was to screen antiviral activity of marine sponges extracts and fractions against hepatitis C virus, as well as to investigate the mode of action of the extract. The results showed that extracts of marine sponges *Homaxinella tanitai* dan *Microxina subtilis* inhibited the growth of HCV at IC₅₀ of 27.1 and 40.5 µg/mL, respectively. Three fractions from *H. tanitai* and two fractions from *M. subtilis* gave more than 50% inhibition against HCV at concentration 100 µg/mL. The time of addition experiment revealed that *H. tanitai* and *M. subtilis* extracts both act at entry and post entry steps. These results suggest that *H. Tanitai* and *M. subtilis* extracts could serve as potential candidates for antiviral agents against hepatitis C virus.

Keywords: antiviral, hepatitis C virus, marine sponges, *Homaxineila tanitai*, *Microxina subtilis*

*Corresponding author:



INTRODUCTION

Hepatitis C virus (HCV) is a positive-stranded RNA virus in the genus Hepacivirus of the family Flaviviridae. HCV has been considered as causative agent of both acute and chronic hepatitis leading to the development of hepatic cirrhosis and hepatocellular carcinoma (1,2). Approximately 130-150 million people globally infected by hepatitis C, and estimated 700,000 people die each year from hepatitis C (World Health Organization, 2016). Based on the heterogeneity of the viral genome, HCV is currently classified into seven genotypes (3-9) and more than 67 subtypes (1a,1b, 2a, 2b etc.) (3, 4). The viral genome, which is a singlestranded open reading encoding a polyprotein precursor consisting of 3000 amino acid residues that is processed by virus-encoded and host cellular proteases into structural and nonstructural proteins. The HCV proteins also play essential roles in the pathological processes associated with HCV infection, such as carcinogenesis as well as glucose and lipid metabolic disorders (4, 5). Current standard therapy for hepatitis C is using triple combination of pegylated interferon, ribavirin and specific NS3 serine protease inhibitor, such as telaprevir and boceprevir. The treatment is considered effective with viral clearance rate to >70% (10,11,12). However, it cause severe side effect, such as anemia and skin rashes from the use of telaprevir and boceprevir, and the presence of drug-resistance virus. Therefore the search for effective and safer therapy for hepatitis C by using natural resources, such as marine sponges is still needed.

The oceans, which occupies over 70% of the Earth's surface, have been the habitat of various living creatures, including algae, sponges, cnidarians, molluscs, bryozoans, ascidians and echinoderms as well as microorganisms. This species diversity makes the marine environment one of the most prolific sources of natural products. Many of these marine creatures produce unique and biologically active compounds which may not be found in the terrestrial ecosystem. These metabolites may be produced as a means of self-defence against predation, since many marine organisms have no spine or protective shell (13). Amongst marine resources, sponges have been the focus of study for many years. Marine sponges are simple, multicellular, and sessile invertebrates with approximately 15,000 species have been discovered Worldwide. More than 5300 metabolites have been reported from this organism (14). Many of these metabolites showed pronounced bioactivity including antiviral.

In this study we investigated anti HCV activity of ethyl acetate extracts of *Hemaxinella tanitai* and *Microxina subtilis* collected from Barrang Lompo Island, South Sulawesi, Indonesia. Flash column fractions of the sponge extracts were also subjected to anti HCV assay. In order to determine the mode of action of the extracts time of addition experiments were conducted.

MATERIAL AND METHODS

Biological Material

Sponges were collected by using SCUBA at a depth of 8-10 m from around Barrang Lompo Island, Makassar, South Sulawesi on May 17th 2014. Samples were kept in plastic pack in ice boxes immediately after collection. Sponge specimens were then frozen at -20°C until analysis. Identification of sponges were conducted by Ecology Laboratory, Department of Biology, Faculty of Mathematic and Sciences, Institut Teknologi Sepuluh November, Surabaya. Voucher specimens were kept in ethanol 70% at the Faculty of Pharmacy, Universitas Airlangga under the accession number 17-5-14-3 and 17-5-14-4.

Extraction and fractionation

Fresh sponges (wet weight 100 – 200 gram) were diced into small pieces and filled in glass container. Samples were extracted in dichloromethane: methanol (1:1) by using ultrasonic vibration for 3 x 5 mins each extraction. The solvent was removed by filtration and the residue was then re-extracted using the same procedure twice. The collected filtrate was evaporated under reduced pressure to give an aqueous residue, which was then partitioned with EtOAc (3 x 50 mL). The organic layer was removed, dried over anhydrous Na₂SO₄, and concentrated under reduced pressure to afford brown viscous extracts.



Anti HCV Assay

The protocol as has been described in (15,16,17). Huh7it cells were seeded in 48-well plates (5×10^4 cells/well) a day before infection. The same volume of JFH1a, with multiplication of infection (MOI) of 0.1 focus-forming units (ffu)/cell, was mixed with serial dilutions of the extracts (100, 50, 25, 12.5, 6.25 and 3.1 $\mu\text{g/ml}$) and inoculated to the cells. Single concentration of 100 $\mu\text{g/ml}$ samples were used for flash column fractions, as well as for time of addition experiments. The mixture was incubated at 37°C for 2 h, and the cells were rinsed twice with serum-free medium to remove residual virus, followed by incubation for an additional 46 h with the same medium. At 48 h p.i., the culture supernatants were collected and used for virus titration.

Virus titration was conducted by placing Huh7it-1 cells (2.4×10^5 cells/well) in a 96-well plate 1 day prior to virus infection. Culture supernatants obtained from HCV-infected cells were serially diluted 25-fold in culture medium and inoculated to the cells. The virus was adsorbed to the cells at 37°C for 2 h, followed by incubation for 46 h with a medium containing 0.4% methylcellulose (Sigma-Aldrich). HCV titers were conducted using a focus formation assay. HCV antigen-positive cells were stained with HCV-infected patient's serum and horseradish peroxidase-conjugated goat anti-human IgG (MBL, Tokyo, Japan). A metal enhanced DAB substrate kit (Thermo Fisher Scientific Inc., Rockford, IL, USA) was used to detect the infectious foci, which were then imaged and counted using the katikati counter software.

The mode of action of the extracts were examined by conducting time of addition experiments. Two sets of experiments were done in parallel: (i) to assess the antiviral effect at the entry step, a mixture of HCV and sample was inoculated into the cells. After virus adsorption for 2 hours, the residual virus and the sample were removed. The cells were then refed with fresh medium without sample for 46 hours; (ii) to assess the antiviral effect at the post entry step, HCV was inoculated to the cells in the absence of the sample. After virus adsorption for 2 hours, medium containing samples were then added followed by incubation at 46 hours; (iii) as a positive control, HCV mixed with the sample was inoculated to the cells. After virus adsorption for 2 hours, the residual virus and the sample were removed, and cells were refed with fresh medium containing the sample for 46 hours. Culture supernatants were obtained at 1 and 2 days post-infection (dpi) and titrated for virus infectivity. Virus and cells treated with medium containing 0.1% DMSO served as a control. The percentage of inhibition of the samples against the virus were then calculated by comparing to the control.

Cytotoxicity Assay

The cytotoxicity of the samples was determined by MTT method as has been described previously (17). The method was based on colorimetric reaction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide with enzyme dehydrogenase inside living cells to form a coloured formazan dye, which corresponded to the number of viable cells. In this assay Huh7it cells were mixed serial dilution of the samples or control in 96 well plates. After 46 hours incubation condition of the cells were observed under microscope. The medium was removed from 96 well plates and then 150 $\mu\text{l/well}$ of MTT 10% was added, followed by incubation for 4 hours at 37°C. The MTT solution was removed from 96 well plates and 100 $\mu\text{l/well}$ of DMSO 100% was then added. Reaction mixtures were homogenize by shaking for 0.5 min before measurement of absorbance at 560 nm and 750 nm.

RESULTS

Anti HCV activities of *Homaxinela tanitai* and *Microxina subtilis* extracts

The ethyl acetate extracts of *H. tanitai* and *M. subtilis* showed anti-HCV activities with 50% inhibitory concentration (IC₅₀) of 27.1 $\mu\text{g/ml}$ and 40.5 $\mu\text{g/ml}$, respectively (Table 1). Cytotoxicity of the crude extract against Huh7it-1 cells was determined using MTT assay. Severe cytotoxicity was not observed at the concentrations tested and the CC₅₀ value was > 100 $\mu\text{g/ml}$. The selectivity index was evaluated (SI; CC₅₀/IC₅₀); the SI values were >3.7 and >2.5 respectively.

Table 1. IC₅₀ and CC₅₀ values of *H. tanitai* and *M. subtilis* extracts

| Samples | IC ₅₀ (µg/ml) | CC ₅₀ (µg/ml) | SI ^a |
|--------------------|--------------------------|--------------------------|-----------------|
| <i>H. tanitai</i> | 27.1 | >100 | >3.7 |
| <i>M. subtilis</i> | 40.5 | >100 | >2.5 |

^a SI: Selectivity index (CC₅₀/IC₅₀)

The ethyl acetate extracts of both marine sponges were subjected to flash column chromatography by using Sepacore® flash column chromatography, employing gradient elution of n-hexane, ethyl acetate and methanol in order of increasing polarity to obtain 7 fractions. Each fractions were then submitted to anti HCV assay using single concentration of 100 µg/ml. The percentage of inhibition of each fractions was calculated (Table 2).

Table 2. Anti HCV activities of *H. tanitai* and *M. subtilis* fractions

| Samples | %Inhibition |
|-------------------------------|-------------|
| <i>H. tanitai</i> Fraction 1 | 0 |
| <i>H. tanitai</i> Fraction 2 | 0 |
| <i>H. tanitai</i> Fraction 3 | 0 |
| <i>H. tanitai</i> Fraction 4 | 88.8 ± 3.5 |
| <i>H. tanitai</i> Fraction 5 | 96.6 ± 2.2 |
| <i>H. tanitai</i> Fraction 6 | 69.2 ± 11.2 |
| <i>H. tanitai</i> Fraction 7 | 49.4 ± 4.7 |
| <i>M. subtilis</i> Fraction 1 | 0 |
| <i>M. subtilis</i> Fraction 2 | 0 |
| <i>M. subtilis</i> Fraction 3 | 0 |
| <i>M. subtilis</i> Fraction 4 | 0 |
| <i>M. subtilis</i> Fraction 5 | 36.8 ± 3.5 |
| <i>M. subtilis</i> Fraction 6 | 88.9 ± 6.9 |
| <i>M. subtilis</i> Fraction 7 | 80.8 ± 8.7 |

Samples were tested at 100 µg/ml; data represented as mean ± SD of two independent experiment each performed in duplicate.

Time of addition experiments were performed to determine the possible mode of action of the extracts. Single concentration of extracts were used in this experiment. Tested samples were added to the cell only during viral inoculation or only after viral inoculation for the remaining culture period until virus harvest. The results (Table 3) suggested that both extracts inhibited HCV activities at both entry and post-entry steps.

Table 3. Time of addition analysis of *H. tanitai* and *M. subtilis* extracts

| Samples ^a | %Inhibition | | | Mode-of-Action |
|----------------------|--------------------|------------------|-------------------------|---------------------------------|
| | During inoculation | Post inoculation | During+Post inoculation | |
| <i>H. tanitai</i> | 88.7 | 84.2 | 99.6 | Entry and post-entry inhibition |
| <i>M. subtilis</i> | 38.1 | 45.4 | 86.9 | Entry and post-entry inhibition |

^aSamples tested at 100 µg/ml

DISCUSSION

Marine sponges have been the fruitful source of many bioactive metabolites, including antiviral. Several marine sponges have been reported to have anti HCV activity, such as *Amphimedon sp.* (18) and undescribed species of the genus *Latrunculia* (19). The ethyl acetate extract of *Amphimedon sp.* showed inhibition of both protease and helicase activities of hepatitis C virus NS3. Anti HCV metabolites have also been successfully isolated from marine sponges, such as halisulfate 3 and sultanine both obtained from unidentified Demospongiae sponge derived from Okinawa Japan. Both compounds inhibited NS3 helicase-activity in a dose dependent manner with IC₅₀ values of 4 and 3 µM, respectively (12). Another sponge metabolite reported to have anti HCV activity is a series of alkaloid discorhabdins, i.e. discorhabdins A, C and dihydrodiscorhabdin C with EC₅₀ value < 10 µM (19).



In this study ethyl acetate extracts of *H. tanitai* and *M. subtilis* exhibited anti HCV activity with IC_{50} values of 27.1 and 40.5 $\mu\text{g/mL}$, respectively. In addition, both extract did not display cytotoxicity at the highest concentration tested 100 $\mu\text{g/mL}$. Flash column chromatography on the extracts were carried out, and obtained seven fractions, which then subjected to anti-HCV assay. The results (Table 2) showed that three fractions from *H. tanitai* and two fractions from *M. subtilis* gave > 50% inhibitions against HCV at concentration of 100 $\mu\text{g/mL}$. In addition, these results also indicated that the possible anti HCV metabolites of *H. tanitai* and *M. subtilis* are semi polar or polar compounds, in which eluted in combination of ethyl acetate and methanol from flash column chromatography.

In order to determine the mode of action of extracts, whether it exerted the effect on the entry or post entry steps, time of addition experiments were carried out. To examine the HCV effect at the entry step, sample was only added at the first 2 h of experiment, followed by incubation of cells with fresh media without sample for 46 h. Meanwhile, to determine whether sample act the post entry step, virus was inoculated into the cell in the absence of sample, after virus adsorption for 2 h, sample was added and incubated for 46h. The results (Table 3) showed that extract of *H. tanitai* exhibited 88.7 and 84.2 % inhibition against HCV at 99.6 $\mu\text{g/mL}$ during inoculation and post inoculation, respectively. However, better inhibition (100%) of extract was obtained when it was added through out experiment. This result suggested that *H. tanitai* extract act at both entry and post entry steps. At the same concentration extract of *M. subtilis* showed lower inhibition against HCV compare to extract of *H. tanitai* at 38.1 and 45.4 % during inoculation and post inoculation, respectively. However, the extract gave higher inhibition at 86.9% when it was added in both steps (during and after inoculation), which indicated that *M. subtilis* extract exerted the activity at entry and post entry steps.

CONCLUSION

The results of this study suggest that the ethyl acetate extracts and fractions of marine sponges *Homaxinella tanitai* and *Microxina subtilis* exhibit antiviral activity against hepatitis C virus. Both extracts exerted the effect at entry and post entry steps.

ACKNOWLEDGEMENT

Authors acknowledge the Ministry of Research, Technology and Higher Education of the Republic of Indonesia for research grant PUPT under contract number 018/SP2H/LT/DRPM/II/2016.

REFERENCES

- [1] Baldo V, Baldovin T, Trivello R, Floreani A. *Curr Pharm Des* 2008; 14: 1646–1654.
- [2] Seeff LB. *Hepatology* 2002; 36: S35–46.
- [3] Gottwein JM, Scheel TK, Jensen TB, Lademann JB, Prentoe JC, Knudsen ML, Hoegh AM, Bukh J. *Hepatology* 2009; 49: 364–77.
- [4] Smith DB, Bukh J, Kuiken C, Muerhoff AS, Rice CM, Stapleton JT, Simmonds P. *Hepatology* 2013; 59: 318–27.
- [5] Moradpour D, Penin F, Rice CM *Nat Rev Microbiol* 2007; 5: 453–6.
- [6] Shoji I, Deng L, Hotta H. *Front Microbiol* 2011; 2: 278.
- [7] Arzumanyan A, Reis HM, Feitelson MA. *Nat Rev Cancer* 2013; 13: 123–35.
- [8] Shepard CW, Finelli L, Alter MJ. *Lancet Infect Dis* 2005; 5: 558–67.
- [9] Mohd Hanafiah K, Groeger J, Flaxman AD, Wiersma ST. *Hepatology* 2013; 57: 1333–42.
- [10] Ghany MG, Nelson DR, Strader DB, Thomas DL, Seeff LB. *Hepatology* 2011; 54: 1433–1444.
- [11] Liang TJ, Ghany MG. *N Engl J Med* 2013; 368: 1907 – 1917.
- [12] Furuta A, Salam KA, Hermawan I, Akimitsu N, Tanaka J, Tani H, Yamasita A, Moriishi K, Nakoshi M, Tsubuki M, Peng PW, Suzuki Y, Yamamoto N, Sekiguchi Y, Tsuneda S, Noda N. *Mar Drugs* 2014; 12: 462 – 476.
- [13] Whitehead R. *Annu. Rep. Prog. Chem. Sect. B* 1999; 95: 183-205.
- [14] Ye J, Zhou F, Al-Kareef AMQ, Wang H. *J Asian Nat Prod Res* 2015; 17: 1 – 23.
- [15] Adianti M, Aoki C, Komoto M, Deng L, Shoji I, Wahyuni TS, Lusida MI, Soetjipto, Fuchino H, Kawahara N, Hotta H. *Microbiol Immunol* 2014; 58: 180 – 187.



ISSN: 0975-8585

- [16] Aoki C, Hartati S, Santi MR, Lydwina, Firdaus R, Hanafi M, Kardono LBS, Shimizu Y, Sudarmono P, Hotta H. *Int J Pharm Pharm Sci* 2014; 6: 211 – 215.
- [17] Tumewu L, Apriyani E, Santi MR, Wahyuni TS, Permanasari AA, Adianti M, Aoki C, Widyawaruyanti A, Hafid AF, Lusida MI, Soetjipto, Hotta H. *Procedia Chemistry* 2016; 18: 169 – 173.
- [18] Fujimoto Y, Salam KA, Furuta A, Matsuda Y, Fujita O, Tani H, Ikeda M, Kato N, Sakamoto N, Maekawa S, Enomoto N, de Voogd NJ, Nakakoshi M, Tsubuki M, Sekiguchi Y, Tsuneda S, Akimitsu N, Noda N, Yamashita A, Tanaka J, Moriishi K. *Plos One* 2012; 7: 1 – 12.
- [19] Na MK, Ding Y, Wang B, Tekwani BL, Schinazi RF, Franzblau S, Kelly M, Stone R, Li, XC, Ferreira D, Hamann MT. *J Nat Prod* 2010; 73: 383 – 387.

Antiviral Activity of Marine Sponges *Homaxinella tanitai* and *Microxina Subtilis* against Hepatitis C Virus

ORIGINALITY REPORT

26%

SIMILARITY INDEX

20%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

| | | |
|----------|--|------------|
| 1 | www.ijppsjournal.com Internet Source | 10% |
| 2 | virologyj.biomedcentral.com Internet Source | 2% |
| 3 | www.mdpi.com Internet Source | 2% |
| 4 | Ken W. L. Yong, I. Wayan Mudianta, Karen L. Cheney, Ernesto Mollo, Joanne T. Blanchfield, Mary J. Garson. " Isolation of Norsesterterpenes and Spongian Diterpenes from (=) ", Journal of Natural Products, 2015 Publication | 1% |
| 5 | www.ncbi.nlm.nih.gov internet Source | 1% |
| 6 | Chowdhry, M. Makroo, R. Singh, M. Agrawa. "Human leucocyte antigen Class I and II alleles associated with anti-hepatitis C virus-positive patie", Indian Journal of Medical Microbiology, | 1% |

July-Sept 2016 Issue

Publication

-
- | | | |
|----------|--|-----------|
| 7 | www.jove.com Internet Source | 1% |
|----------|--|-----------|
-
- | | | |
|----------|--|-----------|
| 8 | "Marine Sponges: Chemicobiological and Biomedical Applications", Springer Nature, 2016 Publication | 1% |
|----------|--|-----------|
-
- | | | |
|----------|--|---------------|
| 9 | academicjournals.org Internet Source | <1% |
|----------|--|---------------|
-
- | | | |
|-----------|---|---------------|
| 10 | www.humanfirewall.com Internet Source | <1% |
|-----------|---|---------------|
-
- | | | |
|-----------|---|---------------|
| 11 | www.neurobio.pitt.edu Internet Source | <1% |
|-----------|---|---------------|
-
- | | | |
|-----------|--|---------------|
| 12 | www.malariaworld.org Internet Source | <1% |
|-----------|--|---------------|
-
- | | | |
|-----------|---|---------------|
| 13 | Darong Yang, Xianghe Meng, Qinqin Yu, Li Xu, Ying Long, Bin Liu, Xiaohong Fang, Haizhen Zhu. "Inhibition of Hepatitis C Virus Infection by DNA Aptamer against Envelope Protein", Antimicrobial Agents and Chemotherapy, 2013 Publication | <1% |
|-----------|---|---------------|
-
- | | | |
|-----------|--|---------------|
| 14 | "The Committee of 3rd International Biology Conference (IBOC) 2016 & 10th Korea-ASEAN Biomass Symposium", AIP Publishing, 2017 Publication | <1% |
|-----------|--|---------------|
-

15 Wu, Qiaozhen, Ying Tang, Xiaoyun Hu, Qin Wang, Wei Lei, Linfu Zhou, and Jianan Huang. <1%
"Regulation of Th1/Th2 balance through OX40/OX40L signalling by glycyrrhizic acid in a murine model of asthma : GA regulation of Th1/Th2 via OX40/OX40L", *Respirology*, 2015.
Publication

16 Furuta, Atsushi, Kazi Abdus Salam, Nobuyoshi Akimitsu, Junichi Tanaka, Hidenori Tani, Atsuya Yamashita, Kohji Moriishi, Masamichi Nakakoshi, Masayoshi Tsubuki, Yuji Sekiguchi, Satoshi Tsuneda, and Naohiro Noda. <1%
"Cholesterol sulfate as a potential inhibitor of hepatitis C virus NS3 helicase", *Journal of Enzyme Inhibition and Medicinal Chemistry*, 2014.
Publication

17 www.wjgnet.com <1%
Internet Source

18 *Natural Products*, 2013. <1%
Publication

19 Calland, Noémie, Jean Dubuisson, Yves Rouillé, and Karin Séron. "Hepatitis C Virus and Natural Compounds: A New Antiviral Approach?", *Viruses*, 2012. <1%
Publication

20 edepot.wur.nl <1 %
Internet Source

21 bio-protocol.org <1 %
Internet Source

22 www.lib.kobe-u.ac.jp <1 %
Internet Source

23 Sa-ngiamsuntorn, Kharit, Adisak
Wongkajornsilp, Phetcharat Phanthong,
Suparerak Borwornpinyo, Narisorn Kitiyanant,
Wasun Chantratita, and Suradej Hongeng. "A
robust model of natural hepatitis C infection
using hepatocyte-like cells derived from human
induced pluripotent stem cells as a long-term
host", Virology Journal, 2016.
Publication

24 "Biomarkers in Liver Disease", Springer Nature,
2017 <1 %
Publication

25 xagenic.com <1 %
Internet Source

26 Matthew Cryer, Kyle Lane, Mary Greer, Rex
Cates et al. " Isolation and identification of
compounds from having human
alphaherpesvirus and vaccinia virus antiviral
activity ", Pharmaceutical Biology, 2017
Publication

27

rome.setac.org

Internet Source

<1%

28

Karamenderes, Canan, Shabana Khan, Babu L. Tekwani, Melissa R. Jacob, and Ikhlas A. Khan. "Antiprotozoal and Antimicrobial Activities of Centaurea. Species Growing in Turkey", *Pharmaceutical Biology*, 2006.

Publication

<1%

29

El-Bitar, Alaa MH, Moustafa MH Sarhan, Chie Aoki, Yusuke Takahara, Mari Komoto, Lin Deng, Mohsen A Moustafa, and Hak Hotta. "Virocidal activity of Egyptian scorpion venoms against hepatitis C virus", *Virology Journal*, 2015.

Publication

<1%

30

Suciati, James A. Fraser, Lynette K. Lambert, Gregory K. Pierens, Paul V. Bernhardt, Mary J. Garson. "Secondary Metabolites of the Sponge-Derived Fungus", *Journal of Natural Products*, 2013

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Antiviral Activity of Marine Sponges *Homaxinella tanitai* and *Microxina Subtilis* against Hepatitis C Virus

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
