



DECEMBER 2015
VOLUME 4, ISSUE 6

Journal of Spine





OMICS International organises 3000+ Global Conferences series Events every year across USA, Europe & Asia with support from 1000 more scientific Societies and Publishes 700+ Open Access Journals which contains over 50000 eminent personalities, reputed scientists as editorial board members.

Open Access Journals gaining more Readers and Citations
 700 Journals and 15,000,000 Readers Each Journal is getting 25,000+ Readers
 This Readership is 10 times more when compared to other Subscription Journals (Source: Google Analytics)

Publication Policies and Ethics

- Author Role
- Editor Role
- Reviewer Role
- Publisher Role

Useful Links

- Aims and Scope
- Article Processing Charges
- Advertise With Us
- Benefits of Publishing
- Citations Report
- Indexing and Archiving
- NIH Funded Work
- Peer Review Process
- Most Viewed Articles
- Track Your Paper
- eBooks

Recommended Journals

- Neuroinfectious Diseases Journals
- Neurological Disorders Journals
- Neurorehabilitation Journals
- Neuroscience Journals
- Neurosurgery Journals
- Orthopedic Journals
- Rheumatology Journals
- Spine & Neurosurgery Journals
- Spine Research Journals

[View More >](#)

Related Subjects

- Health Care Journals
- Medicine Journals
- Neurology Journals
- Neuroscience & Psychology Journals
- Orthopaedics Journals

[View More >](#)

Share This Page



December 2015 | Volume 4, Issue 6

Moving Away from Fusion by Treating the Pain Generator: The Secrets of an Endoscopic Master

Anthony T. Yeung

Editorial: J Spine 2015, 4: e121

DOI: 10.4172/2165-7939.1000e121

[Abstract](#) |
 [Peer-reviewed Full Article](#) |
 [Peer-reviewed Article PDF](#) |
 [Mobile Full Article](#)



Tubercle Bacilli in Spinal Tuberculosis - Morphology, Cell Wall Features, Behaviour and Drugs

Myung-Sang Moon, Hanlim Moon and Sung-Soo Kim

Review Article: J Spine 2015, 4: 265

DOI: 10.4172/2165-7939.1000265

[Abstract](#) |
 [Peer-reviewed Full Article](#) |
 [Peer-reviewed Article PDF](#) |
 [Mobile Full Article](#)



Injectable Bone Substitute Paste Based on Hydroxyapatite, Gelatin and Streptomycin for Spinal Tuberculosis

Hendita Nur Maulida, Dyah Hikmawati and Aniek Setiya Budiati

Research Article: J Spine 2015, 4: 266

DOI: 10.4172/2165-7939.1000266

[Abstract](#) |
 [Peer-reviewed Full Article](#) |
 [Peer-reviewed Article PDF](#) |
 [Mobile Full Article](#)



Cauda Equina Lymphoma - Case Report and Review of Literature

Pedro R Biasi, Luan Lucena, Rafael Augusto Espanhol, Timóteo Abrantes de Lacerda Almeida, Matheus Pintos Brunet, Wellington César de Souza, Eduardo Felipe Martinelli Baldissera, Laura Regyna Toffoli Roso, Luídia Varrone Giacomini, Fernando Luiz Giacomini, Adroaldo Baseggio Mallmann and Charles André Carazzo

Case Report: J Spine 2015, 4: 267

DOI: 10.4172/2165-7939.1000267

[Abstract](#) |
 [Peer-reviewed Full Article](#) |
 [Peer-reviewed Article PDF](#) |
 [Mobile Full Article](#)



Non-Traumatic Anterior Atlas Fracture Following C1 Laminectomy - Case Report

Keishi Tsunoda, Wataru Haraguti, Naoki Kitagawa and Takayuki Matsuo

Research Article: J Spine 2015, 4: 268

DOI: 10.4172/2165-7939.1000268

[Abstract](#) |
 [Peer-reviewed Full Article](#) |
 [Peer-reviewed Article PDF](#) |
 [Mobile Full Article](#)



Relevant Topics

- ADHD
- Alcohol Addiction
- Arthritic Disease
- Arthroplasty
- Backache
- Bacteria Induced Neuropathies
- Biomechanics of Spine
- Bipolar Affective Disorders
- Bone Grafting
- Brain Infection
- Brain Injury Rehabilitation
- Brain Rehabilitation
- CNS and Neurological Disorders
- Cerebral Palsy
- Cervical Laminectomy
- Chiropractic therapy
- Chronic Inflammatory Arthritis
- Clinical Rheumatology
- Cognition
- Drug Addiction
- Epilepsy
- Epilepsy neurorehabilitation
- Experimental Neurology
- Experimental Rheumatology
- Foot and Ankle Surgery
- Fungal Infection
- Hand Surgery
- Hip Replacement
- Infectious Disease in Children
- Ischemic Stroke
- Ischial Spine
- Joint Replacement: Treatment of Arthritic Disease
- Knee Arthroscopy
- Laminectomy
- Lower Back Pain
- Lower Back Surgery
- Lyme
- Medical Rehabilitation
- Minimal Invasive Spine Surgery
- Modern Orthopedics

Tweets by @JournalSpine



Syn-Frame placed with the retractors to form a [Pott's](#) space directly to the [#pathologic](#) vertebra




Figure 2: Syn-Frame placed with the retractors to form a Pott's space directly to the pathologic vertebra

Embed

[View on Twitter](#)

Adult Scoliosis: A Retrospective Analysis of the Correlation between Radiological Parameters and Clinical Outcomes

Giovanni Barbanti Brodano, Lisa Babbi, Marco Girolami, Alessandro Gasbarrini, Stefano Bandiera, Silvia Terzi, Riccardo Ghemondi, Cristiana Griffoni and Stefano Boriani

Research Article: J Spine 2015, 4: 269

DOI: 10.4172/2165-7939.1000269

[Abstract](#) [Peer-reviewed Full Article](#) [Peer-reviewed Article PDF](#) [Mobile Full Article](#)



A Review of Os Odontoideum in Children

Elvira A Allahverdieva, Ryan A Grant and Michael Diluna

Research Article: J Spine 2015, 4: 270

DOI: 10.4172/2165-7939.1000270

[Abstract](#) [Peer-reviewed Full Article](#) [Peer-reviewed Article PDF](#) [Mobile Full Article](#)



Degenerative Cervical Spondylosis- Is Instability the Primary Point of Pathogenesis?

Atul Goel MCh

Commentary: J Spine 2015, 4: 271

DOI: 10.4172/2165-7939.1000271

[Abstract](#) [Peer-reviewed Full Article](#) [Peer-reviewed Article PDF](#) [Mobile Full Article](#)



Ankylosing Spondylitis in Association with Non-Hodgkin's Lymphoma : Successful Anti-TNF Alpha Treatment

Augusta Ortolan, Anna La Salvia, Managrazia Lorenzin, Paola Frallonardo, Dario Manno, Leonardo Punzi and Roberta Ramonda

Case Report: J Spine 2015, 4: 272

DOI: 10.4172/2165-7939.1000272

[Abstract](#) [Peer-reviewed Full Article](#) [Peer-reviewed Article PDF](#) [Mobile Full Article](#)



Transforaminal Lumbar Interbody Fusion (TLIF) Using Pedicle Screws with Mobility

Hideki Ohta, Yoshiyuki Matsumoto, Yoshikazu Nakayama, Tsubasa Sakai, Youhei Iguchi, Masato Tatsumi, Hirotaka Kida and Yoshiharu Takemitsu

Research Article: J Spine 2015, 4: 273

DOI: 10.4172/2165-7939.1000273

[Abstract](#) [Peer-reviewed Full Article](#) [Peer-reviewed Article PDF](#) [Mobile Full Article](#)



Current Status of Interspinous Posterior Devices in USA and Italy: Finally the Pandoras Vase has been Uncovered

Alessandro Landi, Fabrizio Gregori, Biagia la Pira, Angela Ambrosone and Roberto Delfini

Commentary: J Spine 2015, 4: 274

DOI: 10.4172/2165-7939.1000274

[Abstract](#) [Peer-reviewed Full Article](#) [Peer-reviewed Article PDF](#) [Mobile Full Article](#)



Efficacy of Nucleo-Annuloplasty Using Disc-Fx in Lumbar Disc Herniation

Chan Hong Park and Sang Hb Lee

Research Article: J Spine 2015, 4: 275

DOI: 10.4172/2165-7939.1000275

[Abstract](#) [Peer-reviewed Full Article](#) [Peer-reviewed Article PDF](#) [Mobile Full Article](#)



Biochemical Structure and Functions of the Lumbar Disc after an Autologous Chondrocyte Transplantation (ACT) Evaluated by Studies of Two Operated Cases of Repeated Disc Herniation

Abaaslam F, Marszalek P, Glowacki M, Wluka A, Sierzputowski T and Czernicki Z

Research Article: J Spine 2015, 4: 276

DOI: 10.4172/2165-7939.1000276

Abstract

Peer-reviewed Full Article

Peer-reviewed Article PDF

Mobile Full Article



Special Features of - OMICS International Journals

- 700+ Open Access journals
- 21 Days rapid review process
- 50000 Editorial team
- 15 Million readers
- More than 300000 
- Publication immediately after acceptance
- Quality and quick editorial, review processing

OMICS International Journals

Special Features of - OMICS International Conferences

OMICS International Organises 3000+ Conferences Every Year across USA, Europe & Asia

- Well organized scientific program
- 21 Days rapid review process
- 50000 Editorial team
- Renowned speakers and scientists across the globe
- Poster presentations and world class exhibits
- Panel discussions & interactive sessions
- Perfect platform for global networking

OMICS International Conferences

Peer Reviewed Journals

Make the best use of Scientific Research and information from our 700 + peer reviewed, Open Access Journals

Journals by Subject

Agri and Aquaculture	Geology & Earth Science
Biochemistry	Immunology & Microbiology
Bioinformatics & Systems Biology	Informatics
Biomedical Sciences	Materials Science
Business & Management	Mathematics
Chemical Engineering	Medical Sciences
Chemistry	Nanotechnology
Clinical Sciences	Neuroscience & Psychology
Computer Science	Nursing & Health Care
Economics & Accounting	Pharmaceutical Sciences
Engineering	Physics
Environmental Sciences	Plant Sciences
Food & Nutrition	Social & Political Sciences
General Science	Veterinary Sciences
Genetics & Molecular Biology	

Clinical & Medical Journals

Anesthesiology	Molecular Biology	Psychiatry
Cardiology	Nephrology	Pulmonology
Clinical Research	Neurology	Radiology
Dentistry	Nursing	Reproductive Medicine
Dermatology	Nutrition	Surgery
Diabetes & Endocrinology	Oncology	Toxicology
Gastroenterology	Ophthalmology	
Genetics	Orthopaedics	
Haematology	Pathology	
Healthcare	Pediatrics	
Immunology	Physicaltherapy & Rehabilitation	
Infectious Diseases		
Medicine		
Microbiology		

International Conferences 2017-18

Meet Inspiring Speakers and Experts at our 3000+ Global Annual Meetings



Conferences by Country

USA	Spain	Poland
Australia	Canada	Austria
UAE	China	Turkey

Medical & Clinical Conferences

Microbiology	Oncology & Cancer
Diabetes & Endocrinology	Cardiology
Nursing	Dentistry

Conferences By Subject

Pharmaceutical Sciences
Pharma Marketing & Industry
Agri, Food & Aqua

Italy	France	Finland
Germany	India	Ukraine
UK	Malaysia	Denmark
Japan	Singapore	Mexico
Brazil	South Africa	Norway
South Korea	New Zealand	Romania
Netherlands	Philippines	

Healthcare Management	Physical Therapy Rehabilitation
Neuroscience	Psychiatry
Immunology	Infectious Diseases
Gastroenterology	Medical Ethics & Health Policies
Genetics & Molecular Biology	Palliativecare
Pathology	Reproductive Medicine & Women Healthcare
Alternative Healthcare	Surgery
Pediatrics	Radiology
Ophthalmology	

Nutrition
Physics & Materials Science
Environmental Science
EEE & Engineering
Veterinary
Chemical Engineering
Business Management
Massmedia
Geology & Earth science

Contact Us

Agr & Aquaculture Journals

Dr. Krish
agriaquaculture@omicsonline.com
 1-702-714-7001 Extn: 9040

Biochemistry Journals

Datta A
biochemjournals@omicsonline.com
 1-702-714-7001 Extn: 9037

Business & Management Journals

Ronald
business@omicsonline.com
 1-702-714-7001 Extn: 9042

Chemistry Journals

Gabriel Shaw
chemistryjournals@omicsonline.com
 1-702-714-7001 Extn: 9040

Clinical Journals

Datta A
clinicaljournals@omicsonline.com
 1-702-714-7001 Extn: 9037

Engineering Journals

James Franklin
engineeringjournals@omicsonline.com
 1-702-714-7001 Extn: 9042

Food & Nutrition Journals

Katie Wilson
nutritionjournals@omicsonline.com
 1-702-714-7001 Extn: 9042

General Science

Andrea Jason
generalscience@omicsonline.com
 1-702-714-7001 Extn: 9043

Genetics & Molecular Biology Journals

Anna Melissa
geneticsmolbio@omicsonline.com
 1-702-714-7001 Extn: 9006

Immunology & Microbiology Journals

David Goranti
immunomicrobiol@omicsonline.com
 1-702-714-7001 Extn: 9014

Materials Science Journals

Rachle Green
materials@omicsonline.com
 1-702-714-7001 Extn: 9039

Nursing & Health Care Journals

Stephanie Skinner
nursinghealthcare@omicsonline.com
 1-702-714-7001 Extn: 9039

Medical Journals

Nimmi Anna
medicaljournals@omicsonline.com
 1-702-714-7001 Extn: 9038

Neuroscience & Psychology Journals

Nathan T
neuropsychology@omicsonline.com
 1-702-714-7001 Extn: 9041

Pharmaceutical Sciences Journals

Ann Jose
pharmajournals@omicsonline.com
 1-702-714-7001 Extn: 9007

Social & Political Science Journals

Steve Harry
social_politicalsci@omicsonline.com
 1-702-714-7001 Extn: 9042

Injectable Bone Substitute Paste Based on Hydroxyapatite, Gelatin and Streptomycin for Spinal Tuberculosis

Hendita Nur Maulida^{1*}, Dyah Hikmawati² and Aniek Setiya Budiati³

¹Biomedical Engineering Study Program, Faculty of Science and Technology, The University of Airlangga, Surabaya, Indonesia

²Department of Physics, Faculty of Science and Technology, The University of Airlangga, Surabaya, Indonesia

³Department of Clinical Pharmacy, Faculty of Pharmacy, The University of Airlangga, Surabaya, Indonesia

Abstract

World Health Organization (WHO) in 2005 reported that cases of tuberculosis (TB) in the world occur more than 8 million annually and 5-10% was attacked in spine. The most effective treatment of spinal TB is evacuation of infected bone segments and fills with bone graft. It has been synthesized and characterized of Injectable Bone Substitute (IBS) paste based on hydroxyapatite, gelatin and streptomycin. IBS paste synthesized by mixing hydroxyapatite and gelatin 20% w/v with 75:25, 70:30, 65:35 and 60:40 ratio and streptomycin 10 wt%. The mixture was then added with hydroxypropyl methylcellulose (HPMC) 4% w/v as suspending agent. In vitro characterization performed includes acidity (pH), injectability test, setting time, cytotoxicity (MTT assay) and microbacterium test. Acidity test results indicate a fourth variation of the samples had pH values approaching normal body pH (7.3 to 7.6) and is able to maintain stability when measured in 7 days. Injectability test results indicate IBS paste is injectable with the highest percentage of the injectability value at 97.74% ± 0.19%. IBS paste has been setting within 30 minutes to 1 hour when injected on hydroxyapatite scaffold that resembles the bone cavity and is able to cover the pore scaffold seen from the Scanning Electron Microscope (SEM). Scaffold pore size is smaller from range of 780.8 to 835.4 µm into 225.2 µm. MTT assay results showed that IBS paste is not toxic and experiencing proliferation (viability >100%) that are expected to trigger osteoblast cell growth when applied. Microbacterium test results showed that IBS paste is an antibacterial seen from inhibition zone diameter of *Staphylococcus aureus* and has a high strength-sensitive antibacterial. Thus, hydroxyapatite, gelatin and streptomycin composites had qualified as injectable bone substitute which applied in cases of spinal tuberculosis.

Keywords: Spinal tuberculosis; Injectable bone substitute; Hydroxyapatite; Gelatin; Streptomycin; Hydroxypropyl methylcellulose

Introduction

Tuberculosis (TB) is an infectious disease caused by *Mycobacterium tuberculosis* which are systemic and can manifest in almost all organs of the body with the lungs as the primary infection site [1]. World Health Organization (WHO) in 2005 reported that TB cases in the world occur more than 8 million per year and 34% of them occur in Southeast Asia, including Indonesia [2,3]. Indonesia is the 3rd largest contributor to cases of TB after India and China with every 4 minutes there was one patient who died and transmission occurs every 2 seconds [1]. The incidence of extra-pulmonary TB occurs in 25-30% and 5-10% of which occur in the bones and joints and the most attacking in the spine. Bone TB cases is diagnosed when the symptoms have started after one year [4].

Spinal infection by TB could potentially cause serious morbidity including neurological deficits and permanent spinal deformity. Handling of spinal TB in general is giving antituberculosis drugs, immobilization using gips also orthopedic and neurological surgical intervention [5]. Surgical intervention will be taken when after 3-4 weeks of antituberculosis drug administration and conservative therapy does not give a good response. In this case, the most effective spinal lesions treated with immediate surgery to evacuate TB bacteria, take the infected bone and fill the spinal segment involved with bone graft to promote healing and achieve spinal stability [6].

Commercially, the material that can be used as bone graft is Hydroxyapatite (HA). Hydroxyapatite is a bioceramics material that is biocompatible and bioactive because of its mineral content, both physically and chemically very similar to bone [7]. Hydroxyapatite can be obtained from the bones of mammals, fish bones, shells and other materials based on calcium phosphate. However, hydroxyapatite is brittle so that hard formed according to the required implant materials [8]. Some researchers then developed a composite material of

hydroxyapatite with polymers to improve the mechanical properties, one of which gelatin. Gelatin as a polymer derived from natural materials is biodegradable, biocompatible and non-toxic. Gelatin is widely used as space filler because it is easy to set up [9].

Some use in orthopedics requires injectable biomaterial graft which serves as a filler. Injectable system is ready to use, can follow the shape of bone cavity to be filled and polymerized in-situ after being injected. To produce composite injectable, it is required a gel as matrix material maker. Polymers are widely used came from the class of cellulose such as Hydroxypropyl Methylcellulose (HPMC) [8]. HPMC is a water-soluble polymer and is widely used in the food industry as a thickening due to high viscosity [10]. Manufacture of injectable bone substitute (IBS) has been done by several researchers including Weiss et al. [11] with IBS based on calcium phosphate and Shen et al. [12] with IBS based on alendronate and calcium phosphate.

In this study, we have been synthesized and characterized of IBS paste based on nano-hydroxyapatite, gelatin and streptomycin for spinal TB cases. The use of nano-sized hydroxyapatite is expected to easily fit into the pores of the bones thoroughly. While the streptomycin is antibiotic, the primary antituberculosis drugs produced by the soil fungus *Streptomyces griseus*. This drug is readily soluble in water [13].

*Corresponding author: Hendita Nur Maulida, Biomedical Engineering Study Program, Faculty of Science and Technology, The University of Airlangga, Surabaya, Indonesia, Tel: +62 856-5524-2442; E-mail: henditamaulida@gmail.com

Received September 09, 2015; Accepted October 31, 2015; Published November 02, 2015

Citation: Maulida HN, Hikmawati D, Budiati AS (2015) Injectable Bone Substitute Paste Based on Hydroxyapatite, Gelatin and Streptomycin for Spinal Tuberculosis. J Spine 4: 266. doi:10.4172/2165-7939.1000266

Copyright: © 2015 Maulida HN, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The main target of streptomycin is working mechanism at the level of ribosome [14].

Characterization in this study include acidity test to determine degree of acidity of IBS to reach the body's normal pH and its stability as paste, injectability test to measure the ability of IBS can come out of injections, setting time to determine the time of IBS experience setting when injected, MTT Assay test to determine the cytotoxicity of IBS when tested in fibroblast BHK-21 cells and microbacterium test to determine the IBS resistance against bacterial pathogens using *S. aureus*. These characterizations are to evaluate the ability of hydroxyapatite, gelatin, and streptomycin as injectable bone substitute for spinal TB cases.

Materials and Methods

Materials

Tools used include digital scales Mettler Toledo, magnetic stirrer Yellow MAG HS 7, deep freezer, freeze dryer, thermometer, Terumo Syringe 12 ml and 5 ml, glassware (vial bottles, beaker glass and petri dish), micrometers couplers, stopwatch, viscotester VT-04F RION, pH meter Benchtop OAKTON, Scanning Electron Microscope (SEM) FEI Inspect S50 Japan, a set of MTT Assay and microbacterium test.

Materials used include nano-hydroxyapatite of Barramundi fish from the National Nuclear Energy Agency (BATAN) Jakarta, cow skin gelatin from 150 bloom Rousselot (Guangdong, China), streptomycin sulfate from PT. Meiji Indonesian Pharmaceutical Industries, Hydroxypropyl Methylcellulose (HPMC) from Sigma Aldrich, hydroxyapatite scaffold from Tissue Bank of Dr. Soetomo hospital Surabaya, *Staphylococcus aureus*, Trypticase Soy Broth (TSB) and microbiology nutrient agar from Merck.

Synthesis of injectable bone substitute paste

IBS paste made in 4 variations in the composition of hydroxyapatite-gelatin that is 75:25, 70:30, 65:35 and 60:40 (w/w). Gelatin 20% w/v is dissolved in distilled water at 40°C. Hydroxyapatite with variations in composition put into the gelatin solution then added streptomycin 10 wt% as a local dose [15]. Meanwhile, HPMC 4% w/v is dissolved in distilled water at 90°C. Furthermore HPMC solution is added to a solution of gelatin, hydroxyapatite and streptomycin at 40°C and the mixture was stirred for six hours to produce a white IBS paste.

Acidity (pH) test

Acidity test is used to determine the degree of acidity and the stability of IBS as paste when measured in 7 days. Measurement performed with placing electrodes into the sample and pH will automatically appear on the screen of pH meter and make sure that the samples are measured at room temperature.

Injectability test

Injectability conducted to evaluate the ability of IBS paste can come out of the injection within a certain time. IBS first measured the viscosity to ensure the applicable value as injectable system. Measurement results showing viscosity IBS in units dPa.s. Furthermore, injectability test performed using the reference method reported by Shen et al. [12]. In this test, we used the syringe 12 ml with inner diameter of 1.5 cm and needle with inner diameter of 12 mm. Mass of IBS paste before and after injection within 2 minutes measured and its injectability calculated using the following equation.

$$\text{Injectability \%} = \frac{\text{mass expelled from the syringe}}{\text{total mass of the paste before injecting}} \times 100 \% \quad (1)$$

Each test was repeated five times and the average value was calculated as a result of percentage of injectability.

Setting time

Setting time testing was performed using hydroxyapatite scaffold that has been freeze-dried as a substrate [16]. Hydroxyapatite scaffold mass were measured to observe the changes that occur after setting. Testing was performed by injecting IBS samples into the scaffold vertically. Time counting begins when IBS penetrated into the scaffold pores and stopped when the surface of the scaffold was completely dry and covered by the IBS. The test results also were observed using SEM to determine the surface morphology of microscopic scaffold in sizes below 200 nm.

Cytotoxicity test with MTT assay

Cytotoxicity assay was performed using reagents MTT {3-(4,5-dimethyl-2-thiazolil)-2,5-diphenil-2H-tetrazolium bromide} which comprises the step of culturing fibroblast Baby Hamster Kidney (BHK-21) cells, placing samples and reading the results. Optical cell density can be determined using Elisa Reader and use equation (2) to calculate the percentage of living cells. Material is not-toxic if the percentage of living cells is more than 50% [17].

$$\% \text{ Living Cells} = \frac{\text{OD Treatment} + \text{OD Media Control}}{\text{OD Cells Control} + \text{OD Media Control}} \times 100 \% \quad (2)$$

where : OD Treatment = optical density value of the sample after treatment

OD Cells Control = optical density value of the control cells

OD Media Control = optical density value of media control

% Living Cells = the percentage of the number of cells after treatment

Microbacterium test

Microbacterium test is used to determine the IBS resistance against bacterial pathogens by looking at the profile of bacterial inhibition zone diameter for several days. Culture of one ose *Staphylococcus aureus* (SA) suspended in 9 mL Trypticase Soy Broth (TSB) then cultured in incubator 60°C for 24 hours. Absorbance values of SA bacteria are calculated using UV-Visible Spectrometer. Nutrient agar was prepared as a medium for bacteria. IBS paste put in wells agar medium with four repetitions and observed inhibition zone diameter after incubation for 24 hours. IBS strength levels against bacterial resistance were evaluated by comparing clear zone diameter according to Table 1 [18].

Clear Zone Diameter	Microorganism Resistance
>20 mm	Very Sensitive
10-20 mm	Sensitive
5-10 mm	Less Sensitive
<5 mm	Resistant

Table 1: Strength rate of bacteria.

Results and Discussion

Acidity (pH) test

Degree of acidity (pH) becomes one of the important factors to evaluate the performance of IBS material when applied. From the measurement results, we obtained an average pH of each sample with variation composition of HA-gelatin 75:25, 70:30, 65:35 and 60:40 (w/w) respectively is 7.62; 7.54; 7.41 and 7.35. Graph stability of IBS pH value can be seen in Figure 1.




Figure 1: pH Value stability of IBS Paste when measured in 7 days.




Figure 2: Injectability testing results of IBS paste.




Figure 3: Setting time results of IBS paste.




Figure 4: SEM of Scaffold HA Results (a) Before Injected with IBS Samples, Magnification at 50x (b) Magnification at 100x (c) After Scaffold HA injected with IBS Samples and Setting, Magnification at 50x (d) Magnification at 100x.

IBS material requires pH more than 6 to be setting in bone. pH that is not too far from normal pH of around 7.8 (slightly alkaline conditions) can still be tolerated by the body [19]. pH IBS samples were approaching the body's normal pH (about 6.8 to 7.4) are expected to have no pain effect in the bones when it is applied.

Injectability test

IBS paste viscosity values were measured using viscotester show number of 120 dPa.s and applicable in terms of injectability test results as shown in Figure 2. From the test results, it can be concluded that the four samples of IBS has a very good ability in terms of the percentage results approaching 100%. Best percentage of injectability owned by the sample C with HA-gelatin ratio at 65:35 (w/w) is equal to 97.74% ± 0.19%.

Setting time

Setting time performed using HA scaffold freeze-dried as a simulation of human bone parts. Media or substrates that have the same as the main constituent component of the sample will be able to trigger these components for neat and improve the crystallinity. From the test results, the final setting time raised about 30 minutes to 1 hour as shown in Figure 3.

Measurement results of mass scaffold before and after setting also showed an increase. This change is due to the scaffold HA synthesized by the method of freeze-dried is produce a pore that allows the IBS to infiltrate so that the crystallinity increases and pore seemed closed neatly. Changes in mass of the scaffold before and after setting surface microscopically analyzed using SEM as shown in Figure 4.

From the scanning results, it appears that HA scaffold surface is covered evenly by IBS paste and the pores is smaller. The pore size before tested shows the distribution of values in the range of 780.8-835.4 μm and after scaffold injected by IBS in the range of 225.2 μm. It might be concluded that the IBS paste is able to evenly into the pores of the scaffold and bind hydroxyapatite in the vicinity so that crystallinity increases. Thus, IBS paste is able as bone substitution to fill the infected bone segments and vulnerable to further trigger the growth of new bone cells.

Cytotoxicity test with MTT assay

Each various samples of IBS tested in four repetitions and cell viability was calculated according to the equation (2) and then averaged as shown in Figure 5. From the test results, all sample variation of IBS is not-toxic seen from the percentage of cell viability which exceeds 50% [17]. There are several samples that have a percentage of cell viability of more than 100%. This shows that fibroblast cells capable of undergoing proliferation in the sample so that it becomes greater than the control cells. Thus, it can be concluded that the IBS paste can become new cell growth media are also expected to be a medium for the growth of osteoblast cells in bone when applied.




Figure 5: Cytotoxicity test results of IBS paste.

Microbacterium test

Based on the test results, it is known that the IBS paste is antibacterial seen from the area of inhibition zone around IBS discs with a diameter




Figure 6: (a) Petri without IBS, (b) Petri IBS with bacterial inhibition zone diameter.




Figure 7: The increase in inhibition zone diameter average of *S. aureus* bacteria on IBS.

IBS Paste at Variation of HA:Gel (w/w)	Sample Wells Diameter (mm)	Inhibition Zone Diameter (mm)	Clear Zone Diameter (mm)	Resistance Level
75:25	9.0	28.5	19.5	Sensitive
70:30	9.0	28.5	19.5	Sensitive
65:35	9.0	28.5	19.5	Sensitive
60:40	9.0	28.0	19.0	Sensitive

Table 2: Clear Zone Diameter of IBS to Resistance Levels Against *S. aureus* After 24 Hours.

of 28-33 mm (Figure 6). This inhibition zone diameter increased when incubated up to 3 days as shown in Figure 7.

The level of material strength against bacterial resistance can be evaluated by comparing the clear zone diameter by calculating the difference in bacterial inhibition zone diameter with a sample wells diameter. Obtained clear zone diameter test results in Table 2. In addition to a bone filling material, IBS paste also expected to act as drug delivery systems to evacuate the TB bacterium that encourages faster healing and achieve early spinal stability.

Conclusions

1. In the physical characteristics, IBS paste has a pH value close to normal body pH (7.3 to 7.6) and is able to maintain stability when measured in 7 days. IBS paste is injectable with the highest percentage of injectability value at $97.74\% \pm 0.19\%$ and setting within 30 minutes to 1 hour when injected at HA

scaffold that resembles the bone cavity and is able to cover the pore scaffold seen from the SEM results.

2. In the biological characteristics, IBS paste is not-toxic and it is antibacterial seen from inhibition zone diameter of *S. aureus* bacteria.

References

1. Nazar M (2006) Infection of Tuberculosis in Spinal. Medicine magazine Archipelago 39: 279-289.
2. WHO (2009) Global Tuberculosis Control, EPIDEMIOLOGY, STRATEGY, FINANCING. World health organization report.
3. Zuwanda, Janitra R (2013) Diagnosis and Management of Tuberculosis Spondylitis. CDK-208 40:661-673
4. Epi PIG, Purniti PS, Subanada IB, Astawa P (2008) Tuberculosis Spondylitis. Sari Pediatri 10: 177-183.
5. Mehta JS, Bhojraj SY (2001) Tuberculosis of the thoracic spine. A classification based on the selection of surgical strategies. J Bone Joint Surg Br 83:859-863.
6. http://pustaka.unpad.ac.id/wp-content/uploads/2009/05/spondylitis_tuberkulosa.pdf
7. Warastuti Y, Abbas B (2011) Synthesis and Characterization of Irradiated Injectable Bone Substitute Paste Based on Hydroxyapatite. A Scientific Journal for The Applications of Isotopes and Radiation 7: 73-95
8. Bramastya, Yusuf (2013) Effect of Gamma Radiation on Mechanical Properties of UHMWPE-HAp Composite Results from Integrating Mechanical and Felts Heat To Tibial Tray. Bogor Agricultural Institute, Bogor, Indonesia.
9. Nelly S, Sulistiawati F, Fajriani A (2009) Type B Gelatin Gel Strength in Granule Formulation and its Mucoadhesive Characteristic. MAKARA of Health Series 13: 1-4
10. Raymond RC, Sheskey PJ, Quinn ME (2009) Handbook Of Pharmaceutical Excipients. (6thedn), Pharmaceutical Press, USA.
11. Pierre W, Gauthier O, Bouler JM, Grimandi G, Daculsi G (2007) Injectable Bone Substitute Using A Hydrophilic Polymer. BONE 25: 67S- 70S
12. Shen Z, Yu T, Ye J (2014) Microstructure and Properties of Alendronate-Loaded Calcium Phosphate Cement. Mater Sci Eng C Mater Biol Appl 42: 303-311.
13. Martindale (2009) The Complete Drug Reference. (36thedn), Pharmaceutical Press, London.
14. Jawetz, Melnick, Adelberg (2007) Medical Microbiology. (24thedn), The McGraw Hills Companies, USA.
15. Setiya BA, Zaninuddin M, Ferdiansyah, Rachmadani F (2012) Matrix Composites Bovine Hydroxyapatite-Gelatin Effectivity for Gentamicin Delivery System and Regeneration Bone Defect. Airlangga University, Surabaya, Indonesia.
16. Alfian PP (2014) Synthesis and Characterization of Composite Suspension of Hydroxyapatite-Gelatin with Addition Alendromate for Injectable Bone Substitute. Airlangga University, Surabaya, Indonesia.
17. Spielmann H, Seiler A, Bremer S, Hareng L, Hartung T, et al. (2006) The practical application of three validated in vitro embryotoxicity tests. The report and recommendations of an ECVAM/ZEBET workshop (ECVAM workshop 57). Altern Lab Anim 34: 527-538.
18. Theresia M, Eriningsih R, Safitri R (2011) Alginate Membrane as Primary Wound Dressing and Drug Delivery Media of Topical Medications for Infected Wounds. Journal of Industry Research. 2: 161-174.
19. Dorozkhin SV (2011) Medical Application of Calcium Orthophosphate Bioceramics. BIO 1: 1-51.

Citation: Maulida HN, Hikmawati D, Budiati AS (2015) Injectable Bone Substitute Paste Based on Hydroxyapatite, Gelatin and Streptomycin for Spinal Tuberculosis. J Spine 4: 266.doi:10.4172/2165-7939.1000266