

Agrobacterium rhizogenes mediated hairy root induction in Justicia gendarussa Burm.f

by Edy Setiti Wida Utami

Submission date: 06-Apr-2018 03:28PM (UTC+0800)

Submission ID: 942027852

File name: I_ Environ_ Biol_ Sci_ 5_ 4_ 87-93_ 2015_ ISSN_ 2090-4274_ Anggota.pdf (901.21K)

Word count: 3557

Character count: 18573

Agrobacterium rhizogenes Mediated Hairy Root Induction in *Justicia gendarussa* Burm.f

Dwi Kusuma Wahyuni^{1*}, Febri Vidiandi¹, Hery Purnobasuki¹, Tri Muji Ermayanti²,
Bambang Prajoga³, Edy Setiti Wida Utami¹

¹Biology Department, Faculty of Sciences and Technology, Airlangga University

²Center of Biotechnology Research, Indonesian Institute of Sciences

³ Phytochemistry and Pharmacognocny Department, Faculty of Pharmacy, Airlangga University

Received: December 23, 2014

Accepted: February 15, 2015

ABSTRACT

Five strains of *Agrobacterium rhizogenes* (LB510, LB509, YMB072001, A4T, and ATCC 15834) were tested to determine their effects on the induction of hairy root on leaf explants of *Justicia gendarussa* Burm. f.. Leaf explants were sterilized, and then infected within liquid MS medium + sucrose containing *A. rhizogenes* OD600 = 0.1, for 20 minutes while being shaken gently by hand. The explant was co-cultivated on Murashige and Skoog (MS) solid medium at 25°C for 48 hours. After 48 hours of co-cultivation, explants were transferred to MS solid medium with the addition of 250 mg/L cefotaxim, and repeated 1-2 times. Observations were carried out during 6 weeks to observe the specified parameters. Data were analyzed by Kruskal-Wallis Test and followed by Mann-Whitney Test. All strains of *A. rhizogenes* were used in this study were able to produce hairy root. The treatment of LB510 and YMB072001 strain was the best strain in the hairy root induction of *J. gendarussa* Burm. f.. The infection frequency of LB510 strain was highest (100%), longest hairy root (2.23 cm), and highest of hairy root wet weight (0.0243 g). The treatment of YMB072001 strain was the highest number of hairy roots (3.3) per explant and the highest number of hairy roots dry weight (0.0098 g).

KEYWORDS: *Agrobacterium rhizogenes*, hairy root, *Justicia gendarussa* Burm. f.

INTRODUCTION

Gandarusa (*Justicia gendarussa* Burm. f.) is an Indonesian medicinal plant, Acanthaceae family. Gandarusa is used for migraine headaches, fever, hemiplegi, paralysis of facial muscles, swelling, ear pain, inflammation, bronchitis, dyspepsia, eye diseases, bleeding, muscle pain, antirheumatic, antinociception, antihepatotoxic and anti malaria [1, 2]. Based on ethnomedicine study, gandarusa (local name) is used for male birth control drug in the Irian Jaya, Indonesia [3]. Prajogo et al. [4] proved the male antifertility gandarusa efficacious clinically. According Prajoga et al. [5] gandarusa is also useful as antiviral.

Utilizations of gandarusa in the industrialized world are faced with a problem its availability in nature. Gandarusa has not been cultivated and grow wildy. Gandarusa cultivation is a relatively long time traditionally. The level and quality of the desired active compounds are not as expected. Levels of 0.9% of the total flavonoid extraction, while gendarusin A 0.03% [6]. Modern biotechnology interventions are needed to address this problems, one of which is the hairy root culture.

Hairy root culture is the hairy root induction techniques by genetic transformation using *Agrobacterium rhizogenes*. Excess hairy root culture is capacity in the production of secondary metabolites similar to or greater than the parent plant [7]. Trigonelline production in *Trigonella foenum-graecum* hairy root cultures produces three to five times that of plant origin [8]. In industry, the hairy root culture is much more effective than cell culture because of its genetic stability. Culture of the hairy roots can also produce recombinant proteins for potential pharmaceutical industry [9].

A. rhizogenes is a genus of gram-negative soil bacterium, is responsible for the formation of hairy root at the site of infection. These bacteria can transfer the T-DNA, which is in Ri (root inducing) plasmid, the size of few hundred kb, from bacteria to plant cells [10]. Many factors affect the success of the hairy root culture. That are strains of bacteria [11,12,13,14,15], the type of plants [11,13,15,16], parts of plants [11,13,15,16,17], and culture medium [7,11,15,18].

The study of the efficiency of various strains of *A. rhizogenes* on induction on hairy root of gandarusa plants still exist. This study aims to find the right kind of strain for the optimization of the hairy root induction of gandarusa leaf.

*Corresponding Author: Dwi Kusuma Wahyuni, Biology Department, Faculty of Sciences and Technology, Airlangga University. Email: kusumaanwar@yahoo.com

MATERIALS AND METHODS

Rejuvenation of *A. rhizogenes*

A. rhizogenes strain LB510, LB509, YMB072001, A4T, and ATCC15834 was obtained from the Research Center for Biotechnology, Indonesian Institute of Sciences (LIPI) Bogor. The bacteria culture medium were LB and YMB medium of solid and liquid. The bacteria was streaked on solid media. Isolates were stored in the refrigerator for 1 week. The bacteria has been rejuvenated to each 25 mL of YMB medium and liquid LB medium. Then left at room temperature for 24 hours.

Sterilization Leaf Explants

J. gendarussa Burn f. obtained from the Institute of Materia Medica, Batu, Malang. The explants was second leaf of shoots. The explants were washed, then rinsed with running water. Explants were rinsed with water three times. The explants were soaked in 1% fungicide solution for 10 minutes. Explants were rinsed with water three times. The explant was sterilized with 50% Clorox for 5 minutes, then rinsed with sterile distilled water, 3 times.

Induction of hairy roots by *A. rhizogenes*

The sterile explant were infected by inserting into MS medium [19] liquid + sucrose containing *A. rhizogenes* OD600 = 0.1, for 20 minutes. After 48 hours of co-cultivation, the explants were transferred to MS medium with the addition of 250 ppm of cefotaxime, and repeated 1-2 times. Observations were carried out for 6 weeks to observe the parameters determined. Each treatment was repeated 10 times.

Data Analysis

The data were the frequency of infection, length of time the inception of roots, root length, number of roots and root morphology. Quantitative data obtained from this study were analyzed by the Kruskal-Wallis Test at 5% level, followed by The Mann-Whitney Test at 5%. Qualitative data were analyzed descriptively.

RESULTS AND DISCUSSION

The frequency of infection (percentage of explants growing roots compared to the overall number of explants in one treatment) can be determined as shown in table 1. The infection frequency of strain LB510 was highest (100%), 80% of YMB072001 strain, 70% of ATCC15834, 30% of LB509 and 20% of A4T (Table 1). These results was in line with research Manuhara *et al.* [20] that strain LB510 had the optimal frequency of infection for *Talinum paniculatum*, but is not optimal for *Artemisia annua* and *Artemisia cina* [13]. Chandran and Potty [15] showed that different strains affected to different infection frequency of *Ipomoea batatas*, *Solenostemon rotundifolius*, *Vigna vexillata*, and *Canavalia* sp., which were the strain ATCC 15834 had the best results in all species. Strain ATCC 15834 and YMB072001 best for *Artemisia annua* and *Artemisia cina* [13]. Strain 15834 was best for *Solenostemon scutellaroides* [14]. The results showed the frequency of infection of A4T strains was lowest, in contrast to studies of Swain *et al.* [7] showed that the strain A4T was best for *Clitoria ternatea*. According Akramian *et al.* [21] that the different strain of *A. rhizogenes* had the variation of transformation ability. Differences of the virulence of *A. rhizogenes* can be caused by the bacterial plasmids.

Table 1. The frequency of infection of various strains of *A. rhizogenes* on leaf explants *J. gendarussa* Burm. f.

Treatment	Number of infected explant	Replicated	Frequency of infection (%)
Negative Control	0	10	0
LB510	10	10	100
LB509	3	10	30
YMB072001	8	10	80
4T	2	10	20
ATCC15834	7	10	70

Hairy root are formed because of *A. rhizogenes* T-DNA transfered Ri (root inducing) plasmid into the genome of the host plant. The transformation resulted hairy root on the infected part. Opine compounds will be generated and served as a source of nutrients for bacteria [22]. Based on the type of opine, *A. rhizogenes* strains can be separated into five groups, namely octopin, agropin, nopalin, mannopin, and cucumopin [23]. Agropin group was the most commonly used strain [12]. In this study, there were two strains agropin group, including the ATCC15834 and A4T

strains. The LB510, LB509, and YMB072001 strain have not been studied, so that references of all three strains was very little.

Other parameters in this study is the length of hairy root formation time (Figure 1). The hairy root was starting to grow in wound site of the leaf explants *J. gendarussa* Burm. f., on day 14-26 after infected. Based on the average length of the hairy root formation time, the A4T strain was able to induce the growth of hairy root on 14 days, in line to studies on *Amaranthus spinosus*, hairy root emerged on 15-18 days of *A. rhizogenes* strain A4T treatment [24]. This suggested that if the same strain are infected to different species, it will show the difference in the length of hairy root formation time. According to Hu and Du [12] the hairy root will be induced in a short period of time, ranging from one week to over a month, depending on plant species diversity.

Treatment of strains *A. rhizogenes* also affected the number of hairy root formed. The mean number of hairy root was different on each treatment (Figure 2). The average number of hairy root number of strain YMB072001 was 3.3 hairy root (highest), ATCC15834 and LB510 strain was 1.4 hairy root, LB509 strain was 0.5 hairy root and strain A4T was 0.2 hairy root. The results obtained in line with Wahyuni et al. [25] that the number of hairy root of *J. gendarussa* Burm. f. by YMB072001 strain treatment was more than the A4T strain treatment.




Figure 1. Graph of the average length of hairy root formation time on leaf explants *J. gendarussa* Burm. f. with different strains of *A. rhizogenes* treatments (different letters indicate significance at the Mann-Whitney Test with a level of 5%).




Figure 2 Graph of the average number of hairy root on leaf explants of *J. gendarussa* Burm. f. with different strains of *A. rhizogenes* treatment at 6th week (different letters indicate significance at the Mann-Whitney test with a level of 5%).

The observation of hairy root length for six weeks was between 0 to 6.6 cm (Figure 3). The results of these studies was varied in each treatment. LB510 strains induced longest hairy root. In contrast to Puspitaningsih [26] study that strain YMB072001 treatment induced hairy root of *Talinum paniculatum* Gaertn. longer than LB510 strain treatment.




Figure 3. Graph of the average length of hairy root on leaf explants of *J. gendarussa* Burm. f. with different *A. rhizogenes* treatment at 6th weeks (different letters indicate significance at the Mann-Whitney test with a level of 5%).

The growth of the hairy root caused weight gain. Wet weight hairy root obtained is very small, as much as 0.0243 grams LB510, YMB072001 of 0.0241 grams and 0.0112 grams of ATCC15834, whereas LB509 and A4T is zero grams (the wet weight was very small, could not be detected by the balance analytic used in this study). In contrast to the data of wet weight of roots, the highest dry weight was YMB072001 of 0.0098 grams, whereas LB510 lower at 0.0063 grams and 0.0033 grams of ATCC15834.

Table 2. The wet weight and dry weight of hairy root on leaf explants of *J. gendarussa* Burm. f. with different strains of *A. rhizogenes*.

Treatment	Wet weight (gram)	Dry weight (gram)
Negative control	0	0
LB510	0,0243	0,0063
LB509	0	0
YMB072001	0,0241	0,0098
A4T	0	0
ATCC15834	0,0112	0,0033

Hairy root on leaf explants of *J. gendarussa* Burm. f. by LB510 strain treatment was long and branched (Figure 4C). In one explant, there was one or two hairy root. In the treatment of LB509 strain, the roots was thicker at the base (Figure 4B). In the treatment of YMB072001 strain the hairy root was short size, 1-8 hairy root (Figure 4A). In the treatment of A4T strain, the hairy root was different on each explants. The one was length and branched, but the other one was short and unbranched (Figure 4E). In the treatment of ATCC15834 strain, the hairy root was varied on each explants. The one was long and branched but the other one short and unbranched (Figure 4D).




Figure 4 . Hairy root on *J. gendarussa* Burm. f. by *A. rhizogenes* strain treatment. A. YMB072001, B. LB509, C. LB510, D. ATCC15834, E. A4T. hr. hairy root. Bar = 1 cm.

Figure 5 showed that the hairy roots appear on midrib area (Figure 5.1A and 5.2A). It supported Ove et al . [29] study, that hairy root induction genes (*rol* genes) had similar tissue specific expression pattern and are mainly confined to root meristems and the phloem. Danial et al. [30] study showed that the hairy roots of *Eurycoma longifolia* are formed on hypocotyl explant only. In detail analysis showed that whereas the *rolB* promoter seems to be generally expressed in phloem, phloem parenchyma [31]. The *rolC* promoter is specifically expressed in phloem companion cells [32]. The root stem of explants formed roots slightly (Figure 5.2B) was bigger than root stem of the explants formed many roots (Figure 5.1B). The root hair at the ends of branched hairy root (Figure 5.1C) more than unbranched hairy root (Figure 5.2C).




Figure 5. Structure of the hairy root *J. gendarussa* Burm.f. 1. the explants formed many roots. 1A. base of the root, 1B. stem root, 1C. root tip; 2. the explants formed roots slightly. 2A. base of root, 2B. stem root, 2C. root tip, rh. root hair, rb. root branch.

The results of this study shown that *A. rhizogenes* strain of LB510, LB509, YMB072001, A4T, and ATCC154 effected on *J. gendarussa* Burm. f. hairy root on. This is realized from the results of the Kruskal-Wallis statistical analysis that a strain of *A. rhizogenes* influenced of a long time to grow, the number and length of hairy root formed. These results were in line to the study of Lee et al. [10] that the strain of *A. rhizogenes* influenced the induction of hairy root and production of alizarin and purpurin on *Rubia akane* Nakai, hairy root growth and artemisinin production in *Artemisia annua* [27], and the growth of the hairy roots of *Vitis vinifera* [28].

Based on the results, it could be seen that the best strain of *A. rhizogenes* to induce hairy roots on the leaf explants of *J. gendarussa* Burm. f. was LB510 and YMB072001 strain. The infection frequency (100%) and root length (3.3 cm) of LB510 strain was highest. The number of hairy root (3.3) and dry weight of hairy root (0.0098 g) of YMB072001 strain was highest. The length time value of root hair formation, hairy root number and length of hairy root of two of these strains was not differ significantly based on the results of the Mann-Whitney Test. In the other words the ability of both strains to induce hairy root in *J. gendarussa* Burm. f. not significantly different.

7 CONCLUSION

Based on the results and discussion can be concluded that the strain of *A. rhizogenes* strain influenced the induction of hairy root on *J. gendarussa* Burm. f. and the best strains of *A. rhizogenes* in inducing the formation of hairy root in *J. gendarussa* Burm. f. was LB510 and YMB072001 strain.

ACKNOWLEDGEMENT

Thanks to the Ministry of Education of the Republik Indonesia through Dirjend of Higher Education, which has funded this research project through the PUPT research program.

REFERENCES

1. Syamsuhidayat SS. dan J.R. Hutapea, 1991. Inventaris Tanaman Obat Indonesia I. Badan Penelitian dan Pengembangan Kesehatan. Departemen Kesehatan RI.
2. Thomas TD. and H. Yoichiro, 2010. In vitro propagation for conservation of a rare medicinal plant *Justicia gendarussa* Burm.f. by nodal explant and shoot regeneration from callus. *Acta Physiol Plant*, 32:943-950.
3. Suryowinoto M. and A. Pudjoarinto, 1985. Laporan perjalanan ke Jayapura Sentani (Irian Jaya). Fakultas Biologi UGM. Yogyakarta. 19.
4. Prajogo BEW., S. Dudy, dan HS. Mulja, 2007. Analisis kadar gendarusin A pada tanaman budidaya *Justicia gendarussa* Burm. f.. *Jurnal Farmasi Indonesia*, 3: 176 – 180.
5. Prajogo BEW., Nasronudin, P. Widiyanti, dan EB. Aksono, 2010. Pengaruh isolat gendarusin A daun *Justicia gendarussa* Burm. F pada penghambatan reverse transkriptase type I HIV in vitro. Abstrak. DP2M-RM STRATNAS Tahun 2010.
6. Prajogo BEW., S. Dudy, dan HS. Mulja, 2007. Analisis kadar gendarusin A pada tanaman budidaya *Justicia gendarussa* Burm. f.. *Jurnal Farmasi Indonesia*, 3: 176-180.
7. Swain SS, L. Sahu, A. Pal, DP. Barik, C. Pradhan, and PK. Chand, 2012. Hairy root cultures of butterfly pea (*Clitoria ternatea* L.): *Agrobacterium* X plant factor influencing transformation. *World J Microbiol Biotechnol*, 28:729-739.
8. Raheleh A., T. Hasanloo, and M. Khosroshahli, 2011. Evaluation of trigonelline production in *Trigonella foenum-graecum* hairy root cultures of two Iranian masses. *POJ*, 4(7):408-412.
9. Chandran RP. and VP. Potty, 2008. Induction of hairy roots through the mediation of four strains of *Agrobacterium rhizogenes* on five host plants. *Indian Journal of Biotechnology*, 7: 122-128.
10. Lee SY., SG. Kim, WS. Song, YK. Kim, NI. Park, and SU. Park, 2010. Influence of different strains of *Agrobacterium rhizogenes* on hairy root induction and production of alizarin and purpurin in *Rubia akane* Nakai. *Romanian Biotechnological Letters*, 15: 5405-5409.
11. Toruan-Mathius NT., Reflini, N. Haris, J. Santoso, and AP. Roswiem, 2004. Kultur akar rambut *Cinchona ledgeriana* dan *C. succirubra* dalam kultur in vitro. *Menara Perkembangan*, 72(2):72-87.
12. Hu Z. and M. Du, 2006. Hairy roots and its application in plant genetic engineering. *Journal of Integrative Plant Biology*, 48(2): 121-127.
13. Ermayanti TM., E. Al-Hafizh, S. Rahmawati, dan Aryanti, 2009. Transformasi *Artemisia cina* dan *Artemisia annua* dengan *Agrobacterium rhizogenes*. *Journal of Applied and Industrial Biotechnology in Tropical Region*, 2(2):1-5.
14. Saleh NM. and LV. Thuc, 2009. Assessment of hairy roots induction in *Solenostemon scutellarioides* by different strain of *Agrobacterium rhizogenes*. *African Journal of Biotechnology*, 8(15):3519-3523
15. Chandran RP. and VP. Potty, 2011. Different inducer molecules and strains of *Agrobacterium rhizogenes* on enhancing transformation frequency in host plants. *Biotechnology*, 10(2): 203-208.
16. Toruan-Mathius NT., Siti-Chalimah, Muhadiono, L. Aznam, and S. Haran, 2007. Kultur akar rambut in vitro serta pemanfaatan kultur ganda untuk pertumbuhan dan perkembangan endomikoriza (*Gigaspora sp.* dan *Acaulospora sp.*). *Menara Perkebunan*, 75(1): 20-31.
17. Pawar PK. and VL. Maheswari, 2004. *Agrobacterium rhizogenes* mediated hairy root induction in two medicinal important member of family *Solanaceae*. *Indian Journal of Biotechnology*, 3:414-417.

18. John KMM., SD. Joshi, AKA. Mandal, SR. Kumar, and RR. Kumar, 2009. *Agrobacterium rhizogenes*-mediated hairy root production in tea leaves (*Camellia sinensis* (L.) O. Kuntz). Indian Journal of Biotechnology, 8: 430-434.
19. Murashige T. and F. Skoog, 1962. A revised medium for rapid growth and bioassays with tobacco tissue culture. Physiologi Plantarum, 15: 473
20. Manuhara YSW., A. Yachya, AN. Kristanti, 2012. Effect of aeration density on biomass and saponin content of *Talinum paniculatum* Gaertn. hairy root in ballon type bubble bioreactor. J. Pharm. Biomed Sci. 2(4):47-52.
21. Akramian M., SMF. Tabatabaei, and M. Mirmasoumi, 2008. Virulence of different strains of *Agrobacterium rhizogenes* on genetic transformation of four *Hyoscyamus* species. American-Eurasian J. Agric. & Environ. Sci., 3 (5): 759-763.
22. Chilton MD., DA. Tepfer, A. Petit, C. David, F. Casse-Delbart, and J. Tempe, 1982. *Agrobacterium rhizogenes* insert T-DNA into the genome of the host plant roots cells. Nature, 295: 432 – 434.
23. Zhou L., J. Wang, and C. Yang, 1998. Progress on plant hairy root culture and its chemistry. 1. Induction and culture of plant hairy roots. Nat. Product Res. Dev., 10: 87–95.
24. Pal, A., S.S. Swain, K.A. Mukherjee, and K.P. Chand, 2013, *Agrobacterium* pRi TL-DNA *rolB* and TR-DNA opine genes transferred to the spiny amaranth (*Amaranthus spinosus* L.), a nutraceutical crop, Food Technol. Biotechnol, 51 (1): 26–35.
25. Wahyuni DK., A. Prabandari, T.M. Ermayanti, B. Prajoga, dan YSW. Manuhara, 2003. Induksi akar rambut pada eksplan daun gandarusa (*Justicia gendarussa* burm.f.) dengan berbagai konsentrasi bakteri *Agrobacterium rhizogenes*. Proceeding Seminar Nasional Biodiversitas. Universitas Sebelas Maret. Solo.
26. Puspitaningsih, E., 2011, Induksi akar dari eksplan batang ginseng jawa (*Talinum paniculatum* Gaertn.) melalui transformasi gen dengan perantara *Agrobacterium rhizogenes*, Skripsi, Departemen Biologi, Fakultas Sains dan Teknologi, Universitas Airlangga.
27. Giri A. and MJ. Narasu, 2000. Transgenic hairy root: recent trends and application. Biotech Adv., 18: 1-22.
28. Martins TM., A. Domingos, C. Novo, and PML. Lourenço, 2003. Effect of *Agrobacterium rhizogenes* infection on *in vitro* rooting of *Vitis Vinifera*. Vitis, 42 (3): 159–161
29. Ove NN., ONO. Olof, and O. Olsson, 1997. Getting to the root: The role of *Agrobacterium rhizogenes* *rol* genes in the formation of hairy roots. Physiologica Plantarum, 100(3):463-473
30. Danial, M., CL. Keng, SSRS. Alwee, and S. Subramaniam, 2012. Hairy roots induction from difficult-to-transform pharmacologically important plant *Eurycoma longifolia* using wild strains of *Agrobacterium rhizogenes*. Journal of Medical Plants Research, 6(3):479-487

Agrobacterium rhizogenes mediated hairy root induction in Justicia gendarussa Burm.f

ORIGINALITY REPORT

13%

SIMILARITY INDEX

9%

INTERNET SOURCES

11%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

academicjournals.org

Internet Source

2%

2

www.rombio.eu

Internet Source

2%

3

lrd.yahooapis.com

Internet Source

1%

4

Ooi, Chai Theam, Ahmad Syahida, Johnson Stanslas, and Mahmood Maziah. "Efficiency of different Agrobacterium rhizogenes strains on hairy roots induction in Solanum mammosum", World Journal of Microbiology and Biotechnology, 2013.

Publication

1%

5

Recent Trends in Biotechnology and Therapeutic Applications of Medicinal Plants, 2013.

Publication

1%

6

www.scialert.net

Internet Source

1%

7	www.stikes-bth.ac.id Internet Source	1%
8	Samadi, Afsaneh; Carapetian, Jirair; Heidari, Reza; Jafari, Morad and Gorttapeh, Abdollah Hassanzadeh. "Hairy Root Induction in <i>Linum mucronatum</i> ssp. <i>mucronatum</i> , an Anti-Tumor Lignans Producing Plant", <i>Notulae Botanicae Horti Agrobotanici Cluj-Napoca</i> , 2012. Publication	1%
9	www.pomics.com Internet Source	1%
10	<i>Biotechnology in Agriculture and Forestry</i> , 1999. Publication	1%
11	eprints.fbme.utm.my Internet Source	<1%
12	Gatica Arias, Andres Mauricio. "Metabolic engineering of flavonoid biosynthesis in hop (<i>Humulus lupulus</i> L.) for enhancing the production of pharmaceutically active secondary metabolites", <i>Universität Hohenheim</i> , 2013. Publication	<1%
13	media.neliti.com Internet Source	<1%

14

www.frontiersin.org

Internet Source

<1%

15

www.rroij.com

Internet Source

<1%

16

Biotechnology for Medicinal Plants, 2013.

Publication

<1%

17

www.bioversityinternational.org

Internet Source

<1%

18

Jae-An Chun. "Optimization of abiotic factors for improved growth and extracellular production of recombinant fungal phytase in sesame hairy root cultures", *Biotechnology and Bioprocess Engineering*, 06/2007

Publication

<1%

19

Hücherig, Stephanie. "RNAi-Untersuchungen und Überexpression von Genen der Rosmarinsäurebiosynthese", *Philipps-Universität Marburg*, 2010.

Publication

<1%

20

Halder, Mihir, and Sumita Jha. "Enhanced trans-resveratrol production in genetically transformed root cultures of Peanut (*Arachis hypogaea* L.)", *Plant Cell Tissue and Organ Culture (PCTOC)*, 2015.

Publication

<1%

21

"Transgenesis and Secondary Metabolism",
Springer Nature, 2017

Publication

<1%

22

Orhan, E., A. Esitken, S. Ercisli, and F. Sahin.
"Effects of indole-3-butyric acid (IBA), bacteria
and radicle tip-cutting on lateral root induction
in Pistacia vera", The Journal of Horticultural
Science and Biotechnology, 2007.

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Agrobacterium rhizogenes mediated hairy root induction in Justicia gendarussa Burm.f

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
