

The Effect of Organic Nutrient and Growth Regulators on Seed Germination, Embryo and Shoots Development of *Dendrobium antennatum* Lindl. Orchid by In Vitro

by Edy Setiti Wida Utami

Submission date: 10-Apr-2018 12:16PM (UTC+0800)

Submission ID: 944121434

File name: 2_165-171_2016._p-ISSN_2085-191X,_e-ISSN_2338-7610._Ketua.pdf (438.25K)

Word count: 4245

Character count: 21283



5
The Effect of Organic Nutrient and Growth Regulators on Seed Germination, Embryo and Shoots Development of *Dendrobium antennatum* Lindl. Orchid by *In Vitro*

✉ **Edy Setiti Wida Utami, Sucipto Hariyanto**

DOI: 10.15294/biosaintifika.v8i2.5165

32

Department of Biology, Faculty of Science and Technology, Airlangga University, Indonesia

History Article

Received 21 February 2016
Approved 11 June 2016
Published 1 September 2016

Keywords:

coconut water; *Dendrobium antennatum*; 1-Naphthylacetic acid; seed germination; embryo and shoot development

Abstract

Dendrobium antennatum has high economic value as cut flowers and flowerpots. Like orchid seeds in general, *D. antennatum* is difficult to germinate under natural conditions. This study aimed to determine the effect of coconut water on seed germination and embryo development, as well as the effect of NAA on shoots development of *D. antennatum*. This study consisted of two stages. In the first stage, the 12 weeks-old seeds after pollination were sown on MS medium containing 2 g/L peptone + 0%; 5%; 10%; and 20% coconut water. After 8 weeks of culture, the seeds germinated and the shoot formed were recorded. The highest in seed germination (92.2%) and the formation of shoots (51.4%) were obtained when seeds were cultured on MS medium containing 2 g/L peptone + 20% coconut water. In the second stage, the shoots were sub-cultured on MS medium containing 1 mg/L thidiazuron + 0 mg/L; 1 mg/L; 2 mg/L; and 3 mg/L NAA. After 16 weeks of sub-culture, the height of plantlets, the length of the roots and leaves, number of leaves and roots formed were recorded. MS medium containing 1 mg/L thidiazuron + 1 mg/L NAA was the most suitable for the shoots development of *D. antennatum*. The embryo development of *D. antennatum in vitro* begins with the enlargement of embryo, with further it emerges from the seed coat (germinated) followed by the formation of the apical meristems to form the shoots and the roots.

How to Cite

Utami, E., & Hariyanto, S. (2016). The Effect of Organic Nutrient and Growth Regulators on Seed Germination, Embryo and Shoots Development of *Dendrobium antennatum* Lindl. Orchid by *In Vitro*. *Biosaintifika: Journal of Biology & Biology Education*, 8(2), 165-171.

© 2016 Semarang State University

✉ Correspondence Author:
Campus C, Mulyorejo, Surabaya, 60115
E-mail: edysetiti@yahoo.com

p-ISSN 2085-191X
e-ISSN 2338-7610

INTRODUCTION

Dendrobium antennatum Lindl. is known as the antelope orchid, commonly used as an ornamental plant in the pot, complementary bouquets, and cut flowers. The flower is fragrant thus has a potential as a producer of fragrances and medicinal plants (Fanfani and Rossi, 1989). Moreover, *D. antennatum* is flowering more frequent and more durable. Its flowers appear between March and December with the type inflorescence of 9-21 flowers on each stalk (Jones, 2006).

The biological aspects of *D. antennatum* is not yet widely known, especially in terms of seed germination, embryo and shoots development *in vitro*. The knowledge in seed germination, embryo development and regeneration may reflect the degree of ease in propagation of plants that could affect the development of the population. This considered as very useful for conservation and cultivation for ornamental plant, especially for hobbyists and entrepreneurs, in order to estimate the length of time needed to produce *plantlets* ready to acclimatize. The study of embryo development and regeneration of *plantlets D. antennatum* from embryo was expected to provide useful scientific information for its cultivation and conservation. Therefore, this study aimed to understand the seed germination, embryo and shoots development of *D. antennatum in vitro*.

In general, the success in plant tissue culture is influenced by growth regulators and nutrients that are added to the media. The medium used for propagation of orchids, in general contains salt, vitamins, minerals, carbon source, and growth regulators (Bektas *et al.*, 2013). Plant growth regulators which are frequently used in plant tissue culture are a group of auxin and cytokinin. The combination of the type and concentration of these growth regulators added to the culture media have been widely used to enhance the development of shoots in many species of hybrid orchid. Wu *et al.* (2014) reported that the MS medium containing banana extract 100g/L, peptone 1g/L and 10% coconut water added NAA 1mg/L was found to be the best for the formation of *Renanthera imschootiana plantlets*. Similarly, Long *et al.* (2010) from his study reported that Knudson medium containing BA 5mg/L added NAA 0.5 mg/L was able to increase the frequency organogenesis of shoots of *Paphiopedilum villosum var. densissimum* and *Paphiopedilum insigne*. Another important component in plant tissue culture is its organic nutrient (Murdad *et al.*, 2010; Zhang *et al.*, 2013; Zeng *et al.*, 2013). The addition of organic nutrient such as apple extract, ba-

nana extract, potatoes extract, and coconut water have been reported capable to increase germination in some species of orchids (Long *et al.*, 2010; Zeng *et al.*, 2012; Shekarriz *et al.*, 2014; Kaur and Bhutani, 2012). This study evaluated the effect of the addition of coconut water on seed germination and development of the embryo as well as the effect of 1-Naphthylacetic acid (NAA) on the development of *D. antennatum* shoot.

METHODS

Preparation and sterilization of green capsules

The 12 weeks-old capsules after pollination (Figure 1) were washed using a detergent solution for 5 minutes to remove dust particles and were rinsed three times with sterile distilled water. The capsules were put in a laminar flow and they were soaked with 1% sodium hypochloride solution for 10-15 minutes, while they were shaken during this process, and then they were rinsed three times with sterile distilled water. The capsules surfaces sprayed with 70% alcohol and they were flamed and repeated two times. The capsules were put in sterile Petri dish and they were cut into 4 parts. Seeds released from the capsules and were collected with the help of a sterile spatula.

The effect of organic nutrient on seed germination and embryo development of *D. antennatum* Lindl (First stage: Experiment I)

To determine the effect of coconut water in the media on seed germination and embryo development, the seeds were sown on Murashige and Skoog (MS: Murashige and Skoog, 1962) which contained 2g/L peptone (Difco Laboratories Detroit, USA), and was being treated with 0%; 5%; 10%; and 20% coconut water. Sucrose (30g/L) was added into all media (Merck, Made in Germany) and the medium was adjusted to pH 5.6. Media was solidified with 2g/L gellan gum (Phytigel: Sigma Chemical Co., St. Louis, MO) and was sterilized at 120°C for 15 minutes. Each treatment consisted of about 200 seeds, were cultured in flasks contained medium. All treatments were repeated for three times with 5 culture bottles for in each repetition. After 8 weeks of culture, observations of seed germination and embryo development using Stereomicroscope (SM-1 Nikon, Japan). Process in germination seed until the embryo developed into shoot was classified into 6 phases according to the phases of embryo development *D. antennatum* observed, namely: Phase 0, seed, the growth of the embryo was not yet visible. Phase 1, the embryo enlarged. Phase

2, the embryo enlarged and filled the seed coat. Phase 3, ruptured seed coat, embryo emerged from the seed coat (germination). Phase 4, the embryo separated from the seed coat, the apical meristems was visible. Phase 5, the first leaves and/or second leaves appeared (shoots). Percentage at any phase of embryo development were calculated by dividing the number of embryos in each phase divided by the total number of seeds were observed and multiplied by 100% (including seeds that were viable and not viable).

19 Evaluation the effect of growth regulators on the shoot development of *D. antennatum* Lindl (Second stage: Experiment II)

After 8 weeks of culture, shoots obtained from the first experiment were subcultured to new media. Shoots with length 3-5 mm and with 1-2 leaves grown on MS medium containing 1mg/L thidiazuron (Phyto Technology Laboratories, United States) was treated with different concentrations of 0; 1; 2; and 3mg/L 1-Naphthylacetic acid (NAA, MERCK-Schuchardt). Each treatment repeated three times with four culture bottles for each replication. Each culture bottles consisted of 4-5 shoots. The development of shoot covers two aspects: growth and differentiation. For growth, height of the *plantlets*, the length of root and the leaves were observed. For differentiation, the number of roots and leaves were observed. After 16 weeks of culture, *plantlets* were released from the flask and washed with running water to remove the agar. The *plantlets* height, leaf length, root length, number of leaves and roots were recorded.

Data analysis

29 These experiments were designed using Completely Randomized Design. The data were analyzed using analysis of variance with SPSS (Version 14), then the mean values were separated by Duncan's Multiple Range Test (DMRT) at level of significance $\alpha = 0.05$ (Duncan, 1955).

RESULTS AND DISCUSSION

Phases in seed germination and embryo development of *D. antennatum* Lindl .

The seeds of *Dendrobium antennatum* used as explants in this study had an elongated-oval shape with a narrow end and the other is rounded end, 663 μ m in length, and 153 μ m in wide, composed of embryos which were protected by the seed coat. Seed coat looked white, thin, patterned like arrangement of nets. Embryos *D. antennatum* located at the end of the narrowed seed, yellow-

colored, small and undifferentiated (Figure 2A). Phases of seed germination began with changes in the embryo; embryo that was originally yellow-colored and was in size of about 80 μ m (Figure 2A) turned green and shiny and growing embryos enlarged from the previous size which was about 87 μ m (Figure 2B). After 4 weeks of culture, the embryo still growing (Figure 2C), as observed from the size of the embryo increased (188 μ m) and as it filled the seed coat. Furthermore, the embryo was continuously enlarged, followed by the ruptured of the seed coat and embryo emerged from the seed coat (germination phase, Figure 2D). The embryo continued to grow and increased in length followed by the appearance of a protrusion as apical meristems at one end of the embryo (Figure 2E). Once the embryo has reached 5mm in length, the first leaf and the second leaf appeared (Figure 2F).



Figure 1. The 12 weeks-old capsules of *D. antennatum* Lindl after pollination. Bar = 1.4 cm

36 Evaluation the effect of organic nutrient on seed germination and embryo development of *D. antennatum* Lindl

The effect of the addition of organic nutrient (coconut water) on seed germination and embryo development of *D. antennatum* Lindl for 8 weeks after cultured, presented in Table 1.

The results of this study (Table 1) showed that adding coconut water and its concentration have effect on seed germination and embryo development of *D. antennatum*. The highest percentage of total seed germination (92.2%) was obtained when the seeds germinated on MS media given in combination of organic nutrient, namely 2g/L peptone + 20% coconut water and was significantly different with the control (2g/L peptone only) and with other treatments. The lowest seed germination percentage (77.33%) was obtained when the seeds germinated on MS media given with only a single organic nutrient that was 2g/L




Figure 2. Seed culture and embryo development of *Dendrobium antennatum* Lindl. *in vitro*. (A) Phase 0, seed, the growth of the embryo was not yet visible. (B) Phase 1, the embryo enlarged. (C) Phase 2, the embryo enlarged and filled the seed coat. (D) Phase 3, ruptured seed coat and embryo emerged from the seed coat (germinated). (E) Phase 4, the embryo separated from the seed coat and the apical meristems was visible. (F) Phase 5, the first leaves and second leaves appeared (shoots). (G) Shoot growth on MS medium containing peptone 2g/L + 20% coconut water. e: Embryo, sc: Seed coat, sm: Shoot meristem, d₁: First leaf, d₂: Second leaf. Bars: (A) 80µm, (B) 87µm, (C) 188µm, (D) 210µm, (E) 350µm, (F) 500µm, (G) 0.6 cm.

Table 1. The effect of coconut water with different concentration on seed germination and embryo development of *D. antennatum* Lindl at MS medium containing 2g/L peptone, 8 weeks after culture.

Coconut water concentration (%)	Phases of embryo development					Total germinated (Phase 3-5)
	Phase 1	Phase 2	Phase 3 (germinated)	Phase 4	Phase 5 (shoot)	
0	12,1±2,6 ^d	10,6±1,2 ^{bc}	42,5±2,4 ^c	19,8±2,0 ^c	14,9±1,8 ^a	77,3±2,1 ^a
5	10,6±1,2 ^c	9,7±1,4 ^b	35,3±2,0 ^b	20,4±2,3 ^c	23,9±1,6 ^b	79,7±2,1 ^b
10	8,3±1,4 ^b	11,6±2,6 ^c	27,8±2,0 ^a	17,1±1,5 ^b	35,3±2,0 ^c	80,2±2,8 ^b
20	2,7±1,6 ^a	5,1±1,6 ^a	26,3±1,1 ^a	14,4±1,9 ^a	51,4±1,5 ^d	92,2±1,5 ^c

The mean ± SD followed by different letters in the same column indicate significant difference according to Duncan's test at a significance level $\alpha = 0.05$.

peptone (control). Similarly, the observation in the percentage of shoot formed (phase 5), which indicated that there was an effect of coconut water as organic nutrient at various concentrations on the shoot development of *D. antennatum*. The highest percentage of shoot formation (51.44%) was found when the seeds germinated on MS medium given with a combination of organic nutrient 2g/L peptone + 20% coconut water and was significantly different from the control treatment

(2g/L peptone). The lowest percentage of shoot formation (14.99%) was found when the seeds germinated on MS medium were given only with a single organic nutrient that was 2g/L peptone, this shows that the seeds were able to germinate and to grow into shoots on MS medium using only a single organic nutrient, but the percentage is lower when compared with provided with combination of organic nutrient. The same results had been reported as well by Utami *et al.*,

(2015) that the seeds of *Paphiopedilum liemianum* cultured in VW media given only with 2g/L of peptone were also able to germinate and to develop into *plantlets*. This is possible because peptone contains vitamin ie: biotin, pyridoxine, thiamin, nitrogen (Dutra *et al.*, 2008) and amino acids, proteins (Nhut *et al.*, 2008) which are able to increase the growth and development of the explants. The results also indicated that MS medium provided with 2g/L peptone + 20% coconut water as organic nutrient was suitable for seed germination and for further embryo development in order to form shoot. Hasanah *et al.*, (2014) reported that the addition of 15% coconut water into the MS medium containing Hyponex as fertilizers is able to increase the growth of *Dendrobium kelemense plantlets*. Enhancement in percentage of seed germination and embryo development on such treatment is possible because in MS medium other than by peptone, was also being provided with coconut water that contains carbohydrates, vitamins, amino acids, organic acids, enzymes that play an important roles for the development of plant cells (Yong *et al.*, 2009; Nambiar *et al.*, 2012). The percentage of germinated seeds (Phase 3) was 26.3% and 14.4% (Phase 4), where it was found that the seeds cultured in the media provided with 2g/L peptone + 20% coconut water was lower than other treatments. This is because the treatment caused the embryo further developed into phase 5 (shoot).

Evaluation the effect of growth regulators (NAA) on shoot development of *D. antennatum* Lindl

To determine the effect of various concentrations of NAA as growth regulators on the growth and differentiation of shoot, 70 days-old shoots obtained from the first experiment with, 3-5 mm long, consisted 1-2 leaves, and without root were being transferred on MS medium containing 1mg/L thidiazuron + NAA with different concentration: 0mg/L; 1mg/L; 2mg/L; 3mg/L. The result (Figure 3 and 4) showed that the addition of NAA in MS medium resulted to a positive response compared to the control treatment (1mg/L thidiazuron without NAA). The maximum of growth and differentiation response was observed when the shoots were sub-cultured on MS medium containing thidiazuron at a concentration of 1mg/L added with low concentrations of NAA (1mg/L), which gained an average of the height of plantlets 44,5mm, length of the leaf 19mm, length of the root 18mm, consisted of 7.2 leaves and 8 roots/*plantlets*. This was possible because the medium, provided with cytokinin (thidiazuron) and auxin (NAA), which plays roles in differentiation of cell and plant tissue as well as in the formation of xylem and roots (Vogel and Mucedo, 2011). The results (Figure 3 and 4) also showed that the presence of plant growth regulator cytokinin, ie. 1mg/L thidiazuron with-




Figure 3. The effect of growth regulators NAA on shoot differentiation of *D. antennatum* at 16 weeks after the culture. P0 = MS medium containing 1mg/L thidiazuron, without NAA, P1 = MS medium containing 1mg/L thidiazuron + 1mg/L NAA, P2 = MS medium containing 1mg/L thidiazuron + 2mg/L NAA, P3 = MS medium containing 1mg/L thidiazuron + 3mg/L NAA. The mean number of leaves in treatment P0 = 7.1 ± 1.0; P1 = 7.2 ± 0.6; P2 = 6.5 ± 0.3; and P3 = 5.8 ± 1.1. The mean number of roots in the treatment of P0 = 1.1 ± 0.2; P1 = 8.0 ± 1.0; P2 = 6.7 ± 1.3; and P3 = 5.4 ± 0.9. The mean ± SD followed by different letters indicate significant difference according to Duncan's test at a significance level $\alpha = 0.05$




Figure 4. The effect of growth regulators NAA on shoot growth of *D. antennatum* at 16 weeks after the culture. P0 = MS medium containing 1mg/L thidiazuron, without NAA, P1 = MS medium containing 1mg/L thidiazuron + 1mg/L NAA, P2 = MS medium containing 1mg/L thidiazuron + 2mg/L NAA, P3 = MS medium containing 1mg/L thidiazuron + 3mg/L NAA. Average length of leaves (mm) in the treatment P0 = 10.9 ± 0.4 ; P1 = 19.0 ± 1.3 ; P2 = 14.4 ± 1.6 ; and P3 = 13.8 ± 0.9 . The mean of length of the roots (mm) in the treatment P0 = 4.6 ± 0.5 ; P1 = 18.0 ± 1.6 ; P2 = 11.8 ± 1.2 ; and P3 = 12.7 ± 0.9 . The mean of the height of *plantlets* (mm) in the treatment P0 = 38.3 ± 0.5 ; P1 = 44.5 ± 1.0 ; P2 = 39.8 ± 1.5 ; and P3 = 39.9 ± 1.5 . The mean \pm SD followed by different letters indicate significant difference according to Duncan's test at a significance level $\alpha = 0.05$.




Figure 5. The shoots development of *D. antennatum* Lindl. (A). 2 weeks-old shoots were cultured on MS medium containing 1mg/L thidiazuron + 1mg/L NAA. (B). *Plantlets* resulted from the growth and differentiation of shoots on MS medium containing 1mg/L thidiazuron + 1mg/L NAA after 16 weeks of culture. The roots (arrow) appeared at the bottom of the *plantlet*. Bars:(A) 0.75cm, (B) 0.83cm.

out NAA, can increase the shoot differentiation, forming the leaf which was 7.1 leaves/*plantlet* in average; higher when compared with using combination of 1mg/L thidiazuron + 2mg/L NAA (6.5 leaves/*plantlet*) and 1mg/L thidiazuron + 3mg/L NAA (5.8 leaves/*plantlet* in average). This was possible because thidiazuron plays roles in

cell division and organ regeneration (Guo *et al.*, 2011; Oluk and Orthan, 2010). Enhanced shoot formation of *Astragalus cariensis* with thidiazuron had been reported by Erisen *et al.*, (2011) as well as in growth in seedling *Paphiopedilum liemianum* (Utami *et al.*, 2015).

CONCLUSION

An increase in the percentage of seed germination and embryo development of *D. antennatum* on MS medium treated with the combination of organic nutrient compared to a single treatment. The percentage of seed germination was 92.22 % and the highest shoot formation 51.44% was found when the seeds were cultured on MS medium containing 2g/L peptone + 20% coconut water. The addition of NAA in MS medium containing 1mg /L thidiazuron increased the differentiation and growth of shoot. The highest formation of roots (8 root/plantlet), the formation of leaves (7 leaves/plantlet) as well as in term of the length of roots (18mm), height of plantlets (44.5mm), and length of leaves (19mm) obtained when shoots were sub-cultured on MS medium containing 1mg/L thidiazuron + low concentration of NAA (1mg/L).

REFERENCES

- Bektas, E., Cuce, M., & Sokmen, A. (2013). *In vitro* germination, protocorm formation, and plantlet development of *Orchis colliophora* (Orchidaceae), a naturally growing species in Turkey. *Turkish Journal of Botany*, 37(2), 336-342.
- Duncan, D. B. (1955). Multiple range and multiple F tests. *Biometrics*, 11: 1-42.
- Dutra, D., Johnson, T. R., Kauth, P. J., Stewart, S. L., Kane, M. E., & Richardson, L. (2008). Asymbiotic seed germination, *in vitro* seedling development, and greenhouse acclimatization of the threatened terrestrial orchid *Bletia purpurea*. *Plant Cell, Tissue and Organ Culture*, 94(1), 11-21.
- Erisen, S., Atalay, L., & Yorgancilar, M. (2011). The effect of thidiazuron on the *in vitro* shoot development of endemic *Astragalus cariensis* in Turkey. *Turkish Journal of Botany*, 35(5), 521-526.
- Fanfani, A., & Rossi, W. (1989). *Guide to orchid*. Simon & Schuster Inc., New York.
- Guo, B., Abbasi, B. H., Zeb, A., Xu, L. L., & Wei, Y. H. (2011). Thidiazuron: A multi-dimensional plant growth regulator. *African Journal of Biotechnology*, 10(45), 8984-9000.
- Hasanah, U., Suwarsi, R. S., & Sumadi. (2014). Utilization of fertilizer leaves, coconut water and banana porridge as component *Dendrobium kelemense* orchid plantlet growth medium. *Biosaintifika: Journal of Biology & Biology Education*, 6(2), 161-168.
- Jones, D.L. (2006). *A complete to native orchids of Australia*. Reed New Holland, Sydney.
- Kaur, S., & Bhutani K.K. (2012). Organic growth supplement stimulants for *in vitro* multiplication of *Cymbidium pendulum* (Roxb.) Sw. *Hort. Sci*, 39(1), 47-52.
- Long, B., Niemiera, A.X., Cheng, Z.Y., & Long, C.L. (2010). *In vitro* propagation of four threatened *Paphiopedilum* species (Orchidaceae). *Plant Cell, Tissue and Organ Culture*, 101(2), 151-162.
- Murashige, T., & Skoog, G. (1962). A revised medium for rapid growth and bioassay with tobacco tissue culture. *Physiology Plant*, 15(3), 473-479.
- Murdad, R., Latip, M.A., Aziz, Z.A., & Ripin, R. (2010). Effect of carbon source and potato homogenate on *in vitro* growth and development of Sabah'Endangered orchid: *Phalaenopsis gigantea*. *Asia-Pac J. Mol. Biol. Biotechnol* 18 (1): 199-202.
- Nambiar, N., Tee, C.S., & Maziah M. (2012). Effect of organic additives and different carbohydrate source on proliferation of protocorm-like bodies in *Dendrobium* Alya Pink. *Plant Omics Journal*, 5(1), 10-18.
- Nhut, D. T., Thi, N. N., Khiet, B. L. T., & Luan, V. Q. (2008). Peptone stimulates *in vitro* shoot and root regeneration of Avocado (*Persea americana* Mill). *Scientia Horticulturae*, 115(2), 124-128.
- Oluk, E. A., & Orthan, S. (2010). Thidiazuron induced micropropagation of *Hypericum triquetrifolium* Turra. *Afr. J. Biotechnol*, 8(15), 3506-3510.
- Shekarriz, P., Kafi, M., Deilamy, S. D., & Mirmasounii, M. (2014). Coconut water and peptone improve seed germination and protocorm like body formation of hybrid *Phalaenopsis*. *Agriculture Science Development*, 3(10), 317-323.
- Utami, E.S.W., Purnobasuki, H., Soedarti, T., & Hariyanto, S. (2015). Asymbiotic seed germination and *in vitro* seedling development of *Paphiopedilum liemianum* Fowlie, an endangered terrestrial orchid in northern Sumatra, Indonesia. *Journal of Plant Sciences*, 10(1), 25-34.
- Vogel, I.N., & Maceo, A.F. (2011). Influence of IAA, TDZ and light quality on asymbiotic germination, protocorm formation and plantlet development of *Cyrtopodium glutiniferum* Raddi., a medicinal orchid. *Plant Cell, Tissue and Organ Culture*, 104(2), 147-155.
- Wu, K., Zeng, S., Lin, D., Teixeira da Silva, J.A., Bu, Z., Zhang, J., & Duan, J. (2014). *In vitro* propagation and reintroduction of the endangered *Renanthera imshootiana* Rolfe. *PLOS* 9(10), 1-12.
- Yong, J. W. H., Ge, L., Fei, Ng Y., & Tan, S. N. (2009). The chemical composition and biological properties of coconut (*Cocos nucifera* L.) water. *Molecules*, 14(12), 5144-5164.
- Zeng, S., Wu, K., Teixeira da Silva, J.A., Zhang, J., Chen, Z., Xia, N., & Duan, J. (2012). Asymbiotic seed germination, seedling development and reintroduction of *Paphiopedilum wardii* Sumerh., an endangered terrestrial orchid. *Scientia Horticulturae*, 138, 198-209.
- Zeng, S., Wang, J., Wu, K., Teixeira da Silva, J.A., Zhang, J., & Duan, J. (2013). *In vitro* propagation of *Paphiopedilum hangianum* Perner & Gruss. *Scientia Horticulturae*, 151, 147-156.
- Zhang, Y., I Lee, Y., Deng, L., & Zhao, S. (2013). Asymbiotic germination of immature seeds and seedling development of *Cypripedium macranthos* Sw., an endangered lady's slipper orchid. *Scientia Horticulturae*, 164, 130-136.

The Effect of Organic Nutrient and Growth Regulators on Seed Germination, Embryo and Shoots Development of *Dendrobium antennatum* Lindl. Orchid by In Vitro

ORIGINALITY REPORT

25%

SIMILARITY INDEX

21%

INTERNET SOURCES

21%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

kb.psu.ac.th

Internet Source

2%

2

biologi.fst.unair.ac.id

Internet Source

2%

3

W. Udomdee, P.J. Wen, S.W. Chin, F.C. Chen.

"EFFECT OF CARBON SOURCE ON PROTOCORM-LIKE BODY INDUCTION, PROLIFERATION AND REGENERATION IN DENDROBIUM SNOWFLAKE 'RED STAR'",
Acta Horticulturae, 2015

Publication

1%

4

dergipark.ulakbim.gov.tr

Internet Source

1%

5

www.scilit.net

Internet Source

1%

6

indianforester.co.in

Internet Source

1%

7	lankesteriana.org Internet Source	1%
8	www.kew.org Internet Source	1%
9	world-food.net Internet Source	1%
10	Otieres Cirino de Carvalho, Vespasiano Borges de Paiva Neto, Daly Roxana Castro Padilha, Tomás Gomes Reis Veloso et al. "Cyrtopodium paludicolum germination with two Tulasnella isolates", Acta Botanica Brasilica, 2017 Publication	1%
11	Aasim, M.. "In vitro micropropagation from plumular apices of Turkish cowpea (Vigna unguiculata L.) cultivar Akkiz", Scientia Horticulturae, 20091001 Publication	1%
12	www.academicjournals.org Internet Source	1%
13	waprogramming.com Internet Source	1%
14	Abdullah , Ghassan Rashid Al-Khateeb , Abdullatif Ali Layous , Laurene Najib. "Response of the Strawberry Cv. Elsanta Micro Propagation in Vitro to Different Carbon	1%

Sources and Concentrations \ Jordan Journal of Agricultural Sciences .- 2013 , Vol. 9 , No. 1 , pp. 1 - 11.", , 2013.

Publication

15

www.transectscience.org

Internet Source

1%

16

ir.library.osaka-u.ac.jp

Internet Source

1%

17

Yao Ouyang, Yulu Chen, Jinfeng Lü, Jaime A. Teixeira da Silva, Xinhua Zhang, Guohua Ma. "Somatic embryogenesis and enhanced shoot organogenesis in *Metabriggsia ovalifolia* W. T. Wang", *Scientific Reports*, 2016

Publication

1%

18

Bhattacharyya, Paromik, Suman Kumaria, Nikhil Job, and Pramod Tandon. "Phyto-molecular profiling and assessment of antioxidant activity within micropropagated plants of *Dendrobium thyrsiflorum*: a threatened, medicinal orchid", *Plant Cell Tissue and Organ Culture (PCTOC)*, 2015.

Publication

<1%

19

www.revistas.unal.edu.co

Internet Source

<1%

20

agriculturejournals.cz

Internet Source

<1%

- 21 Winarto, Budi, Fitri Rachmawati, Anggraeni Santi, and Jaime A. Teixeira da Silva. "Mass propagation of Dendrobium 'Zahra FR 62', a new hybrid used for cut flowers, using bioreactor culture", *Scientia Horticulturae*, 2013.
Publication <1%
-
- 22 pgmp.agro.ufg.br
Internet Source <1%
-
- 23 Parthibhan, S.. "Influence of nutritional media and photoperiods on in vitro asymbiotic seed germination and seedling development of *Dendrobium aqueum* Lindley", *African Journal of Plant Science*, 2012.
Publication <1%
-
- 24 ARDA, Hayati, DAYAN, Sergun, KARTAL, Çiler and GÜLER, Necmettin. "In vitro CONSERVATION OF CRITICALLY ENDANGERED *Dianthus ingoldbyi* Turritt UNDER SLOW GROWTH CONDITIONS", *Trakya Üniversitesi Fen Bilimleri Enstitüsü*, 2016.
Publication <1%
-
- 25 academicjournals.org
Internet Source <1%
-
- 26 Anuradha Parale. "Influence of organic <1%

supplements on production of shoot and callus biomass and accumulation of bacoside in *Bacopa monniera* (L.) Pennell.", *Physiology and Molecular Biology of Plants*, 04/2010

Publication

27

eprints.uns.ac.id

Internet Source

<1%

28

journalarticle.ukm.my

Internet Source

<1%

29

media.neliti.com

Internet Source

<1%

30

Guimarães, Fernanda Aparecida Rodrigues, Marlon Corrêa Pereira, Cristiane da Silva Felício, Denise Pereira Torres, Sabrina Feliciano Oliveira, Tomás Gomes Reis Veloso, and Maria Catarina Megumi Kasuya. "Symbiotic propagation of seedlings of *Cyrtopodium glutiniferum* Raddi (Orchidaceae)", *Acta Botanica Brasilica*, 2013.

Publication

<1%

31

biosains.mipa.uns.ac.id

Internet Source

<1%

32

www.ibnusina.utm.my

Internet Source

<1%

33

www.plantpath.cornell.edu

Internet Source

<1%

34

S., Anjusha, and Gangaprasad A.. "Callus culture and in vitro production of anthraquinone in *Gynochthodes umbellata* (L.) Razafim. & B. Bremer (Rubiaceae)", *Industrial Crops and Products*, 2016.

Publication

<1%

35

[insightsociety.org](https://www.insightsociety.org)

Internet Source

<1%

36

staging.vivo.ufl.edu

Internet Source

<1%

37

Zhang, Yu, Yung-I Lee, Lian Deng, and Shiwei Zhao. "Asymbiotic germination of immature seeds and the seedling development of *Cypripedium macranthos* Sw., an endangered lady's slipper orchid", *Scientia Horticulturae*, 2013.

Publication

<1%

38

www.dtic.mil

Internet Source

<1%

39

Zeng, Songjun, Weichang Huang, Kunlin Wu, Jianxia Zhang, Jaime A. Teixeira da Silva, and Jun Duan. "In vitro propagation of *Paphiopedilum* orchids", *Critical Reviews in Biotechnology*, 2015.

Publication

<1%

40

Wiszniewska, Alina, Barbara Nowak, Anna

Kołton, Ewa Sitek, Krzysztof Grabski, Michał Dziurka, Olga Długosz-Grochowska, Kinga Dziurka, and Zbigniew Tukaj. "Rooting response of *Prunus domestica* L. microshoots in the presence of phytoactive medium supplements", *Plant Cell Tissue and Organ Culture (PCTOC)*, 2016.

Publication

<1%

41

www.lfbentonite.com

Internet Source

<1%

42

Thakur, Julie, Mayank D. Dwivedi, Pragya Sourabh, Prem L. Uniyal, and Arun K. Pandey. "Genetic Homogeneity Revealed Using SCoT, ISSR and RAPD Markers in Micropropagated *Pittosporum eriocarpum* Royle- An Endemic and Endangered Medicinal Plant", *PLoS ONE*, 2016.

Publication

<1%

43

tigerprints.clemson.edu

Internet Source

<1%

44

BEKTAŞ, Ersan and SÖKMEN, Atalay. "In vitro seed germination, plantlet growth, tuberization, and synthetic seed production of *Serapias vomeracea* (Burm.f.) Briq.", TÜBİTAK, 2016.

Publication

<1%

45

Kaur, Saranjeet, and Kamlesh Kumar Bhutani. "Asymbiotic seed germination and

<1%

multiplication of an endangered orchid –
Paphiopedilum venustum (Wall. ex Sims.)",
Acta Societatis Botanicorum Poloniae, 2016.

Publication

46

Teixeira da Silva, Jaime A., Tikam Singh Rana,
Diganta Narzary, Nidhi Verma, Deodas
Tarachand Meshram, and Shirish A. Ranade.
"Pomegranate biology and biotechnology: A
review", *Scientia Horticulturae*, 2013.

Publication

47

digital.csic.es

Internet Source

48

Khamchatra, N., K.W. Dixon, S. Tantiwivat, and
J. Piapukiew. "Symbiotic seed germination of
an endangered epiphytic slipper orchid,
Paphiopedilum villosum (Lindl.) Stein. from
Thailand", *South African Journal of Botany*,
2016.

Publication

49

"Date Palm Biotechnology Protocols Volume I",
Springer Nature, 2017

Publication

50

Dace Grauda, Lita Lapiņa, Biruta Jansone,
Aldis Jansons, Isaak Rashal. "Recovering
Genetic Resources of Some Legume Species
of Latvian Origin by Plant Tissue Culture",
Proceedings of the Latvian Academy of

<1%

<1%

<1%

<1%

<1%

Sciences. Section B. Natural, Exact, and Applied Sciences, 2013

Publication

51

Kunlin Wu, Songjun Zeng, Danni Lin, Jaime A. Teixeira da Silva, Zhaoyang Bu, Jianxia Zhang, Jun Duan. "In Vitro Propagation and Reintroduction of the Endangered *Renanthera imschootiana* Rolfe", PLoS ONE, 2014

<1%

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

The Effect of Organic Nutrient and Growth Regulators on Seed Germination, Embryo and Shoots Development of *Dendrobium antennatum* Lindl. Orchid by In Vitro

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
