

Volume : 3 | Issue : 11 | Nov 2014 | ₹ 250/-

PEER REVIEWED & REFERRED INTERNATIONAL JOURNAL
ISSN - 2250-1991 | IMPACT FACTOR - 1.6714



PARIPEX - INDIAN JOURNAL OF RESEARCH

Journal DOI : 10.15373/22501991

Listed & Indexed in

International ISSN Directory, Paris

Journal for All Subjects

www.paripeex.in



Paripex - Indian Journal of Research

Journal for All Subjects

Editor-In-Chief

Dr A Kumar

Director, College Development Council (CDC)
Director, Internal Quality Assurance Cell (IQAC)
Professor in Management,
Department of Business Administration, Faculty of Management,
Bhavnagar University,

Editorial Advisory Board

Dr.(Prof) Vijay Kumar Soni
Rajasthan

Dr. Kaushal Kishore
Chhattisgarh

Dr. R. Ramachandran
Tamilnadu

Dr. Ashok S. Pawar
Aurngabad

Dr. C. D. Balaji
Chennai

Dr. Ravikant Swami
Himachal Pradesh

Dr. A.R. Saravankumar
Tamilnadu

Prof. Jothi M
Kattankulathur

Dr. Amit Bandyopadhyay
Calcutta

Prof. C M Maran
Tamilnadu

Dr. V. Kumaravel
Namakkal

Dr. Anukrati Sharma
Jaipur

Dr. R. Ganapathi
Alagappa University

Dr. K. Prabhakar,
Chennai

Advertisement Details

Position	B/W (Single Color)	Fore Color
Full Inside Cover	₹ 6250	₹ 12500
Full Page (Inside)	₹ 5000	-

Subscription Details

Period	Rate	Discount	Amount Payable
One Year (12 Issues)	₹ 2400	Nil	₹ 2400
Two Year (24 issues)	₹ 4800	₹ 200	₹ 4600
Three Year (36 issues)	₹ 7200	₹ 300	₹ 6900
Five Year (60 issues)	₹ 12000	₹ 600	₹ 11400

You can download the Advertisement / Subscription Form from website www.paripex.in. You will require to print the form. Please fill the form completely and send it to the **Editor, Paripex, Indian Journal of Research** along with the payment in the form of Demand Draft/Cheque at Par drawn in favour of **Paripex** payable at Ahmedabad.

- Thoughts, language vision and example in published research paper are entirely of author of research paper. It is not necessary that both editor and editorial board are satisfied by the research paper. The responsibility of the matter of research paper/article is entirely of author.
- Editing of the Paripex is processed without any remittance. The selection and publication is done after recommendations of atleast two subject expert referees.
- In any condition if any National/International University denies accepting the research paper published in PIJR, then it is not the responsibility of Editor, Publisher and Management.
- Only the first author is entitle to receive the copies of all co-authors
- Before re-use of published research paper in any manner, it is compulsory to take written permission from the Editor-PIJR, unless it will be assumed as disobedience of copyright rules.
- All the legal undertaking related to Paripex is subject to Ahmedabad Jurisdiction.
- The research journal will be send by normal post. If the journal is not received by the author of research papers then it will not be the responsibility of the Editor and publisher. The amount for registered post should be borne by author of the research paper in case of second copy of the journal.

Editor,

Paripex - Indian Journal of Research

8-A, Banans, Opp. SLU Girls College, New Congres Bhavan, Paldi,
Ahmedabad-380006, Gujarat, INDIA

Contact.: +91-8866003636 E-mail : editor@paripex.in

INDEX

Sr. No.	Title	Page No
1	Mineral composition of urinary stones - Quantitative analysis by FTIR Spectroscopy - Mujeebu Rahiman M, Vidya Bernhardt G, Janita D'Souza R T, Muhammed Manzoor. AP, Vipin C, Nischith D'souza	1-3
2	Struggling with pain and disability: a qualitative study of the experiences of pregnant women living with pelvic girdle pain - Mrs M V Smitha, Dr Molly Babu,	4-6
3	Laws of Thought - Alope Bhunia	7-8
4	LEARNING DISABILITY THE NEGLECTED CLASSROOM REALITY - Dr. Manoj Kumar Choudhary, Dr. Sandeep Kumar	9-12
5	The Effect of Gamma Radiation on Optical Properties and Hall Effect of thin ZnS Film - Shatha S.M. Alazzawi	13-16
6	Quality of Work Life : A Review - MR.SANTOSH L.PATIL, DR.S.Y.SWADI	17-19
7	A Study on Job Involvement With Special Reference to Marketing Professionals in Coimbatore - R.Anitha, Dr.K.Muralidharan	20-22
8	Sangeet Sadhna/Abhyas - Prof. Alaknanda palnitkar	23-26
9	Effect of SAQ Training and Plyometric Training on Selected Motor Fitness and Physiological Variables Among Junior Basketball Players - RAKESH PRASAD, Dr. P. K. SUBRAMAINIAM	27-28
10	Effect of Progersterone Dose and Intravaginal Sponge Diameter on Reproductive Performance of Induced-Estrous Local Ewes in West Java - Soeparna Soeparna, Ranga Setiawan, Siti Darodjah, Tita Damayanti Lestari	29-31
11	Combined Effect of Yoga and Endurance Exercises on Lean Body Mass Among Players - C.KARTHIKEYAN, Dr A. UTHIRAPATHY	32-33
12	Legal Protection of Child Labour in India: Problems and Perspective - Dr. Anand Pawar, Rakesh Vaidya	34-35
13	A Gender-Wise Analysis on Metacognition And Learning Styles on Problem Solving Skill of B.Ed Trainees - T.Yogaraj, Dr. R. Selvaraju	36-37
14	Prevalence of Viral infections in pregnant females in Northern India - Dr Neha Gupta	38-39
15	Banking Practices among Petty Shop Owners with Special Reference In Coimbatore City - J.Thiravia Mary Gloria, Dr.P.Santhi	40-43
16	HR in Krishna Grameena Bank in Gulbarga - Prof. HEMANTHKUMAR S. KOLHAPUR, Dr. GIRAM S. J.	44-45
17	Domestic Violence and Subjective Well Being Among The Women Professionals And House Wives - K.Selvi, Dr.K.Muralidharan	46-48
18	Spread of Buddhism in the West - Ravi chanra Reddy. Bh	49-51
19	Comparison of Speed and Explosive Power Among Different Levels of Handball Players - Dr. B. Chittibabu	52-53
20	Effect of High Intensity Interval Training on Aerobic Power and Anaerobic Power of Male Handball Players - Dr. B. Chittibabu	54-55

Sr. No.	Title	Page No
21	An Empirical Evaluation Of Foreign Exchange Market Efficiency Of Chfjpy Currency Pair Dr. P.Krishnaveni, Dr.S.Varadaraj, Mrs. P. Karthika	56-58
22	Human Right Education for Developing the Personality of the Student - DR. R.N DINESH	59-61
23	Integrated Child Development Services (Icids): a Stepping Stone for Universal Elementary Education - Dr.S.S.Swain, Sri Binod Bihari Das	62-63



Effect of Progesterone Dose and Intravaginal Sponge Diameter on Reproductive Performance of Induced-Estrous Local Ewes in West Java

Soeparna Soeparna

Laboratory of Animal Reproduction, Faculty of Animal Husbandry, Universitas Padjadjaran Jl. Raya Bandung Sumedang KM 21. Sumedang West java. Indonesia

Rangga Setiawan

Laboratory of Animal Reproduction, Faculty of Animal Husbandry, Universitas Padjadjaran Jl. Raya Bandung Sumedang KM 21. Sumedang West java. Indonesia

Siti Darodjah

Laboratory of Animal Reproduction, Faculty of Animal Husbandry, Universitas Padjadjaran Jl. Raya Bandung Sumedang KM 21. Sumedang West java. Indonesia

Tita Damayanti Lestari

Laboratory of Animal Reproduction, Faculty of Animal Husbandry, Universitas Padjadjaran Jl. Raya Bandung Sumedang KM 21. Sumedang West java. Indonesia

ABSTRACT

The use of estrous induction technique is able to synchronize estrous of each ewe in order to increase the reproductive efficiency of livestock. However, the success rate of this technique is very dependent on the progesterone dose and its absorbance in the body. Therefore, this study was aimed to figure out reproductive performance of local sheep induced by the combination of progesterone doses and vaginal sponge diameters. 48 local ewes were used in this study and were grouped into 12 treatment groups. The treatments consist of the combination of progesterone doses (20, 30, and 40 mg) with vaginal sponge diameters (1, 2, 3, and 4 cm). The method used a randomized block design and the measured variables were: estrous response, pregnancy rate, litter size, and birth weight. The result showed that the highest estrous responses of the ewes were from the combination of sponge diameter 4 cm in all variation of progesterone doses. Meanwhile, the best pregnancy rate was found in the group of 20 and 30 mg progesterone dose in the diameter of 2 cm. However, this study did not find any difference among groups for litter size and birth weight. In conclusion, estrous induction using progesterone is effectively to synchronize estrous phase and pregnancy period of ewes, although it does not effect on litter size and birth weight of lamb.

KEYWORDS

Local ewes, Estrous Synchronization, Progesterone, Sponge diameter

INTRODUCTION

Estrous induction is an effective and efficient technique to increase pregnancy rate in ewes. Through this technique, pregnancy and lambing time, even age of the lamb could be uniformed, therefore the farmer can easily plan the management and production. Some researchers stated that pregnancy rate after estrous induction reached 75-80% (Koyuncu and Alticekic 2010; Santos et al., 2011; Zonturlu et al., 2011).

In principle, estrous induction using hormones is to manipulate estrous phase either by prolonging or shortening luteal phase (Iida et al., 2004; Wildeus, 1999). In the present study, progesterone was used to induce estrous by prolonging luteal phase. Based on several studies (Robinson, 1976) reported that the success of using progesterone for estrous induction yielded > 90 % within 48 hours after sponge removal. However, the report also stated that the dose of progesterone must be fit to body condition of the animal. A lower dose of progesterone tends to an un-uniformity of estrous. Another study also reported that level of progesterone has a positive correlation with estrous occurrences (Allison and Robinson, 1970). They stated that it might be due to the effect of body weight physiologically responses the induction.

Indonesian local sheep has body weight ranged of 25-30 kg. This body weight is lower if compared to sheep in subtropical area which is has body weight around 40 – 45 kg (Simone-

tti et al., 1999). Furthermore, a study from (Rubianes et al., 1999) stated that the use of progesterone dose in sub-tropical sheep approximately 60 mg. This implies that an appropriate dose of progesterone is an important factor in the effectiveness of estrous induction and it is differ from animal to animal dependent on the body condition and their environmental condition. Therefore, this study was to figure out a dose of progesterone in estrus induction which is fit to a local sheep and variation of vaginal sponge diameter was included in this study to find out the best absorbance of hormone during the induction.

MATERY AND METHOD

In this study, 48 non-pregnant ewes were used and grouped into three different groups of progesterone dose, i.e. 20, 30, and 40 mg. Each group was then divided into four different of vaginal sponge diameter groups (1,2,3, and 4 cm of diameter), so the experimental design was a 3 x 4 factorial to study the effect of the treatments on estrous response, pregnancy rate, litter size, and birth weight. The sponge contained different progesterone doses were inserted into vagina and kept for 14 days. After 14 days of treatment, the sponges were removed. The method of this study used a randomized block design and was followed by Fisher's LSD (Least Significance Different). The measured variables were estrous response, pregnancy rate, litter size, and birth weight. Estrous respond was monitored every day and measured by using Draminsky's

estrous detector for sheep. Estrous occurrence also defined based on cervical opening and this is indicator for artificial insemination as well (Kershaw et al., 2005). Pregnant ewes were defined using Draminsky's pregnancy detector.

RESULT AND DISCUSSION

Estrous Response is showed in Figure 1.




Figure 1. Estrous response on the progesterone doses and vaginal sponge diameters

According to Figure 1. Each group showed estrous signs after the removal of vaginal sponge. The signs is based on vulva characteristics, vaginal mucus measured by estrous detector, and cervix dilatation as an indicator for insemination according to (Kershaw et al., 2005). The result showed that estrous begins from day-1 to day-3 after the removal of vaginal sponge. The percentage of estrous at day-1 is relatively low for all groups, however, this finding is higher than that finding of (Ciptadi, 2010) who found no estrous animal at day-1 after the induction. The reason for this result might be due to the level of progesterone in the blood circulation is still relatively high. This condition leads to the negative feedback mechanism inhibiting Follicle Stimulating Hormone (FSH). The further effect of this inhibition causes retardation of follicle growth and estrogen production, therefore the female does not show estrous signs (Robinson and Kendrick, 1992).

Estrous response is increased at day-2 and day-3 after removal, in the ranged of 50-70% and 25-75%, respectively. This results are lower compared to the result of (Zonturlu et al., 2011) who reported that the responses at day-2 is 81-92%. (Luther et al., 2007) also reported that in general, estrous response using vaginal sponge is 75%. A hundred percentages of estrous response is reported in another finding, but the estrous induction used PGF_{2α} and 400 IU of PMSG after progesterone treatment (Ataman et al., 2006). These findings indicate that the level of progesterone in the blood stream has been decreased which furthermore leads to the growth of follicle caused by the release of FSH. Along with follicle growth, estrogen secretion increase due to the proliferation of granulose cell causing the female estrous (Hafez, 1993).

Pregnancy rate is showed in Figure 2.




Figure 2. Pregnancy rate on progesterone levels and vaginal sponge diameters

The pregnancy rate of the ewes in all groups was recorded to compare the effect of the treatments (Fig. 2). Fig. 2 shows the combination of progesterone 20 mg and sponge diameter 2 cm; and progesterone 30 mg and sponge diameter 3 cm give 100% of pregnancy rate. These indicate that both of progesterone levels of 20 and 30 mg are optimum doses to stimulate hormonal mechanism in relation to estrous induction and ovulation. Combination of sponge diameter 2 cm and 3 cm are ideal combination as the size is fit to the vaginal space capacity of local sheep, therefore these sponges give best hormone absorption.

The lowest rate of pregnancy belongs to the group with dose 40 mg of progesterone. This result is slightly similar with the previous research used 60 mg of progesterone yielded 45% of pregnancy rate (Yadi et al., 2011). A similar result also found in (Awel and Eshetu, 2009) who used a high level of progesterone with a low pregnancy rate. The reason to address these facts that the use of high level of progesterone gives side effect on the semen transportation in the female reproductive tract (Dogan and Nur, 2006), and can caused embryonic death (Cam and Kuran, 2004). Furthermore, (Dixon et al., 2006) reported that a low dose of progesterone generate good ovulatory follicle, and pregnancy rate has negative correlation with follicle growth and progesterone dose. However, biological architecture underlying this mechanism needs to be revealed to give a better understanding.

Litter size.

Table 1. litter size of induced-estrous local sheep

Progesterone level (mg)	Diameter of vaginal sponge			
	1 cm	2 cm	3 cm	4 cm
20	1,0 ± 0,0	1.7 ± 0,47	1,0 ± 0,0	1,0 ± 0,0
30	2,0 ± 0,0	1,0 ± 0,0	1.7 ± 0,9	1.3 ± 0,47
40	1,0 ± 0,0	2,0 ± 0,0	1,0 ± 0,0	1.7 ± 0,47

Overall, out of 23 estrous-induced ewes giving a birth, 15 ewes have singletons, seven ewes have double, and one has triple. However, litter size in this study did not differ significantly among all groups. This result is in accordance with those reported by (Bitaraf et al., 2007) that estrous induction using CIDR, vaginal sponge, and cloprostenol do not significantly effect on litter size.

Birth weight

Table 2. Birth weight of induced-estrous local sheep

Progesterone level (mg)	Diameter of vaginal sponge			
	1 cm	2 cm	3 cm	4 cm
20	2.8 ± 0.25	2.4 ± 0,12	3,0 ± 0,00	2.9±0,00
30	1.8 ± 0,00	2.5 ± 0,37	2.5 ± 0, 25	2.5±0,70
40	2.8 ± 0,00	2.8 ± 0,00	NA	2.2 ± 0,27

In general, birth weight of this study ranged 1.8 to 3 kg (Table 2.). The lowest of birth weight belongs to the group of progesterone 30 mg combined with vaginal sponge 1 cm, and the highest of litter weight belongs to the group of progesterone 20 mg combined with vaginal sponge 3 cm, however, there were no significant differences among all groups. This result is in accordance with those resulted by (Anilkumar et al., 2010) that estrous induction does not effect on birth weight of lamb.

Table 3. Birth weight based on type of birth

	Total	Birth weight (kgs)
n of ewes	23	
Type of birth		
Single	65% (15/23)	2,6 ± 0,43
Double	30% (7/23)	2,2 ± 0,34
Triplet	4,1% (1/23)	2,17 ± 0,43

Lambing weight based on type of birth (single, twins, and triplets) of estrous-induced local sheep is listed in Table 3. Twins and triplets have lighter birth weights (2.2 ± 0.34 and 2.17 ± 0.43 kg, respectively) compared to single (2.6 ± 0.43 kg). This finding is lower compared to the finding by (Yilmaz and Altin, 2011) in local sheep crossed. This implies that litter size has a negative correlation with birth weight of lamb.

CONCLUSION

1. Estrous synchronization using progesterone in vagina sponge is able to synchronize estrous at the same time in local ewes
2. Vagina sponge with 4 cm diameter give the best estrous response in all doses of progesterone combination
3. Progesterone doses of 20 and 30 mg give the best result in pregnancy rate
4. Combination of progesterone dose and sponge diameter does not affect to the litter size and birth weight of local sheep

ACKNOWLEDGMENT

The research was accomplished within the research project of Universitas Padjadjaran financed by Directorate of Higher Education Ministry of National Education and Culture Republic of Indonesia.

REFERENCES

- Allison, A. J., and T. J. Robinson. 1970. The effect of dose level of intravaginal progestagen on sperm transport, fertilization and lambing in the cyclic merino ewe. *J. Reprod. Fert.* 22: 515-531. | Anikumar, R., C. Chandrahasan, M. Iyue, M. Selvaraju, and A. Palanisamy. 2010. Reproductive and economic efficiency in nilagiri and sandyno ewes treated with pmng. *Livestock Research for Rural Development* 22. | Ataman, M. B., M. Aköz, and O. Akman. 2006. Induction of synchronized oestrus in akkaraman cross-bred ewes during breeding and anestrus season: The use of short-term and long-term progesterone treatments. *Revue Méd Vét* 157: 257-260. | Awel, H., and L. Eshetu. 2009. Estrus synchronization in sheep with synthetic progestagens. *Trop Anim Health Prod* 41: 1521-1524. | Bitaraf, A., M. J. Zamiri, M. Kafi, and J. Izadifard. 2007. Efficacy of cidr, fluogestone acetate sponges and cloprostenol for estrous synchronization of nadooshani goats during the breeding season Iranian Journal of Veterinary Research 8: 218-224. | Cam, M. A., and M. Kuran. 2004. Effects of a single injection of hcg or gnrh on day 12 post mating on fetal growth and reproductive performance of sheep. *Animal Reproduction Science* 80: 81-90. | Ciptadi, G. 2010. Pemanfaatan larutan iodine povidon sebagai hormon stimulan gertak berahi kambing secara alamiah. *J. Ternak Tropika* 11, No. 2: 74-81. | Dixon, A. B., M. Knights, J. L. Pate, P. E. Lewis, and E. K. Inskip. 2006. Reproductive performance of ewes after 5-day treatment with intravaginal inserts containing progesterone in combination with injection of prostaglandin f2alpha. *Reprod Domest Anim* 41: 142-148. | Dogan, I., and Z. Nur. 2006. Different estrus induction methods during the non-breeding season in kivircik ewes. *Veterinary Medicine Czechoslovakia* 51: 133-138. | Hafez, E. S. E. 1993. *Reproduction in farm animals*. 6th Ed. Lea and Febiger. Philadelphia. | Iida, K., N. Kobayashi, H. Kohno, A. Miyamoto, and Y. Fukui. 2004. A comparative study of induction of estrus and ovulation by three different intravaginal devices in ewes during the non-breeding season. *J. Reprod. Dev* 50: 63-69. | Kershaw, C. M. et al. 2005. The anatomy of the sheep cervix and its influence on the transcervical passage of an inseminating pipette into the uterine lumen. *Theriogenology* 64: 1225-1235. | Koyuncu, M., and S. O. Alticekic. 2010. Effects of progestagen and pmng on estrous synchronization and fertility in kivircik ewes during natural breeding season Asian-Aust. *J. Anim. Sci.* Vol. 23, No. 3: 308 - 311. | Luther, J. S. et al. 2007. The effect of gnhrh, ecg and progestjn type on estrous synchronization on following laparoscopic ai in ewes. *Small Rumin Res* 72: 227-231. | Robinson, J. E., and K. M. Kendrick. 1992. Inhibition of luteinizing hormone secretion in the ewe by progesterone. Associated changes in the release of gamma-aminobutyric acid and noradrenaline in the preoptic area as measured by intracranial microdialysis. *J. Neuroendocrinol* 4: 231-236. | Robinson, T. J. 1976. Controlled breeding of sheep and goats. *Sheep breeders' cong.* In: Tomes, G.L., d.E. Robertson, r.J. Lightfoot. Institute of Technology, W. Australia: 423-437. | Rubianes, E., R. Ungerfeld, and T. De Castro. 1999. Inducción y sincronización de celo en ovejas y cabras (induction and synchronization of estrus in ewe and goats). III Simposio Internacional de Reproducción Animal, Montevideo, Uruguay: 109-131. | Santos, G. M. G. et al. 2011. Reproductive performance of ewes treated with an estrus induction/synchronization protocol during the spring season. *Anim. Reprod.* 8: 3-8. | Simonetti, L., G. Ramos, and J. C. Gardon. 1999. Estrus presentation and distribution in ewes treated with intravaginal sponges impregnated with medroxyprogesterone acetate (map) in combination with pregnant mare serum gonadotropin (pmng). *Braz. J. Vet. Res. Anim. Sci.* 36: | Wildeus, S. 1999. Current concepts in synchronization of estrus: Sheep and goats. *Proceedings of the American Society of Animal Science.* | Yadi, J., M. F. Mehran Farhoudi Moghaddam, S. Khalajzadeh, and A. A. Solati. 2011. Comparison of estrus synchronization by pgf2B, cidr and sponge with pmng in kalk uhi ewes on early anestrus season. *International Conference on Asia Agriculture and Animal* 13: 61-65. | Yilmaz, M., and T. Altin. 2011. Growth characteristics in lambs of estrus synchronized ewes in grower conditions. *Turk. J. Vet. Anim. Sci.* 35(6): 412-429. | Zonturlu, A. K., N. Özyurtlu, and C. Kacar. 2011. Effect of different doses pmng on estrus synchronization and fertility in awassi ewes synchronized with progesterone during the transition period *Kafkas Univ Vet Fak Derg* 17 (1): 125-129. |