


- Editorial • *Open access*  
Editorial: Current Issues on Future Researches and Applications of Natural Product Medicines  
Elfahmi  
Pages 1-2  
 [Download PDF](#)   [Article preview](#) 
- Research article • *Open access*  
Development of Palladium-catalyzed Cycloalkenylation and its Application to Bioactive Natural Product Synthesis  
Masahiro Toyora  
Pages 3-12  
 [Download PDF](#)   [Article preview](#) 
- Research article • *Open access*  
*Boesenbergia Pandurata* Roxb., An Indonesian Medicinal Plant: Phytochemistry, Biological Activity, Plant Biotechnology  
Agus Chahyadi, Rika Hartati, Komar Ruslan Wirasutisna, Elfahmi  
Pages 13-37  
 [Download PDF](#)   [Article preview](#) 
- Research article • *Open access*  
Biotransformation of Protoberberine Alkaloids by the Endophytic Fungus *Coelomyces* AFKR-3 Isolated from Yellow Moonseed Plant (*Archangelisia Flava* (L.) Merr.)  
Andria Augusta, Dewi Wulansari, Praptiwi, Arif Nurkanto, Ahmad Fathoni  
Pages 38-43  
 [Download PDF](#)   [Article preview](#) 
- Research article • *Open access*  
Recent Study of Turmeric in Combination with Garlic as Antidiabetic Agent  
Elin Yulinah Sukandar, Primal Sudjana, I. Ketut Adnyana, Ame Suciati Setiawan, Umi Yuniarni  
Pages 44-56  
 [Download PDF](#)   [Article preview](#) 

- Research article • *Open access*  
Changes in Secondary Metabolite Contents Following Crude Drug Preparation  
Sukrasno  
Pages 57-62  
 [Download PDF](#) [Article preview](#) 
- Research article • *Open access*  
Molecular Docking Studies of Chemical Constituents of *Tinospora cordifolia* on  
Glycogen Phosphorylase  
Rina Herowati, Gunawan Pamudji Widodo  
Pages 63-68  
 [Download PDF](#) [Article preview](#) 
- Research article • *Open access*  
Assessing Heterologous Expression of Hyoscyamine 6 $\beta$ -Hydroxylase – A  
Feasibility Study  
Nils J.H. Aversch, O. Kayser  
Pages 69-78  
 [Download PDF](#) [Article preview](#) 
- Research article • *Open access*  
Cytotoxic Activities of Fractions and Two Isolated Compounds from Sirih  
Merah (Indonesian red betel), *Piper Crocatum* Ruiz & Pav.  
Emrizal, Armon Fernando, Riska Yuliandari, Kamal Rullah, ... Dayar Arbain  
Pages 79-84  
 [Download PDF](#) [Article preview](#) 
- Research article • *Open access*  
Effect of Terbinafin and DMSO on The Gene Expression Level of Squalene  
Synthase (Sqs) and Amorpha-4,11-Diene Synthase (Ads) in *Artemisia Annuua* L  
Elfahmi, Luthfia Nurhalim, Tati Kristanti, Agus Chahyadi, Sony Suhandono  
Pages 85-91  
 [Download PDF](#) [Article preview](#) 
- Research article • *Open access*

## Encapsulation of Risperidone into Chitosan-based Nanocarrier via Ionic Binding Interaction

Diky Mudhakhir, Caroline Wibisono, Heni Rachmawati

Pages 92-100

 [Download PDF](#) [Article preview](#) 

Research article • *Open access*

## In vivo Antimalarial Activity of *Andrographis Paniculata* Tablets

Aty Widyawaruyanti, Muhammad Asrory, Wiwied Ekasari, Dwi Setiawan, ... Achmad Fuad Hafid

Pages 101-104

 [Download PDF](#) [Article preview](#) 

Research article • *Open access*

## Development of Immunonutrient from Pumpkin (*Cucurbita Moschata* Duchense Ex. Lamk.) Seed

Maria Immaculata Iwo, Muhamad Insanu, Caryn Anne Santhana Dass

Pages 105-111

 [Download PDF](#) [Article preview](#) 

Research article • *Open access*

## Analysis of Secondary Metabolite Production in Somatic Embryo of Pasak Bumi (*Eurycoma Longifolia* Jack.)

Iriawati, Andira Rahmawati, Rizkita R. Esyanti

Pages 112-118

 [Download PDF](#) [Article preview](#) 

Research article • *Open access*

## Influence of $\beta$ -cyclodextrin on Cefixime Stability in Liquid Suspension Dosage Form

Jessie Sofia Pamudji, Rachmat Mauludin, Nurhabibah

Pages 119-127

 [Download PDF](#) [Article preview](#) 

Research article • *Open access*

## Effect of Steeped Black Tea (*Camellia sinensis* (L.) var. assamica) on Immunoglobulin Titer and Lymphocyte Proliferation in Responses to

☰ Outline

Download full text in PDFDownload

Export ▾

Search ScienceDirect

Advanced

## Outline

Abstract

Keywords

References



Procedia Chemistry

Volume 13, 2014, Pages 101-104

open access



# In vivo Antimalarial Activity of *Andrographis Paniculata* Tablets ☆

Aty Widyawaruyanti <sup>a, b</sup> ✉, Muhammad Asrory <sup>a, b</sup>, Wiwied Ekasari <sup>a</sup>, Dwi Setiawan <sup>a</sup>, Achmad Radjaram <sup>a</sup>, Lidya Tumewu <sup>b</sup>, Achmad Fuad Hafid <sup>a, b</sup>

☒ Show more

<https://doi.org/10.1016/j.proche.2014.12.012>

Get rights and content

Under a Creative Commons license

## Abstract

The formulation of three phytopharmaceutical products of *Andrographispaniculata* fractions (AP fraction A and B) containing diterpene lactones as an active substance were developed and their antimalarial activities against *Plasmodium berghei* has been examined. *In vivo* antimalarial assay on *P. berghei* infected mice was carried out by oral administration, twice a day for four consecutive days of the AP fractions product, which were Tablet I: wet granulated formula of AP fraction A; Tablet II: wet granulated formula of AP fraction B; Tablet III: solid dispersion formula of AP fraction B.. The results revealed that three phytopharmaceutical products of *A.paniculata* were inhibited parasite's growth with inhibition range of 70.15% to 80.35%. There was no significant difference of antimalarial activities

between Tablet II and III, meanwhile there was significant difference among Tablet I with Tablet II and Tablet III. It was concluded that antimalarial activity depending on raw material form of *A. paniculata* active substance.

### Keywords

*Andrographis paniculata* Ness; tablets; *in vivo* antimalarial activity; *Plasmodium berghei*; inhibition of parasite's growth



International Seminar on Natural Product Medicines, ISNPM 2012

## In vivo Antimalarial Activity of *Andrographis paniculata* Tablets

Aty Widyawaruyanti<sup>a,b\*</sup>, Muhammad Asrory<sup>a,b</sup>, Wiwied Ekasari<sup>a</sup>, Dwi Setiawan<sup>a</sup>,  
Achmad Radjaram<sup>a</sup>, Lidya Tumewu<sup>b</sup>, Achmad Fuad Hafid<sup>a,b</sup>

<sup>a</sup>Faculty of Pharmacy, Universitas Airlangga, Surabaya 60286, Indonesia

<sup>b</sup>Institute of Tropical Disease, Universitas Airlangga, Surabaya 60115, Indonesia

### Abstract

The formulation of three phytopharmaceutical products of *Andrographispaniculata* fractions (AP fraction A and B) containing diterpene lactones as an active substance were developed and their antimalarial activities against *Plasmodium berghei* has been examined. *In vivo* antimalarial assay on *P. berghei* infected mice was carried out by oral administration, twice a day for four consecutive days of the AP fractions product, which were Tablet I : wet granulated formula of AP fraction A; Tablet II : wet granulated formula of AP fraction B; Tablet III : solid dispersion formula of AP fraction B.. The results revealed that three phytopharmaceutical products of *A.paniculata* were inhibited parasite's growth with inhibition range of 70.15% to 80.35%. There was no significant difference of antimalarial activities between Tablet II and III, meanwhile there was significant difference among Tablet I with Tablet II and Tablet III. It was concluded that antimalarial activity depending on raw material form of *A. paniculata* active substance.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

Peer-review under responsibility of the School of Pharmacy, Bandung Institute of Technology

**Keywords:** *Andrographis paniculata* Ness; tablets; *in vivo* antimalarial activity; *Plasmodium berghei*; inhibition of parasite's growth

### 1. Introduction

Malaria was one of the oldest recorded diseases in the world. Each year from 300 to 500 million new cases were diagnosed and approximately 1.5 million people died because the disease; majority were children<sup>1</sup>. The re-emerging of malaria in many parts of the world was due to the rapid increasing of resistance to most of the available antimalarial drugs, as well as resistance of vectors to insecticides<sup>2,3</sup>. Drug resistant for strains of *Plasmodium falciparum* has been found in many endemic areas of the world and many of conventional anti-malarial drugs have

\* Corresponding author. Tel.: +62-31-503-3710; fax: +62-31-502-0514  
E-mail address: [aty\\_ww@yahoo.com](mailto:aty_ww@yahoo.com)

been associated with treatment failure. Furthermore, the difficulty of creating efficient vaccines and also adverse side-effects of the existing antimalarial drugs highlight the urgent need for novel and well-tolerated antimalarial drugs<sup>2</sup> for both prophylaxis and treatment of malaria.

*Andrographis paniculata* Nees which known as sambiloto was traditionally used for antimalaria in Indonesia. The previous research had shown that *A. paniculata* extract have antimalarial activity both in vitro and in vivo<sup>4,5</sup>. Methanol extract of *A. paniculata* exhibited antimalarial activity against *P. falciparum* with IC<sub>50</sub> value of 7.2 µg/ml. Furthermore, the results of in vivo toxicity assay indicating no toxicity associated with the use of this extract in mice system<sup>5</sup>. Andrographolide was the main content of sambiloto which reported have antimalarial activity. Andrographolide has IC<sub>50</sub> value of 9.1 µM<sup>6</sup>. Andrographolide was considered to be active compound and has good antimalarial activity based on its IC<sub>50</sub> value which was less than 20 µM<sup>7</sup>. Our previous research also found that andrographolide shown antiplasmodial activity as blood schizontocidal activity and also gametocytocidal with IC<sub>50</sub> value of 12.16 and 3.61 µg/ml, respectively<sup>8</sup>.

Regarding to the potential antimalarial activity of *A. paniculata*, Widyawaruyanti et al<sup>9</sup> was developed phytopharmaceutical products by conducted formulation study of two fractions from ethanol extract of this plant which were AP fraction A and B. AP fraction A was obtained from fractionated ethanol extract using ethyl acetate, while further purification of AP fraction A resulted as AP fraction B. Although AP fractions have different physicochemical properties, both of them were contain andrographolide (diterpene lactone compound) as an active substance. The formulation study of AP fractions A and B was performed based on two methods which were wet granulation and solid dispersion, therefore three phytopharmaceutical products were obtained which were Tablet I: wet granulated formula of AP fraction A; Tablet II: wet granulated formula of AP fraction B; and Tablet III: solid dispersion formula of AP fraction B.

The aim of this study was to observe in vivo antimalarial activities of Tablet I, II and III based on The Peter's test (The 4-days suppressive test) by oral administration of products at a dose of 12.55 mg andrographolide/kg mice body weight, twice a day for four consecutive days..

## 2. Experiments

### 2.1. Material

*A. paniculata* dry herbs powder was obtained from pharmaceutical company PT. Kimia Farma Tbk, Bandung, Indonesia. *A. paniculata* was extracted by maceration method using ethanol 96% as a solvent. Ethanol extract of *A. paniculata* was further fractionated by liquid-liquid fractionation method using ethyl acetate and water to obtain AP fraction A. Dried AP fraction A was further purified so that AP fraction B was obtained. AP fraction A was dark green and sticky, meanwhile AP fraction B was greenish amorphous powder.

Tablet I contained of 75 mg of AP fraction A per 400 mg tablet formulated by wet granulation method. Tablet II contained of 15 mg of AP fraction B per 300 mg tablet formulated by wet granulation method. Tablet III contained of 60 mg of AP fraction B per 150 mg tablet formulated by solid dispersion method.

The chloroquine sensitive *Plasmodium berghei* ANKA strain was obtained from Institute of Biomolecular Eijkmann, Jakarta and maintained on mice at Malaria Laboratorium, Faculty of Pharmacy, Universitas Airlangga. The inoculum of *P. berghei* parasitized erythrocytes was prepared by determining the percentage parasitemia of mice donor and diluting the blood with alceivers solutions in proportion indicated by the determination. The mice were infected intraperitoneally with 200 µL of 5% parasite blood stock from frozen deposits. Once the percent of parasitemia on this donor mice reached 20%, the mice blood taken intracardially and diluted with PBS or alceivers solutions (1:3) up to 5% parasitemia. The test mice has been infected by 200 µL of this diluted parasitized blood that contain 5% parasite.

Male albino Swiss mice Balb-C strain (20-30 g weight) were obtained from Animal Laboratory, Faculty of Pharmacy, Universitas Airlangga. They were maintained on standard animal pellets and water ad libitum. Permission and approval for animal studies were obtained from Faculty of Veterinary Medicine, Universitas Airlangga.

## 2.2. Methods

### 2.2.1. Preparation of samples

Tablet I, Tablet II and Tablet III of which were equal to 12.55mg andrographolide/kg mice body weight were powdered and suspended in 0.5% CMCNa solution. Sample solutions were administered per oral (1 ml/mouse) twice a day.

### 2.2.2. In vivo antimalarial assay

Antimalarial activity of Tablet I, II and III were evaluated using the Peter's test methods (The 4-days suppressive test). Each mouse was inoculated intraperitoneally on the first day (day 0) with 0.2 ml of infected blood containing *P. berghei* parasitized erythrocytes (5% parasitemia). The animals were divided into four groups of five mice each and administered orally with suspension of Tablet I, Tablet II and Tablet III containing andrographolide 12.55 mg/kg body weight twice a day and an equivalent volume of CMC-Na solutions (as negative control) at one day after inoculation, for four consecutive days (day 0 to day 3). On the fifth day (day 4), thin films were made from the tail blood of each mouse and the parasitemia level was determined by counting the number of parasitized erythrocytes out of 1000 erythrocytes in random fields of the microscope. Average percentage parasite's inhibition was calculated as following:

$$100\% - \left( \frac{Xe}{Xk} \times 100\% \right)$$

Xe: % parasitemia growth of experimental group

Xk: % parasitemia growth of negative control

## 3. Results and Discussion

Tablet I, II and III produced inhibition effect at a dose employed in this study. The parasite's inhibition were 70.35%, 78.16% and 80.35% for sample equal to andrographolide 20.10 mg/kg/day dose of Tablet I, II and III respectively. The inhibition produced by the products were significant ( $P < 0.05$ ) compared to control (Table 1).

Tabel 1. Antiplasmodial Activity of Tablet I, II, and III during 4-days Test

Sample	Active substance	Andrographolide dose (mg/kg bw/day)	Average parasitemia (%)	Average inhibition (%)
Tablet I	AP fraction A	25.10	3.37±0.29	70.15
Tablet II	AP fraction B	25.10	2.85±0.48	78.16
Tablet III	AP fraction B	25.10	2.96±0.41	80.35
Negative control	-	-	10.85±1.54	-

Data were expressed as mean±SD for five animals per group F=53.789

\*P<0.001 compared to control

The data was analyzed by one-way analysis of variance (one way ANOVA) at 95% confidence limit ( $\alpha = 0.05$ ) to detect the significant differences in inhibition of parasite's growth of each test group. ANOVA and Post Hoc LSD (Least Significant Difference) test resulted the calculating F(53.789) and  $P < (0.001)$ . It concluded that there were significant difference among Tablet I and Tablet II, Tablet I and Tablet III, but no significant difference between Tablet II and Tablet III. Tablet I and II displayed different antimalarial activity, they were formulated by the same



method, but different in active substance. Meanwhile, Tablet II and Tablet III displayed similar antimalarial activity, in which they were formulated by different method but contain the same active substance.

The antimalarial activities of Tablet I, II and III were mainly due to the active substances which contained in tablets instead of formulation method which conducted to produce tablets. Regarding to the fact that andrographolide was one of the active substances which proven active as an antimalarial and assumed to be responsible to the antimalarial activity of *A. paniculata*, then tablets were produced based on the content of andrographolide. Tablet I, II and III were containing the same andrographolide level as an effort to reach the same antimalarial activity level. Tablet I contain AP fraction A which was crude ethyl acetate fraction of *A. paniculata* extract with multi-components content and lower concentration of andrographolide, meanwhile tablet II and III contain AP fraction B which was diterpene lactone fraction obtained from purification of AP fraction A, more concentrate with higher andrographolide concentration. AP fraction B has shown better antimalarial activity than AP fraction A at the same level of andrographolide. Physicochemical properties of AP fraction B was greenish amorphous powder shown different form with AP fraction A which was dark green and sticky. There was a possibility that the sticky form of AP fraction A affected the dissolution profile of andrographolide as an active substance and produced the low bioavailability in the in vivo system. The results shown that the form of raw material of *A. paniculata* fractions were influenced the antimalarial activities of tablets.

#### 4. Conclusion

*A. paniculata* formulated products containing AP fraction B as an active substance in Tablet II and III showed higher parasite growth's inhibition than product containing AP fraction A in Tablet I. The different formulation method was not influenced the antimalarial activity. The antimalarial activity products depending on the raw material form of *A. paniculata* fraction which contained in the products.

#### Acknowledgement

This research was supported by the Fund of the Directorate of Higher Education, Republic of Indonesia Ministry of National Education.

#### References

1. Greenwood BM, Bojang K, Whitty CJ, Targett GA: Malaria. *Lancet* 2005; **365**: 1487-98.
2. Ridley RG. Medical need, scientific opportunity and the drive for antimalarial drugs. *Nature* 2002; **415**: 686-93.
3. Zirihi GN, Mambu L, Guédé-Guina F, Bodo B, Grellier P. In vitro antiplasmodial activity and cytotoxicity of 33 West African plants used for treatment of malaria. *J Ethnopharmacol* 2005; **98**: 281-85.
4. Rahman NNNA, Furuta T, Kojima S, Takane K, Ali Mohd M. Antimalarial Activity of extracts of Malaysian medicinal plants. *J Ethnopharmacol* 1999; **64** (3), 249–54.
5. Mishra K, Dash AP, Swain BK, Dey N. Antimalarial activities of *Andrographis paniculata* and *Hedyotis corymbosa* extracts and their combination with curcumin. *Malaria J* 2009; **8** (26), 1-9
6. Mishra K, Dash AP, Dey N. Andrographolide: A novel antimalarial diterpene lactone compound from *Andrographis paniculata* and its interaction with curcumin and artesunate. *J Trop Med* 2011; 1-6.
7. Nogueira C, Lopes LMX. Review: Antiplasmodial natural products. *Molecules* 2011; **16**, 2146-90.
8. Widyawaruyanti A. Antimalarial activity test of diterpene lactone compound isolated from *Andrographis paniculata* Nees. Project Grand-QUE Research Report, Faculty of Pharmacy, Universitas Airlangga, 2001 (unpublished).
9. Widyawaruyanti A. The Development of Antimalarial Phytopharmaceutical Product from diterpene lactone fraction of *Andrographis paniculata* Nees, Research Report of Hibah Kompetitif Penelitian Strategis Nasional Batch I, Faculty of Pharmacy, Universitas Airlangga, 2009 (unpublished)