

New N-allylthiourea derivatives: Synthesis, molecular docking and in vitro cytotoxicity studies

by Tri Widiandani

Submission date: 19-Sep-2019 10:45AM (UTC+0800)

Submission ID: 1175530035

File name: TJPR_17_8_Tri_Widiandani.pdf (304.09K)

Word count: 4059

Character count: 22385

Original Research Article

New N-allylthiourea derivatives: Synthesis, molecular docking and *in vitro* cytotoxicity studies

Tri Widiandani^{1*}, Siswandono¹, Edy Meiyanto², Melanny Ika Sulistyowaty^{1,3}, Bambang Tri Purwanto¹, Suko Hardjono¹

¹Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Universitas Airlangga, Surabaya, ²Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, Indonesia, ³Graduate School in Biomedical and Health Sciences, Hiroshima University, Hiroshima, Japan

*For correspondence: **Email:** tri-w@ff.unair.ac.id; **Tel:** +62-81803022660

Sent for review: 12 March 2018

Revised accepted: 26 July 2018

Abstract

30 **pose:** To synthesise derivatives of N-allylthiourea and evaluate their anticancer activities against epidermal growth factor receptor (EGFR) using *in silico* and *in vitro* methods. **27**

Methods: Four compounds were synthesized using the Schotten-**10**mann reaction. The structures of the synthesized compounds were confirmed using infrared (IR), proton nuclear magnetic resonance (¹H-NMR), carbon nuclear magnetic resonance (¹³C-NMR) and electrospray ionization mass spectrometry (ESI-MS) methods. Molecular modeling was carried out with Molegro Virtual Docker version 5.5 through docking of the compounds onto the protein binding site of EGFR, with protein data bank (PDB) codes 1M17, 1XKK, and 3POZ. *In vitro* cytotoxicity was evaluated in MCF-7 cell lines using MTT assay.

Results: The synthesized compounds showed lower Rerank Scores, relative to N-allylthiourea and hydroxyurea. The low Rerank Score values implied stable molecular bonds, and hence higher biological activities. In addition, the derivatives showed cytotoxicities against MCF-7 cell line (IC₅₀: 0.21 – 0.38 mM) which were superior to those of N-allylthiourea (IC₅₀: 5.22 mM) and hydroxyurea (IC₅₀: 2.89 mM).

Conclusion: The predicted anticancer activities of the synthesized compounds are consistent with results from *in silico* studies and assays of cytotoxicity against MCF-7 cell lines. Thus, N-allylthiourea derivatives can potentially be developed as anticancer drugs.

Keywords: N-allylthiourea derivatives, *In silico*, Synthesis, Cytotoxicity, MCF-7, EGFR

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited.

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

INTRODUCTION

Chemotherapy is one method of cancer treatment that works by inhibiting the growth of cancer cells or killing them. It is known that chemotherapy has many adverse side effects, such as hair loss, diarrhea, dry mouth, nausea,

vomiting, and fatigue [1]. Therefore, it is important to evolve newer, more effective and more selective anticancer agents [2,3]. In the present study, thiourea derivatives were developed as potential anticancer agents by modifying the structure of N-allylthiourea (ATU) [3-8]. The choice of N-allylthiourea was due to

its thiol group, a pharmacophore which is responsible for its biological activity. In addition, *N*-allylthiourea possesses a primary amine group (-NH₂). These groups allow for possible modifications that may lead to the development of ATU as an anticancer agent.

The chemical modification of the structure of *N*-allylthiourea was effected by reacting it with derivatives of benzoyl chlorides, resulting in generation of *N*-benzoyl-3-allylthiourea (A – D, Figure 1) using the Schotten-Baumann reaction [7]. In the first step, the amine group of *N*-allylthiourea reacted with the C-carbonyl of benzoyl chloride forming an amide, with the release of a proton and a chloride ion. The addition of the weak base, triethylamine (TEA) was necessary to neutralize the acidic proton formed, otherwise, the reaction would not have proceeded. Moreover, the addition of TEA blocked the protonation of the amine, a reaction which would have made it impossible for the base to react as a nucleophile [7].




Figure 1: (i) Steps in the synthetic process according to the Schotten-Baumann procedure. (ii) Chemical structures of the four derivatives of *N*-allylthiourea: **A**, 3,4-dichlorobenzoyl-3-allylthiourea; **B**, 2,4-dichlorobenzoyl-3-allylthiourea; **C**, 4-fluorobenzoyl-3-allylthiourea; **D**, 4-methylbenzoyl-3-allylthiourea

Molecular modeling is usually conducted to examine the affinity of a ligand to its docking site through evaluation of the energy of drug-receptor binding. It has been shown that the evaluation of the interaction between a molecule and its docking protein involved in signal transduction may aid in identifying potential biological activity, as well as possible mechanisms of action [8,10,11]. Studies indicate that the 1M17, 293K, and 3POZ receptors are used as models of the epidermal growth factor receptor, EGFR [1,12]. The standard ligands for these receptors are erlotinib, lapatinib, and TAK-285, respectively (TAK-285 is a selective EGFR inhibitor) [12]. In addition, these receptors resemble thiourea derivatives (EGFR/HER-2 inhibitors) which act by inhibiting the tyrosine kinase receptor (RTKs) in

the intracellular region [13-15]. As inhibitors of EGFR, thiourea derivatives exhibit anti-proliferation activities against tumor cells, including human breast cancer cells [3,4,14].

One important parameter in molecular docking is the Rerank Score (RS). It is the energy of the ligand-receptor interaction, and a measure of bond energy between ligand and receptor [7,8,10]. The smaller the RS value, the more stable the bond between ligand and receptor. Therefore, a strong interaction implies a high biological activity [8].

In the present study, the cytotoxicities of *N*-allylthiourea and its derivatives against MCF-7 breast cancer cell line were investigated using MTT assay [6,16,18].

EXPERIMENTAL

Chemicals and reagents

N-Allylthiourea was purchased from Merck; 3,4-dichlorobenzoyl chloride; 2,4-dichlorobenzoyl chloride; 4-fluorobenzoyl chloride, and 4-methylbenzoyl chloride were products of Sigma-Aldrich. The reactions were monitored by Thin Layer Chromatography (TLC) carried out on silica gel 60 F₂₅₄ plates (Merck). Spot detection was performed at UV 254 nm [19]. Purification of the products was carried out by recrystallization [7,19].

Infrared spectra were measured using a Perkin-Elmer Spectrum One spectrophotometer and major absorptions were listed in cm⁻¹. ¹H-NMR (600 MHz) and ¹³C-NMR (150 MHz) spectra were measured on JEOL JNM-ECS 600 instrument with DMSO-d₆ as solvent and chemical shifts reported in ppm on the δ-scale. Positive ion HR-ESI-MS was accomplished by an Applied Biosystems QSTAR XL Nanospray™ system. *In silico* studies were carried out using Molegro® Virtual Docker version 5.5; 1M17, 1XKK, and 3POZ receptors were obtained from Protein Data Bank (www.pdb.org). The 2-D and 3-D structures of *N*-allylthiourea derivatives and the ligand were measured using ChemBio® Office Ultra 11.0 and MMFF94 for minimization of the energy [8,11].

Cell culture

MCF-7 cell lines were obtained from the Laboratory of Gene Regulation at Graduate School of Biological Sciences, Nara Institute of Science and Technology (NAIST), Japan. The cells were grown on high-glucose Dulbecco's modified Eagle's medium (DMEM; Nacal

Tesque, Japan) containing 10 % fetal bovine serum (FBS) (Sigma, USA) and 25 % penicillin-streptomycin (Sigma, USA). The cells were harvested from culture dish using Trypsin-EDT (34 0.25 % (Sigma, USA). The MTT reagent (5 mg/ml) was prepared by dissolving MTT (Nacalai, Japan) in phosphate-buffered saline (PBS).

Docking

Receptor and ligand were downloaded from Protein Data Bank. The ligand-receptor complexes were measured in the cavity. The 3-D structures of derivatives of *N*-allylthiourea were replaced to the fixed cavity. The ligands were docked towards the 1M17, 1XKK, and 3POZ receptors [12]. In the next step, Docking Score, Rerank Score, RMSD value, and the state of the environment of each compound derivative, such as hydrogen bonds, hydrophobic interactions, and electronic bonds, were determined [11,16,20].

Synthesis of *N*-allylthiourea derivatives

N-Allylthiourea (0.025 moles) was dissolved in 20 mL of tetrahydrofuran. Benzoyl chloride derivatives (1 eq) and triethylamine (7 mL) were added to the solution, stirred for 30 min at room temperature, and refluxed for 3 - 4 h. Thereafter, the mixture was evaporated and washed with saturated aqueous sodium bicarbonate solution. The crude product was purified using column chromatography, with hexane: acetone (3:2); chloroform: hexane (5:2) and chloroform: hexane: ethyl acetate (1:6:2) as mobile phases at the indicated volume ratios. Recrystallization was done using ethanol: water at a volume ratio of 1:1 [7].

EVALUATION OF CYTOTOXICITY

The cytotoxic effects of the derivatives against MCF9 cells were evaluated using MTT assay. The cells were seeded into 96-well plates at a density of 8×10^3 cells/well and incubated at 37 °C for 24 h in an atmosphere containing 5 % CO₂. The synthesized compounds were applied to the wells separately at different concentrations. After a 24-h incubation, MTT (5 mg/mL in PBS) was added to each well, and incubation was continued for 4 h at 37 °C to allow for formation of formazan crystals. Then, the reaction in each well was stopped by addition of the stopper reagent, a solution of 10 % sodium dodecylsulphate (SDS, Merck) in 0.01 N HCl (Merck), which also served to solubilize the formazan crystals. The plates were kept overnight in the dark at room temperature, after

which the absorbance of the formazan solution in each well was read at 570 nm in an ELISA plate reader [2,17,18].

RESULTS

Derivatives synthesized from *N*-allylthiourea

The desired derivatives were synthesized through nucleophilic substitution of *N*-allylthiourea with benzoyl chloride derivatives i.e. 3, 4-dichlorobenzoyl chloride; 2, 4-dichlorobenzoyl chloride; 4-fluorobenzoyl chloride, and 4-methylbenzoyl chloride, yielding compounds A, B, C and D, respectively (Figure 1). The detailed physicochemical and spectral data of the compounds are described below:

Compound (A) was obtained as a white crystal (49 % yield) with the following spectral characteristics: IR (cm⁻¹) = 1672; 3467; 1548; and 1447; ¹H-NMR δ_H (ppm, *J* in Hz) = 4.30 (t, *J* = 5.2, 2H), 5.20 (d, *J* = 12, 1H), 5.25 (d, *J* = 3, 1H), 6.00 - 5.93 (m, 1H), 7.88 (d, *J* = 6, 2H), 7.89 (d, *J* = 6, 2H), 8.21 (s, 1H), 10.83 (s, 1H), and 11.61 (s, 1H); ¹³C-NMR δ_C (ppm) = 46.86, 116.69, 128.81, 130.57, 130.63, 131.43, 132.71, 132.96, 135.64, 165.80, and 180.09; HR-ESI-MS *m/z* = 288.9222 [M + H]⁺ (calculated value for C₁₁H₁₂ON₂Cl₂S = 288.9262). These spectral data agreed with the structure of 3,4-dichlorobenzoyl-3-allylthiourea (A).

The synthesized compound (B) was a white crystal obtained at a yield of 75 %. Its spectral data were: IR (cm⁻¹) = 1684, 3467, 2885 and 1425; ¹H-NMR δ_H (ppm, *J* in Hz) = 11.81 (s, 1H), 10.64 (s, 1H), 7.75 (d, *J* = 2, 1H), 7.62 (d, *J* = 8.3, 1H), 7.54 (d, *J* = 5.1, 1H), 6.02-5.92 (m, 1H), 5.25 (d, *J* = 10.3, 1H), 5.20 (d, *J* = 1.6, 1H), and 4.29 (t, *J* = 3.6, 2H); ¹³C-NMR δ_C (ppm) = 180.24, 167.09, 136.17, 133.88, 133.45, 134.6, 131.03, 129.54, 127.80, 117.33 and 47.31; HR-ESI-MS *m/z* = 288.9966 [M + H]⁺ (calculated value for C₁₁H₁₁ON₂Cl₂S = 288.9969). These spectral data agreed with the structure of 2,4-dichlorobenzoyl-3-allylthiourea (B).

Compound (C) was a white crystalline salt (48 % yield) with the following spectral data: IR (cm⁻¹) = 1674; 3429; 1568 and 1519; ¹H-NMR δ_H (ppm, *J* in Hz) = 4.30 (t, *J* = 13.2, 2H), 5.20 (d, *J* = 12, 1H), 5.26 (d, *J* = 18, 1H), 6.00-5.94 (m, 1H), 7.36 (t, *J* = 18Hz, 2H), 8.03 (t, *J* = 12, 2H), 10.93 (s, 1H), and 11.43 (s, 1H); ¹³C-NMR δ_C (ppm) = 46.83, 115.42, 116.65, 128.73 and 131.55 (2C); 133.05, 164.00, 166.99, 180.30; HR-ESI-MS *m/z* = 239.0650 [M + H]⁺ (calculated value for C₁₁H₁₂ON₂FS = 239.0654). These spectral data

agreed with the structure of 4-fluorobenzoyl-3-allylthiourea (C).

The derivative compound (D) was a white crystal (35 % yield). Its spectral features were: IR (cm^{-1}) = 1663, 3467, 1609 and 1448; $^1\text{H-NMR}$ δ_{H} (ppm, J in Hz) = 11.26 (s, 1H), 11.00 (s, 1H), 7.87 (d, J = 8.2, 2H), 7.33 (d, J = 7.9, 2H), 6.02-5.93 (m, 1H), 5.28 (d, J = 42, 1H), 5.20 (d, J = 12, 1H), 4.30 (t, J = 3.6, 2H), and 2.39 (s, 3H); $^{13}\text{C-NMR}$ δ_{C} (ppm) = 180.176, 166.847, 144.736, 131.941, 129.929, 128.861 (2C), 127.574 (2C), 117.801, 48.160, 21.769. HR-ESI-MS m/z = 235.0900 [$\text{M} + \text{Na}$] $^+$ (calculated value for $\text{C}_{12}\text{H}_{15}\text{ON}_2\text{S}$ = 235.0905). These were consistent with the structure of 4-methylbenzoyl-3-allylthiourea (D).

Docking results




Figure 2: The amino acids involved in the interaction process between ligands in epidermal growth factor receptor. (1) 2-D interaction of compound A with 1M17; (2) H-bond interaction with 1XKK in 3-D illustration; and (3) 2-D interaction of ligand standard TAK-285 with 3POZ

Table 1: Interactions between the synthesized compounds (ligands) and amino acid residues in the EGFR receptor

Code	1M17		3POZ		1XKK	
	H-bond	Steric bond	H-bond	Steric bond	H-Bond	Steric bond
A	Met769	Met769, Thr766, Leu820, Leu768, Ile765, Gln767, Ala719	Met793, Thr854	Met793, Gly796, Gln791, Thr790, Val726, Ile744, Ala743, Ile789, Leu788	Met793, Thr854	Leu792, Thr790, Gln791, Ala743, Ile744, Ile789, Gly796, Met793
	Met769	Met769, Thr766, Leu820, Leu768, Leu764, Ile765, Ile720, Gln767, Ala719	Met793, Thr854	Met793, Gly796, Gln791, Thr790, Val726, Ile744, Ala743, Ile789, Leu788	Met793, Thr854	Leu792, Leu788, Thr790, Gln791, Ala743, Ile789, Gly796, Met793
C	Met769	Met769, Thr766, Leu820, Lau764, Leu768, Ile765, Gln767, Ala719	Met793, Thr854	Met793, Gly796, Gln791, Thr790, Ala743, Ile789, Leu788	Met793, Thr854	Leu788, Thr854, Thr790, Gln791, Ala743, Ile789, Gly796, Met793
D	Met769	Met769, Thr766, Leu820, Lau764, Leu768, Ile765, Gln767, Ala719	Met793, Thr854	Met793, Gly796, Gln791, Thr790, Ala743, Ile789, Leu788	Met793, Thr854	Leu788, Thr854, Thr790, Gln791, Ala743, Ile789, Gly796, Met793
ATU	-	Met769, Leu768, Gln767, Leu820	-	Met793, Thr790, Gln791, Leu844	-	Met793, Thr790, Gln791, Leu844
HU	-	Met769, Leu768, Gln767	-	Met793, Gln791	-	Met793, Thr790, Gln791
SL	Met769	Met769, Gln767, Gly772, Pro770, Asp831	Met769, Arg841	Phe856, Cys775	Met793	Gln791, Arg776, Thr790, Lau788, Asp800, Cys797

Note: ATU = Allylthiourea; HU = hydroxyurea; SL = standard ligand

The interaction between the synthesized compounds (ligands) and amino acid residues are shown in Table 1 and illustrated in 2D in Figure 2.

The docking scores with respect to interaction with epidermal growth factor receptors were obtained using the Molegro Virtual Docker as shown in Table 2.

Cytotoxicity of compounds A - D

The cytotoxic activities of the four derivatives were assessed using MTT assay, and the results are shown in Table 2. The viabilities of the cells obtained from the measurement of the absorbance of formazan formed after treatment with MTT reagent, and representative morphologies of MCF-7 cell line after treatment are shown in Figure 3.

DISCUSSION

In drug development and discovery it is important to ensure that the drug-like physicochemical properties are maintained as described by Lipinsky's rule [9]. Most orally-administered drugs are relatively small and moderately lipophilic molecules.

Table 2: Docking (RS) and IC₅₀ data

Compound	Substituent	Docking Score (kcal/mol)*			IC ₅₀ (mM) ±SD
		1M17	1XKK	3POZ	MCF-7
A	3,4-Cl ₂	-94.7885	-90.8922	-81.6869	0.213±1.61
B	2,4-Cl ₂	-94.1892	-88.2606	-75.4107	0.231±1.34
C	4-F	-91.2717	-89.1355	-80.7994	0.261±1.92
D	4-CH ₃	-68.5424	-84.8147	-76.6435	0.230±1.05
ATU	-	-42.2023	-55.9170	-56.4322	5.219±1.81
HU	-	-44.2384	-44.2922	-41.3393	2.793±1.66

*Rerank Score is the value of energy after re-ranking by considering RMSD factor and chemical bonds involved [15].






Figure 3: (i) Effect of compounds A-D on viability of MCF-7 cells. (1a): Hydroxyurea (HU) and *N*-allylthiourea (ATU); (1b) Effect on serial concentrations of HU and ATU on viability of MCF-7 cells; (ii) Effect of compound A (200 μ M) on viability of MCF-7 cells, relative to control cells. Treatment with the synthesized compounds changed the morphology of MCF-7 cell line after 24-h incubation. Cell morphology was observed using an inverted microscope at a magnification of x100.

Chemdraw[®] is a helpful tool for predicting the lipophilicity of organic compounds [8]. The log P values of *N*-benzoyl-3-allylthiourea, *N*-allylthiourea and hydroxyurea are 2.0450, 0.0799, and -1.8000, respectively [7]. The higher the log P value, the more lipophilic the compound. Based on this parameter, the derivatives of *N*-allylthiourea are more lipophilic than *N*-allylthiourea and hydroxyurea. Thus, theoretically, the derivatives of *N*-allylthiourea can penetrate the cell membrane more easily, and so possess higher biological activities than the parent compound. The infrared spectra of the synthesized compounds A-D showed peaks above 1663 - 1684 cm^{-1} , indicating the presence of C=O carbonyl groups in their structures. Furthermore, the chemical structures of compounds A-D were confirmed through several parameters such as ratio of protons as seen from integrations and splitting patterns of the peaks in ¹H-NMR spectra, number of peaks in ¹³C-NMR spectra, and the exact mass of molecular ion measured by HR-MS.

Based on *in silico* studies on the interactions between the ligands and EGF receptor, compound A interacted with amino acid residues such as Met769, Met793, and Thr854 through steric bonds and H-bonds. The interactions of other compounds (B, C and D) followed a similar pattern. The synthesized compounds exhibited the same ATP binding site for the H-bond binding of the amino acid residues Met793 and Met769 to the EGFR. However, *N*-allylthiourea and hydroxyurea bound amino acid residues only through steric bonds: H-bonds were not involved.

The presence of the lipophilic benzoyl group in *N*-benzoyl-3-allylthiourea resulted in enhanced biological activity due to stabilization of the ligand-receptor bond. This is consistent with the Rerank Score values of the *N*-allylthiourea derivatives (compounds A - D) which were lower than those of *N*-allylthiourea (lead compound) and hydroxyurea (clinically-used anticancer drug). The addition of substituents with certain lipophilic, electronic and steric properties improves the stability of ligand-receptor bonds.

The variation of substituents on *N*-benzoyl-3-allylthiourea caused differences in the number of interaction with amino acids. However, it did not bring about significant differences in Rank Score values. From the results of *in silico* studies, some derivatives of *N*-allylthiourea were predicted to have greater anticancer activities than hydroxyurea. The results of *in vitro* cytotoxicity performed on MCF-7 cell lines using the MTT method revealed that the IC₅₀ of compounds **A - D** (0.21 – 0.38 mM) were less than that of *N*-allylthiourea (IC₅₀ = 5.22 mM). These results indicate that the four derivatives of *N*-allylthiourea were more cytotoxic than the lead compound, *N*-allylthiourea. In addition, cytotoxicity studies revealed that all derivatives of *N*-allylthiourea were more cytotoxic against MCF-7 cell lines when compared to the commercial anticancer drug, hydroxyurea (IC₅₀ = 2.79 mM).

CONCLUSION

Molecular docking data have predicted that *N*-allylthiourea derivatives are epidermal growth factor receptor inhibitors. This is supported by results from cytotoxicity studies using MCF-7 cancer cell lines. Therefore, *N*-allylthiourea derivatives possess potential anticancer properties that can be harnessed for the development of new anticancer drugs.

DECLARATIONS

Acknowledgement

This research was supported by the Ministry of Research Technology and Higher Education, The Republic of Indonesia. The authors are grateful for access to conduct NMR and ESI-MS measurements at the Natural Science Center for Basic Research and Development (N-BARD) of the Graduate School of Biomedical and Health Sciences, Hiroshima University. The authors are grateful to Professor Yasumasha Bessho and Associate Professor Norihiro Ishida-Kitagawa for access to cell culture facility at the Laboratory of Gene Regulation, Department of Biological Science, Nara Institute of Science and Technology, Japan.

Conflict of Interest

No conflict of interest associated with this work.

Contribution of Authors

The authors declare that this work was done by the authors named in this article and all liabilities

pertaining to claims relating to the content of this article will be borne by them.

REFERENCES

- Goldwein JW, Vachani C. *Chemotherapy: The Basic*. Onkolink.com [homepage on the Internet]. [Update 2016 April 12, cited 2018 February 11] Available from: <https://www.onkolink.org/cancer-treatment/chemotherapy/overview/chemotherapy-the-basics>.
- Rudyanto M, Widiandani T, Syahrani A. 6-Allyl-3-(4-Methoxybenzyl)-8-Methoxy-3,4-Dihydro-2H-Benzo[e][1,3]Oxazine and 4-Allyl-2-Methoxy-6-(4-Methoxybenzyl) Aminomethylphenol: Synthesis and cytotoxicity test on MCF-7 cells. *Int J Pharm Clin Res* 2016; 8(5): 387-391.
- Li HQ, Yan T, Yang Y, Shi L, Zhou CF, Zhu HL. Synthesis and structure-activity relationships of *N*-benzyl-*N*-(*X*-2-hydroxybenzyl)-*N'*-phenylureas and thioureas as antitumor agents. *Bioorganic Med Chem* 2010; 18(1): 305-313.
- Liu W, Zhou J, Zhang T, Zhu H, Qian H, Zhang H, Huang W, Gust H. Design and synthesis of thiourea derivatives containing a benzo [5,6] cycloheptal [1,2-*b*]pyridine moiety as potential antitumor and anti-inflammatory agents. *Bioorganic Med Chem Lett* 2012; 22: 2701-2704.
- Huang XC, Wang M, Pan YP, Yao GY, Wang HS, Tiang XY, Qin JK, Zhang Y. Synthesis and antitumor activities of novel thiourea α -aminophosphonates from dehydroabietic acid. *Eur J Med Chem* 2013; 69: 508-520.
- Widiandani T and Siswandono S. Structure modification and anticancer activity prediction of Allylthiourea derivatives based on *In silico*, *Bikfar* 2015; 4(1): 33-40.
- Widiandani T, Arifianti L, Siswandono S. Docking, Synthesis and Cytotoxicity Test on Human Breast Cancer Cell Line (t47d) of *N*-allylcarbamothioylbenzamide. *Int J Pharm Clin Res* 2016; 8 (5): 372-376.
- Siswandono. *Medicinal Chemistry*. Book 1. 2nd ed. Surabaya: Airlangga University Press; 2016. 25 p.
- Lipinski CA, Lombardo F, Dominy BW, Feeney PJ. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. 2001; *Adv. Drug Del Rev* 46(1-3): 3-26.
- Young DC. *Computational Drug Design, A Guide for Computational and Medicinal Chemists*. New York: John Wiley and Sons; 2009.
- CLCBio. *Molegro Virtual Docker User Manual*, MVD 2012. 5.5 for Windows, Linux, and Mac OS X, Molegro – A CLC Bio Company. 2012.
- Aertgeerts K, Skene R, Yano J, Sang BC, Zou H, Snell G, Jennings A, Iwamoto K, Habuka N, Hirokawa A, et al. Structural Analysis of the Mechanism of Inhibition and Allosteric Activation of the Kinase Domain of HER2 Protein. *J Biol Chem* 2011; 286(21): 18756-18765.

13. Laskin JJ and Sandler AB. Epidermal growth factor receptor: a promising target in solid tumours. *Can Treat Rev* 2004; 30(1): 1-17.
14. Manjula SN, Noolvi NM, Parihar KV, Reddy SAM, Ramani V. Synthesis and antitumor activity of optically active thiourea and their 2-aminobenzothiazole derivatives: a novel class of anticancer agents. *Eur J Med Chem* 2009; 44(7): 2923-2929.
15. Wieduwilt MJ, Moasser MM. The epidermal growth factor receptor family: biology driving targeted therapeutics. *Cell Mol Life Sci* 2008; 65(10): 1566-1584.
16. Siswandono S, Widiandani T, Hardjono S. Docking and Cytotoxicity Test on Human Breast Cancer Cell Line (T47d) of N-(Allylcarbamothioyl)-3-chlorobenzamide and N-(Allylcarbamothioyl)-3,4-dichlorobenzamide. *Res J Pharm Biol Chem Sci* 2017; 8(2): 1909-1914.
17. Rudyanto M, Widiandani T, Syahrani A. Some Benzoxazine and aminomethyl derivatives of Eugenol: Cytotoxicity on MCF-7 cell line. *Int J Pharm Pharm Sci* 2015; 7(5): 229-232.
18. Meiyanto E, Putri DDP, Susidarti RA, Murwanti R, Sardjiman, Aditya FA, Husna U, Pumomo H, Kawaichi M. Curcumin and its Analogues (PGV-0 and PGV-1), Enhance Sensitivity of Resistant MCF-7 Cells to Doxorubicin through Inhibition of HER2 and NF- κ B Activation. *Asian Pac J Cancer Prev* 2014; 15(1): 179-184.
19. Rudyanto M, Ekowati J, Widiandani T, Honda T. Synthesis and brine shrimp lethality test of some benzoxazine and aminomethyl derivatives of eugenol. *Int J Pharm Pharm Sci* 2014; 6: 96-98.
20. Yang W, Hu Y, Yang YS, Zhang F, Zhang YB, Wang XL, Tang JF, Zhong WQ, Zhu HL. Design, modification and 3D QSAR studies of novel naphthalin-containing pyrazoline derivatives with/without thiourea skeleton as anticancer agents. *Bioorg Med Chem* 2013; 21: 1050-1063.

New N-allylthiourea derivatives: Synthesis, molecular docking and in vitro cytotoxicity studies

ORIGINALITY REPORT

13%

SIMILARITY INDEX

9%

INTERNET SOURCES

11%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	www.ijppsjournal.com Internet Source	1%
2	www.google.com Internet Source	1%
3	Meiyanto, Edy, Dyaningtyas Dewi Pamungkas Putri, Ratna Asmah Susidarti, Retno Murwanti, Sardjiman Sardjiman, Aditya Fitriasaki, Ulfatul Husnaa, Hari Purnomo, and Masashi Kawaichi. "Curcumin and its Analogues (PGV-0 and PGV-1) Enhance Sensitivity of Resistant MCF-7 Cells to Doxorubicin through Inhibition of HER2 and NF-kB Activation", Asian Pacific Journal of Cancer Prevention, 2014. Publication	1%
4	www.tandfonline.com Internet Source	1%
5	en.wikipedia.org Internet Source	1%

6	146.230.128.141 Internet Source	1%
7	legacy.impactjournals.com Internet Source	1%
8	Yasunori Sato, Naoto Sasaki, Megu Saito, Nao Endo, Fumihiko Kugawa, Akemichi Ueno. "Luteolin Attenuates Doxorubicin-Induced Cytotoxicity to MCF-7 Human Breast Cancer Cells", Biological & Pharmaceutical Bulletin, 2015 Publication	1%
9	biomedicaloptics.spiedigitallibrary.org Internet Source	1%
10	link.springer.com Internet Source	1%
11	www.nrcresearchpress.com Internet Source	<1%
12	Cesarini, Sara, Angelo Ranise, and Andrea Spallarossa. "A Successful Application Of Parallel Synthesis To Computer-assisted Structural Optimization Of New Leads Targeting Human Immunodeficiency Virus-1 Reverse Transcriptase The Case Of Acylthiocarbamates And Thiocarbamates", Critical Reviews in Combinatorial Chemistry, 2008. Publication	<1%

- 13 baadalsg.inflibnet.ac.in Internet Source <1%
-
- 14 www.freepatentsonline.com Internet Source <1%
-
- 15 Bertus, P.. "Pentadienyl transfer reagents based on zirconium: preparation and reactions with carbonyl compounds", Tetrahedron, 20040202 Publication <1%
-
- 16 Rohit Singh, Gurubasavaraj V. Pujar, Madhusudan N. Purohit, V. M. Chandrashekar. "Synthesis, in vitro cytotoxicity, and antibacterial studies of new asymmetric bis-1,2,4-triazoles", Medicinal Chemistry Research, 2012 Publication <1%
-
- 17 Genevieve D. Paderes. "Structure-Based and Property-Compliant Library Design of 11 β -HSD1 Adamantyl Amide Inhibitors", Methods in Molecular Biology, 2011 Publication <1%
-
- 18 Hans P. Märki. "Optically Pure Isoproterenol Analogues With Side Chains Containing an Amide Bond: Synthesis and biological properties", Helvetica Chimica Acta, 03/16/1988 Publication <1%
-
- 19 Madhusudan Purohit, Yegeri C. Mayur. "Synthesis, in vitro cytotoxicity, and anti-

microbial studies of 1,4-bis(4-substituted-5-mercapto-1,2,4-triazol-3-yl)butanes", Medicinal Chemistry Research, 2010

Publication

20

Maralinganadoddi P. Sadashiva, Raghavendra Gowda, Xianzhu Wu, Gajanan S. Inamdar et al. "A non-cytotoxic N-dehydroabietylamine derivative with potent antimalarial activity", Experimental Parasitology, 2015

Publication

21

www.jcps.ac.cn

Internet Source

22

Hans-Joachim Gais, C. Venkateshwar Rao, Ralf Loo. "Anionic Cross-Coupling Reaction of α -Metallated Alkenyl Sulfoximines and Alkenyl Sulfoximines with Cuprates Featuring a 1,2-Metal-Ate Rearrangement of Sulfoximine-Substituted Higher Order Alkenyl Cuprates and an α -Metallation of Alkenyl Sulfoximines by Cu", Chemistry - A European Journal, 2008

Publication

23

patents.google.com

Internet Source

24

John D. Williams, Alaa E. Mourad, John C. Drach, Leroy B. Townsend. " Design and Synthesis of Acyclic Nucleoside Analogs with Chlorinated Imidazo[1,2-]pyridine Bases ",

<1%

<1%

<1%

<1%

<1%

Nucleosides, Nucleotides and Nucleic Acids, 2003

Publication

25

repository.hkbu.edu.hk

Internet Source

<1%

26

Leo Havemann. "Chapter 218-1 Open Educational Resources", Springer Science and Business Media LLC, 2016

Publication

<1%

27

www.connectjournals.com

Internet Source

<1%

28

Yujia Dai, Yan Guo, Robin R. Frey, Zhiqin Ji et al. "Thienopyrimidine Ureas as Novel and Potent Multitargeted Receptor Tyrosine Kinase Inhibitors", Journal of Medicinal Chemistry, 2005

Publication

<1%

29

Hwei-Ru Tsou, Nellie Mamuya, Bernard D. Johnson, Marvin F. Reich et al. "6-Substituted-4-(3-bromophenylamino)quinazolines as Putative Irreversible Inhibitors of the Epidermal Growth Factor Receptor (EGFR) and Human Epidermal Growth Factor Receptor (HER-2) Tyrosine Kinases with Enhanced Antitumor Activity", Journal of Medicinal Chemistry, 2001

Publication

<1%

30

Kovacs, Erika, Julie Anne Zorn, Yongjian Huang, Tiago Barros, and John Kuriyan. "A

<1%

Structural Perspective on the Regulation of the Epidermal Growth Factor Receptor", Annual Review of Biochemistry, 2015.

Publication

31

Engler, T.A.. "Stereoselective syntheses of substituted pterocarpan with anti-HIV activity, and 5-aza-/5-thia-pterocarpan and 2-aryl-2,3-dihydrobenzofuran analogues", Bioorganic & Medicinal Chemistry, 199610

Publication

<1%

32

Nicolaou, K.C.. "New synthetic technology for the construction of N-hydroxyindoles and synthesis of nocathiacin I model systems", Tetrahedron, 20070702

Publication

<1%

33

Yoshizawa, K.. "Siloxallenes revisited. A useful functional intermediate for the synthesis of (Z)-@b-branched Morita-Baylis-Hillman type adducts and (Z)-chalcones", Tetrahedron, 20070702

Publication

<1%

34

"Pharmaceutical Bioassays", Wiley, 2009

Publication

<1%

Exclude bibliography On

New N-allylthiourea derivatives: Synthesis, molecular docking and in vitro cytotoxicity studies

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
