



2016 COUNCIL  
FORUM, 2016

# Asian Pacific Journal of Tropical Biomedicine



7  
2017

## Editorial Board

[About the journal](#)

### Editors-in-Chief

Prof. Dr. Jeffrey M. Bethony

Department of Microbiology, Immunology & Tropical Medicine, School of Medicine and Health Sciences, The George Washington University, Washington DC, USA

E-mail: [jbethony@gwu.edu](mailto:jbethony@gwu.edu)

Prof. Dr. Santiago Mas-Coma

Departamento de Parasitología, Facultad de Farmacia, Universidad de Valencia, Valencia, Spain

E-mail: [S.Mas.Coma@uv.es](mailto:S.Mas.Coma@uv.es)

Prof. Dr. Malcolm K. Jones

School of Veterinary Science, Faculty of Science, The University of Queensland, Queensland, Australia

E-mail: [m.jones@uq.edu.au](mailto:m.jones@uq.edu.au)

Prof. Dr. Vanessa Steenkamp

Department of Pharmacology, University of Pretoria, Pretoria, South Africa

E-mail: [vanessa.steenkamp@up.ac.za](mailto:vanessa.steenkamp@up.ac.za)

Nazni Bte Hj. Wasi Ahmad

Senior Research Officer, Medical Entomology Unit, Institute for Medical Research, Kuala Lumpur, Malaysia

E-mail: [nazni@imr.gov.my](mailto:nazni@imr.gov.my)

### President Executive-Editor-in-Chief

Prof. Shunhai Qu

Hainan Medical University Journal Publisher, Hainan Medical University, Haikou, China

E-mail: [qu.sh@163.com](mailto:qu.sh@163.com)

### Deputy Editors-in-Chief

Prof. Dr. Jong-Yil Chai

Department of Parasitology and Tropical Medicine, Seoul National University College of Medicine, Seoul, Korea

Dr. Pierre Roques

ImmunoVirology Division, Institute for Emerging Diseases and Innovative Therapies, Fontenay-aux-Roses, France

Dr. Alcides Troncoso

Department of Infectious Diseases, School of Medicine, Buenos Aires University, Buenos Aires, Argentina

Dr. Stephen Munga

Centre for Global Health Research, Kenya Medical Research Institute, Nairobi, Kenya

### Associate Editors-in-Chief

Prof. Giuseppe La Torre

Department of Public Health and Infectious Diseases, Sapienza University of Rome, Rome, Italy

Prof. Odir Antônio Dellagostin

Biotechnology Unit, Center for Technology Development, Federal University of Pelotas, Pelotas, Brazil

Prof. Sung-Jong Hong

Department of Medical Environmental Biology, Chung-Ang University College of Medicine, Seoul, Korea

Prof. Hassan Vatandoost

Department of Entomology and Vector Control, School of Public Health and National Institute of Health Research, Teheran University of Medical Sciences, Teheran, Iran

Prof. Maria José Ferreira

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Lisbon, Lisbon, Portugal

Dr. Leonard E. G. Mboera

National Institute for Medical Research, Dar es Salaam, Tanzania

Prof. Dr. Iván Darío Vélez

Facultad de Medicina, Universidad de Antioquia, Medellín, Colombia

[SUBMIT ARTICLE](#)[SUBSCRIBE](#)[POPULAR ARTICLES](#)[JOIN AS REVIEWER](#)[GET EMAIL ALERTS](#)[RECOMMEND](#)

## **Editorial Directors**

Prof. Dr. Viroj Wiwanitkit  
Department of Laboratory Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

Assoc. Prof. Dr. Saravanan Thangamani  
Department of Pathology, University of Texas Medical Branch (UTMB), Galveston, USA

Prof. Dr. Laura Bongiovanni  
Faculty of Veterinary Medicine, University of Teramo, Teramo, Italy

## **Statistic Editors**

Assoc. Prof. Khaled Khatab  
Health Economics Modelling and Medical Statistics, Faculty of Health and Wellbeing, Sheffield Hallam University, Sheffield, UK

Assist. Prof. Sarath Chandra Janga  
Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, USA

## **Executive Editors**

Prof. Dr. Salah Akkal  
Department of Chemistry, University of Constantine, Constantine, Algeria

Dr. Hervé Hoste  
French National Institute for Agricultural Research, Université de Toulouse, Toulouse, France

Prof. Thomas N. Jr. Tully  
Department Veterinary Clinical Sciences, Louisiana State University School of Veterinary Medicine, Baton Rouge, USA

Prof. Dr. Barbara R Conway  
Department of Pharmacy, University of Huddersfield, Huddersfield, UK

## **Executive Members of Editorial Board**

Assoc. Prof. Vittorio Sambri  
Unit of Microbiology, Department of Specialistic, Diagnostic and Experimental Medicine, University of Bologna, Bologna, Italy

Prof. Dr. Qiushui He  
Department of Microbiology and Immunology, University of Turku, Turku, Finland

Assist. Prof. Dr. Bindu Sukumaran  
Emerging Infectious Diseases Programme, Duke-NUS Medical School, Singapore

Prof. Dr. Suhail Ahmad  
Department of Microbiology, Kuwait University, Kuwait City, Kuwait

Prof. Indra Vythilingam  
Department of Parasitology, University of Malaya, Kuala Lumpur, Malaysia

Prof. Dr. Wael Mohamed Abou EL-Makarem EL-Deeb  
Department of Clinical Studies, College of Veterinary Medicine and Animal Resources, King Faisal University, Al-Ahsa, Saudi Arabia

Assoc. Prof. Dr. Ben Slama Karim  
Département de Biotechnologie, Université de Tunis El Manar, Tunis, Tunisia

Prof. Bing Huang  
Department of Animal Parasitology, Shanghai Veterinary Research Institute, Chinese Academy of Agricultural Science, Shanghai, China

Prof. Xiao-Nong Zhou  
National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention, Shanghai, China

Dr. Nahid Einollahi  
Department of Lab Medical Sciences, Faculty of Allied Medical Sciences, Tehran University of Medical Sciences, Tehran, Iran

Prof. Dr. Munir Aktas  
Department of Parasitology, College of Veterinary Medicine, Firat University, Elazig, Turkey

Dr. José Manuel Lorenzo Rodríguez  
Researcher in Meat Technology Centre of Galicia, Meat Technology Centre of Galicia, Galicia, Spain

Dr. Giovanni Benelli  
Insect Behaviour Group, Department of Agriculture, Food and Environment University of Pisa, Pisa, Italy

Prof. Dr. Jeong Hwan Shin  
Department of Laboratory Medicine, Busan Paik Hospital, Inje University College of Medicine, Busan, Korea

Luis Ignacio González Granado  
Teaching assistant, Immunodeficiencies Unit, Hospital 12 de Octubre, Madrid, Spain

Dr. Abhishek Mathur  
Executive Director (R&D), NCS Group, Nagpur, Maharashtra, India



Dr. Ahmed Ismail  
Consultant Clinical Pathologist & Chief Lab Services, Supreme Council of Health, Doha, Qatar

Dr. Américo David Rodríguez Ramírez  
Centro Regional de Investigación en Salud Pública, Chiapas, Mexico

Prof. Xue-Jie Yu  
Department of Pathology, University of Texas Medical Branch, Texas, USA

Prof. Arun Kumar  
Department of Biochemistry, Shri Shankaracharya Institute of Medical Sciences, Bhilai, India

Prof. Dr. Chia-Kwung Fan  
Department of Parasitology, Taipei Medical University College of Medicine, Taipei

Assist. Prof. Dr. Cyrille Bisseye  
Centre de Recherche Biomoléculaire Pietro Annigoni (CERBA)/LABIOGENE, UFR/SVT, Ouagadougou, Burkina Faso

Prof. Dr. Dongmi Kwak  
College of Veterinary Medicine, Kyungpook National University, Daegu, Korea

Dr. Donovan Anthony McGrowder  
Chemical Pathology Department, Faculty of Medical Sciences, University Hospital of The West Indies, Kingston, Jamaica

Dr. Farouk El Allaki  
Epidemiology and Surveillance Section, Canadian Food Inspection Agency, Quebec, Canada

Prof. Dr. Gabriel Trueba  
Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito, Quito, Ecuador

Assoc. Prof. Gordana J. Dragovic Lukic  
Department of Pharmacology, Clinical Pharmacology and Toxicology, School of Medicine, University of Belgrade, Belgrade, Serbia

Dr. Hanspeter Marti  
Swiss Tropical and Public Health Institute, Basel, Switzerland

Prof. Dr. Herbert Auer  
Department of Medical Parasitology, Medical University of Vienna, Vienna, Austria

Hetron Mweemba Munang'andu  
Department of Basic Sciences and Aquatic Medicine, Norwegian School of Veterinary Science, Oslo, Norway

Dr. Irma Khachidze  
Department of Behavior, Cognition Functions and Human Psychophysiology, Beritashvili Institute of Physiology, Tbilisi, Georgia

Dr. Jean-François Faucher  
Praticien hospitalier en maladies infectieuses et tropicales, CHU de Besançon, Service des maladies infectieuses et tropicales, Hôpital Minjoz, Besançon cedex, France

Dr. José A. Oteo  
Infectious Diseases Department, Center of Rickettsioses and Vector-borne Diseases, Hospital San Pedro-CIBIR, Logroño, Spain

Dr. Juraj Majtán  
Section of Molecular and Applied Zoology, Institute of Zoology, Slovak Academy of Sciences, Bratislava, Slovakia

Dr. Kalman Imre  
Department of Animal Production and Veterinary Public Health, Faculty of Veterinary Medicine, Banat University of Agricultural Sciences and Veterinary Medicine Timisoara, Timisoara, Romania

Dr. Kingsley Badu  
Noguchi Memorial Institute for Medical Research, University of Ghana, College of Health Science, Accra, Ghana

Lala Ravaomanarivo  
Département d'Entomologie, de la Faculté des Sciences d'Antananarivo, University of Antananarivo, Antananarivo, Madagascar

Prof. Dr. Liwang Cui  
Department of Entomology, The Pennsylvania State University (PSU), Pennsylvania, USA

Assist. Prof. Dr. Luís R. Silva  
Laboratory for Process Engineering, Environment, Biotechnology and Energy (LEPABE), Faculty of Engineering (FEUP)  
University of Porto, Porto, Portugal

Dr. M. Fawzi Mahomoodally  
Department of Health Sciences, Faculty of Science, University of Mauritius, Reduit, Mauritius

Mamadou C Baldé  
Institut Pasteur de Guinée, Cheikh Anta Diop University, Kindia, Guinée

Prof. Dr. Milkyas Endale  
Department of Chemistry, Hawassa University, Hawassa, Ethiopia

Dr. Mochammad Hatta

Molecular Biology and Immunology Laboratory for Infectious Diseases, Department Microbiology, Faculty Medicine  
Hasanuddin University, Makassar, Indonesia

Moses Samje

Department of Medicine, Faculty of Health Sciences, University of Bamenda, Bamili, Cameroon

Assist. Prof. Muftah A. M. Shushni

Department of Pharmacognosy, University of Tripoli, Tripoli, Libya

Dr. Muhammad Akram

Department of Eastern Medicine and Surgery, Faculty of Medical and Health Sciences, The University of Poonch, Azad Jammu and  
Kashmir, Pakistan

Dr. Musso Munyeme

Department of Disease Control, School of Veterinary Medicine, University of Zambia, Lusaka, Zambia

Prof. Dr. Nguyen Hoang Loc

Department of Biotechnology, College of Sciences, Hue University, Hue, Vietnam

Dr. Prof. Obembe Olawole

Department of Biological Sciences, Covenant University, Ota, Nigeria

Prof. Dr. Pius Tshimankinda Mpiana

Department of Chemistry, Faculty of science, University of Kinshasa, Kinshasa, Democratic Republic of the Congo

Prof. Dr. Polrat Wilairatana

Department of Clinical Tropical Medicine, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand

Dr. Pravin Malla Shrestha

Energy Biosciences Institute, University of California, Berkeley, USA

Dr. Rebuma Firdessa

Institute for Molecular Infection Biology, University of Wuerzburg, Wuerzburg, Germany

Prof. Rinaldo Poncio Mendes

Faculdade de Medicina de Botucatu, Universidade Estadual Paulista Júlio de Mesquita Filho, Botucatu, Brazil

Dr. Robin A J Nicholas

Mycoplasma Group, Department of Statutory and Exotic Bacteria, Veterinary Laboratories Agency (Weybridge), Surrey, United Kingdom

Dr. Assoc. Prof. Samath D Dharmaratne

Faculty of Medicine, University of Peradeniya, Peradeniya, Sri Lanka

Dr. Assist. Prof. Song Gao

Department of Pharmacological and Pharmaceutical Sciences, The University of Houston, Houston, USA

Dr. Assist. Prof. Tarig Mohamed Saad Alnour

Department of Microbiology, Alzaiem Alazhari University, Khartoum, Sudan

Dr. Tashi Tobgay

Vector-Borne Disease Control Programme, Ministry of Health Royal Government of Bhutan, Thimphu, Bhutan

Dr. Thomas Dorlo

Department of Pharmaceutical Biosciences, Uppsala University, Uppsala, Sweden

Prof. Tohru Gono

Medical Mycology Research Center, Chiba University, Chiba, Japan

Dr. Vania Giacomet

Department of Paediatrics, University of Milan, Luigi Sacco Hospital, Milano, Italy

Dr. Veasna Duong

Virology Unit, Institut Pasteur du Cambodge, Phnom Penh, Cambodia

Assist. Prof. Dr. Weaam N. E. Ebrahim

Department of Pharmacognosy and Medicinal Plants, Faculty of Pharmacy, Mansoura University, Mansoura, Egypt.

Dr. Xuanming Shi

Department of Physiology, University of Texas Southwestern Medical Center, Dallas, TX, USA

Assist. Prof. Dr. Yong Song

Faculty of Medicine, Dentistry and Health Sciences, The University of Western Australia, Perth, Australia

Dr. Yu Pang

National Center for Tuberculosis Control and Prevention,  
Chinese Center for Disease Control and Prevention, Beijing, China

Assist. Prof. Yuki Eshita

Department of Infectious Disease Control, Faculty of Medicine, Oita University, Oita, Japan

[Editorial and Ethics Policies](#)



[Open Access](#)

[\\$ No Fee](#)



[View mobile site](#)

ISSN: Print -2221-1691, Online - 2588-9222

Keywords

Author name

Asian Pacific Journal of Tropical Biomedicine

Volume

Issue

Pages

## Asian Pacific Journal of Tropical Biomedicine

Latest issue   All issues

### Volume 7, Issue 7

Pages 591-674 (July 2017)

[< Previous vol/issue](#)

[Next vol/issue >](#)

☐ ☒ Expand all article previews [Download PDFs](#) [Export](#)

## Review

### ☐ Red propolis: Chemical composition and pharmacological activity

[Open access](#) - Review article

Pages 591-598

Luciane Corbellini Rufatto, Denis Amilton dos Santos, Flávio Marinho, João Antonio Pêgas  
Henriques, ... Sidnei Moura



Download PDF

Article preview

### ☐ An overview on application of phage display technique in immunological studies

[Open access](#) - Review article

Pages 599-602

Abbas Rami, Mahdi Behdani, Najmeh Yardehnavi, Mahdi Habibi-Anbouhi, Fatemeh Kazemi-  
Lomedasht



Download PDF

Article preview

### ☐ *Euphorbia tirucalli* L.: Review on morphology, medicinal uses, phytochemistry and pharmacological activities

[Open access](#) - Review article

Pages 603-613

Prashant Y. Mali, Shital S. Panchal



Download PDF

Article preview

## Basic research

### ☐ Antihypertensive and antioxidant activity of *Cassipourea filiformis* L.: A correlative study

[Open access](#) - Original research article

Pages 614-618









Yori Yuliandra, A. Armenia, Helmi Arifin



Download PDF

Article preview



- ☐ Antihyperglycemic effect of methanol extract of *Tamarix aphylla* L. Karst (Saltcedar) in streptozocin–nicotinamide induced diabetic rats  
Open access - Original research article  
Pages 619-623  
Rooh Ullah, Shafiq Ahmed Tariq, Naeem Khan, Nawaz Sharif, ... Khalid Mansoor  
 Download PDF   Article preview 
- ☐ *Candida albicans* isolated from urine: Phenotypic and molecular identification, virulence factors and antifungal susceptibility  
Open access - Original research article  
Pages 624-628  
Laura Wiebusch, Adriana Araújo de Almeida-Apolonio, Luana Mireli Carbonera Rodrigues, Bruna de Paula Bicudo, ... Kelly Mari Pires de Oliveira  
 Download PDF   Article preview 
- ☐ Antimicrobial activity of Tunisian *Euphorbia paralias* L.  
Open access - Original research article  
Pages 629-632  
Malek Besbes Hlila, Kaouther Majouli, Hichem Ben Jannet, Mahjoub Aouni, ... Boulbaba Selmi  
 Download PDF   Article preview 
- ☐ Antiviral activity of the dichloromethane extracts from *Artocarpus heterophyllus* leaves against hepatitis C virus  
Open access - Original research article  
Pages 633-639  
Achmad Fuad Hafid, Chie Aoki-Utsubo, Adita Ayu Permanasari, Myrna Adianti, ... Hak Hotta  
 Download PDF   Article preview 
- ☐ Toxicological evaluation and protective effect of ethanolic leaf extract of *Launaea taraxacifolia* on gentamicin induced rat kidney injury  
Open access - Original research article  
Pages 640-646  
Lydia Enyonam Kuatsienu, Charles Ansah, Michael Buenor Adinortey

[Download PDF](#)[Article preview](#)

- ☐ The protective effect of rutin and quercetin on 5-FU-induced hepatotoxicity in rats

[Open access](#) - Original research article

Pages 647-653

Volkan Gelen, Emin Şengül, Semin Gedikli, Gözde Atila, ... Mustafa Makav

[Download PDF](#)[Article preview](#)

- ☐ Chemical constituents, *in vitro* antimicrobial and cytotoxic potentials of the extracts from *Macaranga barteri* Mull-Arg

[Open access](#) - Original research article

Pages 654-659

Akintayo Ogundajo, Benjamin Okeleye, Anofi Omotayo Ashafa

[Download PDF](#)[Article preview](#)

- ☐ Optimization of ionic liquid-based microwave-assisted extraction of polyphenolic content from *Peperomia pellucida* (L) kunth using response surface methodology

[Open access](#) - Original research article

Pages 660-665

Islamudin Ahmad, Arry Yanuar, Kamarza Mulia, Abdul Mun'im

[Download PDF](#)[Article preview](#)

- ☐ Dysregulation of Notch signaling related genes in oral lichen planus

[Open access](#) - Original research article

Pages 666-669

Nunthawan Nowwarote, Thanaphum Osathanon

[Download PDF](#)[Article preview](#)

- ☐ Antibiotic susceptibility and molecular characterization of resistance genes among *Escherichia coli* and among *Salmonella* subsp. in chicken food chains

[Open access](#) - Original research article

Pages 670-674



---

ISSN: 2221-1691

Copyright © 2018 Hainan Medical University. Production and hosting by Elsevier B.V.

---

**ELSEVIER**

[About ScienceDirect](#) [Remote access](#) [Shopping cart](#) [Contact and support](#) [Terms and conditions](#)  
[Privacy policy](#)

Cookies are used by this site. For more information, visit the [cookies page](#).

Copyright © 2018 Elsevier B.V. or its licensors or contributors. ScienceDirect® is a registered trademark of Elsevier B.V.

 **RELX Group™**



## Publish Open Access

Make an impact.

tandfonline.com

OPEN

# Asian Pacific Journal of Tropical Biomedicine

Country [Netherlands](#) -

Subject Area and Category [Biochemistry, Genetics and Molecular Biology](#)  
[Biochemistry, Genetics and Molecular Biology \(miscellaneous\)](#)

Publisher [Wolters Kluwer Medknow Publications](#)

Publication type [Journals](#)

ISSN [22211691, 25889222](#)

Coverage [2011-2020](#)

Scope The journal will cover technical and clinical studies related to health, ethical and social issues in field of biology, bacteriology, biochemistry, biotechnology, cell biology, environmental biology, microbiology, medical microbiology, pharmacology, physiology, pathology, immunology, virology, toxicology, epidemiology, vaccinology, hematology, histopathology, cytology, genetics and tropical agriculture. Articles with clinical interest and implications will be given preference.

[Homepage](#)

[How to publish in this journal](#)

[Contact](#)

[Join the conversation about this journal](#)

54

H Index


BRIDGE GROUP




# Quality Supplements at iHerb® iHerb

Quartiles


Biochemistry, Genetics and Molecular Biology (miscellaneous)




SJR




Citations per document




Total Cites Self-Cites



External Cites per Doc Cites per Doc



% International Collaboration




Citable documents Non-citable documents

Cited documents Uncited documents

1.2k

1.2k

Asian Pacific Journal of  
Tropical Biomedicine



← Show this widget in  
your own website

Just copy the code below  
and paste within your html  
code:

```
<a href="https://www.scimagojr.com" data-bbox="260 194 407 207">
```



Metrics based on Scopus® data as of April 2020

**Nafis** 3 months ago

Why is the Country of Publication mentioned China at [https://www.ncbi.nlm.nih.gov/nlmcatalog?term=2221-1691\[ISSN\]](https://www.ncbi.nlm.nih.gov/nlmcatalog?term=2221-1691[ISSN]) ?

reply



**Melanie Ortiz** 3 months ago

SCImago Team

Dear Nafis,

Thank you for contacting us.

SJR is a portal with scientometric indicators of journals indexed in Scopus and all the metadata (including the Country) have been provided by Scopus /Elsevier.

Unfortunately, we can not help you regarding your query, we suggest you contact the NCBI Support team.

Best Regards, SCImago Team

**Bo Cui** 9 months ago


The journal is indexed for SCI and has a current IF of about 1.7

reply



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: [www.elsevier.com/locate/apjtb](http://www.elsevier.com/locate/apjtb)Original article <http://dx.doi.org/10.1016/j.apjtb.2017.06.003>

## Antiviral activity of the dichloromethane extracts from *Artocarpus heterophyllus* leaves against hepatitis C virus



Achmad Fuad Hafid<sup>1,2\*</sup>, Chie Aoki-Utsubo<sup>3</sup>, Adita Ayu Permanasari<sup>1</sup>, Myrna Adianti<sup>1</sup>, Lydia Tumewu<sup>1</sup>, Aty Widyawaruyanti<sup>1,2</sup>, Sri Puji Astuti Wahyuningsih<sup>6</sup>, Tutik Sri Wahyuni<sup>1,2</sup>, Maria Inge Lusida<sup>1,4</sup>, Soetjipto<sup>1,5</sup>, Hak Hotta<sup>7</sup>

<sup>1</sup>Institute of Tropical Disease, Airlangga University, Surabaya 60115, Indonesia

<sup>2</sup>Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Airlangga University, Surabaya 60286, Indonesia

<sup>3</sup>Department of International Health, Kobe University Graduate School of Health Sciences, 7-10-2, Tomogaoka, Suma-ku, Kobe 654-0142, Japan

<sup>4</sup>Department of Microbiology, Faculty of Medicine, Airlangga University, Surabaya 60131, Indonesia

<sup>5</sup>Department of Biochemistry, Faculty of Medicine, Airlangga University, Surabaya 60131, Indonesia

<sup>6</sup>Department of Biology, Faculty of Science and Technology, Airlangga University, Surabaya 60115, Indonesia

<sup>7</sup>Department of Oral Vaccine and Drug Development, Kobe University Graduate School of Health Sciences, 1-5-6 Minatojima-minamimachi, Chou-ku, Kobe 650-0047, Japan

### ARTICLE INFO

#### Article history:

Received 22 May 2017

Accepted 19 Jun 2017

Available online 23 Jun 2017

#### Keywords:

Hepatitis C virus

*Artocarpus* sp.

*Artocarpus heterophyllus*

Antiviral

### ABSTRACT

**Objective:** To determine anti-viral activities of three *Artocarpus* species: *Artocarpus altilis*, *Artocarpus camansi*, and *Artocarpus heterophyllus* (*A. heterophyllus*) against Hepatitis C Virus (HCV).

**Methods:** Antiviral activities of the crude extracts were examined by cell culture method using Huh7it-1 cells and HCV genotype 2a strain JFH1. The mode of action for anti-HCV activities was determined by time-of-addition experiments. The effect on HCV RNA replication and HCV accumulation in cells were analyzed by quantitative reverse transcription-PCR and western blotting, respectively.

**Results:** The dichloromethane (DCM) extract of *A. heterophyllus* exhibited strong anti-HCV activity with an inhibitory concentration (IC<sub>50</sub>) value of (1.5 ± 0.6) µg/mL without obvious toxicity. The DCM extracts from *Artocarpus altilis* and *Artocarpus camansi* showed moderate anti-HCV activities with IC<sub>50</sub> values being (6.5 ± 0.3) µg/mL and (9.7 ± 1.1) µg/mL, respectively. A time-of-addition studies showed that DCM extract from *A. heterophyllus* inhibited viral entry process though a direct virucidal activity and targeting host cells. HCV RNA replication and HCV protein expression were slightly reduced by the DCM treatment at high concentration.

**Conclusions:** The DCM extract from *A. heterophyllus* is a good candidate to develop an antiviral agent to prevent HCV reinfection following liver transplantation.

## 1. Introduction

Hepatitis C Virus (HCV) infection is a major health problems that lead to liver diseases such as chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma. Approximately 170 million people of world population are chronically infected with HCV [1–3]. HCV exhibits high genetic diversity and different genotypes which are classified into seven (1–7) genotypes with 67 confirmed and 20 provisional subtypes [4]. In

\*Corresponding author: Achmad Fuad Hafid, Institute of Tropical Disease, Airlangga University, Surabaya 60115, Indonesia.

Tel: +62 8121644180

Fax: +62 31 5992445

E-mail: [achmadfuad@yahoo.com](mailto:achmadfuad@yahoo.com) (A.F. Hafid).

Peer review under responsibility of Hainan Medical University. The journal implements double-blind peer review practiced by specially invited international editorial board members.



HCV-positive patients, the cumulative risk of developing hepatocellular carcinoma in the 40–74 age group is 21.6% among males and 8.7% among females [5].

HCV is a small enveloped virus with a positive-sense, single-stranded RNA genome that encodes a large polyprotein consisting of three structural proteins and seven nonstructural proteins. The structural proteins of enveloped glycoproteins E1 and E2 are responsible for virus binding to the receptor molecules on cell surface, such as scavenger receptor class B type I (SR-BI), CD81, claudin 1, and occludin [1]. Meanwhile, the nonstructural proteins of p7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B are responsible for viral RNA replication and viral particle construction [6–8].

The current treatment of HCV infection has markedly changed in the recent years. The direct acting antiviral agents (DAAs) combined with Interferon (IFN) have been approved and new IFN-free regimen combinations are recently available in many countries. DAAs targets nonstructural proteins of HCV resulted in the disruptions of the viral replication and infection. Currently approved DAAs consist of NS3 protease inhibitors such as simeprevir, asunaprevir and vaniprevir; NS5A inhibitors such as daclatasvir and ledipasvir; and NS5B RNA-dependent RNA polymerase (RdRp) inhibitors such as sofosbuvir [9,10]. The current treatment regimen using DAAs has dramatically improved sustained virological response (SVR) in most patients of different HCV genotypes. However, the emergence of drug resistance virus, safety for long usage, expensive cost of DAAs therapy, and limited access to the treatment, especially for patients in countries with relatively low income remain major barriers to HCV treatment. Thus, development of effective and inexpensive anti-HCV agents is still required.

Tropical rainforests exhibit a vast diversity in plants and those plants are sources for potential drug development. It has been previously reported that anti-HCV activities of Indonesian medicinal plants, in which 4 out of 21 plants extracts revealed anti-HCV activity against the HCV J6/JFH1-P47. One of the four plants was *Ficus fistulosa* known as Moraceae family [11]. Moraceae family consists of 60 genera and includes 1400 species. The important genus of the Moraceae family is *Artocarpus* which is composed of 50 species [12]. *Artocarpus* is known to have wide bioactivities against virus [13,14], bacterial [15,16], malarial [17,18], and fungi [19,20].

In this study, three species of *Artocarpus* from Purwodadi Botanical Gardens, East Java, Indonesia, namely *Artocarpus altilis* (*A. altilis*) (breadfruit), *Artocarpus camansi* (*A. camansi*) (breadnut), and *Artocarpus heterophyllus* (*A. heterophyllus*) (jackfruit) were screened for anti-HCV activities.

## 2. Materials and methods

### 2.1. Cells and viruses

A clone of human hepatocellular carcinoma-derived Huh7 cells, Huh7it-1 [21], was cultivated in Dulbecco's Modified Eagle Medium (GIBCO Invitrogen, Carlsbad, CA, USA) supplemented with 10% Fetal Bovine Serum (Biowest, Nualle, France), 0.15 mg/mL Kanamycin (Sigma-Aldrich, St. Louis, MO, USA) and non-essential amino acids (GIBCO-Invitrogen) in 5% CO<sub>2</sub> at 37 °C. A cell culture-adapted HCV variant was propagated as described previously [9]. In brief, Huh7it-1 cells ( $1.8 \times 10^7$  cells) were infected with JFH1  $1.8 \times 10^7$  focus-forming unit (ffu) for 4 h with agitation in every 30 min. The HCV-infected cells were incubated for 5 d. The supernatants at

day 3 post-infection were collected and used for antiviral experiments.

### 2.2. Preparation of crude extracts

The leaves of *A. altilis*, *A. camansi*, and *A. heterophyllus* were obtained from Purwodadi Botanical Garden, Indonesia and verified by a licensed botanist. These *Artocarpus* leaves were extracted with several solvents (ethanol 80%, hexane, dichloromethane and methanol). The leaves were extracted using *n*-hexane and ethanol 80%. Meanwhile, the residue from *n*-hexane extract was further extracted using dichloromethane (DCM). Thereafter, the residue from dichloromethane was extracted using methanol. All extracts were dissolved in dimethyl sulfoxide (DMSO) at a concentration of 100 mg/mL and then stored at –30 °C.

### 2.3. Virus titration and immunostaining

Virus titration and immunostaining was performed as described previously [9,21,22]. In brief, virus supernatants diluted in the medium and inoculated onto the Huh7it-1 cells. After virus absorption for 4 h, the cells were cultured with medium containing 0.4% methyl cellulose (Sigma-Aldrich) for 41 h. The cells were fixed with 10% formaldehyde solution and permeabilized with 0.5% triton X-100 in PBS. The cells were stained with anti-HCV patient anti-serum and HRP-goat anti-human Ig antibody (MBL). The HCV antigen positive cells were visualized with Metal Enhanced DAB substrate kits (Thermo Fisher Scientific, Rockford, USA) and infectious foci were counted under microscope.

### 2.4. Antiviral activity assay

Antiviral activity assay was performed as described previously [9,21,22]. Huh7it-1 cells ( $5.2 \times 10^4$ ) were inoculated with HCV at multiplication of infection (MOI) of 0.1 in the presence of different concentrations of plant extracts (100, 50, 25, 12.5, 6.3 and 3.1 µg/mL). After virus absorption for 2 h, the cells were rinsed with the medium and were further incubated in the medium containing the same extracts for 46 h. For time-of-addition experiments, the cells were treated with the medium containing extracts only during viral inoculation (entry event) or only after viral inoculation (post entry event). Culture supernatants at 48 h post-infection were collected for virus titration. The 50% inhibitory effect (IC<sub>50</sub>) was calculated by SPSS probit analysis.

### 2.5. Virucidal activity assay

Virucidal activity test was performed as described previously [9]. In brief, the HCV suspension ( $10^6$  ffu/mL, 75 µL) was mixed with an equal volume of DCM extract and incubated for 2 h at 37 °C. Following by inoculated the virus suspension to the cells and incubated for 4 h at 37 °C. After removing viral inoculum, the cells were overlaid with 0.5% methyl cellulose-containing medium and incubated for 41 h.

### 2.6. Effect of host expression assay

The extract of DCM from *A. heterophyllus* was preincubated with cell ( $5.2 \times 10^4$ ) for 2 h at 37 °C. Then, cells were inoculated with HCV (MOI of 0.1) for 4 h. After viral absorption, cells were replaced with medium and incubated for 41 h. The culture supernatant was collected for virus titration and immunostaining.



## 2.7. Immunoblotting

The cells were lysed in a sodium dodecyl sulfate (SDS) sample buffer and the protein concentrations were determined using a bicinchoninic protein assay kit (Thermo Fisher Scientific). Equal amounts of proteins were separated in SDS-polyacrylamide gel electrophoresis and transferred onto a polyvinylidene difluoride membrane (Millipore, Bedford, MA, USA). The membranes were probed with an HCV NS3 mouse monoclonal antibody (clone H23; Abcam, Cambridge, MA, USA), an HCV NS5A mouse monoclonal antibody (clone 7B5; Biofront, Tallahassee, FL) or glyceraldehyde-3-phosphate dehydrogenase (GAPDH) antibody (MBL, Nagoya, Japan) as primary antibodies followed by HRP-conjugated goat anti-mouse immunoglobulin (MBL) as the secondary antibody. Target proteins were visualized using enhanced chemiluminescence detection system (Biorad; GE healthcare, UK).

## 2.8. Quantitative reverse transcription-polymerase chain reaction (qRT-PCR)

RNA extraction, cDNA synthesis, and qRT-PCR were performed as described previously [9]. In brief, RNA was extracted from cells using TRIzol reagent (Invitrogen, Carlsbad, CA, USA). One  $\mu\text{g}$  of total RNA was transcribed using a ReverTra Ace qPCR RT Kit (Toyobo, Osaka, Japan) with random primers, and cDNA was amplified for real-time quantitative PCR using SYBR Premix Ex Taq (Takara Bio, Shiga, Japan) in a MicroAmp 96-well plate. PCR was performed using ABI 7300 Real-Time PCR system with specific primers to amplify an NS3 region of the HCV genome 5'-CTTTGACTCCGTGATCGACT-3' (sense) and 5'-CCCTGTCTTCCTCTACCTG-3' (antisense).

## 2.9. MTT assay

The cytotoxicity of the samples was assessed by 3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide (MTT) assay as described previously [9]. In brief, cells in 96 well plates were treated with various concentrations of crude extracts for 48 h. The medium was replaced with MTT containing medium and incubated for 4 h. Insoluble precipitates were dissolved with DMSO and the absorbance at 560 nm was measured using a microplate reader. The percentages of cell viability were compared to the control and calculated for 50% cytotoxic concentration ( $\text{CC}_{50}$ ) values.

## 2.10. Data analysis

Results were expressed as mean  $\pm$  SD. Differences between two data sets were evaluated by Student's two-tailed *t*-test. A *P*-value of  $< 0.05$  was considered as statistically significant.

## 3. Results

### 3.1. Anti-HCV activities of *A. altalis*, *A. heterophyllus*, and *A. camansi*

We prepared crude extract samples from three *Artocarpus* species (*A. altalis*, *A. heterophyllus*, and *A. camansi*) using four various solvents (80% ethanol, *n*-hexane, DCM, and methanol) and obtained a total of 12 samples. Those samples were used

Table 1

Anti-HCV activity ( $\text{IC}_{50}$ ) and cytotoxicity ( $\text{CC}_{50}$ ) of *A. altalis*, *A. heterophyllus* and *A. camansi*.

Sample		$\text{IC}_{50}$ ( $\mu\text{g/mL}$ )	$\text{CC}_{50}$ ( $\mu\text{g/mL}$ )	SI
<i>A. altalis</i>	80% Ethanol	$8.9 \pm 0.3$	$> 50$	$> 5.7$
	Hexane	$> 100$	$> 500$	NA
	DCM	$6.5 \pm 0.3$	$> 50$	$> 7.8$
	Methanol	$10.7 \pm 1.6$	$> 200$	$> 18.6$
<i>A. heterophyllus</i>	80% Ethanol	$12.9 \pm 2.6$	$> 800$	$> 62.1$
	Hexane	$> 100$	$> 400$	NA
	DCM	$1.5 \pm 0.6$	$> 200$	$> 134.8$
	Methanol	$6.8 \pm 0.8$	$> 600$	$> 88.6$
<i>A. camansi</i>	80% Ethanol	$6.7 \pm 0.9$	$> 50$	$> 7.5$
	Hexane	$> 100$	$> 500$	NA
	DCM	$9.7 \pm 1.1$	$> 50$	$> 5.2$
	Methanol	$13.0 \pm 0.7$	$> 100$	$> 7.8$


Data represent mean  $\pm$  SD of data from triplicate experiments. NA, not applicable, SI, selectivity index.

for antiviral screening against HCV (JFH1 strain). HCV was inoculated onto Huh7it-1 cells in the presence of the samples and after viral adsorption for 2 h, the cells were extensively rinsed and further incubated in the same extracts containing medium for 46 h. The 50% HCV inhibition concentration ( $\text{IC}_{50}$ ) and the 50% cytotoxic concentration ( $\text{CC}_{50}$ ) and selectivity indexes (SI:  $\text{CC}_{50}/\text{IC}_{50}$ ) of tested samples are shown in Table 1. The results showed that 6 of the 12 samples possessed strong anti-HCV activities with  $\text{IC}_{50}$  value of  $< 10 \mu\text{g/mL}$  and 3 extracts moderate activities with  $\text{IC}_{50}$  value of  $(10\text{--}20) \mu\text{g/mL}$ . Among samples possessing anti-HCV activity, the DCM extract of *A. heterophyllus* exhibited the strongest activity with an  $\text{IC}_{50}$  value of  $1.5 \mu\text{g/mL}$  and  $\text{CC}_{50} > 200 \mu\text{g/mL}$  (SI:  $> 134.8$ ). Methanol extract of *A. heterophyllus* and 80% ethanol extract of *A. heterophyllus* showed anti-HCV activities with  $\text{IC}_{50}$  values of  $6.8 \mu\text{g/mL}$  and  $12.9 \mu\text{g/mL}$ , respectively without any cytotoxicity effect with  $\text{CC}_{50}$  values  $> 600 \mu\text{g/mL}$  and  $> 800 \mu\text{g/mL}$  and SI values of 88.6 and 62.1, respectively. The DCM extracts of *A. altalis* and *A. camansi* revealed stronger anti-HCV activity with  $\text{IC}_{50}$  values of  $6.5 \mu\text{g/mL}$  and  $9.7 \mu\text{g/mL}$ , respectively, with  $\text{CC}_{50}$  value of  $> 50 \mu\text{g/mL}$ . On the other hand, the hexane extracts of the three *Artocarpus* species did not exhibit significant anti-HCV activities at the concentration of  $100 \mu\text{g/mL}$ . Dose-dependent inhibition of HCV infection and cell viability of each sample was shown in Figure 1.

### 3.2. Mode of action of extracts of *A. altalis*, *A. heterophyllus* and *A. camansi*

To determine the inhibitory stage(s) of extracts possessing anti-HCV activities in HCV life cycle, we conducted time-of-addition experiments, in which three sets of experiments were done in parallel: First, the extract(s) and virus were co-added onto the cells for 2 h and after virus adsorption, the cells were further incubated in the presence of the same extracts for 46 h (treatment during entry and post-entry steps). Second, the extract(s) and virus were co-added onto the cells for 2 h and after virus adsorption, the cells were further incubated in the absence of the extracts for 46 h (treatment during entry step). Third, the HCV virus without sample was inoculated onto the culture cells. After virus adsorption for 2 h, the infected cells were incubated





**Figure 1.** Dose-dependent inhibition of HCV infection and cell viability from *A. heterophyllum*, *A. camansi*, and *A. altilis*.

Percent inhibition of HCV infection and cell viability by the extracts compared to the untreated control are shown. Data represent mean ± SD from three independent experiments. DCM: dichloromethane, MeOH: methanol EtOH: 80% ethanol.

with media containing same extracts for 46 h (treatment after virus entry). The results showed that all the three extracts (80% ethanol, DCM, methanol) of *A. altilis* and *A. camansi* exerted anti-HCV activity mainly at the post-entry event (Table 2). In contrast, the extracts of 80% ethanol, DCM and methanol of *A. heterophyllum* exhibited HCV inhibition principally at viral entry, while the post entry steps showed the lesser extent (Table 2).

### 3.3. Thin layer chromatography (TLC) analysis of extracts of *A. altilis*, *A. heterophyllum*, and *A. camansi*

Bioactivities of medicinal plants were influenced by the chemical contents of the plants.

Screenings of the bioactive components in the extracts were performed by TLC analysis and the result was demonstrated in Figure 2. The DCM, methanol and 80% ethanol extracts of

*A. altilis* and *A. camansi* showed to contain with flavonoid component which is indicated by one major orange spot. On the other hand, TLC of methanol and DCM extracts of *A. heterophyllum* showed the presence of terpenoid and steroid which are indicated by purple and blue spots, respectively. DCM extract of *A. heterophyllum* also contained chlorophyll-related compounds with detection of red spots under detection of UV irradiation 365 nm. While the hexane extracts of *A. altilis*, *A. heterophyllum*, and *A. camansi* showed the presence of terpenoids as one major spots.


### 3.4. DCM extract of *A. heterophyllum* inhibits HCV infection through a direct virucidal effect and affecting host cells

Since the strong inhibition of DCM extract, further analysis to conduct the mechanism of action was performed for DCM

**Table 2**

Mode of action of crude extracts from *A. altilis*, *A. heterophyllum*, and *A. camansi*.


Sample	Solvent	Conc. (µg/mL)	% Inhibition			Mode of action
			During + post inoculation	During inoculation	Post inoculation	
<i>A. altilis</i>	80% Ethanol	30	97.5	3.1	98.1	Post-entry inhibition
	DCM	30	99.0	7.7	98.8	Post-entry inhibition
	Methanol	30	92.4	9.7	90.1	Post-entry inhibition
<i>A. heterophyllum</i>	80% Ethanol	25	89.4	56.7	15.0	entry inhibition
	DCM	6	91.0	81.1	22.5	entry inhibition
	Methanol	25	95.5	72.3	19.9	entry inhibition
<i>A. camansi</i>	80% Ethanol	30	97.1	10.6	93.9	Post-entry inhibition
	DCM	30	97.0	12.9	96.4	Post-entry inhibition
	Methanol	30	94.0	17.3	95.3	Post-entry inhibition



**Figure 2.** TLC analysis of extracts of *A. altilis*, *A. heterophyllum* and *A. camansi*.

Silica gel F254 TLC was used as stationary phase and chloroform:methanol (9:1, v/v) as mobile phase. Detection under (A) UV 254 nm, (B) UV 365 nm, (C) heating TLC plate at 105 °C for 5 min after spraying 10% sulfuric acid and (D) observed under UV 365 nm after using spray reagent and heated. Sample: (1) hexane extract of *A. heterophyllum*, (2) DCM extract of *A. heterophyllum*, (3) methanol extract of *A. heterophyllum*, (4) 80% ethanol extract of *A. heterophyllum*, (5) hexane extract of *A. altilis*, (6) DCM extract of *A. altilis*, (7) methanol extract of *A. altilis*, (8) 80% ethanol extract of *A. altilis*, (9) hexane extract of *A. camansi*, (10) DCM extract of *A. camansi*, (11) methanol extract of *A. camansi*, (12) 80% ethanol extract of *A. camansi*.

extract of *A. heterophyllum*. Time-of-addition experiment to determine the effect of extracts in the entry or post entry steps of HCV life cycle showed that the DCM extract of *A. heterophyllum* exerts anti-HCV activity at mainly in the entry-step. To evaluate whether the DCM extract acted on the target of HCV or host cells, the infectivity of the extract-treated HCV virions were determined. HCV virions were treated with the DCM extract or medium as the control for 2 h at 37 °C and then pretreated-HCV was inoculated onto Huh7it-1 cells for virus titration. As shown in Figure 3A, pretreatment of HCV inoculum with the DCM extract (6.3 µg/mL) was significantly decreased the HCV infectivity upon to 18.2% ( $P < 0.00001$ ) compared with the untreated control (81.8% reduction). We next assessed the effect of DCM-pretreated host cells on HCV infection. Huh7it-1 cells were pretreated with the DCM extract for 2 h and then rinsed extensively to remove the extract. The pre-treated cells were challenged with HCV infection in the absence of the DCM extract. The result showed that the pretreatment of the cells with DCM extract (6.3 µg/mL) was significantly decreased HCV infectivity upon 9.9% ( $P < 0.00001$ ) compared with the untreated control (90.1% reduction) (Figure 3B). These results suggested that the DCM extract of *A. heterophyllum* targeted both HCV virion and host cells. Since treatment of cells with the DCM extracts (6.3 µg/mL) revealed the effect at the post-viral adsorption step and also somehow inhibited HCV infection (Table 2), further confirmation of the effect on HCV RNA replication and HCV protein accumulation in the cells was evaluated. Real-time RT-PCR and immunoblotting analysis demonstrated that low concentration of DCM extract did not clearly suppress HCV replication and HCV protein accumulation, however high concentrations of the DCM ( $> 12.5$  µg/mL) acted to inhibit HCV replication (Figure 3C and D).



**Figure 3.** (A) Analysis of virucidal activity. HCV suspension was mixed with the DCM extract of *A. heterophyllum* for 2 h at 37 °C before inoculation onto the cells. (B) Effect of pretreatment of cells with DCM extracts of *A. heterophyllum* on HCV infection.

Cells were preincubated with the DCM extracts for 2 h and then challenged with HCV infection. (C) The HCV-infected cells were treated with the DCM extracts of *A. heterophyllum*. HCV protein accumulation in the cells was analyzed by western blotting against NS3, NS5A or GAPDH as a loading control. (D) The level of HCV RNA in the cells was measured by qRT-PCR. Data represent means from triplicate experiments  $\pm$  SD. \* $P < 0.001$  compared to the untreated control; \*\* $P < 0.00001$  compared to the untreated control.







#### 4. Discussion

Medicinal plants are potential resources for various bioactivities. Several components from medicinal plants have been reported to possess potential bioactivity including anti-HCV. Plants of *Artocarpus* genus have been used as traditional medicine in Indonesia for the treatment of fever, dysentery, and malaria. The genus of *Artocarpus* is rich in phenolic compounds, including flavonoid, stilbenoids, arylbenzofurans, and Jacalin (a lectin) [23–25] that were reported to possess a wide range of biological activities including anticancer, anti-inflammatory, antihypertensive, antibacterial, and antiviral [26].

In the present study, we screened crude extracts of three *Artocarpus* species: *A. altilis*, *A. camansi*, and *A. heterophyllus* for anti-HCV activities. A plant of *A. altilis* is known as breadfruit, which was used traditionally to treat liver disorders, hypertension, and diabetic. A total of 130 compounds were identified from *A. altilis*, of which more than 70 are derived from the prenylpropanoid pathways [25]. Meanwhile *A. camansi* is known with local name breadnut. The morphology of *A. altilis* and *A. camansi* is similar for leaves, fruits, and stems [25]. *A. camansi* is also believed to have similar medicinal properties to *A. altilis* [24].

Plant of *A. heterophyllus* has a local name Jackfruit and the leaves are usually entire (without lobes), much smaller than breadfruit and breadnut leaves. *A. heterophyllus* was known active as antibacterial activity against 24 species of bacteria [27]. Jacalin, a Jackfruit lectin from *A. heterophyllus* was reported to inhibit DNA viruses such as herpes simplex virus type II (HSV-2), varicella-zoster virus (VZV), and cytomegalovirus (CMV) [28], however, there is no reported yet about its anti-HCV activities.

Anti-HCV activities were demonstrated that 80% ethanol, DCM, and methanol extract of *Artocarpus* species mediated strong inhibition against HCV with  $IC_{50}$  value less than 15 g/mL, while the hexane extracts did not access any anti-HCV activities in the concentration of 100 g/mL (Table 1 and Figure 1). Further analysis was demonstrated that extracts of *A. altilis* and *A. camansi* exhibited HCV inhibition mainly at the post-entry step with percentage inhibition higher than 90%, while extracts of *A. heterophyllus* inhibit HCV in the entry step with percentage inhibition higher than 80% (Table 2). Common constituent(s) present in the extracts of *A. camansi* and *A. altilis* may exert similar anti-HCV activities.

To confirm the mechanism of *A. heterophyllus* how to inactivate the virus, virucidal activity and effect of host expression test were conducted. The result showed that pre-treatment of the HCV virion or the Huh7it-1 cells with DCM extract of *A. heterophyllus* strongly reduced HCV infection. It suggested that DCM extract of *A. heterophyllus* exerts antiviral activities through direct virucidal activity and effecting host cells (Figure 2A and B) which may interfere the interaction with some receptors in the host cells. Some host cell molecules have been reported to be important entry factors or (co)receptors for HCV, such as glycosaminoglycan (GAG), low density lipoprotein receptor (LDLR), the scavenger receptor class B member I (SR-BI), the tetraspanin CD81, claudin-1 (CLDN1), and occludin (OCLN) which play as necessary keys in the attachments process of HCV to the host cells [29]. Since antiviral drug(s) targeting host factor(s) is generally known to lower emergence rate of drug resistance compared to the direct-acting antiviral drugs, the DCM extract of *A. heterophyllus* may be useful as a new drug development for the treatment of HCV especially to prevent HCV re-infection following liver transplantation.

Further analysis to confirm the effect of DCM extract of *A. heterophyllus* in the post entry step of HCV life cycle, western blot analysis was performed to examine the expressions of NS3 and NS5A HCV protein levels which played the important role in the replication of HCV. The result demonstrated mild inhibition of NS3 and NS5A protein expression level in DCM extract of *A. heterophyllus*-treated cells. Consistently, the HCV RNA levels were slightly inhibited by DCM extract of *A. heterophyllus*, however significant inhibition of RNA level was observed when the concentration was increased up to 12.5 and 25  $\mu$ g/mL. These results were suggested the possible inhibition process attachment, assembly, release of virions and replication steps.

The bioactivities of components were influenced by the biochemical constituents among the plants. Our studies have not yet identified the compound(s) responsible for anti-HCV activity from *A. altilis*, *A. heterophyllus*, and *A. camansi* in this study. Other study was reported that *A. heterophyllus* contains lectin, artocarpine, artocarpetin, cycloheterophyllin, artonins A, morin, oxydihydroartocarpesin, cynomacurin, isoartocarpin, cyloartocarpin, artocarpesin, norartocarpetin, cycloartinone and artocarpone. The leaves and stem are also reported to contain sapogenins, cycloartenone, cycloartenol, sitosterol and tannins [24].

Our TLC profiles were identified terpenoid and steroid components which served as major components of the DCM extracts of *A. heterophyllus* (Figure 2). The DCM, methanol and 80% ethanol extracts of *A. altilis* and *A. camansi* contained flavonoid as a major component. Some terpenoids have been reported to inhibit HCV infection such as Saikosaponin b2 from *Bupleurum kaoi* [30]; oleanic acid and ursolic acid [31]; Platycodin D, D2, D3, deapioplatycodin D, D2, platyconic acid A [32]; and andrographolide, a diterpenoid lactone from *Andrographis paniculata* [33]. Many flavonoids have also been reported to exert anti-HCV activity: Epigallocatechin-3-gallate (EGCG) [34], quercetin, luteolin, apigenin and ladanin [35], naringenin and silymarin/silibinin. The grapefruit flavonoid naringenin was reported to inhibit HCV assembly and release. Silibinin, the major component of silymarin, was reported to exert anti-HCV activity by blocking HCV entry, HCV fusion, replication and production of new progeny virus. These compounds are currently in phase I and phase 2/3 clinical trial studies, respectively [36,37]. As for the *A. altilis*, *A. heterophyllus*, and *A. camansi*, further analyses is required to determine the responsible compound(s) for anti-HCV activities in their extracts.

Extracts of *A. altilis*, *A. camansi*, and *A. heterophyllus* possess anti-HCV activity. The DCM extract of *A. heterophyllus* exhibits strong anti HCV activity through a direct virucidal activity and effecting host cells. The DCM extract of *A. heterophyllus* was a good candidate to develop a new antiviral agent to treat HCV infection and to prevent HCV re-infection following liver transplantation.

#### Conflict of interest statement

The authors declare no conflict of interests.

#### Acknowledgements

We thank to Dr. Takaji Wakita (National Institute of Infectious Diseases, Tokyo, Japan) for providing pJFH-1 and





Dr. Yohko Shimizu for providing Huh7it-1 cells. This work was supported in part by Mandat Project Airlangga University and Science and Technology Research Partnerships for Sustainable Development (SATREPS) program from Japan Science and Technology Agency (JST) and Japan International Cooperation Agency (JICA).

## References

- [1] Dubuisson J, Cosset FL. Virology and cell biology of the hepatitis C virus life cycle: an update. *J Hepatol* 2014; **61**(Suppl 1): S3-13.
- [2] Lange CM, Jacobson IM, Rice CM, Zeuzem S. Emerging therapies for the treatment of hepatitis C. *EMBO Mol Med* 2014; **6**(1): 4-15.
- [3] Pawlotsky JM. Treatment of chronic hepatitis C: current and future. *Curr Top Microbiol Immunol* 2013; **369**: 321-42.
- [4] Messina JP, Humphreys I, Flaxman A, Brown A, Cooke GS, Pybus OG, et al. Global distribution and prevalence of hepatitis C virus genotypes. *Hepatology* 2015; **61**(1): 77-87.
- [5] Tanaka J, Kumagai J, Komiya Y, Yoshizawa H. Area and age specific prevalence rate of HCV infection in Japan. *Nippon Rinsho* 2004; **62**(Suppl 7(Pt 1)): 253-7.
- [6] Penin F, Dubuisson J, Rey FA, Moradpour D, Pawlotsky JM. Structural biology of hepatitis C virus. *Hepatology* 2004; **39**(1): 5-19.
- [7] Moradpour D, Penin F, Rice CM. Replication of hepatitis C virus. *Nat Rev Microbiol* 2007; **5**(6): 453-63.
- [8] Rupp D, Bartenschlager R. Targets for antiviral therapy of hepatitis C. *Semin Liver Dis* 2014; **34**(1): 9-21.
- [9] Apriyanto DR, Aoki C, Hartati S, Hanafi M, Kardono LB, Arsianti A, et al. Anti-hepatitis c virus activity of a crude extract from longan (*Dimocarpus longan* Lour.) leaves. *Jpn J Infect Dis* 2016; **69**(3): 213-20.
- [10] Wahyuni TS, Aoki C, Hotta H. Promising anti-virus hepatitis C compounds from natural resources. *Nat Prod Commun* 2016; **11**(8): 1193-200.
- [11] Wahyuni TS, Tumewu L, Permanasari AA, Apriani E, Adianti M, Rahman A, et al. Antiviral activities of Indonesian medicinal plants in the east java region against hepatitis C virus. *Virol J* 2013; **10**: 259.
- [12] Verheij EWM, Coronel RE. *Plant resources of south-east Asia*. Bogor, Indonesia: Prosea; 1992.
- [13] Likhitwitayawuid K, Sritularak B, Benchanak K, Lipipun V, Mathew J, Schinazi RF. Phenolics with antiviral activity from *Milletia erythrocalyx* and *Artocarpus lakoocha*. *Nat Prod Res* 2005; **19**(2): 177-82.
- [14] Sasivimolphan P, Lipipun V, Likhitwitayawuid K, Takemoto M, Pramyothin P, Hattori M, et al. Inhibitory activity of oxyresveratrol on wild-type and drug-resistant varicella-zoster virus replication in vitro. *Antivir Res* 2009; **84**(1): 95-7.
- [15] Khan MR, Omoloso AD, Kihara M. Antibacterial activity of *Artocarpus heterophyllus*. *Fitoterapia* 2003; **74**(5): 501-5.
- [16] Sato M, Tsuchiya H, Miyazaki T, Fujiwara S, Yamaguchi R, Kureshiro H, et al. Antibacterial activity of hydroxychalcone against methicillin-resistant *Staphylococcus aureus*. *Int J Antimicrob Agents* 1996; **6**(4): 227-31.
- [17] Widyawaruyanti A, Surya S, Kalauni K, Awale S, Nindatu M, Zaini NC, et al. New prenylated flavones from *Artocarpus champe-den*, and their antimalarial activity in vitro. *J Nat Prod* 2007; **61**: 410-3.
- [18] Boonlaksiri C, Oonanant W, Kongsaree P, Kittakoop P, Tanticharoen M, Thebtaranonth Y. An antimalarial stilbene from *Artocarpus integer*. *Phytochemistry* 2000; **54**(4): 415-7.
- [19] Jayasinghe L, Balasooriya BA, Padmini WC, Hara N, Fujimoto Y. Geranyl chalcone derivatives with antifungal and radical scavenging properties from the leaves of *Artocarpus nobilis*. *Phytochemistry* 2004; **65**(9): 1287-90.
- [20] Trindade MB, Lopes JL, Soares-Costa A, Monteiro-Moreira AC, Moreira RA, Oliva ML, et al. Structural characterization of novel chitin-binding lectins from the genus *Artocarpus* and their anti-fungal activity. *Biochim Biophys Acta* 2006; **1764**(1): 146-52.
- [21] Aoki C, Hartati S, Santi MR, Lydwina T, Firdaus R, Hanafi, et al. Isolation and identification of substances with anti-hepatitis C virus activities from *Kalanchoe pinnata*. *Int J Pharm Pharm Sci* 2014; **6**(2): 211-5.
- [22] Wahyuni TS, Widyawaruyanti A, Lusida MI, Fuad A, Soetjipto, Fuchino H, et al. Inhibition of hepatitis C virus replication by chalepin and pseudane IX isolated from *Ruta angustifolia* leaves. *Fitoterapia* 2014; **99**: 276-83.
- [23] Jagtap UB, Bapat VA. *Artocarpus*: a review of its traditional uses, phytochemistry and pharmacology. *J Ethnopharmacol* 2010; **129**(2): 142-66.
- [24] Hari A, Revikumar KG, Divya D. *Artocarpus*: a review of its phytochemistry and pharmacology. *J Pharma Search* 2014; **9**(1): 7-12.
- [25] Sirkawar MS, Hui BJ, Subramaniam K, Valeisamy BD, Yean LK, Balaji K. A review on *Artocarpus altilis* (Parkinson) Fosberg (breadfruit). *J Appl Pharm Sci* 2014; **4**(8): 91-7.
- [26] Swami SB, Thakor NJ, Haldankar PM, Kalse SB. Jackfruit and its many functional components as related to human health: a review. *Compr Rev Food Sci Food Saf* 2012; **11**(6): 565-76.
- [27] Septama AW, Panichayupakaranant P. Antibacterial assay-guided isolation of active compounds from *Artocarpus heterophyllus* heartwoods. *Pharm Biol* 2015; **53**(11): 1608-13.
- [28] Wetprasit N, Threesangsri W, Klamklai N, Chulavatnatol M. Jackfruit lectin: properties of mitogenicity and the inhibition of herpesvirus infection. *Jpn J Infect Dis* 2000; **53**(4): 156-61.
- [29] Qian XJ, Zhu YZ, Zhao P, Qi ZT. Entry inhibitor: new advances in HCV treatment. *Emerg Microbes Infect* 2016; **5**: e3.
- [30] Lin C. HCV NS3-4A serine protease. In: Tan SL, editor. *Hepatitis C viruses: genomes and molecular biology*. UK: Norfolk; 2006.
- [31] Kong L, Li S, Liao Q, Zhang Y, Sun R, Zhu X, et al. Oleanolic acid and ursolic acid: novel hepatitis C virus antivirals that inhibit NS5B activity. *Antivir Res* 2013; **98**(1): 44-53.
- [32] Kim JW, Park SJ, Lim JH, Yang JW, Shin JC, Lee SW, et al. Triterpenoid saponins isolated from *Platycodon grandiflorum* inhibit hepatitis C virus replication. *Evid Based Complement Altern Med* 2013; **2013**: 560417.
- [33] Lee JC, Tseng CK, Young KC, Sun HY, Wang SW, Chen WC, et al. Andrographolide exerts anti-hepatitis C virus activity by up-regulating haeme oxygenase-1 via the p38 MAPK/Nrf2 pathway in human hepatoma cells. *Br J Pharmacol* 2014; **171**: 237-52.
- [34] Ciesek S, Hahn TV, Colpitts CC, Schang LM, Friesland M, Steinmann J, et al. The green tea polyphenol, Epigallocatechin-3-gallate, inhibits hepatitis C virus entry. *Hepatology* 2011; **54**(6): 1947-55.
- [35] Calland N, Dubuisson J, Rouille Y, Seron K. Hepatitis c virus and natural compound: a new antiviral approach? *Viruses* 2012; **4**: 2197-217.
- [36] Wagoner J, Morishima C, Graf TN, Oberlies NH, Teissier E, Pecqueur EI, et al. Differential in vitro effects of intravenous versus oral formulations of silibinin on the HCV life cycle and inflammation. *PLoS One* 2011; **6**(1): e16464.
- [37] Zeisel MB, Crouchet E, Baumert TF, Schuster C. Host-targeting agents to prevent and cure hepatitis c virus infection. *Viruses* 2015; **7**(11): 5659-85.