

# INTERNATIONAL JOURNAL OF PHARMACY AND PHARMACEUTICAL SCIENCES

Online ISSN: 0975-1491 | Print ISSN: 2656-0097

[Home](#) [About](#) [Current](#) [Archives](#) [Editorial Board](#) [Submissions](#)  
[Instructions to Authors](#) [Contact Us](#)

Search

[HOME](#) / [ARCHIVES](#) / Vol 7, Issue 1, 2015

## LETTER TO EDITOR

**DRUG PROMOTION IN IRAQ AND ITS NEGATIVE EFFECT ON BOTH PERSONAL AND COUNTRY BUDGET**

EHAB MUDHER MIKHAEL

Pages 1

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

## REVIEW ARTICLE(S)

**CURRENT STATUS OF EPILEPSY IN MALAYSIA AND WAY AHEAD**

ALINA ARULSAMY, BEY HING GOH, MOHD FAROOQ SHAIKH

Pages 2-5

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

**PHARMACOLOGICAL PROFILE OF MANGROVE ENDOPHYTES - A REVIEW**

SALINI G

Pages 6-15

[View PDF](#) [Abstract](#) [Download PDF](#) [View HTML](#)

 [View PDF](#)  [Abstract](#)  [Download PDF](#)  [View HTML](#)

---

**ANTIOXIDANT AND ANTIINFLAMMATORY EFFECTS OF VITEX NEGUNDO ON FREUNDÄ™S COMPLETE ADJUVANT INDUCED ARTHRITIS**

HARSHA LAD, DEEPTI DIXIT, ANKITA JOSHI, DEEPAK BHATNAGAR

Pages 81-85

 [View PDF](#)  [Abstract](#)  [Download PDF](#)  [View HTML](#)

---

**ANTIPROLIFERATIVE EFFECT AND SELECTIVITY OF SODIUM DICHLORO-BIS [N-PHENYL-5-CHLORO-SALICYLIDENEIMINATO-N, O] RUTHENATE (III) IN VITRO**

EMIRA KAHROVIÄ†, SANDRA KRALJEVIÄ† PAVELIÄ†

Pages 86-88

 [View PDF](#)  [Abstract](#)  [Download PDF](#)  [View HTML](#)

---

**EVALUATION OF THE ANTIOXIDANT POTENTIAL FOR DIFFERENT EXTRACTS OF AL-TAIF POMEGRANATE (PUNICA GRANATUM L) INDUCED BY ATRAZINE AND MALATHION PESTICIDES IN LIVER OF MALE ALBINO MICE**

REHAM Z. HAMZA, ASMAA M. ABD EL-AZEEZ, NAHED A. HUSSIEN

Pages 89-94

 [View PDF](#)  [Abstract](#)  [Download PDF](#)  [View HTML](#)

---

**DEVELOPMENT AND EVALUATION OF ORAL CONTROLLED RELEASE MATRIX TABLETS OF LAMIVUDINE: OPTIMIZATION AND IN VITRO-IN VIVO STUDIES**

NELSON KENNETH, VARADARAJAN PARTHASARATHY, CHIKKANNA NARENDRA, PRAKASAM KALYANI

Pages 95-101

 [View PDF](#)  [Abstract](#)  [Download PDF](#)  [View HTML](#)

---

**EFFECTS OF OLIVE DRYING AND STORAGE ON THE OXIDATIVE STATUS, AROMA, CHLOROPHYLL AND FATTY ACIDS COMPOSITION OF OLIVE OIL**

WISSAL DHIFI, MAHER BEN KHEDHER, SANA BELLILI, CARMEN SADAKA, LARA WAKIM, MARC EL BEYROUTHY, BRAHIM MARZOUK, WISSEM MNIF

Pages 102-108

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

---

### ANTIBACTERIAL ACTIVITY OF FRESH WATER CRAB AND SNAIL AND ISOLATION OF ANTIBACTERIAL PEPTIDES FROM HAEMOLYMPH BY SDS PAGE

PACKIA LEKSHMI N C J, VIVEKA S, ANUSHA S, JEEVA S, RAJA BRINDHA J, SELVA BHARATH M  
Pages 109-114

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

---

### THE INFLUENCE OF SODIUM ORTHOVANADATE ON P85 AND GSK-3 EXPRESSIONS TO THE BLOOD GLUCOSE REGULATION OF TYPE 2 DIABETIC MICE (MUS MUSCULUS) MODEL

RIMA HAYANTY RITONGA, BUDI SUPRPTI, JUNAIDI KHOTIB  
Pages 115-119

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

---

### DIURETIC EFFECT OF THE ETHANOL EXTRACTS OF PHYLLANTHUS ACIDUS L (SKEELS) LEAVES IN WISTAR RATS

SUCI NAR VIKASARI, ELIN YULINAH SUKANDAR, AFIFAH BAMBANG SUTJIATMO, SORAYA RIYANTI  
Pages 120-123

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

---

### FORMULATION OF ORODISPERSIBLE TABLET OF LUFFA ACUTANGULA (L) ROXB USING NOVEL CO-PROCESSED VIA SPRAY DRIED EXCIPIENT

RINA ADRIANY, EFFIONORA ANWAR, RETNOSARI ANDRAJATI, M. HANAFI  
Pages 124-129

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

---

### ANTIPROLIFERATIVE EFFECT OF FLOWER EXTRACTS OF SPILANTHES PANICULATA ON HEPATIC CARCINOMA CELLS

AKANKSHA MISHRA, SUBHADEEP ROY, SIDDHARTHA MAITY, RAJNISH KUMAR YADAV, AMIT KUMAR KESHARI, SUDIPTA SAHA  
Pages 130-134

[View PDF](#)[Abstract](#)[Download PDF](#)[View HTML](#)

---



## THE INFLUENCE OF SODIUM ORTHOVANADATE ON P85 AND GSK-3 EXPRESSIONS TO THE BLOOD GLUCOSE REGULATION OF TYPE 2 DIABETIC MICE (*MUS MUSCULUS*) MODEL

RIMA H. RITONGA<sup>1\*</sup>, BUDI SUPRAPTI<sup>2</sup>, JUNAIIDI KHOTIB<sup>2\*</sup>

<sup>1</sup>Post-Graduate Program, Faculty of Pharmacy, Airlangga University, <sup>2</sup>Department of Clinical Pharmacy, Faculty of Pharmacy, Airlangga University Jln. Dharmawangsa Dalam Surabaya 60286, Indonesia.  
Email: rimahayantyrionga@yahoo.com, junaidi-k@ff.unair.ac.id

Received: 23 Sep 2014 Revised and Accepted: 25 Oct 2014

### ABSTRACT

**Objective:** The present study was designed to investigate the influence of sodium orthovanadate on the P85 (Regulatory Subunit of PI3-Kinase) and GSK-3 (Glycogen Synthase Kinase 3) expressions in skeletal muscle tissue to the decreasing of blood glucose levels of alloxan-induced diabetic mice.

**Methods:** Mice were divided into 5 groups i. e. (1) naive group, (2) diabetic group and (3-5) orthovanadate-treated diabetic groups at the dose of 16, 32 and 64 mg/kg BW respectively. Diabetic mice model was induced by intraperitoneal administration of alloxan monohydrate at the dose of 200 mg/kgBW. Diabetic state was occurred on day 3 after alloxan injection and then started the treatment of sodium orthovanadate for 7 days. The muscular tissue was harvested on day 10 after treatment and was stained using routine histology staining, haematoxylin-eosin for morphological qualitative analysis and immunohistochemical approaches to observe the expressions of P85 and GSK-3 in skeletal muscle.

**Results:** Diabetes condition was shown by the increasing of fasting blood glucose levels from  $59.1 \pm 11.2$  mg/dL to  $310.6 \pm 107.2$  mg/dL on day 3. Administration of sodium orthovanadate at the dose of 16, 32 and 64 mg/kg BW reduced the fasting blood glucose levels after 7 days treatment ( $p < 0.001$ ) at diabetic mice significantly. The results of histology staining showed that sodium orthovanadate improved a necrosis in skeletal muscle cells alloxan-induced diabetic mice. On immunohistochemical approaches, sodium orthovanadate might decreased the P85 expressions and increased the GSK-3 expressions in skeletal muscle cells alloxan-induced diabetic mice ( $p < 0.001$ ).

**Conclusion:** This study reveals that the administration of sodium orthovanadate on diabetic mice led to the attenuation of the PTPase activity. The inhibition cause a decreasing the expression of P85 and increasing of GSK-3 that cause the decreasing of blood glucose levels and an improvement of insulin target cells in the muscle cells.

**Keywords:** Sodium orthovanadate, Diabetes mellitus, P85 (Regulatory Subunit of PI3-Kinase), GSK-3 (Glycogen Synthase Kinase 3).

### INTRODUCTION

Diabetes mellitus (DM) is the most common endocrine disorder in man, currently affecting over 170 million people world-wide and, potentially, over 365 million in the year 2030 [1]. Type 2 diabetes mellitus (T2DM) is the most common form of diabetes worldwide accounting for 90% of cases globally and affecting approximately 4% of the world's adult population [2]. Besides  $\beta$  cell failure, the major pathophysiological event contributing to the development of T2DM is the resistance of target tissues to insulin [1].

Molecular mechanism that may potentially lead to insulin resistance is a disruption in the balance between the amounts of the phosphoinositide 3 kinase (PI3-kinase) subunits, consisting of a regulatory subunit P85, which is tightly associated with a catalytic subunit, P110. Normally, a pool of free P85 monomers exists in stoichiometric excess to the P110 catalytic subunit, with the P85-P110 heterodimer being responsible for the PI3-kinase activity [1].

In insulin-resistant state, significantly increasing the expression of the free P85 monomers and bind to the phosphorylated IRS-1 proteins in insulin signaling pathway, blocking access to P85-P110 heterodimers and subsequently affecting the ability of insulin to stimulate the association of the P85-P110 heterodimer with IRS-1, thus reducing the PI3-kinase insulin signaling. On previous study, the excess in total P85 in skeletal muscle insulin resistance was entirely accounted for by an increase in the free P85 $\alpha$ -specific isoform [3].

Other mechanisms involved in the development of insulin resistance is a reduction of the cells capacity to synthesize and store glycogen by glycogen synthase kinase-3 (GSK-3) activation. GSK-3 was originally identified as a regulator of glycogen synthase (GS), a rate-limiting enzyme that promotes glycogen deposition. In basal condition, stimulation of cells with insulin causes inactivation of GSK-3 through a PI 3-kinase-dependent mechanism. PI 3-kinase-induced activation of PKB (also termed Akt) results in PKB

phosphorylation of both GSK-3 isoforms (Ser9 of GSK-3 $\beta$ ; Ser21 of GSK-3 $\alpha$ ), which inhibits GSK-3 activity.

This leads to the dephosphorylation of substrates including GS, resulting in their functional activation and consequent increased glycogen synthesis. In insulin-resistant state, in the absence of insulin, GSK-3 was activated by tyrosine phosphorilation.

The active GSK-3 will phosphorylate four serine residues in the C-terminal domain of GS and negatively regulate its activity, reduce the capacity of cells to synthesize and store glycogen [4, 5].

Current antidiabetes therapies are generally safe and effective but have certain limitations, i. e. require the functional insulin receptor and non insulin oral antidiabetes therapies additionally require minimum endogenous levels of insulin.

Therefore, insulin-mimetic compounds that act independently of insulin and the insulin receptor may thus represent a novel therapeutic strategy for severe insulin-resistant patients, T1DM, and T2DM [6].

Vanadium compounds act in an insulin-mimetic manner both *in vitro* and *in vivo* have been well established. Vanadium is an ultra trace element, widely distributed in the nature and its compounds would closely mimic the physiological action of insulin by initial entry into the portal system [7].

In this regard and supporting a possible therapeutic use of vanadium, the present study aimed to investigate sodium orthovanadate (SOV), the inorganic salt compound of vanadium, influence in the reduction of P85 expression and the elevation of GSK-3 expression to the reduction of blood glucose levels of alloxan-induced diabetic mice. Therefore, this mechanism will be the basis of new therapeutic strategies in patients with T2DM.



## MATERIALS AND METHODS

### Animals and experimental groups

A group of 25-40 male mice of Balb/C strain, weighing between 20-30 g and 6-8 weeks of age, was obtained from Animal Laboratory at the Pharmacy Faculty, Airlangga University, Indonesia. Animals were maintained in the climatically controlled animal house facility of Animal Laboratory at the Pharmacy Faculty, for one week before the initiation of the experimentation.

The animals were fed *ad libitum* with mice feed and tap water. All of mice were divided into five groups. Group 1 was non-diabetic control mice (n=5). Group 2 was diabetic-untreated control mice (n=5). Group 3 (n=5), Group 4 (n=5) and Group 5 (n=8) were diabetic-treated mice. The methodology of this experiment was performed after the approval by Airlangga University Animal Care and Use Committee (ACUC).

### Animal model of diabetes mellitus

Diabetes was induced by a single intraperitoneal injection of alloxan monohydrate (Sigma Aldrich Inc.) at the dose of 200 mg/kgBW, was dissolved freshly in cold normal saline. Non-diabetic control mice received an injection of the vehicle. After 3 days, mice (n=23) with marked hyperglycemia (fasting blood glucose levels exceeding 140 mg/dL) were selected and used for the study [8].

### Experimental protocol

Group 1 and Group 2 were administered orally with aquadest. Group 3 (n=5), Group 4 (n=5) and Group 5 (n=8) was forced-fed orally with sodium orthovanadate (Calbiochem Inc.) at the dose of 16, 32 and 64 mg/kg BW/day for 7 days respectively. Fasting blood glucose levels were determined on day 0 (before the induction of diabetes), day 3 (start of treatment), and day 10 (end of treatment). At the end of treatment (day 10), mice were anesthetised and skeletal muscle tissue were collected and fixed in 10% neutral buffered formalin for histochemistry and immunohistochemistry analysis.

### Determination of blood glucose levels

Blood samples were collected from the tail vein of 8-hours-fasted mice for determination of blood glucose levels using On-Call® Plus Blood Glucose Monitoring System.

### Analysis of skeletal muscle histology

Skeletal muscle tissue were processed to any treatment of tissue necessary to impregnate them within a solid medium (paraffin wax) to facilitate the production of sections for microscopy, then were checked histochemically by haematoxylin-eosin (HE) staining [9]. Observed using a light microscope with a magnification of 100 and 400x.

### Analysis of P85 and GSK-3 expressions

Skeletal muscle tissue sections were checked immunohistochemically using PI3-kinase P85 $\alpha$  antibody ([A01524-100] GenScript Inc.) for P85 expressions and p-GSK-3 $\beta$  (Ser 9) antibody ([sc-11757] Santa Cruz Biotechnology Inc.) for GSK-3 expressions, at 1:700 dilution. For the evaluation of P85 and GSK-3 expressions, a modified semiquantitative IRS scale of Remmele was applied (Table 1). Semiquantitative IRS scale taking into account both percentage of positive cells (A) and intensity of the reaction colour (B), with the final score representing product of the two variables (A x B), and ranges from 0 to 12 pt [10].

### Statistical analysis

Analysis of the research data were processed with SPSS V.17.0 for windows with significance level  $p < 0.05$ . All data were expressed as mean  $\pm$  Standard Deviation (SD). Differences of fasting blood glucose levels between day 0 and 3 on non-diabetic control group were analyzed with unpaired t-test. Differences of fasting blood glucose levels between day 0 and 3 on diabetic-untreated control group were analyzed with Mann-Whitney test. Differences of fasting blood glucose levels between groups on day 10 were analyzed with one-way ANOVA. Differences of P85 or GSK-3 expression between groups were analyzed with Kruskal-Wallis test and followed by Mann-Whitney test.

Table 1: Semiquantitative IRS scale

A	B
0 pt - no cells with positive reaction	0 pt - no colour reaction
1 pt - to 10% cells with positive reaction	1 pt - low intensity of colour reaction
2 pt - 11-50% cells with positive reaction	2 pt - moderate intensity of colour reaction
3 pt - 51-80% cells with positive reaction	3 pt - intense colour reaction
4 pt - > 80% cells with positive reaction	

## RESULTS

### Effect of alloxan monohydrate administration in naive mice

Blood glucose levels of mice after the administration of alloxan monohydrate were summarized in table 2. In the naive mice group which only received vehicle injection intraperitoneally, no elevating in fasting blood glucose levels from day 0 to 3 significantly ( $p = 0.274$ ). While in the mice group that received alloxan monohydrate injection, there was an elevating of fasting blood glucose levels significantly from  $59.1 \pm 11.2$  mg/dL on day 0 to  $310.6 \pm 107.2$  mg/dL on day 3 ( $p < 0.001$ ).

### Histology of skeletal muscle cells after alloxan monohydrate administration

The muscle cells in the sectional of skeletal muscle tissue of non-diabetic mice looks solid, the shape and size of the cells were homogeneous, and had a nucleus at the cell edge (Fig. 1, sign  $\rightleftarrows$ ). While in the muscle cells of diabetic mice, the shape and size of the cells were not homogeneous, there was nucleus in the middle, swelling of the cell nucleus and aggregation of the cells that led to muscle cell necrosis (Fig. 1, sign  $\rightarrow$ ).

### Immunohistochemistry of P85 expressions in skeletal muscle cells

The sectional of skeletal muscle tissue of non-diabetic mice, P85 was expressed in a sufficient quantities in the cytoplasm (Fig. 2, sign  $\rightleftarrows$ ).

While in the muscle cells of diabetic mice, P85 was more expressed than non-diabetic mice, that described an excessive P85 (Fig. 2, sign  $\rightarrow$ ).

### Immunohistochemistry of GSK-3 expressions in skeletal muscle cells

The sectional of skeletal muscle tissue of non-diabetic mice, GSK-3 was expressed in a quite strongly in the cytoplasm (Fig. 3, sign  $\rightleftarrows$ ). While in the muscle cells of diabetic mice, GSK-3 was expressed weaker than non-diabetic mice, that described a reduction of GSK-3 phosphorylation (Fig. 3, sign  $\rightarrow$ ).

### Effect of sodium orthovanadate (SOV) administration in DM mice

Blood glucose levels of mice after the administration of sodium orthovanadate were summarized in table 3. Administration of sodium orthovanadate on diabetes condition for 7 days reduced fasting blood glucose levels on day 10. The higher dose of sodium orthovanadate, the greater reduction in fasting blood glucose levels.

### Histology of skeletal muscle cells after SOV administration

After administration of sodium orthovanadate for 7 days, there was an improvement in skeletal muscle cells (Fig. 4). Cell nucleus in skeletal muscle cells was clearly visible, oval, located at the edge of the cell (Fig. 4, sign  $\rightleftarrows$ ), looks no further aggregation of cell




nucleus (Fig. 4, sign  $\rightarrow$ ) and inflammatory cells, the boundary of perimysium and endomysium seems clearly and closely so that the

tissue looks more solid by muscle cells. The higher dose of sodium orthovanadate, the greater improvement in skeletal muscle cells.


**Table 2: Fasting blood glucose levels on day 0 and 3**

Group	Total mice	Blood glucose levels on day $\pm$ SD (mg/dL)		Sig.
		0	3	
Naive	5	71.6 $\pm$ 10.2	78.4 $\pm$ 8.0	p = 0.274
DM	23	59.1 $\pm$ 11.2	310.6 $\pm$ 107.2*	p < 0.001

Values are statistically significant at \*p < 0.05 vs day 0.



**Fig. 1: Appearance of skeletal muscle by HE staining. Naive state was shown by fig. A (magnification 100x) dan C (magnification 400x). DM state was shown by fig. B (magnification 100x) dan D (magnification 400x).**




**Fig. 2: Expressions of P85 in skeletal muscle cells reacted with PI3-kinase P85 $\alpha$  antibody with magnification 400x. Naive state was shown by fig. A. DM state was shown by fig. B.**


**Immunohistochemistry of P85 and GSK-3 expressions in skeletal muscle cells**

The results of IRS scoring of the muscle cells that expressed the P85 and GSK-3 were summarized in table 4. Administration of sodium orthovanadate for 7 days, reduced the excessive P85 expressions

and increased GSK-3 expressions in skeletal muscle tissue. The higher dose of sodium orthovanadate, the greater reduction in P85 expressions, was characterized by fewer brown color produced (Fig. 5). The higher dose of sodium orthovanadate, the greater enhancement of GSK-3 expressions, was characterized by the growing strength of brown color produced (Fig. 6).



**Fig. 3: Expressions of GSK-3 in skeletal muscle cells reacted with p-GSK-3 $\beta$  antibody with magnification 400x. Naive state was shown by fig. A. DM state was shown by fig. B.**



**Fig. 4: Appearance of skeletal muscle by HE staining with magnification 400x (A) DM. (B) DM+sodium orthovanadate 16 mg/kgBW. (C) DM+sodium orthovanadate 32 mg/kgBW. (D) DM+sodium orthovanadate 64 mg/kgBW.**

**Table 3: Fasting blood glucose levels on day 10**

Group	Fasting blood glucose levels average on day 10 $\pm$ SD (mg/dL)
Naive	64.6 $\pm$ 18.3*
DM	406.4 $\pm$ 64.1
DM+Sodium orthovanadate 16 mg/kgBW	303.0 $\pm$ 126.8*
DM+Sodium orthovanadate 32 mg/kgBW	231.8 $\pm$ 57.1*
DM+Sodium orthovanadate 64 mg/kgBW	75.6 $\pm$ 40.8*

Values are statistically significant at \*p < 0.05 vs DM group.



Table 4: P85 and GSK-3 expressions

Group	Protein expressions average per field of	
	P85	GSK-3
Naive	2.7 ± 1.4*	6.2 ± 2.2*
DM	7.6 ± 1.6	3.2 ± 1.6
DM+Sodium orthovanadate 16 mg/kgBW	6.4 ± 2.1*	6.9 ± 2.2*
DM+Sodium orthovanadate 32 mg/kgBW	4.5 ± 2.1*	7.4 ± 2.3*
DM+Sodium orthovanadate 64 mg/kgBW	3.9 ± 2.0*	9.2 ± 2.4*

Values are statistically significant at \* $p < 0.05$  vs DM group.




Fig. 5: Expressions of P85 in skeletal muscle cells reacted with PI3-kinase P85 $\alpha$  antibody with magnification 400x (A) DM. (B) DM+sodium orthovanadate 16 mg/kgBW. (C) DM+sodium orthovanadate 32 mg/kgBW. (D) DM+sodium orthovanadate 64 mg/kgBW




Fig. 6: Expressions of GSK-3 in skeletal muscle cells reacted with p-GSK-3 $\beta$  antibody with magnification 400x (A) DM. (B) DM+sodium orthovanadate 16 mg/kgBW. (C) DM+sodium orthovanadate 32 mg/kgBW. (D) DM+sodium orthovanadate 64 mg/kgBW

## DISCUSSION

The main finding of our study was that sodium orthovanadate reduced the excess of P85 $\alpha$  expressions (Table 4; Fig. 5) and increased phosphorylation at serin-9 of GSK-3 $\beta$  expressions (Table 4; Fig. 6), would lead to the reduction of blood glucose levels (Table 3) and improved a necrosis in skeletal muscle cells alloxan-induced diabetic mice that given sodium orthovanadate treatment for 7 days (Fig. 4).

Sodium orthovanadate adopts a trigonal bipyramidal structure that mimics the transition state of the phosphoryl transfer reaction, thereby acting as a competitive inhibitor of PTP-1B [11], allowing the phosphoester bond (which also forms an autophosphorylation) and thus the insulin signal transduction remain intact [12]. Inhibition of PTP-1B activity effectively raises the concentration of phosphorylated insulin receptor and IRS-1 [13], thereby allowing the tyrosine-phosphorylated IRS-1 to dock with the p85 regulatory

subunit of phosphatidylinositol-3-kinase (PI3-kinase). This interaction unregulated the p110 catalytic subunit of PI3-kinase, which catalyzes the production of phosphoinositide moieties that subsequently activate 3-phosphoinositide-dependent kinases (PDK), including PDK1. One downstream target of PDK1 is the serine/threonine kinase Akt. Interestingly, one substrate for phosphorylation by Akt action is GSK-3. Akt phosphorylates specific serine residues on GSK-3 [14], inhibit GSK-3 via Ser 9/21 phosphorylation, results in the dephosphorylation and activation of GS, leading to increased rates of glycogen synthesis [5], thereby improved a necrosis in skeletal muscle cells in diabetic state. The activation of these steps up to and including PI3-kinase and Akt ultimately results in the translocation of a specific glucose transporter protein isoform (GLUT-4) to the membranes of the sarcolemma and the t-tubules, where glucose transport takes place via a facilitative diffusion process. The amount of GLUT-4 protein incorporated into the sarcolemmal membrane correlates closely with the degree of insulin-stimulated glucose transport [14], thereby reduced the elevation of blood glucose levels in diabetic state.

Based on that finding, the reduction of P85 expressions and the elevation of GSK-3 expressions can be used for new therapeutic strategy to reduce the elevating blood glucose levels and to improve a necrosis that may happened in patients with T2DM.

## CONCLUSION

Vanadium compounds improved metabolic disorders in models of type 2 diabetes mellitus. In type 2 diabetes mellitus, sodium orthovanadate treatment normalized hyperglycemia by reduced P85 expressions and increased GSK-3 expressions in insulin signaling pathway. Hence, this compound is a potential candidate for oral therapy in diabetes as substitutes for insulin.

## CONFLICT OF INTERESTS

Declared None

## ACKNOWLEDGEMENT

We thank to Managing Trustee, Department of Clinical Pharmacy, Faculty of Pharmacy, Airlangga University for their support in the implementation of this research.

## REFERENCES

- Saini V. Molecular mechanisms of insulin resistance in type 2 diabetes mellitus. *World J Diabetes* 2010;1(3):68-75.
- Sudagani J, Hitman GA. Diabetes Mellitus: etiology and epidemiology. *encyclopedia of human nutrition* (third edition). London: Elsevier Inc; 2013.
- Barbour LA, Rahman SM, Gurevich I, Leitner JW, Fishcer SJ, Roper MD, *et al.* Increased P85 $\alpha$  is a potent negative regulator of skeletal muscle insulin signaling and induces *in vivo* insulin resistance associated with growth hormone excess. *J Biol Chem* 2005;280(45):37489-94.
- Doble BW, Woodgett JR. GSK-3: tricks of the trade for a multi-tasking kinase. *J Cell Sci* 2003;116(7):1175-86.
- Patel S, Doble BW, MacAulay K, Sinclair EM, Drucker DJ, Woodgett JR. Tissue-specific role of glycogen synthase kinase 3 $\beta$  in glucose homeostasis and insulin action. *Mol Cell Biol* 2008;28(20):6314-28.
- Vicente SG, Yraola F, Marti L, Munoz EG, Barrado MJG, Canto C, *et al.* Oral insulin-mimetic compounds that act independently of insulin. *Diabetes* 2007;56:486-93.

7. Arya GS, Hedaytullah MD, Raghvendra, Yadav A, Sachan K. Treating diabetes mellitus with vanadium salts-a future prospectus: a review. *Int J Pharm Sci Rev Res* 2011;8(2):183-5.
8. Jiang S, Du P, An L, Yuan G, Sun Z. Anti-diabetic effect of coptis chinensis polysaccharide in high-fat diet with STZ-induced diabetic mice. *Int J Biol Macromol* 2013;55:118-22.
9. Fawcett DW. *A textbook of histology*, 12<sup>th</sup> ed. USA: Chapman and Hall Inc; 2002.
10. Nowak M, Madej JA, Dziegiel P. Intensity of COX-2 expression in cells of soft tissue fibrosarcomas in dogs as related to grade of tumour malignancy. *Bull Vet Inst Pulawy* 2007;51:275-9.
11. O'Connor JC, Freund GG. Vanadate and rapamycin synergistically enhance insulin-stimulated glucose uptake. *Metab* 2003;52(6):666-74.
12. Rehder D. Biological and medicinal aspect of vanadium. *Inorg Chem Commun* 2003;6:604-17.
13. Tracey AS, Willsky GR, Takeuchi E. *Vanadium: Chemistry, biochemistry, pharmacology, and practical application*. New York: CRC Press; 2007.
14. Henriksen EJ, Dokken BB. Role of glycogen synthase kinase-3 in insulin resistance and type 2 diabetes. *Curr Drug Targets* 2006;7(10):1-7.



## International Symposium 2020

international journal indexed by Scopus, Wos, Esci, Doaj, Google Scholar, Copernicus, etc

[isasem.com](#)

OPEN

# International Journal of Pharmacy and Pharmaceutical Sciences

**Country** India -  SIR Ranking of India

**Subject Area and Category** Pharmacology, Toxicology and Pharmaceutics  
Pharmaceutical Science  
Pharmacology

# 37

H Index

**Publisher** International Journal of Pharmacy and Pharmaceutical Sciences

**Publication type** Journals

**ISSN** 09751491

**Coverage** 2009-2016 (cancelled)

**Scope** International Journal of Pharmacy and Pharmaceutical Sciences (Int J Pharm Pharm Sci) is peer-reviewed, monthly (Onward April 2014) open access Journal. IJPPS publishes original research work that contributes significantly to advance the scientific knowledge in pharmacy and pharmaceutical sciences including Pharmaceutical Technology, Pharmaceutics, Novel Drug Delivery, Biopharmaceutics, Pharmacokinetics Pharmacognosy and Natural Product Research Pharmaceutical/Medicinal Chemistry, Computational Chemistry and Molecular Drug Design, Pharmaceutical Analysis Pharmacology, Pharmacy Practice, Clinical and Hospital Pharmacy Cell Biology, Genomics and Proteomics, Pharmacogenomics, Bioinformatics, Pharmacoeconomics Research outcomes from medical sciences/case study and biotechnology of pharmaceutical interest are also considered. IJPPS publishes original research work either as an Original Article or as a Short Communication. Review articles on the current topic under mentioned scopes are also considered for publication. In addition, a hypothesis is also invited now for the publication onwards March 2016.



[Homepage](#)

[How to publish in this journal](#)

[Contact](#)



[Join the conversation about this journal](#)

## International Symposium 2020

international journal indexed by Scopus, Wos, Esci, Doaj, Google Scholar, Copernicus, etc


[isasem.com](#)

OPEN


Pharmaceutical Science

Pharmacology


SJR



Citations per document




Total Cites Self-Cites




Cites / Doc. (4 years)  
 Cites / Doc. (3 years)  
 Cites / Doc. (2 years)


External Cites per Doc Cites per Doc




% International Collaboration



Citable documents Non-citable documents




Cited documents Uncited documents



International Journal of Pharmacy and...

Show this widget in your own website



Just copy the code below and paste within your html code:



<a href="https://www.scima

powered by scimagojr.com



Search

## Editorial Board

### Editor-in-Chief

Prof. M. S. Bhatia, India  
(Bharati Vidyapeeth College of Pharmacy, Kolhapur, India)  
Email: manish.bhatia@bharativedyapeeth.edu  
Email: editor@ijppsjournal.com

### Associate Editors

Dr. Avijeet Jain, India  
(Shri Sathya Sai Institute of Pharmaceutical Sciences, Bhopal, India)  
Email: avijeet\_9826275340@rediffmail.com

Diah Ayu Maharani  
Department of Preventive and Public Health, Dentistry Faculty of Dentistry, University of Indonesia  
Email: raniabdillah@gmail.com

Dr. Subhash C Mandal, India  
(Department of Pharmaceutical Technology, Jadavpur University, Kolkata, India)  
Email: subhashmandal@yahoo.com

Dr. Lokesh Deb, India  
(Medicinal Plants and Horticultural Resources Division, Institute of Bioresources and Sustainable Development (IBSD), Department of Biotechnology, Government of India, Takyelpat Institutional Area, Imphal, Manipur, India)  
Email: lokesh\_deb@rediffmail.com

Dr. Wong Tin Wui, Malaysia  
(Non-Destructive Biomedical and Pharmaceutical Research Center, University of Technology MARA, Malasiya)  
Email: wongtinwui@yahoo.com

### Assistant Editors

Dr. Idress Hamad Attittala, Libya  
(Omar El-Mukhtar University, Faculty of Science, Botany Department, El-Beida, Libya)  
Email: idressattitalla2004@yahoo.com

Dr. Alok Nahata, Malaysia  
(Alor Star, Malaysia)  
Email: aloknahata@gmail.com

### Executive Editor

Mr. Niranjan Pathak, India  
(Dept. of Pharmaceutical Sciences, Dr H S Gour Central University, Saugor, India)  
Email: niranjanpathaklib@gmail.com

### Editorial Board Members

Dr. Furhan Iqbal  
Bahauddin Zakariya University Multan, Pakistan

Dr. Ebtessam Ahmed Mohammed Essa  
Department of Pharmaceutical Technology, Faculty  
of Pharmacy, Tanta University, Tanta, Egypt

Dr. Syed Muhammad Farid Hasan  
Faculty of Pharmacy, University of Karachi, Karachi,  
Pakistan

Dr. Abdel Raheim Mohammed Ahmed Donia  
College of Pharmacy – Salman Bin Abdul Aziz  
University, Egypt

Dr. Pranay Jain  
Institute of Engineering & Technology, Kurukshetra  
University, Kurukshetra, Haryana, India

Dr. Dr. Niaz Ali  
Institute of Basic Medical Sciences, Khyber Medical  
University, Peshawar, Pakistan

Dr. M. Saeed Arayne  
Chairman, Department of Chemistry, University of  
Karachi, Pakistan

Dr. Wanzala Wycliffe  
School of Pure and Applied Sciences, South Eastern  
Kenya University, Kenya

Dr. Mayuree Tangkiatkumjai  
Faculty of Pharmacy, Srinakharinwirot University,  
Ongkharak, Nakhonnayok, Thailand

Dr. Anup Naha  
Dept. of Pharmaceutic, MCOPS, Manipal, Karnataka,  
India

Dr. Jagdish Labhubhai Kakadiya  
Indubhai Patel College of Pharmacy and Research  
Centre, Petlad-Khambhat Road, Dharmaj, Anand,  
Gujarat, India

Dr. Javed Intekhab  
G. F. College (Rohilkhand University), Shahjahanpur,  
U.P., India

Dr. Manish P. Patel  
Dept. of Pharmaceutics and Pharmaceutical  
Technology, Nootan Pharmacy College, Visnagar,  
Gujarat, India

Dr. Narendra Babu Shivanagere Nagojappa  
J.N. Medical College, KLE University, Belgaum,  
Karnataka, India

Dr. C. Chellaram  
Vel Tech Multi Tech Engg. College, Chennai, India

Dr. Mehdi Shafiee Ardestani  
Department of Medicinal Chemistry and  
Radiopharmacy, Tehran University of Medical  
Sciences, Tehran, Iran

Dr. Amal Amin Mohamed  
Plant Biochemistry Department; Agriculture  
Division -National Research Center, Dokki, Cairo,  
Egypt

Dr. Rabab Kamel Mahmoud  
National Research Center, Cairo, Egypt

Dr. Syed Adnan Ali Shah  
Universiti Teknologi MARA (UiTM), Puncak Alam  
Campus, Bandar Puncak Alam, Selangor D. E.,  
Malaysia

Dr. Sat Pal Singh Bisht  
Roland Institute of Pharmaceutical Sciences  
Berhampur, Orissa, India.

Dr. Shahu Ingole  
Smt. Kashibai Navale Medical College & Hospital,  
Pune India

Dr. Gina Samy El-Feky  
Pharmaceutics Department, Faculty of Pharmacy,



- Dr. Maha Ali Eissa Ahmed  
Department of Pharmacology, Faculty of Pharmacy,  
MISR University for Science and Technology  
(MUST), Giza Governorate, Egypt.
- Dr. Rajesh Mukthavaram  
Health Science Drive, University of California, San  
Diego, Lajolla, CA, California, USA
- Dr. Saifullah Khan  
International Islamic University, Malaysia
- Norhaniza Aminudin  
Institute of Biological Sciences, Faculty of Science,  
University of Malaya, Kuala Lumpur, Malaysia.
- Dr. Shazia Jamshed  
Kulliyah of Pharmacy, International Islamic  
University Malaysia (IIUM), Kuantan, Pahang,  
Malaysia
- Dr. Mayuree Tangkiatkumjai  
Department of Clinical Pharmacy and Social  
Pharmacy Faculty of Pharmacy, Srinakharinwirot  
University, Nakhonnayok, Thailand
- Dr. İsmail Murat Palabiyik  
Faculty of Pharmacy, University of Ankara,  
Tandoğan, Ankara, Turkey
- Nadeem A. Kizilbash  
Faculty of Medicine, Northern Border University  
Arar, Saudi Arabia
- Dr. Debajit Kalita  
Department of Botany, Morigaon College, Assam,  
India
- Dr. Seyed Mohammad  
Department of Biology, Faculty of Basic Sciences,  
University of Mazandaran, Babolsar, Iran
- Dr. Shalini Sivadasan  
Faculty of pharmacy, AIMST University, Semeling,  
Kedah, Malaysia
- Dr. Sujimon Tanvichien  
Srinakharinwirot University Nakornayok Rd. A.  
Ongkarak T.Ongkarak, Nakornayok, Thailand
- Prof. Dr.-Ing. habil. Dr. h. c. Lothar Mörl  
Institute für Apparate- und Umwelttechnik Otto-  
von-Guericke-Universität Magdeburg, Germany
- Dr. V. Ravichandran  
Faculty of Pharmacy, AIMST University, Semeling,  
Kedah, Malaysia
- Dr. Zahid Hussain  
Department of Pharmaceutics, Faculty of Pharmacy,  
Modern Science and Arts University, Egypt
- Dr. Abdalla Ahmed Elbashir Ahmed  
Khartoum University, Sudan
- Dr. Yesudass Dominic Ravichandran  
School of Advanced Sciences, VIT University,  
Vellore, Tamil Nadu, India
- Dr. Seema Akbar  
Research Institute of Unani Medicine, The  
University of Kashmir Campus, Srinagar, J. & K.,  
India
- Dr. P. Thillai Arasu  
Department of Chemistry, Wollega University,  
Nekemta, Ethiopia
- Dr. Sooraj S. Nath  
Safi Institute of Advanced Study, Kozhikode, India
- Dr. Erum Shireen  
Dept. of Biochemistry, University of Karachi,  
Pakistan
- Dr. M. M. Gupta  
School of Pharmacy, Faculty of Medical Sciences,  
The University of The West India, India
- Dr. Najma Sultana  
United Biotechnologies, Karachi, Pakistan
- Dr. Sivakumar P  
Department of Petroleum Engineering, JCT College  
of Engineering and Technology, Pichanur, Tamil  
Nadu, India
- Dr. Evren Algin Yapar  
Department in Ministry of Health, Turkish  
Medicines and Medical Devices Agency, Ankara,  
Turkey
- Dr. Vishal Vijay Pandey  
Jayawantrao Sawant College of Pharmacy &  
Research, Hadapsar, Pune, India
- Dr. Shamkuwar Prashant Babarao  
Government College of Pharmacy, Thiba Palace,  
Ratnagiri, India
- Dr. S. K. Starling  
Department of Chemistry, Mewar University  
Chittorgarh, Rajasthan, India
- Dr. Syed Sajjad Hussen  
Manipal College of Pharmaceutical Sciences,  
Manipal University, India
- Dr. Ahmed Osman  
Department of Psychology, Faculty of Education,  
Assiut University, Malaysia

Universiti Teknologi MARA, Puncak Alam Campus,  
Malaysia

Dr. Ajay Kumar Meena  
Department of AYUSH, Ministry of Health & Family  
Welfare, Government of India, India

Dr. Ashish C. Suthar  
Herbal R & D, Piramal Life Sciences Ltd., Mumbai,  
India

Dr. Manish A. Rachchh  
Pharmacological Research and IPR University road,  
Rajkot, Gujarat, India

Kiran Kumar Chereddy  
Manager at Novartis Pharma AG  
Postfach 4002, Basel  
Switzerland

Dr. Fahd M. Abd Al Galil  
Department of Zoology, Faculty of Applied  
Science, Thamar University, Yemen

Dr. Priyanka Bhatt  
Department of Pharmaceutical Sciences, College of  
Pharmacy, University of South Florida, USA

Dr. Beril Anilanmert  
Istanbul University-Cerrahpasa Institute of Forensic  
Sciences, Cerrahpasa/Istanbul

Dr. Abdel-Tawab Halim Mossa Abd El-Aziz  
Department of Pesticide Chemistry, National  
Research Centre (NRC) Dokki, Cairo, Egypt

Dr. V. Vinod Prabhu  
Department of Biochemistry, University of Madras,  
Guindy campus, Tamil Nadu, India

Dr. Ching Siew Mooi  
University Putra, Malaysia

Dr. Asif Husain  
Jamia Hamdard University, New Delhi, India

Dr. Muhammad Shahzad Aslam  
School of Bioprocess Engineering, Universiti  
Malaysia Perlis, Kompleks Pusat Pengajian  
Jejawi, Arau, Perlis

Dr. Gláucio Diré Feliciano  
State University Center Foundation of the West  
Zone / Rio de Janeiro, Brazil

Dr Dilipkumar Pal  
Department of Pharmaceutical Sciences, Guru  
Ghasidash Vishwavidyalaya, Bilaspur, C.G., 495 009,  
India

Ali Abdullah Ali Al-yahawi  
Assistant Professor of Clinical Pharmacy &  
Therapeutics, Yemen



Online ISSN: 0975-1491

Print ISSN: 2656-0097

Journal Metrics 2018



Source Normalized Impact per Paper (SNIP): 2.029

---


SCImago Journal Rank (SJR): 0.23

---

ISSN: 0975-1491

---

**International Journal of  
Pharmacy and...**



---

Visitor No. 06472

---

Our Journals || Open Access Policy || Publication & Peer Review Policy || Publication Ethics

The publication is licensed under a Creative Commons License (CC BY) View Legal Code

Copyright © 2018 All Rights Reserved. Innovare Academic Sciences | Powered By CyberDairy