

DAFTAR PUSTAKA

- Batista, I. 1999. Recovery of proteins from fish waste products by alkaline extraction. European Food Research Technology, 210(2) :84–89.
- Benjakul, S and Visessanguan, W. 2010. Impacts of freezing and frozen storage on quality changes of seafoods. In Devahastin, S. (Ed.), Physicochemical Aspects of Food Engineering and Processing. Boca Raton. FL: CRC Press, 283-301.
- Chaijan, M., Panpipat, W., and Benjakul, S. 2010. Physicochemical and gelling properties of short-bodied mackerel (*Rastrelliger brachysoma*) protein isolate prepared using alkaline-aided process. Food and Bioproducts Processing, 88: 174-180.
- Chanarat, S and Soottawat, B. 2013. Impact of microbial transglutaminase on gelling properties of Indian mackerel fish protein isolates. food chemistry, 136 : 929-937
- Chang, R. 2000. Macromolecules. In Physical Chemistry for the Physical and Biological Sciences. R. Chang, Ed. Sausalito. University Science Books, 895-940.
- Chen, Y. C., and Jaczynski, J. 2007. Protein recovery from rainbow trout (*Oncorhynchus mykiss*) processing by-products via isoelectric solubilization/ precipitation and its gelation properties as affected by functional additives. Journal of Agricultural and Food Chemistry, 55(22): 9079–9088.
- Choi J. Y, and Kim J. S. 2005. Fish protein recovered using pH shifting method and its physicochemical properties. Ocean University of China, 4(3): 224–228
- Deman, M. J. 1997. Kimia Makanan. Bandung. Penerbit ITB
- Feng, Y and Hultin H.O. 2001. Effect of pH on the rheological and structural properties of gels of water-washed chicken-breast muscle at physiological ionic strength. Journal Agricultural and Food Chemistry, 49: 3921–35.
- Foh, M B K., Xia, W., Issoufou, A., and Qixing J. 2012. Influence of pH Shift on Functional Properties of Protein Isolated of Tilapia (*Oreochromis niloticus*) Muscles and of Soy Protein Isolate. Food and Bioprocess Technology, 5: 2192-2200.
- Gehring, C. K., Gigliotti, J. C., Moritz, J. S., Tou, J. C., and Jaczynski, J. 2011. Functional and nutritional characteristics of proteins and lipids recovered

- by isoelectric processing of fish byproducts and low-value fish: A review. *Food Chemistry*, 124(2): 422–431.
- Gotanco, R.G.B and Menez, M.A.J. 2004. Population Genetic Structure of Milkfish, *Chanos chanos*, Based on PCR-RFLP Analysis of the Mitochondrial Control Region. *Marine Biology*, 145: 789-801.
- Guo, Y., Kong, B., Xia, X., Yu, T., and Liu, Q. 2014. Changes in physicochemical and protein structural properties of common carp (*Cyprinus carpio*) muscle subjected to different freeze-thaw cycles. *Journal of Aquatic Food Product Technology*, 23(6): 579-590.
- Hafiludin, H. 2015. Analisis Kandungan Gizi Pada Ikan Bandeng Yang Berasal Dari Habitat Yang Berbeda. *Jurnal Kelautan. Indonesian Journal of Marine Science and Technology*, 8(1): 37-43.
- Hapsari, N dan Dedin, F R. 2014. Efektivitas Metode Pemisahan Dalam Produksi Isolat Protein Nabati Berbahan Baku Lokal. *Jurnal Ilmiah Teknik Lingkungan*, 6(1)
- Huidobro, A., Alvlrez, C., and Tejada, M. 1998. Hake Muscle Altered by Frozen Storages Affected by Added Ingredients. *Journal of Food Science*, 63(4):638–43.
- Huidobro, A., Mohamed, G. F., and Tejada, M. 1998. Aggregation of myofibrillar proteins in hake, sardine and mixed minces during frozen storage. *Journal of Agricultural and Food Chemistry*, 46: 2601–2608.
- Hultin, H.O., Kelleher, S.D., Feng, Y.M., Kristinsson, H.G., Richards, M.P., Undenand, I.A. 2000. High efficiency alkaline protein extraction. US Patent
- Hultin, H. O., Kristinsson, H. G., Lanier, T. C., and Park, J. W. 2005. Process for recovery of functional proteins by pH shift. USA. CRC Press, 107-140
- Ingadottir, B. 2004. The use of acid and alkali aided solubilization and precipitation methods to recover functional proteins from tilapia white muscle (MS thesis). Gainsville. University of Florida.
- Kim, Y. S., Park, J. W., and Choi, Y. J. 2003. New approaches for the effective recovery of fish proteins and their physicochemical characteristics. *Fisheries Science*, 69(6): 1231–1239
- Kinsella, J. E. 1976. Functional Properties of Protein in Foods: a Survey. *Critical Reviews in Food Science and Nutrition*, 7: 219-280

- Kobayashi, Y., Jae, W.P. 2016. Biochemical and physical characterizations of fish protein isolate and surimi prepared from fresh and frozen whole fish. *LWT - Food Science and Technology*
- Kristensen, L and Purslow, P. P. 2001. The Effect of Ageing on the Water Holding Capacity of Pork : Role of Cytoskeletal Proteins. *Meat Science*, 58(1): 17-23
- Kristinsson, H. G and Hultin, H. O. 2003. Changes in conformation and subunit assembly of cod myosin at low and high pH and after subsequent refolding. *Journal of Agricultural and Food Chemistry*, 51(24): 7187–7196.
- Kristinsson, H. G., and Hultin, H. O. 2003. Effect of Low and High pH Treatment on the Functional Properties of Cod Muscle Proteins. *Journal of Agricultural and Food Chemistry*, 51(17): 5103–5110
- Kristinsson, H.G and Ingadottir, B. 2005. The recovery and gel forming ability of proteins recovered from tilapia muscle with acid and alkali aided solubilization and precipitation. In: IFT Annual Meeting; 2005 July 16–20. Institute of Food Technologists Abstract
- Kristinsson, H. G and Liang, Y. 2006. Effect of pH-shift processing and surimi processing on Atlantic croaker (*Micropogonias undulates*) muscle proteins. *Journal of Food Science*, 71: 304-312.
- Kristinsson, H. G., Theodore, A. E., Demir, N., and Ingadottir, B. A. 2005. Comparative study between acid- and alkali-aided processing and surimi processing for the recovery of proteins from channel catfish muscle. *Journal of Food Science*, 70: 298–306.
- Kudo, G., Okada, M., Miyauchi, D. 1973. Gel-forming capacity of washed and unwashed flesh of some Pacific coast species of fish. *Marine Fish Rev*, 32:10–5
- Lee, J and Jae, W P. 2016. Pacific Whiting Frozen Fillets as Affected by Postharvest Processing and Storage Conditions. *Food Chemistry*, 201 : 177-184
- Lehninger, L Albert. 1982. Dasar-dasar biokimia. Jakarta. PT Gelora Aksara Pratama
- Leygonie, C., Britz, T. J., and Hoffman, L. C. 2012. Impact of freezing and thawing on the quality of meat: Review. *Meat Science*, 91(2): 93-98.
- Madsen, A. 1984 . Forbedring af Processen Til Fremstilling af Frossen Fish ud Fra Mindre Fiskearter og Biprodukter fra Filetproduktionen, Ph.D. thesis. Denmark. Akademiet for de tekniske videnskaper

- Morissey, M.T., Park, J.W., Huang, L. 2000. Surimi processing waste: its control and utilization. New York. Marcel Dekker.
- Nelson, J. S. 2006. Fishes of the World. Fourth Edition. USA. John Wiley and Sons. 601
- Ngapo, T. M., Babare, I. H., Reynolds, J., and Mawson, R. F. 1999. Freezing and thawing rate effects on drip loss from samples of pork. Meat Science, 53: 149–158.
- Nolsoe, H and Undeland, I. 2009. The acid and alkaline solubilization process for the isolation of muscle proteins: State of the art. Food and Bioprocess Technology. 2: 1-27
- Nopianti R, Huda N, Fazilah A, Ismail N, Easa AM. 2012. Effect of different types of low sweetness sugar on physic chemical properties of threadfin bream surimi (*Nemipterus spp.*) during frozen storage. Int Food Research, 19: 1011-1021
- Panpipat, W and Chaijan, M. 2017. Functional properties of pH-shifted protein isolates from bigeye snapper (*Priacanthus tayenus*) head by-product. International Journal of Food Properties, 20(3): 596-610.
- Pramono, H., D, Y Pujiastuti., A, M Sahidu. 2017. Biochemical and Physicochemical Analysis of Fish Protein Isolate Recovered from Red Snapper (*Lutjanus sp*) by-Product Using Isoelectric Solubilization/ Precipitation Method. IOP Conf. Series : Earth and Enviromental Series 137
- Rawdkuen, S., Sai-Ut, S., Khamsorn, S., Chaijan, M., and Benjakul, S. 2009. Biochemical and gelling properties of tilapia surimi and protein recovered using an acid-alkaline process. Food Chemistry, 112: 112-119.
- Rosenberg, I.M. 2005, Protein Analysis and Purification: Benchtop Techniques, Second Edition. Boston. Birkhauser
- Sano, T., Ohno, T., Otsuka-fuchino, H., Matsumoto, J. J., and Tsuchiya, T. 1994. Carp natural actomyosin: thermal denaturation mechanism. Journal of Food Science, 59(5): 1002-1008.
- Shabanpour, B and Etemadian, Y. 2016. Chemical changes and shelf-life of conventional surimi and proteins recovered using pH change method from common carp (*Cyprinus carpio*) muscle during 5 months storage at -18°C. Iranian Journal of Fisheries Sciences, 15(1) 311-332

- Shaviklo, G.R., Arason, S., Thorkelsson, G., Sveinsdottir, K. and Martinsdottir, M. 2010. Sensory attributes of haddock balls affected by added fish protein isolate and frozen storage. *Journal of Sensory Studies*, 3: 316-331
- Shaviklo, G.R., Thorkelsson, G., and Arason, S. 2010. The Influence of Additives and Frozen Storage on Functional Properties and Flow Behaviour of Fish Protein Isolated from Haddock (*Melanogrammus aeglefinus*). *Turkish Journal of Fisheries and Aquatic Science*, 10(3): 333-340
- Sikorski, Z. E and Kolakowska, A. 1990. Seafood: Resources, nutritional composition, and preservation Boca Raton, FL: CRC Press, Inc. 111-124
- Sun, X D and Richard, A H. 2011. Factors Influencing Gel Formation by Myofibrillar Proteins of Muscle Foods. *Comprehensive Reviews in Food Science and Food Safety*. 10
- Surasani, V. K. R., Khatkar, S. K., and Singh, S. 2017. Effect of process variables on solubility and recovery yields of proteins from pangas (Pangasius pangasius) frames obtained by alkaline solubilization method; Characteristics of isolates. *Food and Bioproducts Processing*. 106: 137-146.
- Suwetja, I. K. 2007. Biokimia Hasil Perikanan. Jilid III. Rigormortis, TMAO, dan ATP. Manado. Fakultas Perikanan dan Ilmu Kelautan. Uversitas Sam Ratulangi
- Tahergorabi, R., Kristen, E. Matak., and Jacek, J. 2014. Fish Protein Isolate: Development of Functional Foods with Nutraceutical Ingredients. *Journal of Functional Foods*
- Thorkelsson, G., Sigurgisladottir, S., Geirsdottir, M., Jóhannsson, R., Guérard, F., Chabeaud, A., Bourseau, P., Vandanon, L., Jaouen, P., Chaplain-Derouiniot, M., Fouchereau-Peron, M., Martinez-Alvarez, O., Le, Gal Y., Ravallec-Ple, R., Picot, L., Berge, J.P., Delannoy, C., Jakobsen, G., Johansson, I., Batista, I and Pires C. 2008 . Mild Processing Techniques and Development of Functional Marine Protein and Peptide Ingredients. in Børresen, Improving seafood products for the consumer. Woodhead Publishing Limited.
- Undeland, I., Kelleher, S. D., Hultin, H. O., McClements, J., and Thongraung, C. 2003. Consistency and solubility changes in herring (*Clupea harengus*) light muscle homogenates as a function of pH. *Journal of Agricultural and Food Chemistry*. 51: 3992–3998
- Vareltzis, K., Soultos, N., Zetou, F and Tsiaras I. 1990 . Proximate Composition and Quality of Hamburger-type Product Made from Minced Beef and Fish protein Concentrate. *Lebensm Wiss Technol*. 23:112-6

- Wang, W., Nema, S., and Teagarden, D. 2010. Protein aggregation: Pathways and influencing factors. International Journal of Pharmaceutics, 390(2): 89-99.
- Winarno, G. 2004. Kimia Pangan dan Gizi. Jakarta. PT Gramedia Pustaka
- Xiong, Y. L. 1997. Structure-function relationships of muscle proteins. In Damodaran,S. (Ed.), Food proteins and their applications. New York. Marcel Dekker, Inc. 341-392
- Xiong, Y. L and Blanchard, S.P. 1993. Viscoelastic properties of myofibrillar-polysaccharide composite gels. Journal of Food Science, 58:164–7
- Yongsawatdigul, J. and Park, J.W., 2004, Effects of alkali and acid solubilization on gelation characteristics of rockfish muscle proteins. Journal of Food Science, 69: 499–505.