

JOURNAL OF  
STEM CELL &  
REGENERATIVE  
BIOLOGY



OMMEGA PUBLISHERS  
ANSWERING INTELLIGENCE

## EDITORIAL MEMBERS



**Ahmed Abdelbaset-Ismail**

Post-Doctoral Associate  
Stem Cell Institute

Jarvis Graham Brown Cancer Centre, United States

[View Profile](#)



**Martin Morad**

Professor  
Cardiac Signaling Center, United States

[View Profile](#)



**Jianxun Song**

Associate Professor  
Dept. of Microbiology & Immunology  
Penn State University, United States

[View Profile](#)



**Bikash Pattnaik**

Senior Scientist  
Dept of Pediatrics

University of Wisconsin-Madison, United States

[View Profile](#)



**Young-Ho Lee**

Professor  
Dept of Pediatrics

Hanyang University College of Medicine, Korea South

[View Profile](#)



**Sarkar Fazlul**

Professor  
Dept of Pathology & Oncology

Wayne state University, United States

[View Profile](#)



**Hendy Hendarto**

Head  
Dept of Obstetric Gynecology

Medical School Airlangga University, Indonesia

[View Profile](#)



**Herrera Guillermo A**

Professor  
Dept of Pathology

Louisiana State University Health Sciences Center, United States

[View Profile](#)



**Mark Stephen Kindy**  
Professor  
Dept of Neurosciences  
University of South Florida,  
[View Profile](#)



**Johan Bengzon**  
Professor  
Lund University Hospital, Sweden  
[View Profile](#)



**Shuen-Kuei Liao**  
Chief Scientific Officer  
Dept of Cancer Biology and Drug Discovery  
Taipei Medical University, Taiwan  
[View Profile](#)



**Raul A. Urrutia**  
Professor  
Mayo Clinic, United States  
[View Profile](#)



**Haiyang Tang**  
Professor  
Department of Medicine  
University of Arizona, United States  
[View Profile](#)




**JIN-SAN ZHANG**  
Professor  
Wenzhou Medical University, China  
[View Profile](#)



**Irina Conboy**  
Associate Professor  
Dept of Bioengineering  
University of California Berkeley, United States  
[View Profile](#)



**Kuldip Sidhu\***  
Associate professor  
Dept of Medicine  
University of New south wales, Australia  
[View Profile](#)



## ABOUT JOURNAL

Journal of stem cell and regenerative biology is established with a focus to publish high quality articles in all the aspects of stem cells written by the authors internationally. Ommega publishers encourage the scientists and students who are working hard to bring awareness about the advancements in the stem cell related therapies. Our intention is to bring out new discoveries in to light so that the readers will have knowledge on the novel research. Ommega open access publishes the peer review articles in the field of stem cells and regenerative biology so that the readers receive complete information about the topic.

## PROFILE

Ommega Online Publishers is the house for researchers and scientists, who work hard day in and out for findings that benefit mankind and the environment.

Ommega Online Publishers is an open access publishing group of peer reviewed journals in the fields of science, technology and medicine with a sole intention of promulgation of research articles throughout the globe.

Open access model provides the scientists and researchers a platform to share their innovations, discoveries and findings thus removing all the barriers that were imposed in traditional publishing models.

**The Best Part Of Ommega Online Publishers is it Offers its Readers To Access The Content Without Any Charge Through Open Access Policy.**

### VISION

Ommega Online Publishers aims to bring the world under one roof for knowledge sharing for sustainable growth and promotion. Ommega Online Publishers publishes reliable, top tier Scientific, Technical and Medical information that sets new benchmark in the process of observation, research and discovery.

Ommega Online Publishers is committed to serve the scientific, medical communities and the whole society by publishing highest standards of peer-reviewed international quality Journals.

### OUR PHILOSOPHY

**Commitment To Provide High Quality Work**

**Speedy Process**

**Dedicated Expertise Personnel**

**One Stop Solutions**


**Minimum Publication Charges**

### JOURNAL INDEXED AT





## JOURNALS - ARTICLE IN PRESS



### Journal of Stem Cell and Regenerative Biology

JOIN AS EDITOR

SUBMIT MANUSCRIPT

[AUTHOR GUIDELINES](#) [OPEN ACCESS](#) [PEER REVIEW SYSTEM](#) [EDITOR GUIDELINES](#)

## VOLUME 2 ISSUE 1 YEAR 2016

Effectiveness of Insulin Transferrin Selenium Supplementation to Vitrified Mice Using Hemi Straw on Zona Hardening, Expression of Heat Shock Protein 70 and Caspase 3

Widjiati Widjiati

Hendy Hendarto<sup>2</sup>, Sri Pantja Madyawati<sup>1</sup>, Epy Muhammad Luqman<sup>1</sup>, Viski Fitri Hendrawan<sup>3</sup>, Agung Budianto Ahmad<sup>1</sup>  
Research



E-PUB



FULL TEXT



ABSTRACT



PDF

Intra-Bone Marrow-Hematopoietic Stem Cell Transplantation; A Powerful Tool For Hematopoietic Stem Cell Transplantation

Susumu Ikehara

Yasushi Adachi<sup>1,2</sup>, Ming Li<sup>2</sup>, Susumu Ikehara<sup>2\*</sup>  
Commentary



E-PUB



FULL TEXT



ABSTRACT



PDF

Thinking Out Loud On Endometriosis And Stem Cell Relationship

Hendy Hendarto

Editorial



E-PUB



FULL TEXT



ABSTRACT



PDF

Autologous Corneal Repair Using *in-vitro* Adult Stem Cell Expansion

Charles McGhee

Peter J Wilson, Jeremy J Mathan, Jennifer J McGhee, Salim Ismail, Trevor Sherwin  
Mini-Review



E-PUB



FULL TEXT



ABSTRACT



PDF

## Effectiveness of Insulin Transferrin Selenium Supplementation to Vitrified Mice Using Hemi Straw on Zona Hardening: Expression of Heat Shock Protein 70 and Caspase 3

Widjiati Widjiati<sup>1\*</sup>, Hendy Hendarto<sup>2</sup>, Sri Pantja Madyawati<sup>1</sup>, Epy Muhammad Luqman<sup>1</sup>, Viski Fitri Hendrawan<sup>3</sup>, Agung Budiarto Ahmad<sup>1</sup>

<sup>1</sup>Faculty of Veterinary Medicine, Airlangga University, Indonesia

<sup>2</sup>Faculty of Medicine, Airlangga University, Indonesia

<sup>3</sup>Faculty of Veterinary Medicine, Brawijaya University, Indonesia

**Corresponding Author:** Widjiati, W, Faculty of Veterinary Medicine, Airlangga University, Campus C, Jl Mulyorejo Surabaya-60115, Tel: (031)5992785; E-mail: [widjiati1962@gmail.com](mailto:widjiati1962@gmail.com)

### Abstract

Addition of Insulin Transferrin Selenium in vitrification medium can scavenge free radicals caused by temperatures stressors due to the freezing. Insulin Transferrin Selenium is complex protein that can stimulate cell growth, prevent cell damage due to the role of antioxidants inside, so it can maintain the viability of the embryo after thawing. Insulin Transferrin Selenium can improve the quality and viability of in-vitro blastocyst culture results. The purpose of this study is to prove the effectiveness Insulin Transferrin Selenium supplementation in vitrification medium can reduce the hardening zone of mouse embryo post thawing, proving the effectiveness of Insulin Transferrin Selenium supplementation in vitrification medium on the expression of Heat Shock Protein 70 in mouse embryos post thawing, and prove the effectiveness of Insulin Transferrin Selenium supplementation the vitrification medium can decrease the expression of caspase 3 in mouse embryos post thawing. This study covered the stages of superovulation and collection eggs, in-vitro fertilization (IVF), modification vitrification medium with supplementation of Insulin Transferrin Selenium (ITS), embryo vitrification with hemi straw, embryo culture, examination of the hardening zone of mouse embryo, examination of Heat Shock Protein 70 expression and examination of caspase 3 expression in embryos post thawing. The results showed that there were differences between the addition or supplementation of Insulin Transferrin Selenium and without addition of Insulin Transferrin Selenium in the vitrification medium to hardening zone and caspase 3, but it was no differentiation in the expression of Heat Shock Protein 70. The conclusion is the supplementation of Insulin Transferrin Selenium in vitrification medium decreases the hardening zone of the mouse embryos post thawing, decreases the caspase 3 expression in the mouse embryos post thawing and there is no difference in the expression of Heat Shock Protein 70 in the mouse embryos post thawing.

**Received Date:** October 23, 2015

**Accepted Date:** November 25, 2015

**Published Date:** November 30, 2015

**Citation:** Widjiati, W. Effectiveness of Insulin Transferrin Selenium Supplementation to Vitrified Mice Using Hemi Straw on Zona Hardening: Expression of Heat Shock Protein 70 and Caspase 3. (2015) J Stem Cell Regen Bio 1(1): 1- 8.



**Keywords:** Transferrin Selenium; Vitrification; Heat Shock Protein 70 (HSP70); Hemi straw; Caspase 3

### Introduction

Vitrification method is where material is frozen and placed in hyperosmolarity media or media with a high concentration of cryoprotectant. Then the material is directly immersed in liquid nitrogen so that the frozen solution seems to be as glass. Technically, this method can reduce cell damage due to freezing temperatures. In addition this method can reduce embryo damage due to freezing temperatures because the critical temperature can be exceeded by very fast and can avoid the formation of ice crystals that can damage the cells<sup>[1,2]</sup>.

Vitrification method has some problems, which make declining quality of embryos post thawing due to the thickening of pellucida zone. The thickening of pellucida zone will affect the attachment of the embryo to endometrium at the time of



**Copy rights:** © 2015 Widjiati, W. This is an Open access article distributed under the terms of Creative Commons Attribution 4.0 International License.

implantation. Hardening zone will affect the attachment of endometrium tropoblast while implantation<sup>[3]</sup>. Loss of viability of embryos also caused many blastomere cells under going apoptosis as a result of the low temperature stressor to high temperature while thawing. Improvement of Heat Shock Protein70 during freezing will result in increased caspase 3 that trigger cell apoptosis.

A decrease in viability of embryo post thawing greatly affects to the successful rate of embryo implantation, more over will decrease the pregnancy rate. This is possible, because the formation of hardening zone or thickening of pellucida zone in frozen embryos after thawing, making it difficult for implantation. Therefore, a study is required to optimize the vitrification medium, in order to optimize the role of cryoprotectant to protect the embryo from temperature stressors of vitrification method. Addition of Insulin Transferrin Selenium in vitrification medium can scavenge free radicals caused temperature stressors due to freezing<sup>[4]</sup>. Insulin Transferrin Selenium is a complex protein that can stimulate cell growth, prevent cell damage due to the role of antioxidants inside, so it can maintain the viability of embryos post thawing. According to<sup>[3]</sup>, Insulin Transferrin Selenium can improve the quality and viability of in-vitro blastocyst culture results.

Based on the background, research is needed to prove the effectiveness of Insulin Transferrin Selenium on the thickness of the pellucida zone, Heat Shock Protein70 and Caspase3 in the mouse embryos vitrified with hemistraw.

## Materials and Method

This study was experimental laboratory and the design was Completely Randomized Design (CRD), assuming all treatments from sampling to implementation as well as laboratory conditions were same.

### Superovulation and collection of eggs

Female mice were injected with Pregnant Mare Serum Gonadotropin (PMSG; Foligon, Interved) at dosage of 5 IU. Forty-eight hours later they were injected with Human Chorionic Gonadotropin (HCG or Chorulon) and directly mated with castrated male monomatingly. Seventeen hours later, the vaginal plug was examined. Female mice with positive vaginal plug means that the egg can be collected. Then the female mice with positive vaginal plug, to removed and collected the fallopian tubes. Furthermore, the fallopian tubes were flushed with PBS in the petri dish, and examined under inverted microscope by tearing fertilization bags. Finally, the flushed eggs were washed and prepared for *in-vitro* fertilization.

### In-Vitro fertilization

The collected eggs were washed three times in PBS and MEM medium. The washed eggs were transferred to the medium of fertilization. Spermatozoa were taken from cauda epididymal of male mice, then immersed in the fertilization medium where the eggs already in it. The eggs were mixed with spermatozoa then incubated in 5% CO<sub>2</sub> incubator at a temperature of 37° C for 7 hours, then the granulosa cells were dropped to observe 2 pn.

### Culture of embryos to reach morula stage

Once 2 pn was formed, the zygote was transferred into

culture medium and incubated in 5% CO<sub>2</sub> incubator at 37°C. Culture medium was replaced every 2 days until the embryos reached the morula stage.

### Modification of the vitrification medium with supplementation Insulin Transferrin Selenium(ITS)

The vitrification medium contain Phosphate Buffer-Saline(PBS) added with intracellular cryoprotectant Ethylen-glycol (EG) 30% and Sucrose 1M. The modification medium was the same as mentioned above but it was added by adding Insuline Transferrin Selenium with dosage of 5 µg/ml, 10 µg/ml and 15µg/ml<sup>[5]</sup>. So overall this study consist of 4 groups. Those groups were control group (PBS + 30% EG + Sucrose 1M), group1 (PBS + 30% EG + Sucrose1M + ITS 5 µg/ml, group2 (PBS + 30% EG + Sucrose1M + ITS 10 µg/ml) and group3 (PBS + 30% EG + Sucrose1M + ITS 15 µg/ml).

### Embryo vitrification with hemi straw

This vitrification used<sup>[6]</sup> modified method. Morula embryo stage entered to the vitrification medium 1 containing PBS for 2 minutes, then transferred to vitrification medium 2 containing 30% EG + ITS for 2 minutes, then transferred medium 3 containing 1M sucrose for 30 seconds, then the embryo was place that the end of the hemistraw. Further more hemistraw that has been revealed by liquid nitrogen dipped in liquid nitrogen and put in a largestraw, so the embryo will be remain at the end of the hemi straw. Then the big straw fixed on each tips and inserted into the cassette straw. Further more cassette inserted in container goblet of liquid nitrogen.

### Thawing frozen embryos

Thawing embryos used<sup>[6]</sup> modified method. Embryo was removed from the straw, then dropped into the vitrification medium 4 containing PBS+1M sucrose for 2 minutes, then transferred to medium vitrification 5 containing PBS + 0.5M sucrose for 2 minutes, then transferred to PBS medium for 2 minutes. Further more morula embryos cultured in CO<sub>2</sub> incubator for 24 hours to observe the development of embryo searching the blastocyst stage.

### Examination of hardening zone

Examination of hardening zone was performed with image moticroprograme using inverted microscope. Measurements were performed 3times for each embryo, then the results of the thickness zone is the average value of the measurement results.

### Examination of heat shock protein70 expression by immunocytochemistry

After thawing embryos were fixed on objects glass, then rehydrated with multi level alcohol concentration, washed with PBS, and then soaked respectively in 3% hydrogen peroxide (in DIwater) for 20 minutes, 1% BSA in PBS 30 minutes at room temperature, Primary Antibodies (anti-HSP 70) 1: 1000 overnight at 4°C, biotin-labeled secondary antibody(anti-Rat IgG Biotin Labelled) and primary antibody anti-HSP 70, 1hour at room temperature, the SA-HRP (Sterp Avidin-Hoseradish Peroxidase), 60 minutes at room temperature, chromogen DAB (3,3-diaminobenzidine tetrahydrochloride), 20 min at room temperature, counter stain (Acetoorcein), 3 minutes at room tem-

perature and finally examined under a light microscope.

### Examination of the expression of caspase3 with immunocytochemistry

After thawing embryos were fixed on objects glass, then rehydrated with multi level alcohol concentration, washed with PBS, and then soaked respectively in 3% hydrogen peroxide (in DI water) for 20 minutes, 1% BSA in PBS 30 minutes at room temperature, Primary Antibodies (anti-HSP 70) 1: 1000 overnight at 4°C, biotin-labeled secondary antibody (anti-Rat IgG Biotin Labelled) and primary antibody anti-HSP 70 1 hour at room temperature, the SA-HRP (StreptAvidin-Horse Radish Peroxidase) 60 minutes at room temperature, chromogen DAB (3,3-diaminobenzidine tetrahydrochloride), 20 minutes at room temperature, counter stain (Acetocein) 3 minutes at room temperature and finally examined under a light microscope.

### Data Analysis

The data was conducted in a same controlled environment and conditions. Data obtained from the picture of hardening zone, the expression of HSP70, Caspase3 was tested by ANOVA and if there is a difference between treatment followed by Duncan test<sup>[6]</sup>.

### Results

#### Viability of mouse embryos after thawing

The viability of the thawed embryos have tested by ANOVA, and showed significant difference ( $p < 0.05$ ), followed by the Tukey test to see the difference in each treatment. The results can be seen in (Table 1).

Insulin Transferrin Selenium was given in vitrification medium could increase the viability of mouse embryos after thawing. It can be seen from the results of the analysis between the control group without Insulin Transferrin Selenium and the treatment with Insulin Transferrin Selenium. The viability of the mouse embryos after thawing in the control group was not significantly different from the group treated with the addition of Insulin Transferrin Selenium with a dosage of 5 µg/ml, but there is significantly different in dosage of 10 µg/ml and 15 µg/ml Insulin Transferrin Selenium group respectively. Moreover, the addition of Insulin Transferrin Selenium with a dosage of 5 µg/ml was significantly different from the group treated with the addition of Insulin Transferrin Selenium with a dosage of 10 µg/ml and 15 µg/ml, but the group treated with addition of Insulin Transferrin Selenium with a dosage of 10 µg/ml and 15 µg/ml were not significantly different. The results of statistical tests in the (Table 1) showed that the addition of Insulin Transferrin Selenium with a dosage of 10 µg/ml and 15 µg/ml contributed the best results of the mouse embryo viability after thawing.

**Table 1.** Average Viability of Mouse Embryos after Thawing

Treatment Group	X ± SD
Control group	4.33 <sup>a</sup> ± 1,2
Treatment group 1	5.00 <sup>a</sup> ± 1,8
Treatment group 2	7.33 <sup>b</sup> ± 0,5
Treatment group 3	7.56 <sup>b</sup> ± 0,7

Notes : Different superscripts in the same column is significant different ( $p < 0,05$ )

Control group : PBS + EG 30 % + Sukrosa 1 M  
 Treatment group 1 : PBS + EG 30 % + Sucrose 1 M + ITS 5 µg/ml  
 Treatment group 2 : PBS + EG 30 % + Sucrose 1 M + ITS 10 µg/ml  
 Treatment group 3 : PBS + EG 30 % + Sucrose 1 M + ITS 15 µg/ml

#### Hardening zone of the mouse embryos after thawing

Based on the results of the calculation on the thickness of pellucida zone (hardening zone) on the embryos after thawing, tested by ANOVA, showed significant difference ( $p < 0.05$ ), followed by Duncan test to see the difference among the treatment. The results showed in (Table 2).

The ANOVA analysis on the thickness of the zone above showed  $F_{hitung} = 3.362$  with significance of  $0.025 < 0.05$ , so there is a noticeable difference in treatment, further more testing was done with Duncan test in order to see the difference in each treatment. The hardening zone control group was not significantly different compared with the group added with Insulin Transferrin Selenium with a dosage of 5 µg/ml and 10 µg/ml, but significantly different from the treatment group with Insulin Transferrin Selenium dosage of 15 µg/ml. Moreover, the addition of Insulin Transferrin Selenium with a dosage of 15 µg/ml was not significantly different from the group treated with the addition of Insulin Transferrin Selenium with a dosage of 10 µg/ml, but significantly different from the group treated with the addition of Insulin Transferrin Selenium with a dosage of 5 µg/ml. The statistical test resulting in (Table 2) showed that the addition of Insulin Transferrin Selenium with a dosage of 15 µg/ml gave the best results due to reducing of hardening zone, so the embryos can hatch and implant.

**Table 2:** Mean and standard deviation of the thickness of the pellucida zone on mouse Embryos after thawing

Group	Mean ± SD
Control group	14,53 <sup>b</sup> ± 1,72
Treatment group 1	14,37 <sup>b</sup> ± 1,83
Treatment group 2	13,70 <sup>ab</sup> ± 1,98
Treatment group 3	12,67 <sup>a</sup> ± 1,57

Notes : different superscript in the same column is significant different ( $p < 0,05$ )

Control group : PBS + EG 30 % + Sukrosa 1 M  
 Treatment group 1 : PBS + EG 30 % + Sucrose 1 M + ITS 5 µg/ml  
 Treatment group 2 : PBS + EG 30 % + Sucrose 1 M + ITS 10 µg/ml  
 Treatment group 3 : PBS + EG 30 % + Sucrose 1 M + ITS 15 µg/ml

#### Expression of Heat Shock Protein70 (HSP 70) in mouse embryos after thawing

Based on the calculation of HSP70 expression of mouse embryos after thawing, was tested using Kruskal-Wallis test, the result showed not significantly ( $p > 0.05$ ). The result shows in the (table 3) below. The result of Kruskal-Wallis test on HSP70 data showed not significantly of  $0.919 > 0,05$ , means there was not significant difference on the treatment. Therefore it was not continue with Mann-Whitney test. HSP 70 expression of mouse embryos after thawing in the control group was not significantly different from treated with the addition of Insulin Transferrin Selenium with a dosage of 5 µg/ml, 10 µg/ml, and 15 µg/ml. Similarly, the addition of Insulin Transferrin Selenium showed no significant difference among treatment groups. The ineffectiveness of the addition Insulin Transferrin Selenium to vitrification medium on the expression of Heat Shock Protein70 in the mouse embryos in post thawing was probably there is no



cold shock during post thawing. The increased expression of Heat Shock Protein70 during the vitrification to protect against cold shock.

**Table 3:** Results of statistic analysis HSP 70 expression of mouse embryos after thawing

Group	Mean	Asymp. Sig.
Control group	177,50 <sup>a</sup>	0,919
Treatment group1	177,50 <sup>a</sup>	
Treatment group 2	181,50 <sup>a</sup>	
Treatment group3	185,50 <sup>a</sup>	

Notes : Same superscript in the different column is not significant ( $p > 0.05$ )

Control group : PBS + EG 30 % + Sukrosa 1 M

Treatment group 1 : PBS + EG 30 % + Sucrose 1 M + ITS 5  $\mu\text{g/ml}$

Treatment group 2 : PBS + EG 30 % + Sucrose 1 M + ITS 10  $\mu\text{g/ml}$

Treatment group 3 : PBS + EG 30 % + Sucrose 1 M + ITS 15  $\mu\text{g/ml}$

### Expression of caspase 3 in mouse embryos after thawing

Based on the calculation of caspase 3 expression of mouse embryos after thawing, tested using Anova test, the result obtained was significantly very different ( $p < 0.05$ ), then continued with Tukey test to find out differences of each treatment. The result shows in the (table 4).

**Table 4:** Average and standard deviation of expression of caspase 3 in mouse embryos after thawing

Treatment Group	X $\pm$ SD
Control group	5.89 <sup>a</sup> $\pm$ 1.2
Treatment group 1	5.11 <sup>ab</sup> $\pm$ 1.7
Treatment group 2	4.00 <sup>bc</sup> $\pm$ 1.2
Treatment group 3	3.22 <sup>c</sup> $\pm$ 0.9

Notes : Different superscript in the same coloumn is significantly different ( $p < 0.05$ )

Control Group : PBS + EG 30 % + Sucrose 1 M

Treatment Group 1 : PBS + EG 30 % + Sucrose 1 M + ITS 5  $\mu\text{g/ml}$

Treatment Group 2 : PBS + EG 30 % + Sucrose 1 M + ITS 10  $\mu\text{g/ml}$

Treatment Group 3 : PBS + EG 30 % + Sucrose 1 M + ITS 15  $\mu\text{g/ml}$

Data showed significance of  $0.01 < 0.05$ , therefore there was significantly difference on the treatment. Further test using Tukey test was carried out, to see the difference of each treatment. The expression of caspase 3 in mouse embryos after thawing of control group was not significantly different from that of treatment group supplemented with Insulin Transferrin Selenium with a dosage of 5  $\mu\text{g/ml}$  dan 10  $\mu\text{g/ml}$ , but it was significantly different from that of treatment group supplemented with Insulin Transferrin Selenium with a dosage of 10  $\mu\text{g/ml}$  and that 15  $\mu\text{g/ml}$  respectively. Moreover, the expression of caspase 3 in mouse embryos after thawing of treatment group supplemented with Insulin Transferrin Selenium with a dosage of 5  $\mu\text{g/ml}$  was not significantly different from that of treatment group supplemented with Insulin Transferrin Selenium with a dosage of 10  $\mu\text{g/ml}$ , but it was significantly different from that of treatment group supplemented with Insulin Transferrin Selenium with a dosage of 15  $\mu\text{g/ml}$ . More over, the expression of caspase 3 in mouse embryos after thawing of treatment group supplemented with Insulin Transferrin Selenium with a dosage of 10  $\mu\text{g/ml}$  was not significantly different from that of treatment group supplemented with Insulin Transferrin Selenium with a

dosage of 15  $\mu\text{g/ml}$ .

### Discussion

Viability of mouse embryos after thawing at least determined the success rate of embryo transfer and their of pregnancy, however many factors can influence the rate of pregnancy, such as hormonal condition in recipient, reproductive tract condition of the recipient, implantation success, and other factor.

The result showed adding Insulin Transferrin Selenium with a dosage of 10  $\mu\text{g/ml}$  and that of 15  $\mu\text{g/ml}$  gave the best result becuse it could decrease apoptosis level of mouse embryos post thawing, therefore when the mouse embryos were re-cultured and the transfer of mouse embryos was done, it was expected successfull rate of embryo growth will be very high. The decrease of caspase 3 related to blastomer cell vitality would increase the viability of the mouse embynos after in-vitro culture, therefore it could increase the success rate of embryo transfer and their pregnancy. *In vitro* culture as culture conditions, the addition of cells to the media, hormone supplements and additional growth factors not only affect embryonic development but also the viability of embryos after freezing and warming<sup>[7-9]</sup>. In addition, the quality and stage of embryo development is also important for survival after freezing and thawing<sup>[10]</sup>. The types of factors cryoprotectant, freezing, and thawing procedures can also affect the viability of embryos after thawing. Glycerol a cryoprotectant is better than the Ethylene glycol on the quality of the blastocyst embryos after thawing<sup>[11]</sup>. According to<sup>[12]</sup>, states that embryo viability after cryopreservation correlated with the level of development of the early embryo frozen<sup>[13]</sup>. Reported the viability embryo morula after frozen is lower than blastosis. Embryonic stage development very sensitive to the freezing and affect the viability of embryos after thawing<sup>[11]</sup>.

The addition of ITS on the freezing spermatozoa can improve sperm motility post-thawing. AFP can act through mechanisms non colligative to inhibit the growth of ice crystals, with thermal hysteresis (lowering the freezing point locally, creating a difference between freezing and melting point), or by interacting directly with the membrane and increase the stability of phospholipids membrane<sup>[4]</sup>. Freezing and thawing back proved can cause significant changes in lipid peroxidation in human sperm membrane, causing damage morphology, viability and decreased motility<sup>[14]</sup>. Addition of Insulin Transferrin Selenium in vitrification medium possible because Insulin Transferrin Selenium binded free radicals through increased glutathione<sup>[15, 16]</sup>. Free radicals are molecules with no electron pairs and are highly toxic to cells as they will damage endogene anti-oxidants, consequently it will decrease glutathione peroxidase. The decrease of glutathione would cause cell damage due to the increase of poly unsaturated fatty acid (PUFA) and the result of the peroxidation lipid process, in the form of Malondeadehid binded with components in cell membranes which were able to damage structure and function of cells<sup>[17, 18]</sup>. Selenium content of the ITS will be conjugated with glutathione peroxidase and catalase containing iron. Transferrin is considered able to protect cell membranes against peroxidation caused by free radicals.

Cell damage due to oxidative stress is related to mitochondria damage, which will released pro-apoptosis molecules. When permeability of mitochondria membranes were damaged, anti-apoptosis molecules in mitochondria membranes were re-

placed by pro-apoptosis ones and the condition would activate caspase 3 as executor of apoptosis<sup>[17]</sup>. Oxidative stress originated from vitrification process of Reactive Oxygen Species (ROS) caused by cold shock. The decrease in embryo viability showed that cryoprotectant was unable to protect embryos optimally due to the oxidative stress. Adding Insulin Transferrin Selenium was able to scavenge free electrons released on targeted cells, therefore it would increase glutathione. The glutathione increase would muffle ROS released during cold shock, therefore it would get better embryo viability.

Hardening zone could decrease the rate of implantation as trophoblast cells and inner cell mass can not get out of pellucida zone, therefore trophoblast could not stick on endometrium of uterus, consequently implantation process would not be happened. The decrease in pellucida zone thickness in this study needs to be proven further if it is equivalent with the increase in pregnancy rate if the embryo from thawing is transferred to recipient. HSP70 was always found in the basal condition or a condition without stress, but the expression of HSP70 increased when the cells were under stress. In the embryo vitrification cold shock occurred. The very drastic changes in temperature cause and increase in HSP because HSP70 properties as chaperone molecules, that is, directing proteins involved in the process of synthesis, translocation and prevention of aggregation<sup>[19]</sup>. In post thawing HSP70 expression did not differ between the groups with and without supplementation of Insulin Transferrin Selenium. This possibility due to the absence of cold shock, so HSP70 did not play a role in the process of protein denaturation induced by temperature stress, so there was no increase in the expression of HSP70.

## Conclusion

The conclusion of this study is that Insulin Transferrin Selenium in the vitrification medium decreases hardening zone and caspase 3 expression in mouse embryos after thawing, but there is no difference in Heat Shock Protein70 expression between the groups and without supplementation of Insulin Transferrin Selenium respectively.

## References

1. Turathum, B., Saikhun, K., Sangsuwan, P., et al. Effects of vitrification on nuclear maturation, ultrastructural changes and gene expression of canine oocytes. (2010) *J Reprod Biol Endocrinol* 8: 70.
2. Wilding, M.G., Capobianco, C., Montanaro, N., et al. Human cleavage-stage embryo vitrification is comparable to slow-rate cryopreservation in cycles of assisted reproduction. (2010) *J Assist Reprod Genet* 27(9-10): 549-554.
3. Amir, G., Rubinsky, B., Kassif, Y., et al. Preservation of myocyte structure and mitochondrial integrity in sub zero cryopreservation of mammalian hearts for transplantation using anti freeze proteins-an electron microscopy study. (2003) *European J Cardio-Thoracic Surg* 24(2): 292-297.
4. Younis, A.I., Rooks, B., Khan, S., et al. The effect of anti-freeze peptide III (AFP) and insulin transferrin selenium (ITS) on cryopreservation of chimpanzee (*Pan troglodytes*) spermatozoa. (1998) *J Andrology* 19(2): 207-214.
5. Jeong, Y., Hossein, M.S., Bhandari, D.P., et al. Effect insulin transferrin selenium in defined and porcine follicular fluid supplemented IVM media on porcine IVF and SCNT embryo production. (2008) *J Anim Reprod Sci* 106(1-2): 13-24.
6. Djuwita, I., Boediono, A., AgungPriyono, S., et al. In Vitro Fertilization and Embryo Development of Vitrified Ovine Oocytes Stressed in Sucrose. (2005) *Hayati Journal of Biosciences* 73-76.
7. Fukui, Y., McGowan, L.T., James, R.W., et al. Factors affecting the in-vitro development to blastocysts of bovine oocytes matured and fertilised in vitro. (1991) *J Reprod Fert* 92(1): 125-131.
8. Massip, A., Mermillod, P., Wilsand, C., et al. Effects of dilution procedure and culture conditions after thawing on survival of frozen bovine blastocysts produced in vitro. (1993) *J Reprod Fert* 97(1): 65-69.
9. Shamsuddin, M., Larsson, B., Gustafsson, H., et al. A serum-free, cell-free culture system for development of bovine one-cell embryos upto blastocyst stage with improved viability. (1994) *Theriogenology* 41(5): 1033-1043.
10. Han, Y., Yamashina, M.H., Koyama, N., et al. Effects of quality and developmental stage on the survival of IVF-derived bovine blastocysts cultured in vitro after freezing and thawing. (1994) *Theriogenology* 42(4): 645-654.
11. Gustafsson, H., Larsson, B., Shamsuddin, M., et al. Factors Affecting the Survival of Frozen Thawed Bovine In Vitro Produced Blastocysts. (2001) *J Anim Sci* 14(1): 7-12.
12. Greve, T., Avery, B., Callesen, H. Viability of in-vivo and in-vitro produced bovine embryos. (1993) *Reprod Dom Anim* 28(3): 164-169.
13. Pollard, J.W., Leibo, S.P. Comparative cryobiology of in vitro and in vivo derived bovine embryos. (1993) *Theriogenology* 39(1): 287.
14. Alvarez, G., Storey, B.T. Evidence that membrane stress contributes more than peroxidation to sublethal cryodamage in cryopreserved human sperm: glycerol and other polyols as sole cryoprotectant. (1993) *J Androl* 14: 199-209.
15. Cordova, B., Morato, R., Izquierdo, D., et al. Effect of the addition of insulin-transferrin-selenium and/or L-ascorbic acid to the in vitro maturation of prepubertal bovine oocytes on cytoplasmic maturation and embryo development. (2010) *J Theriogenology* 74(8): 1341-1348.
16. Yeon, W.J., Hossein, M.S., Bhandari, D.P., et al. Effect of insulin-transferrin-selenium in defined and porcine follicular fluid supplemented IVM medium on porcine IVF and SCNT embryo production. (2008) *J Anim Reprod Sci* 106(1-2): 13-24.
17. McGavin, M.D. *Pathologic Basic of Veterinary medicine*. 4th Edition. (2007) Mosby Elsevier.
18. McKee, T., McKee, J.R. *Aerobic metabolism II: Electron transport and oxidative phosphorylation* In: *Biochemistry the molecular basic of life*. (2003) 3rd Ed McGraw-Hill: 319-326.
19. Kacimi, R., Chentoufi, J., Honb, N., et al. Hypoxia differentially regulates stress proteins in cultured cardiomyocytes. (2000) *Cardiovas Res* 46(1): 139-150.

## EDITORIAL MEMBERS



**Ahmed Abdelbaset-Ismail**  
Post-Doctoral Associate  
Stem Cell Institute  
James Graham Brown Cancer Centre, United States  
[View Profile](#)



**Martin Morad**  
Professor  
Cardiac Signaling Center, United States  
[View Profile](#)



**Jianxun Song**  
Associate Professor  
Dept. of Microbiology & Immunology  
Penn State University, United States  
[View Profile](#)



**Bikash Pattnaik**  
Senior Scientist  
Dept of Pediatrics  
University of Wisconsin-Madison, United States  
[View Profile](#)



**Young-Ho Lee**  
Professor  
Dept of Pediatrics  
Hanyang University College of Medicine, Korea South  
[View Profile](#)



**Sarkar Fazlul**  
Professor  
Dept of Pathology & Oncology  
Wayne state University, United States  
[View Profile](#)



**Hendy Hendarto**  
Head  
Dept of Obstetric Gynecology  
Medical School Airlangga University, Indonesia  
[View Profile](#)



**Herrera Guillermo A**  
Professor  
Dept of Pathology  
Louisiana State University Health Sciences Center, United States  
[View Profile](#)



**Mark Stephen Kindy**  
Professor  
Dept of Neurosciences  
University of South Florida,  
[View Profile](#)



**Johan Bengzon**  
Professor  
Lund University Hospital, Sweden  
[View Profile](#)



**Shuen-Kuei Liao**  
Chief Scientific Officer  
Dept of Cancer Biology and Drug Discovery  
Taipei Medical University, Taiwan  
[View Profile](#)



**Raul A. Urrutia**  
Professor  
Mayo Clinic, United States  
[View Profile](#)



**Haiyang Tang**  
Professor  
Department of Medicine  
University of Arizona, United States  
[View Profile](#)




**JIN-SAN ZHANG**  
Professor  
Wenzhou Medical University, China  
[View Profile](#)



**Irina Conboy**  
Associate Professor  
Dept of Bioengineering  
University of California Berkeley, United States  
[View Profile](#)



**Kuldip Sidhu\***  
Associate professor  
Dept of Medicine  
University of New south wales, Australia  
[View Profile](#)



# Journal of Stem Cell and Regenerative Biology

[JOIN AS EDITOR](#)

[SUBMIT MANUSCRIPT](#)

[AUTHOR GUIDELINES](#) [OPEN ACCESS](#) [PEER REVIEW SYSTEM](#) [EDITOR GUIDELINES](#)

## ABOUT JOURNAL

Journal of stem cell and regenerative biology is established with a focus to publish high quality articles in all the aspects of stem cells written by the authors internationally. Ommega publishers encourage the scientists and students who are working hard to bring awareness about the advancements in the stem cell related therapies. Our intention is to bring out new discoveries in to light so that the readers will have knowledge on the novel research. Ommega open access publishes the peer review articles in the field of stem cells and regenerative biology so that the readers receive complete information about the topic.

## PROFILE

Ommega Online Publishers is the house for researchers and scientists, who work hard day in and out for findings that benefit mankind and the environment.

Ommega Online Publishers is an open access publishing group of peer reviewed journals in the fields of science, technology and medicine with a sole intention of promulgation of research articles throughout the globe.

Open access model provides the scientists and researchers a platform to share their innovations, discoveries and findings thus removing all the barriers that were imposed in traditional publishing models.

**The Best Part Of Ommega Online Publishers Is It Offers Its Readers To Access The Content Without Any Charge Through Open Access Policy.**

### VISION

Ommega Online Publishers aim's to bring the world under one roof for knowledge sharing for sustainable growth and promotion. Ommega Online Publishers publishes reliable, top tier Scientific, Technical and Medical information that sets new benchmark in the process of observation, research and discovery.

Ommega Online Publishers is committed to serve the scientific, medical communities and the whole society by publishing highest standards of peer-reviewed international quality Journals.

### OUR PHILOSOPHY

**Commitment To Provide High Quality Work**

**Speedy Process**

**Dedicated Expertise Personnel**

**One Stop Solutions**

**Minimum Publication Charges**

### JOURNAL INDEXED AT



[HOME](#) [PROFILE](#) [JOURNALS](#) [SUBMIT MANUSCRIPT](#) [CONFERENCES](#) [CONTACT US](#) [REGISTRATION](#)

© 2014 OMMEGA PUBLISHERS ALL RIGHTS RESERVED.

