

The Effectiveness of Spiritual Emotional Breathing Towards Respiratory Function and Immune Response of Tuberculosis Patients

by Kusnanto Kusnanto

Submission date: 12-Mar-2019 12:15AM (UTC+0800)

Submission ID: 1091462765

File name: 17.pdf (587.81K)

Word count: 3330

Character count: 18273



The Effectiveness of Spiritual Emotional Breathing Towards Respiratory Function and Immune Response of Tuberculosis Patients

Kusnanto Kusnanto, Joni Haryanto, Tintin Sukartini, Elida Ulfiana and Made Mahaguna Putra

Faculty of Nursing Universitas Airlangga, Surabaya, Indonesia

ABSTRACT

Introduction: Tuberculosis is one cause of infectious death worldwide. In relation to the healing of pulmonary tuberculosis in Indonesia, there are still certain areas where the cure rate is still low. This study aims to identify the effect of spiritual emotional breathing (SEB) on the quality of respiratory function and the modulation of immune response in tuberculosis patients.

Methods: The study used a quasi-experimental design with two groups of pre-post-test design. The population was 34 patients with tuberculosis in East Perak's primary health care. The independent variable was SEB (spiritual emotional breathing). The dependent variables were peak expiratory flow rate (PEFR), pulse, oxygen saturation, breath frequency, breath sound, stiffness complaints, human IL-2, human cortisol, IgG.

Results: The results showed that there was a significant difference in PEFR, pulse, oxygen saturation, respiratory rate, respiratory sound, stiffness, human IL-2, human cortisol, IgG.

Conclusion: SEB can improve the quality of respiratory function and the modulation of immune response in tuberculosis patients. The emotional spiritual approach is part of the science of energy psychology that aims to turn the negative energy in a person into positive energy that can help the healing process. This therapy is performed as a complementary therapy for TB patients to improve their quality of life and the control of symptoms.

ARTICLE HISTORY

Received: May 19, 2018
Accepted: June 8, 2018

KEYWORDS

immune; spiritual emotional breathing; respiratory function; tuberculosis

CONTACT

Kusnanto Kusnanto
✉ kusnanto@fkp.unair.ac.id
✉ Faculty of Nursing
Universitas Airlangga,
Surabaya, Indonesia

Cite this as: Kusnanto, K., Haryanto, J., Sukartini, T., Ulfiana, E., & Putra, M. (2018). The Effectiveness of Spiritual Emotional Breathing towards Respiratory Function and Immune Response of Tuberculosis Patients. *Jurnal Ners*, 13(1). 93-97. doi:<http://dx.doi.org/10.20473/jn.v13i1.8373>

INTRODUCTION

Tuberculosis is one cause of infectious death worldwide (WHO, 2008). The healing of pulmonary tuberculosis in Indonesia is still low in certain areas. The disease is chronic, and can affect the quality of life of the sufferer. Patients living with tuberculosis (TB) experience the significant disruption of their social life and are exposed to both stigma and discrimination. Pulmonary tuberculosis poses serious problems, relating to the concept of quality of life consisting of the aspects of physical, psychological, social, and environmental health (Kusnanto, Pradanie and Karima, 2016). TB is a common cause of death in people with HIV. The duration of the treatment for tuberculosis is prolonged, and is at least 6 months for drug-prone TB and 18-24 months for multi-drug resistant TB (MDR-TB), which does not respond to the two most effective anti-TB drugs, isoniazid and rifampicin.

Long-term treatment, adverse drug reactions during the TB treatment, stigma and financial burden contribute to poor treatment and treatment outcomes. In addition, ensuring patient adherence to treatment through a direct facility for observation therapy (DOT) competes with patient-centered services, adding to the involved finances. The costs come from out-of-pocket and there are indirect costs associated with the treatment. This is even though anti-TB drugs are given free of charge in most countries. Rapid healing of the symptoms of TB at the onset of treatment also contributes to improper patient care (i.e. people who are lost to follow-up) as competing interests are prioritised. Poor treatment obedience and the negligence of follow-up increases the morbidity, mortality, and risk of developing drug resistance, and can lead to prolonged TB transmission (Hoorn et al., 2016).

EFT (emotional freedom technique) is one of the therapeutic groups that is referred to as

psychological energy (Church et al., 2012). EFT is a mild and non-invasive form of emotional acupressure by using an acupuncture meridian called "acupuncture without needles". Tapping the meridian point on the upper body releases the energy blockage ("energy therapy") and reinforces the patient's cognitive change for them to take full responsibility for their own health and well-being (Bougea et al., 2013). In addition, there is a repetition of self-acceptance statements, which are suggested to contribute to cognitive restructuring, a well-known psychotherapeutic technique, in which individuals identify and correct negative thoughts (Patterson, 2016). The effectiveness of SEFT (Spiritual Emotional Freedom Technique) lies in the merging of Spiritual Power with Energy Psychology. Spiritual Power contains the five main principles of sincerity, confidence, gratitude, and patience. Energy Psychology is a set of principles and techniques utilising the body's energy system to improve the state of the person's mind, emotions, and behaviour. Chemical imbalances and energy disturbances in the human body can cause emotional distress, including depression. SEFT intervention in the body's energy system can change the chemical conditions in the brain (neurotransmitter), which can further change the emotional condition of a person, including the condition of depression. In addition, SEFT is effective, easy, fast, cheap, the effect can be permanent and there are no side effects. It is universal and empowers the individual (so they are not dependent on the therapist), and it can be explained scientifically (Astuti, Yosep and Susanti, 2015).

Spiritual Emotional Breathing (SEB) is a therapy using the Spiritual Emotion Freedom Technique (SEFT), which is then followed by breathing exercises. The emotional spiritual approach is a part of the science of energy psychology that aims to turn the negative energy in a person into positive energy that can help the healing process. (Zainudin, 2006). This approach is done using the body's energy system to cure patients, as in acupuncture or acupressure, but with simple techniques. The process consists of three stages, namely the set-up, the tune-in and the tapping. The set-up aims to ensure the proper flow of bodily energy. The tune-in aims to direct our mind to the pain and to feel the pain that we experience. The tapping is a light tap on certain points that neutralises emotional and pain disorders (Zainudin, 2006).

In improving the immunological factor, our study shows that EFT can increase lymphocyte proliferation. There is the assumption that greater proliferation is associated with more effectiveness when it comes to an immune response (Aggarwal and Gurney, 2002). Furthermore, EFT can increase IL-17 in the peripheral blood of chemically and pulmonary injured veterans. IL-17 is a pro-inflammatory cytokine produced by activated memory T cells, that has a key role in the host's defence against microbial infections such as

mycobacterium tuberculosis (Jovanovic et al., 1998). It has a key role in the initiation and maintenance of inflammatory responses (Zhang et al., 2012). The cells that produce IL-17 have an important role in controlling both immune and inflammatory reactions (Lutgendorf and Costanzo, 2003). Many studies have shown that stress can affect the function and number of immune cells (Carlson et al., 2004), the production of many cytokines, like IL-4, IFN- γ , L-10 and the reduction of lymphocyte proliferation (Lengacher et al., 2013). The main objectives of this study were to analyse changes in breath frequency, PEFR, oxygen saturation, breath sound, congested, cortisol, IL-2, and IG. The second goal was to explain the relationship between the variables.

11

MATERIALS AND METHODS

This research was a quasi-experimental research study with a non-randomised (purposive sampling) Control Group Pre-test and Post-test Design. The sample in this study was made up of 34 people with the inclusion criteria of 1) positive acid-resistant bacteria TB patient, 2) productive age and 3) got standard therapy from primary health care. The exclusion criteria in this study were that they were TB patients with complications. The independent variable in this research was SEB (Spiritual Emotional Breathing), while the dependent variables in this research were cortisol, IL-2, IgM, PEFR, Oxygen Saturation, Respiratory Rate (RR), Pulse, Complaints, and breath sound.

First, we selected patients who met the inclusion criteria from each group in the first part of the study, and we enrolled those patients into the second part of the study. The patients were divided into 2 groups of 17 patients, and they were trained using an original program. The patients were also instructed to practice daily exercise at home. Before the exercises, the patient was trained beforehand and given guidance. Compliance with daily exercise was assessed weekly using an observation sheet filled in with the patients at home. The assessments were repeated 12 weeks after the beginning of the SEB program. SEB involves focusing on the situation identified as causing distress and tapping on specific meridian points of the body with breathing retraining, which consisted of relaxation, pursed lip breathing, and slow-deep breathing with training occurring in both the supine and sitting positions using the classic method of Miller (Miller, 1954), with slight modifications. Intervention was performed once a day. The control group was given treatment according to the standard of care in the primary health care centre. The respondent's sampling was done by the purposive sampling method. The sample was taken by a nurse in primary health care. The samples were then analysed at the Institute of Tropical Diseases.

The place of study was East Perak primary health care, January - February 2015. The instrument used

was a peak flow meter to measure PEFR. Pulse oximetry was used to measure SaO₂. Salivary Cortisol ELISA was used for the cortisol examination. The lowest detectable level of cortisol that can be distinguished from the Zero Standard is 0.537 mg/mL or 0.0537 µg/dL at the 95% confidence limit. Legend Max Human IL-2 Elisa used for the human IL-2 examination, and the minimum detectable concentration of IL-2 is 4 pg/mL. Gen Way Mycobacterium IgG was used for the examination of IgG, with a clinical sensitivity of 100%. The salivary cortisol and IgG concentration were determined using an enzyme-linked immunosorbent assay. The collected data was analyzed using a paired t-test. The instrument used in the research was made in Japan. The analysis was conducted using SPSS with Paired t-test ($p \leq 0.05$). The ethical considerations were met, as the participants signed an informed consent form. Moreover, their privacy, confidentiality, and voluntary participation was ensured. The study was approved by the Health Research Ethics Committee Faculty of Nursing Universitas Airlangga (No. 57-KEPK).

RESULTS

Based on Table 1, the number of the intervention and control group respondents was 17 people. The number of male respondents in the intervention group was 76.47% and in the control, it was 52.94%. The female respondents in the intervention group were 23.52% and 47.05% respectively. The mean age of the respondent intervention group was 41.17 years and in the control group, it was 47.76 years.

The mean human value of IL-2 in the intervention group before intervention was 30.95, and after intervention, it was 45.52. The mean values of the human cortisol group before and after intervention were 156.64 and 137.60 respectively. The mean IgG groups of before and after intervention were respectively 10.75 and 14.58.

Table 1 Demographic and Baseline Characteristics

Characteristics	Intervention	Control
Gender		
Male	13 (76.47%)	9 (52.94%)
Female	4 (23.52%)	8 (47.05%)
Age (year)	41.17	47.76
Breathing Pattern		
Pre test	Dyspnea 12 (70.59%)	10 (58.82%)
	Normal 5 (29.41%)	7 (41.18%)
Post test	Dyspnea 0 (00.00%)	12 (70.59%)
	Normal 17 (100.00%)	5 (29.41%)
Breathing Sound		
Pre test	Vesicular 11 (64.71%)	13 (76.47%)
	Not 6 (35.29%)	4 (23.53)
Post test	Vesicular 17 (100.00%)	15 (88.24)
	Not 0 (00.00%)	2 (11.76)
	vesicular	

DISCUSSION

Based on Table 2, there was an SEB effect on human IL-2, human cortisol, and Ig G. In improving immunologic factors, (Babamahmoodi et al., 2015) shows that EFT may increase lymphocyte proliferation. It is assumed that greater proliferation is associated with a more effective immune response. Furthermore, EFT may increase IL-17 in chemically-injured veteran's peripheral blood. IL-17 is a pro-inflammatory cytokine produced by activated memory T cells and has a key role in host defense against microbial infections, such as Mycobacterium tuberculosis. It has a key role in the initiation and maintenance of the inflammatory response. The cells that produce IL-17 have an important role in controlling immune and inflammatory reactions. Many studies have shown that stress can affect the function and number of immune cells, the production of many cytokines, such as IL-4, IFN-γ, L-10 and the reduction of lymphocyte proliferation. Furthermore, stress management intervention can reduce the immunosuppressive effects of stressors. There are many papers on the effect of other stress management techniques such as mindfulness-based stress reduction (MBSR) on immunological factors, but no studies of EFT and immunity. In Table 2, it can be seen that the decrease of the cortisol in the control group occurred because the respondent was able to manage the stress. Cortisol can be associated with emotions, and can cause long-term physiological effects (Trial, 2012).

The exercise of the respiratory muscles and chest muscles can increase lung capacity. The results showed a strong correlation between the diffusion capacity of vital pulmonary capacity. Exercise in the form of breathing exercises such as deep breathing exercises can be done by healthy people or lung problem sufferers in order to increase lung volume and capacity. The lung function status is identified by the ability of gas exchange to provide oxygen to the alveoli, so it is mutually sustainable to determine the oxygen saturation value. Doing deep breathing exercises can help in the process of oxygenation so that the saturation levels can persist within the normal range (Priyanto, 2010).

Deep breathing will reduce sympathetic reactions to improve respiratory patterns and decrease inspiratory and expiratory muscle contractions (Yadav, Singh and Singh, 2009). Shortness of breath may decrease as the breathing pattern changes from the rapid rate, which is under the control of the involuntary respiratory centre of the brain stem, and the pattern thus becomes more controlled. (Nusdwinringtya, 2000). Our study had some limitations. To elucidate the effectiveness of SEB on non-TBC patients, a prospective, randomised, controlled study is necessary. Another limitation of our study was the difficulty of proving statistical equivalence. In general, to prove statistical equivalence, there needs to be a large sample size.

Table 2 Statistical Analysis

Variable	Intervention (Mean)		P value	Control (Mean)		P value
	Before	After		Before	After	
Human IL-2	30.95	45.52	0.00	32.75	44.06	0.01
Human Cortisol	156.64	137.60	0.01	131.02	43.52	0.79
IgG	10.75	14.58	0.00	8.29	3.48	0.10
PEFR	221.18	254.71	0.00	236.47	242.94	0.28
Heart rate	87.29	79.64	0.00	87.41	91.29	0.00
Saturation O2	98.29	95.70	0.00	87.41	95.94	0.00
Respiratory Rate	22.70	17.94	0.00	21.29	23.11	0.03

CONCLUSION

Spiritual Emotional Breathing (SEB) can decrease human IL-2, human cortisol, and Ig G. Using an emotional spiritual approach is a part of the science of psychological energy that is capable of improving the immune system. This therapy is performed as a complementary therapy for TB patients to improve their quality of life and the control of their symptoms. It is used as a therapy that supports major treatment therapies.

REFERENCES

- Aggarwal, S. and Gurney, A. L. (2002) 'IL-17: prototype member of an emerging cytokine family', *Journal of leukocyte biology*, 71(1), pp. 1–8. Available at: <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=11781375&retmode=ref&cmd=prlinks%5Cnpapers3://publication/uid/C6B73D07-6483-4B7C-B182-BAE723F8A26A>. (Accessed: January 1, 2017).
- Ando, M. *et al.* (1999) 'A study of the long-term effectiveness of an outpatient pulmonary rehabilitation program', *Nihon Kogyaku Gakkai zasshi = the journal of the Japanese Respiratory Society*, 37(9), pp. 680–687.
- Astuti, R., Yosep, I. and Susanti, R. D. (2015) 'Effect of Intervention Spiritual Emotional Freedom Technique toward Decrease the Level of Depression Housewife with HIV', *Jurnal Keperawatan Padjadjaran*, 3(April 2015), pp. 44–56.
- Babamahmoodi, A. *et al.* (2015) 'Emotional Freedom Technique (EFT) Effects on Psychoimmunological Factors of Chemically Pulmonary Injured Veterans', *Journal Allergy Asthma Immunology*, 14(February), pp. 37–47.
- Bougea, A. M. *et al.* (2013) 'Effect Of The Emotional Freedom Technique On Perceived Stress , Quality Of Life , And Cortisol Salivary Levels Intension - Type Headache Sufferers : A Randomized Controlled Trial', *JSCH*. Elsevier Inc., 9(2), pp. 91–99. doi: 10.1016/j.explore.2012.12.005.
- Carlson, L. E. *et al.* (2004) 'Mindfulness-based stress reduction in relation to quality of life, mood, symptoms of stress and levels of cortisol, dehydroepiandrosterone sulfate (DHEAS) and melatonin in breast and prostate cancer outpatients', *Psychoneuroendocrinology*, 29(4), pp. 448–474. doi: 10.1016/S0306-4530(03)00054-4.
- Church, D. *et al.* (2012) 'Single-Session Reduction of the Intensity of Traumatic Memories in Abused Adolescents after EFT : A Randomized Controlled Pilot Study', *Traumatology*, 18(3). doi: 10.1177/1534765611426788.
- Clark, C. J., Cochrane, L. and Mackay, E. (1996) 'Low intensity peripheral muscle conditioning improves exercise tolerance and breathlessness in COPD', *European Respiratory Journal*, 9(12), pp. 2590–2596. doi: 10.1183/09031936.96.09122590.
- Hoorn, R. Van *et al.* (2016) 'The Effects of Psycho-Emotional and Socio- Economic Support for Tuberculosis Patients on Treatment Adherence and Treatment Outcomes – A Systematic Review and Meta- Analysis', pp. 1–27. doi: 10.1371/journal.pone.0154095.
- Jovanovic, D. V. *et al.* (1998) 'IL-17 stimulates the production and expression of proinflammatory cytokines, IL-beta and TNF-alpha, by human macrophages', *Journal of immunology*, 160(7), pp. 3513–21.
- Kusnanto, K., Pradanie, R. and Karima, I. A. (2016) 'Spiritual Emotional Freedom Technique (SEFT) terhadap Kualitas Hidup Penderita Tuberkulosis Paru Spiritual Emotional Freedom Technique (SEFT) and the Quality of Life of People Living with Lung Tuberculosis', *Jurnal Keperawatan Padjadjaran*, 4, pp. 213–224. Available at: <http://jkip.fkep.unpad.ac.id/index.php/jkpe/article/view/284/134>. (Accessed: January 1, 2017).
- Lengacher, C. A. *et al.* (2013) 'Lymphocyte Recovery After Breast Cancer Treatment and Mindfulness-Based Stress Reduction (MBSR) Therapy', *Biological Research for Nursing*, 15(1), pp. 37–47. doi: 10.1177/1099800411419245.
- Lutgendorf, S. K. and Costanzo, E. S. (2003) 'Psychoneuroimmunology and health psychology: An integrative model', *Brain, Behavior, and Immunity*, pp. 225–232. doi: 10.1016/S0889-1591(03)00033-3.
- Miller, W. F. (1954) 'A physiologic evaluation of the effects of diaphragmatic breathing training in patients with chronic pulmonary emphysema', *The American Journal of Medicine*, 17(4), pp. 471–477. doi: 10.1016/0002-9343(54)90122-9.
- Nusdwinuringtya, N. (2000) *Efek latihan otot-otot pernafasan pada penyakit paru obstruksi kronis di instalasi rehabilitasi medik RSUPN Dr.*

- Ciptomangunkusumo* Jakarta. Universitas Indonesia.
- Patterson, S. L. (2016) 'The Effect Of Emotional Freedom Technique On Stress And Anxiety In Nursing Students: A Pilot Study', *Nurse Education Today*. Elsevier B.V. doi: 10.1016/j.nedt.2016.02.003.
- Priyanto (2010) *Pengaruh deep breathing exercise terhadap fungsi ventilasi oksigen paru pada klien post ventilasi mekanik*. Universitas Airlangga.
- Trial, A. R. C. (2012) 'The Effect of Emotional Freedom Techniques on Stress Biochemistry', 200(10), pp. 891-896. doi: 10.1097/NMD.0b013e31826b9fc1.
- WHO (2008) *Global Burden of Disease 2004 Update*. WHO.
- Yadav, A., Singh, S. and Singh, K. . (2009) ' Role of pranayama breathing exercise in rehabilitation of coronary artery disease patient-a pilot study', *Indian Journal of Traditional Knowledge*, 8(3).
- Zhang, L. *et al.* (2012) 'Effects and mechanism of arsenic trioxide on reversing the asthma pathologies including Th17-IL-17 axis in a mouse model', *Iranian Journal of Allergy, Asthma and Immunology*, 11(2), pp. 133-145. doi: 011.02/ijaa.133145.

The Effectiveness of Spiritual Emotional Breathing Towards Respiratory Function and Immune Response of Tuberculosis Patients

ORIGINALITY REPORT

10%

SIMILARITY INDEX

6%

INTERNET SOURCES

8%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Bougea, Anastasia M., Nick Spandideas, Evangelos C. Alexopoulos, Thomas Thomaides, George P. Chrousos, and Christina Darviri. "Effect of the Emotional Freedom Technique on Perceived Stress, Quality of Life, and Cortisol Salivary Levels in Tension-Type Headache Sufferers: A Randomized Controlled Trial", *EXPLORE The Journal of Science and Healing*, 2013.
Publication 1%
- 2** eprints.ners.unair.ac.id
Internet Source 1%
- 3** www.rmdiagnosics.com
Internet Source 1%
- 4** Suprajitno Suprajitno, Khabibah Jannatul Firdaus, Imam Sunarno. "Family Effort in Fulfilling Personal Hygiene for Mental Disorderpeople", *Jurnal Ners dan Kebidanan* 1%

(Journal of Ners and Midwifery), 2017

Publication

-
- | | | |
|----|--|-----|
| 5 | archiv.ub.uni-heidelberg.de
Internet Source | 1% |
| 6 | Susan Librizzi Patterson. "The effect of emotional freedom technique on stress and anxiety in nursing students: A pilot study", Nurse Education Today, 2016
Publication | 1% |
| 7 | www.ncbi.nlm.nih.gov
Internet Source | 1% |
| 8 | Stapleton, Peta, Hannah Chatwin, Mary William, Amanda Hutton, Amanda Pain, Brett Porter, and Terri Sheldon. "Emotional Freedom Techniques in the Treatment of Unhealthy Eating Behaviors and Related Psychological Constructs in Adolescents: A Randomized Controlled Pilot Trial", EXPLORE The Journal of Science and Healing, 2016.
Publication | 1% |
| 9 | theses.bham.ac.uk
Internet Source | 1% |
| 10 | www.reflectionsonnursingleadership.org
Internet Source | <1% |
| 11 | A. Yasmin, A. Aryadi, A. Widya, A. P. Suwirya et al. " Original ResearchHypertension in rural | <1% |

area: The determinants of left ventricular geometric alteration among hypertensive patients at Kubu Village, Karangasem, Bali

Effectiveness comparison between lisinopril and amlodipine in Melanesian patients with hypertension

An experimental study of β -D-Glucans extract of polysaccharide peptides as anti-inflammation and anti-oxidant in patients at high-risk of atherosclerosis

Echocardiographic findings among hypertensive patients in rural area: The descriptive study at Kubu Village, Karangasem, Bali

Addition of left ventricular ejection fraction to the GRACE score in prediction of short term prognosis in patients with non ST-elevation myocardial infarction

Effect of colchicine on HsCRP and mean platelet volume in acute myocardial infarction

Correlation between GRACE score with spatial QRS-T angle in non-ST elevation myocardial infarction patients

Prevalence of Impaired Kidney Function in Hospitalized Hypertensive Patients in Agoesdjam General Hospital, Ketapang, West Borneo, Indonesia

Inappropriate initial treatment in acute coronary syndrome patients: A gap in cardiovascular care

Correlation between plasma histamine level and atherosclerosis severity based on carotid intima media thickness (CIMT) measurement in patients with stable coronary

artery disease
Clinical characteristics, management and outcome of patients with ST elevation myocardial infarction in Pekanbaru city: Preliminary result of single center experience
Phase II cardiac rehabilitation program increases functional capacity in post atrial septal defect surgical closure in adult patients
The identification of human telomerase catalytic subunit in cervical cancer patients and its potential usage as strategy to reawaken hibernating myocardium post myocardial infarction
Could EURO heart risk score predict the severity of coronary lesion?
Pericardial effusion in systemic lupus erythematosus
Intracardiac metastasis of hepatocellular carcinoma
Myocardial bridging assessed by computed tomography scan: A cause of chest pain in patients without obstructed coronary arteries
Correlation between resting heart rate and 6-Minute Walk Test distance in systolic heart failure patients
Severe pulmonary hypertension is a predictor of mortality in pregnant unoperated atrial septal defect
High plasma level of soluble ST2 associated with adverse outcomes during acute phase of ST-elevation myocardial infarction
Effect of extracorporeal shockwave myocardial revascularization on spatial QRS – T angle resolution in stable coronary artery

disease patients
Fragmented QRS study, does it have any structural abnormality relation?
Risk of coronary artery lesions severity to erectile dysfunction incidence in stable coronary heart disease patients
Clinical profiles and outcomes of patients with ST-Elevation Acute Coronary Syndrome (STE-ACS) in non-primary PCI capable hospital
Correlation between QTc and QTcD with all caused in-hospital mortality after thrombolytic therapy in Dr. Moewardi Hospital, Surakarta
Functional capacity improvement after newly-modified cardiac rehabilitation program in post-revascularized coronary artery disease patients
Acute heart failure and diabetes are predictors for mortality in 6 months follow up after acute coronary syndrome
Respiratory training as adjuvant of phase 2 exercise program decreased systolic pulmonary artery pressure in post mitral valve surgery patients with residual pulmonary hypertension
Correlation between Tricuspid Regurgitation Duration with Pulmonary Pressure in Heart Failure Patients
Compatibility of myocardial damage observed by echocardiography to MSCT coronary assessment of arterial stenosis in acute coronary syndrome
Hypolipidemic effect and antioxidant activity of (Tamarind) leaves in hypercholesterolemic-fed rats
Correlation between pulmonary hypertension and functional

tricuspid regurgitation in Cimacan General Hospital
Correlation knowledge and attitude regarding behaviour of controlling blood pressure in elderly hypertension patient in Manggar District
Correlation of lipid profile and high sensitivity C-reactive protein among young people
Increased in-hospital mortality associated with hyponatremia in acute heart failure patients
The association between different criteria of obesity to the severity of coronary artery lesion
Predictor postoperative arrhythmias after pediatric cardiac surgery using aortic cross clamp and cardiopulmonary bypass
Profile of hyperthyroid heart disease patients in National Cardiac Center Harapan Kita : A descriptive study
Effect of cardiac rehabilitation programs on functional capacity of post coronary revascularization patient with decrease ejection fraction using Six Minute Walk-Test
Association of postoperative hyperglycemia with in-hospital mortality in patients undergoing on pump coronary artery bypass grafting
A comparison successful reperfusion with and without fibrinolytic therapy in acute ST-segment elevation myocardial infarct patients
Factors influencing ventricular function in repaired (mid-term evaluation)
Correlation between tricuspid regurgitation duration with right ventricular dysfunction in atrial septal defect patients
The

effect of therapy Spiritual Emotional Freedom Technique (SEFT) changes on blood pressure in elderly hypertension experience in Rejosari Karangas Trenggalek 2015Hyponatremia as a predictor of rehospitalization in patients with acute decompensated heart failureEffect of sildenafil therapy to functional capacity in congenital heart disease patient with severe pulmonary hypertension in Sardjito HospitalEisenmenger syndrome is more prevalence in young female patients with uncorrected atrial septal defect: Result of a single-center registryLeuko-glycemic index as an in-hospital prognostic marker in patients with ST-segment elevation myocardial infarction in Haji Adam Malik General HospitalSerum uric acid levels as predictors of incident complications in acute coronary syndromeAnticoagulant therapy in heart failure patients with sinus rhythmEpicardial fat thickness is related to the blood pressure response abnormalities in exercise stress testingRemote ischemic conditioning prior to primary percutaneous coronary intervention: The effect on Six Minutes Walk Test distance and serum NT-ProBNP levelCorrelation between urinary protein and vascular dysfunction before and after Delivery in preeclamptic womenImpact of high hsCRP level

on adverse cardiovascular events and mortality in ST-elevation acute myocardial infarction undergoing fibrinolysis

Correlation between baseline soluble ST2 and global longitudinal strain 2D Speckle tracking echocardiography with left ventricle remodeling post acute myocardial infarction

Difference of peripheral P level between univalvular and multivalvular of rheumatic heart disease

Correlation between the severity of hyperthyroidism with increased of systolic blood pressure in hyperthyroidism patients

Correlation of red cell distribution width with the severity of coronary artery disease in patients with stable coronary artery disease

Chest pain presentation on acute coronary syndrome patients in Dustira Hospital

Result of exercise stress testing after cardiac rehabilitation program after coronary artery bypass grafting surgery

Relationship between plasma level of Endothelin-1 with major adverse cardiac event in ST-elevation myocardial infarction undergoing percutaneous coronary intervention

Controlled clinic blood pressure lowers the risk of left ventricular hypertrophy: An internship study among hypertensive Indonesians in rural areas

Correlation of CHA₂DS₂-Vasc-HS scores with coronary artery lesions complexity based on SYNTAX score

The correlation between body

mass index and diastolic indices in non-diabetic hypertensive male subjects
Risk of traditional cardiovascular risk factor for the development of erectile dysfunction in stable coronary artery disease patients
Relationship between left ventricle ejection fraction and submaximal exercise capacity in cardiovascular patients
Relationship between type 2 diabetes mellitus with poor heart rate autonomic response in post acute myocardial infarction patients
The ECG features among hypertensive patients in rural area: A descriptive study at Kubu Village, Karangasem, Bali
Chronotropic incompetence in coronary artery bypass graft and heart valve surgery patients: The role of β -blocker
Correlation between fibrinogen and low density lipoprotein (LDL) cholesterol levels with acute myocardial infarction
Cardiac rehabilitation use among patients with coronary artery disease on July–December 2015 in Dr. Hasan Sadikin General Hospital Bandung
Differences of plasma histamine level between acute coronary syndrome and stable coronary artery disease patients
High Endothelin-1 plasma level gives a tendency toward increasing rate of in-hospital major adverse cardiovascular outcomes in ST-elevation myocardial infarction undergoing fibrinolysis
Correlation between serum p53 levels and vascular age determined by carotid intima

media thickness (CIMT) in patients with intermediate cardiovascular risk factor
Clinical significance of precordial ST segment depression on admission electrocardiogram in patients with acute inferior myocardial infarction
Postprocedural high sensitivity C-reactive protein and the risk of recurrent ischemic events after percutaneous coronary intervention in patients with stable angina pectoris
The messenger ribonucleic acid expression of B-type natriuretic peptide, natriuretic peptide receptor type-A and type-C in cardiomyocytes of obese population
The correlation between serum galectin-3 level with wall motion score index in patients with ST-segment elevation myocardial infarction
Risk factors of atrial fibrillation in patients with chronic heart failure
QT interval prolongation after non-ST elevation myocardial infarction in type 2 diabetic compared with nondiabetic patients
The novel effect of polysaccharide peptides of to endothelial dysfunction and dislipidemia in high risk groups of atherosclerosis
Relation of Left Atrial Spontaneous Echocardiographic Contrast to Neutrophil/Lymphocyte Ratio in Patient With Mitral Stenosis
Analysis of characteristic of atrial fibrillation versus not atrial fibrillation in hospitalized heart failure patients
Pulmonary hypertension features in adult atrial septal defect

at RSUP Dr. Sardjito

The association between plasma tryptase with ventricular remodeling after 1 month in patients with acute coronary syndrome

The correlation of plasma histamine and tryptase with ST-elevation acute myocardial infarction in acute coronary syndrome patients

Acute kidney injury after coronary artery bypass graft surgery: Incidence and pre operative risk profile

The effect of diabetes mellitus on in-hospital mortality and major complications after isolated coronary artery bypass grafting surgery

Factor those influence return to work after CABG revascularization on coronary artery disease patients

The correlation of pulmonary vein endothelin-1 level with pulmonary vascular resistance before and after mitral valve surgery with pulmonary hypertension

Mitral valve E-Point septal separation as an independent predictor for in-hospital mortality after acute ST-elevation myocardial infarction

Effects of allopurinol on complications of post coronary artery bypass graft (CABG) surgery in coronary artery disease's patient with left ventricular dysfunction

Maternal and fetal outcomes of pregnancy with valvular heart disease

QRS duration in ST elevation myocardial infarction patients undergoing a primary percutaneous intervention

Knowledge of resident medical

officers on hypertensive emergency: A descriptive study
Hyperuricemia and its association with incident of congestive heart failure in Madurese population
Diagnostic value of Duke Treadmill Score in predicting coronary lesions severity in patients with suspected stable coronary artery diseases
Relationship of plasma glucose levels and troponin-I in patients wi...

Publication

12

jnk.phb.ac.id

Internet Source

<1%

13

D. J. Anderson. "Concentrations and significance of cytokines and other immunologic factors in semen of healthy fertile men", *Human Reproduction*, 09/27/2007

Publication

<1%

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On

The Effectiveness of Spiritual Emotional Breathing Towards Respiratory Function and Immune Response of Tuberculosis Patients

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
