

# Pharmaceutical Sciences

Journal of Faculty of Pharmacy

ISSN 1715-411X


Quarterly

Volume 1

Number 1

2007


No. 1



## Research Article

## Cytoprotective Properties of Carnosine against Isoniazid-Induced Toxicity in Primary Cultured Rat Hepatocytes

(/Article/PHARM\_19891\_20180501171214)

Aziz Eftekhari (/Search/A\_Aziz\_Eftekhari), Reza Heidari (/Search/A\_Reza\_Heidari), Elham Ahmadian (/Search/A\_Elham\_Ahmadian)\*  (https://orcid.org/0000-0002-7230-0489), Mohammad Ali Eghbal (/Search/A\_Mohammad\_Ali\_Eghbal)\*

*Pharm Sci.* 2018;24(4):257-263. doi: 10.15171/PS.2018.38 (https://doi.org/10.15171/PS.2018.38)

 Article (/Article/PHARM\_19891\_20180501171214)

 PDF (/PDF/PHARM\_19891\_20180501171214)


 XML (/XMLs/PHARM\_19891\_20180501171214)

Cited By:

## Research Article

## Effect of Curcumin and N-Acetylcysteine on Brain Histology and Inflammatory Factors (MMP-2, 9 and TNF- $\alpha$ ) in Rats Exposed to Arsenic

(/Article/PHARM\_20987\_20180522190243)

Mostafa Fallah (/Search/A\_Mostafa\_Fallah), Najmeh Moghble (/Search/A\_Najmeh\_Moghble), Iraj Javadi (/Search/A\_Iraj\_Javadi), Hossein Bahadoran (/Search/A\_Hossein\_Bahadoran), Alireza Shahriary (/Search/A\_Alireza\_Shahriary)\*  (https://orcid.org/0000-0003-3240-9591)

*Pharm Sci.* 2018;24(4):264-272. doi: 10.15171/PS.2018.39 (https://doi.org/10.15171/PS.2018.39)

 Article (/Article/PHARM\_20987\_20180522190243)

 PDF (/PDF/PHARM\_20987\_20180522190243)

 XML (/XMLs/PHARM\_20987\_20180522190243)

Cited By:

## Research Article

## Evaluating the Effect of Vitamin C on Myocardial Angiogenesis Under Oxidative Stress Induced by Exhaustive Exercise in Rat

(/Article/PHARM\_20995\_20180225091809)

Ahmad Hemmat\_far (/Search/A\_Ahmad\_Hemmat\_far), Parivash Piraki (/Search/A\_Parivash\_Piraki)\*, Mohammad Ali Samavati Sharif (/Search/A\_Mohammad\_Ali\_Samavati\_Sharif), Naser Behpour (/Search/A\_Naser\_Behpour)

*Pharm Sci.* 2018;24(4):273-279. doi: 10.15171/PS.2018.40 (https://doi.org/10.15171/PS.2018.40)

 Article (/Article/PHARM\_20995\_20180225091809)

 PDF (/PDF/PHARM\_20995\_20180225091809)


 XML (/XMLs/PHARM\_20995\_20180225091809)

Cited By:

## Research Article

## Synergistic Interaction of Fluconazole/Amphotericin B on Inhibition of Enzymes Contributes to the Pathogenesis of *Candida Tropicalis*

(/Article/PHARM\_20955\_20180104092142)

Alireza Khodavandi (/Search/A\_Alireza\_Khodavandi), Fahimeh Alizadeh (/Search/A\_Fahimeh\_Alizadeh)\* 

(<https://orcid.org/0000-0002-7074-0464>), Mahsa Jafarzadeh (/Search/A\_Mahsa\_Jafarzadeh)

*Pharm Sci.* 2018;24(4):280-290. doi: 10.15171/PS.2018.41 (<https://doi.org/10.15171/PS.2018.41>)

 Article (/Article/PHARM\_20955\_20180104092142)

 PDF (/PDF/PHARM\_20955\_20180104092142)

 XML (/XMLs/PHARM\_20955\_20180104092142)

Cited By:

## Research Article

## Quantification of Pimavanserin in Bulk and Tablet Dosage Form Using A Stability Indicating High Performance Liquid Chromatographic Method

(/Article/PHARM\_22094\_20180621063118)

Geetha Bhavani Koduri (/Search/A\_Geetha\_Bhavani\_Koduri), Hari Babu Bollikolla (/Search/A\_Hari\_Babu\_Bollikolla),

Ramachandran Dittakavi (/Search/A\_Ramachandran\_Dittakavi)\*, Srinivasu Navuluri (/Search/A\_Srinivasu\_Navuluri)\*

*Pharm Sci.* 2018;24(4):291-297. doi: 10.15171/PS.2018.42 (<https://doi.org/10.15171/PS.2018.42>)

 Article (/Article/PHARM\_22094\_20180621063118)

 PDF (/PDF/PHARM\_22094\_20180621063118)

 XML (/XMLs/PHARM\_22094\_20180621063118)

Cited By:

## Research Article

## Development of a Micellar Electrokinetic Chromatographic Method with Indirect UV Detection for Pregabalin Determination in Serum Samples

(/Article/PHARM\_1041\_20180608182406)

Hamid Reza Sargazi (/Search/A\_Hamid\_Reza\_Sargazi), Elnaz Tamizi (/Search/A\_Elnaz\_Tamizi), Elaheh Rahimpour

(/Search/A\_Elaheh\_Rahimpour), Abolghasem Jouyban (/Search/A\_Abolghasem\_Jouyban)\*  (<https://orcid.org/0000-0002-4670-2783>)

*Pharm Sci.* 2018;24(4):298-303. doi: 10.15171/PS.2018.43 (<https://doi.org/10.15171/PS.2018.43>)

 Article (/Article/PHARM\_1041\_20180608182406)

 PDF (/PDF/PHARM\_1041\_20180608182406)

 XML (/XMLs/PHARM\_1041\_20180608182406)

Cited By:

## Research Article

## Formulation, Characteristic Evaluation, Stress Test and Effectiveness Study of Matrix Metalloproteinase-1 (MMP-1) Expression of Glutathione Loaded Alginate Microspheres and Gel


(/Article/PHARM\_21011\_20180316093316)

Dewi Melani Hariyadi (/Search/A\_Dewi\_Hariyadi)\*, Noorma Rosita (/Search/A\_Noorma\_Rosita) 

(<https://orcid.org/56074072400>), Fitriia Nugrahaeni (/Search/A\_Fitria\_Nugrahaeni)

*Pharm Sci.* 2018;24(4):304-312. doi: 10.15171/PS.2018.44 (<https://doi.org/10.15171/PS.2018.44>)

 Article (/Article/PHARM\_21011\_20180316093316)

 PDF (/PDF/PHARM\_21011\_20180316093316)

 XML (/XMLs/PHARM\_21011\_20180316093316)

Cited By:

## Research Article

## Transdermal Delivery of Snakehead Fish (*Ophiocephalus striatus*) Nanoemulgel Containing Hydrophobic Powder for Burn Wound

(/Article/PHARM\_22079\_20180622231505)

Robert Tungadi (/Search/A\_Robert\_Tungadi)\* , Widy Susanty

(/Search/A\_Widy\_Susanty), Prisca Wicita (/Search/A\_Prisca\_Wicita), Elvira Pido (/Search/A\_Elvira\_Pido)

*Pharm Sci.* 2018;24(4):313-323. doi: 10.15171/PS.2018.45 (<https://doi.org/10.15171/PS.2018.45>)

 Article (/Article/PHARM\_22079\_20180622231505)

 PDF (/PDF/PHARM\_22079\_20180622231505)

 XML (/XMLs/PHARM\_22079\_20180622231505)

Cited By:

## Research Article

## Apparent Specific Volumes of Sucrose in Different Aqueous Cosolvent Mixtures at 298.2 K

(/Article/PHARM\_4039\_20180727201932)

Dario A Tinjaca (/Search/A\_Dario\_Tinjaca), Maria M Muñoz (/Search/A\_Maria\_Muñoz), Fleming Martinez

(/Search/A\_Fleming\_Martinez)\* , Abolghasem Jouyban

(/Search/A\_Abolghasem\_Jouyban), William E Acree Jr (/Search/A\_William\_Acree\_Jr)

*Pharm Sci.* 2018;24(4):324-331. doi: 10.15171/PS.2018.46 (<https://doi.org/10.15171/PS.2018.46>)

 Article (/Article/PHARM\_4039\_20180727201932)

 PDF (/PDF/PHARM\_4039\_20180727201932)

 XML (/XMLs/PHARM\_4039\_20180727201932)

Cited By:

## Research Article

**Cytotoxic Flavonoids from the Aerial Parts of *Stachys lavandulifolia* Vahl**

(/Article/PHARM\_1842\_20180624125953)

Mohammad-Reza Delnavazi (/Search/A\_Mohammad-Reza\_Delnavazi), Parisa Saiyarsarai

(/Search/A\_Parisa\_Saiyarsarai), Saeedeh Jafari-Nodooshan (/Search/A\_Saeedeh\_Jafari-Nodooshan), Mahnaz Khanavi

(/Search/A\_Mahnaz\_Khanavi), Saeed Tavakoli (/Search/A\_Saeed\_Tavakoli), Hormoz Hadavinia

(/Search/A\_Hormoz\_Hadavinia), Narguess Yassa (/Search/A\_Narguess\_Yassa)\*

*Pharm Sci.* 2018;24(4):332-339. doi: 10.15171/PS.2018.47 (<https://doi.org/10.15171/PS.2018.47>)

 Article (/Article/PHARM\_1842\_20180624125953)

 PDF (/PDF/PHARM\_1842\_20180624125953)

 XML (/XMLs/PHARM\_1842\_20180624125953)

Cited By:

## Short Communication

**Evaluating the Protective Effect of a Combination of Curcumin and Selenium-L-Methionine on Radiation Induced Dual Oxidase Upregulation**

(/Article/PHARM\_23127\_20180719180333)

Peyman Amini (/Search/A\_Peyman\_Amini), Saeed Rezapoor (/Search/A\_Saeed\_Rezapoor), Dheyauldeen Shabeeb

(/Search/A\_Dheyauldeen\_Shabeeb), Ahmed Elejo Musa (/Search/A\_Ahmed\_Elejo\_Musa), Masoud Najafi

(/Search/A\_Masoud\_Najafi)\*, Elahe Motevaseli (/Search/A\_Elahe\_Motevaseli)\*

*Pharm Sci.* 2018;24(4):340-345. doi: 10.15171/PS.2018.48 (<https://doi.org/10.15171/PS.2018.48>)

 Article (/Article/PHARM\_23127\_20180719180333)

 PDF (/PDF/PHARM\_23127\_20180719180333)

 XML (/XMLs/PHARM\_23127\_20180719180333)

Cited By:

**Reviewer Acknowledgement****Acknowledgement of Top Manuscript Reviewers 2018**

(/Article/Pharm-31720)

Ali Shayanfar (/Search/A\_Ali\_Shayanfar)  (<https://orcid.org/0000-0001-7507-9057>), Sanaz Hamedeyazdan

(/Search/A\_Sanaz\_Hamedeyazdan)  (<https://orcid.org/0000-0002-8950-9879>), Alireza Garjani

(/Search/A\_Alireza\_Garjani)\*

*Pharm Sci.* 2018;24(4):346-346. doi: 10.15171/PS.2018.49 (<https://doi.org/10.15171/PS.2018.49>)

 Article (/Article/Pharm-31720)

 PDF (/PDF/Pharm-31720)

 XML (/XMLs/Pharm-31720)

Cited By:

[Submit Your Paper \(/Login\)](#)

[Editorial Board \(/EditorialBoard\)](#)

[Instructions for authors \(/InstructionsforAuthors\)](#)

[Contact editorial office \(/ContactUs\)](#)

[About \(/AimsandScope\)](#)


[Editorial policies \(/EditorialPolicies\)](#)

[Peer Review \(/PeerReview\)](#)

[Archive \(/Archive\)](#)

© 2019 Tabriz University of Medical Sciences; unless otherwise stated





# Formulation, Characteristic Evaluation, Stress Test and Effectiveness Study of Matrix Metalloproteinase-1 (MMP-1) Expression of Glutathione Loaded Alginate Microspheres and Gel

Dewi Melani Hariyadi<sup>1\*</sup>, Noorma Rosita<sup>1</sup>, Fitria Nugrahaeni<sup>1</sup>

<sup>1</sup>Pharmaceutics Department, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia.

## Article Info

### Article History:

Received: 16 March 2018

Revised: 18 August 2018

Accepted: 25 August 2018

ePublished: 30 December 2018

### Keywords:

- Alginate microspheres
- Aerosolization
- Glutathione
- Surfactants
- Mmp-1 expression

## ABSTRACT

**Background:** The present study aimed to formulate and evaluate the stability, characteristics and effectiveness of glutathione-loaded alginate microspheres through increased lipophilicity using surfactant with a Hydrophilic-Lipophilic Balance (HLB) value equal to 7. The selection of glutathione as an antioxidant was based on its prominent role in maintaining intracellular redox balance. Alginate was used as the polymer, while calcium chloride constituted a cross-linking agent and Tween and Span were employed as surfactants.

**Methods:** The study applied an ionotropic gelation-aerosolization method. Microspheres were characterized by their morphology, size, drug loading, entrapment efficiency and yield. Stress testing utilized a forced degradation method, while an effectiveness study of glutathione incorporated a Matrix Metalloproteinase I (MMP-1) parameter on mouse skin. Glutathione-microspheres, to which had been added surfactants with a HLB value equal to 7, were compared to those without surfactants.

**Results:** Microspheres demonstrated both high yield and encapsulation efficiency. From the stability study conducted, it was evident that the glutathione-microspheres with additional surfactant were more stable than glutathione with surfactant, but without microspheres. Similarly, the glutathione-microspheres with additional surfactant were more stable than the glutathione without surfactant. The *in vivo* effectivity showed lipophilic glutathione microspheres were able to decrease MMP-1 expression in the dermis tissue of mice.

**Conclusion:** The results of freeze-dried glutathione-loaded alginate microspheres with surfactant with a HLB value equal to 7 can be utilized as potential glutathione delivery systems.

## Introduction

Glutathione (L- $\gamma$ -glutamyl-L-cysteinyl-glycine) (GSH) is a low molecular weight thiol-tripeptide that plays a prominent role in maintaining intracellular redox balance.<sup>1</sup> Glutathione is a ubiquitous compound of the biologically active sulfhydryl group provided by the cysteine moiety that acts as the active part of the molecule.<sup>2</sup> The sulfhydryl group promotes interaction with a variety of biochemical systems to form glutathione in its predominant intracellular form, which acts as a potent antioxidant and defends against toxic compounds and xenobiotics.<sup>3</sup> However, the biomedical applications of glutathione remain limited due to its relatively short half-life, labile properties and rapid metabolism and elimination.<sup>3</sup> A study was carried out into the lipophilicity increase of glutathione using mixed surfactants of Tween 80 and Span 80.<sup>4</sup> The addition of surfactants with an HLB value equal to 7 affected the lipophilicity of glutathione resulting in similarity to Log P lipophilicity of the skin

(Log P 2-3).<sup>4</sup> Therefore, a surfactant system was needed in order to improve the stability of glutathione.

Microspheres are small spherical particles, with diameters in the micrometer range (typically 1  $\mu$ m to 1000  $\mu$ m (1 mm)). Encapsulation system in the microspheres can be used to protect sensitive materials to environmental conditions such as light, oxygen, water, and temperature, such as glutathione. By using microspheres delivery system it is expected that the active substance will be protected and will be also able to penetrate the dermis layer of the skin. Microspheres can stabilize and protect a drug from degradation, while preserving its biological activity and enhancing its bioavailability.<sup>5</sup> Moreover, they offer prolonged or controlled drug delivery, improved bioavailability and stability.<sup>5-6</sup> Sodium alginate, used in drug delivery systems, is a linear copolymer with a polysaccharide backbone comprising two repeating carboxylated monosaccharide units (mannuronic acid and guluronic acid).<sup>7</sup>

\*Corresponding Author: Dewi Melani Hariyadi, E-mail: dewi-m-h@ff.unair.ac.id

©2018 The Authors. This is an open access article and applies the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

**Table 1.** Formula of Glutathione-Ca Alginate Microspheres with or without surfactant.

Compounds	Function	Concentration of Compound	
		I	II
Glutathione	Active Compound	-	0.5 g
Dried Glutathione+surfactant HLB 7	Active Compound	0.5 g	-
Alginate	Polymer	2% (b/v)	2% (b/v)
CaCl <sub>2</sub> Solution	Crosslinker	1 M	1 M

Formula I: Dried glutathione+surfactant HLB 7 0.5g; Alginate 2% (b/v); 1M CaCl<sub>2</sub> Solution

Formula II: Glutathione; Alginate 2% (b/v); 1M CaCl<sub>2</sub> Solution

Alginate can be cross-linked by external gelation methods allowing the alginate-drug solution to be extruded as microspheres into a CaCl<sub>2</sub> solution. Alginates have guluronate (G) and mannuronate (M) monomer units. Gelling of the alginate occurs when divalent cations participate in the interchain bonding between guluronate units (G-blocks), giving rise to a three-dimensional network in the form of a gel. The “eggbox” model has been formulated to explain the nature of this interaction.<sup>7-8</sup> Sodium alginate forms gel microspheres by crosslinking with Ca<sup>2+</sup> ion.<sup>8-9</sup> This study applied an aerosolization technique, previously used to encapsulate drug and proteins, which crosslinked alginate polymer and CaCl<sub>2</sub> crosslinker to encapsulate a drug model by spraying followed by freeze-drying.<sup>10</sup> The advantages of aerosolization techniques include its ability to produce a simple, rapid, non-toxic and cost-effective method.<sup>10,11</sup> In cases of antioxidant use, denaturation or stability issues of the antioxidants could be avoided.<sup>8,10,11</sup>

The use of alginate microspheres in the field of biotechnology and the pharmaceutical industry is currently widespread due to their unique properties of high biocompatibility and biodegradability.<sup>12</sup> The present study was aimed to formulate and evaluate the characteristics and stability of the glutathione and freeze-dried glutathione-loaded alginate microspheres with surfactants with a HLB value equal to 7. The microspheres were evaluated for size, morphology, encapsulation efficiency, loading and yield. Stress Testing was studied using a forced degradation method<sup>13</sup>. Effectiveness study of glutathione described as MMP-1 parameter will be studied on animal’s skin.

**Materials and Methods**

**Chemical Reagents**

The following pharmaceutical grade chemical reagents were used: Glutathione (Sigma-Aldrich Inc); Sodium alginate (Sigma-Aldrich Inc); CaCl<sub>2</sub>.2H<sub>2</sub>O (Merck); Sodium citrate (Merck); Tween 80 (Merck); Span 80

(Merck); NaH<sub>2</sub>PO<sub>4</sub>.2H<sub>2</sub>O (Merck); Na<sub>2</sub>HPO<sub>4</sub>.12H<sub>2</sub>O (Merck) and Aquadest.

**Formulation of Glutathione-Loaded Alginate Microspheres**

2g of Glutathione was dissolved in 20ml of phosphate buffer solution pH 6±0.05. 0.5g of surfactant (mixed tween 80 and span 80) with a HLB value equal to 7 was added before freeze-drying was conducted for 30 hours at -26 °C. The preparation of alginate microspheres used ionotropic gelation method involving aerosolization. The alginate microspheres formulas were summarized in Table 1.

**Gel Formulation**

The carbomer (0.05 g) was dispersed in a preheated aquadest (5 g) before being cooled and propylene glycol and triethanolamine added through continuous stirring. Carbomer concentration was 1% w/w. The glutathione-alginate microspheres were then added and stirred continuously, while the pH was checked. Gel formulation was shown in Table 2.

**Stress Test of Glutathione-Alginate Microspheres**

The glutathione microspheres’ stress test involved storage in an oven at 50°C, 60°C or 80 °C and 75% RH for five days.<sup>13</sup> Organoleptic observations (color, odor and taste), drug loading and percentage of entrapment were subsequently performed on days 1, 3 and 5.

**Effectiveness Study of MMP-1 of Glutathione-Alginate Microspheres and Gel**

MMP-1, known as collagenase-1, is a zinc and calcium dependent endopeptidase, produced and released by both dermal fibroblasts and keratinocytes, which functions to break down collagens. The effectiveness study of MMP-1 consisted of the following stages: 30 balb/c mice were prepared by first shaving those areas of their backs to be irradiated.

**Table 2.** Gel formula of glutathione-ca alginate microspheres and blank microspheres.

Compound	Formula I (g)	Formula II (g)	Formula III (g)
Glutathione	-	-	-
Glutathione Microspheres	Equal 0.2	Equal 0.2	-
Carbomer	0.05	0.05	0.05
Propylene glycol	0.5	0.5	0.5
Triethanolamine	0.025	0.025	0.025
Aquadest ad	5	5	5

Formula I: Gel Formula Glutathione Microspheres + surfactant HLB 7

Formula II: Gel Formula Glutathione Microspheres -surfactant

Formula III: Formula gel base



A dose of 60 mJ/m<sup>2</sup> UV radiation was administered with interval every two days (at days 1,3,5,7,9,11, and 13), mice were then prepared its skin biopsy for histopathology examination (skin tissue of skin biopsy of mouse diameter 5 mm and depth until sub-cutaneous). Then histopathology examination on fibroblasts which expressing MMP-1 were determined under microscope with measurements taken by means of a calibrated lux meter. The determination of MMP-1 levels (%) was done by counting Fibroblasts expressing MMP-1 divided with total fibroblasts in the field of view.<sup>14</sup> The subjects were divided into three groups: Group I: glutathione microspheres gel with increased lipophilicity, Group II: glutathione gel, Group III: gel base.

#### Characterization of Glutathione-Alginate Microspheres

**Size and morphology:** Size was determined by means of optical microscopy, while morphology was investigated using a Scanning Electron Microscopy (SEM). The 300 particles of wet microspheres were measured using an optical microscope. The particles were firstly grouped to identify the smallest and largest within all the samples, by dividing them into several intervals and classes. The average diameter was then determined using the following equation:

$$D \text{ average} = \frac{\sum nd}{\sum n} \quad \text{Eq. (1)}$$

where:

n = number of particles observed

d = particle size

For freeze-dried microspheres, SEM was used to determine morphology and size by firstly placing them on an adhesive material containing metal grains, for example Platinum (Pt). The gold in the chamber was then evaporated in order to coat the entire surface of the microspheres with its vapor. The surface of the gold-coated microspheres was subsequently observed by means of SEM.

#### Determination of Glutathione-Alginate Microspheres

The drug content of alginate microspheres was quantified by breaking the microspheres formed into 120 mg with 50 ml Na Citrate over seven hours. From standard curves, microspheres were calculated in terms of entrapment efficiency, glutathione content and yield.<sup>15</sup> The results obtained were calculated based on the percentage of glutathione content of each formula using the equation below:

$$\% \text{Drug Loading} = \frac{\text{Weight of glutathione in microspheres}}{\text{Total weight of dry microspheres}} \times 100 \quad \text{Eq. (2)}$$

#### Determination of Glutathione Entrapment Efficiency

Entrapment efficiency was calculated based on the glutathione content in microspheres using the following equation:

$$\text{Entrapment efficiency} = \frac{\text{Weight of glutathione in microspheres}}{\text{Theoretical weight of glutathione}} \times 100 \quad \text{Eq. (3)}$$

#### Determination of Yield

The percentage recovery was calculated from the total number of dry microspheres produced compared to the amount of sodium alginate-glutathione added during the manufacturing process. From the calculation results, it was possible to quantify the yield of microspheres.

$$\% \text{ Yield} = \frac{\text{Total weight of dry microspheres}}{\text{Total weight of glutathione and polymer}} \times 100 \quad \text{Eq. (4)}$$

#### Results

The microencapsulation process can protect the active ingredient against chemical or enzymatic degradation.<sup>16</sup> The formulas were divided into two types of microspheres, namely glutathione-alginate microspheres with increased lipophilicity through the addition of surfactants and glutathione-alginate microspheres to which surfactants had not been added. The production of glutathione-alginate microspheres with increased lipophilicity was achieved by, firstly, adding surfactants to glutathione prior to the encapsulation process by means of alginate and a crosslinker. Another formula of glutathione-alginate microspheres without added surfactants was compared to glutathione-alginate microspheres which had not been subjected to lipophilicity enhancement. The production of microspheres employed an ionotropic gelation method including aerosolization. During microspheres formulation, the excess CaCl<sub>2</sub> that did not react with sodium alginate was removed since it can decrease entrapment efficiency.<sup>9</sup> Maltodextrin, as a lyoprotectant, was intended to stabilize the microspheres against the pressure exerted during the freeze-drying step of any water replacement process.<sup>17</sup> Maltodextrin replaces water molecules by forming hydrogen bonds between maltodextrins and polar groups on microspheres surfaces on conclusion of the drying process. Consequently, microspheres will be protected from mechanical stress and can prevent aggregation during the freeze-drying process. Maltodextrin also plays a role in the formation of microspheres surfaces.<sup>9</sup>

An evaluation of the characteristics of microspheres was performed in terms of their size, shape and surface, IR spectrophotometry, entrapment efficiency, glutathione loadings and degree of yield. The evaluation of the size distribution of wet microspheres involved use of a 300-particle optical microscope. The size of the blank microspheres was confirmed as 1.34 μm and both formulas of microspheres (F1 and F2) showed larger particle sizes of 1.40 μm and 1.58 μm respectively compared to blank microspheres (Table 3). Blank

microspheres were smaller compared to the two formulas of microspheres. Three formulas had a polydispersity index of 0.003. The resulting polydispersity index of less than 0.3 indicated that the sample had a narrow distribution (monodisperse) or uniformly stated size.<sup>18</sup>

A study of the shape and surface of wet microspheres can be seen in Figure 1A, while one of dry microspheres can be seen in Figure 1B. An investigation into the shape and surface of the microspheres by Scanning Electron Microscope (SEM) confirmed that the F1 microspheres had an uneven surface. This was because a certain degree of moisture cannot be eradicated after the sublimation process involving water and surfactant glutathione. This was evidenced from the moisture content level of 8.25% in F1, while that in F2 was lower than 2.65%. The sublimation process of F1 was not optimal because microspheres contained a surfactant with the potential to attract water. F2 microspheres possessed a smooth surface. The F1 and F2 microspheres surfaces became spherical due to the addition of maltodextrin which closes the cavities or pores which had increased in number and

size during the freeze-drying process by forming hydrogen bonds with polar groups on the microspheres surface.<sup>19</sup>

The surface of those microspheres which was not containing surfactants became smooth and spherical as the evaporated water was replaced by maltodextrin. Meanwhile, in F1, which contained surfactant during the lyophilization process, it remained possible for water to be retained in the surfactant.

The results of the overlay of an IR spectrum inspection of the glutathione-alginate microspheres can be seen in Figure 2. The interaction was characterized by shifts in wave numbers, loss of guluronate fingerprint absorption and an uptake of carboxylic salt groups (1614 cm<sup>-1</sup>) of Na alginate due to cross-linked reactions with CaCl<sub>2</sub>. From the results of the second IR-spectra examination of the formulas, the absorption of glutathione-specific groups still existed in all formulas. This means that glutathione was absorbed in the microspheres system without reacting with alginates.

**Table 3.** Particle size distribution of glutathione-Ca alginate microspheres and blank microspheres.


Distance size (µm)	Average of distance size (µm)	Blank Microspheres		F1		F2	
		n	nxd	n	nxd	n	nxd
0.64 – 1	0.76	69	52.44	31	23.56	3	2.28
1.01 – 1.37	0.94	115	136.8	24	22.56	47	44.18
1.38 – 1.74	1.29	76	124.8	124	159.9	97	125
1.75 – 2.11	1.68	34	57.12	98	164.6	85	142.8
2.12 – 2.48	2.22	2	4.6	18	39.96	36	79.9
2.49 – 2.85	2.41	1	2.41	5	12.05	28	67.5
2.86 – 3.22	3.10	2	6.2			1	3.1
3.23 – 3.59	3.39	1	3.39			3	10.2
<b>Average Diameter (µm)</b>			<b>1.34</b>	<b>1.40</b>		<b>1.58</b>	
<b>Polydispersity Index</b>			<b>0.003</b>	<b>0.003</b>		<b>0.003</b>	

The average diameter of Glutathione-Ca Alginate Microspheres was obtained from 300 particles of wet microspheres measured with an optical microscope.


Formula I: Glutathione Microspheres + surfactant HLB 7

Formula II: Glutathione Microspheres - surfactant

nxd: number of particles × average of distance size




**A**



**B**

**Figure 1.** The morphology of the shape and surface of the microspheres (A) F1 (GSH HLB 7), (B) F2 (GSH) observed through 5000x magnification scanning electron microscopy (SEM).



**Figure 2.** IR spectrum of glutathione-alginate microspheres.

On examination of specific Na-alginate group uptake, guluronate fingerprints were absent from all formulas. While the uptake of carboxylate salt groups was still present in all formulas, one of the two absorptions of  $1614\text{ cm}^{-1}$  was absent. The wavelengths of the carboxylic salt group in F1 and F2 were  $1423.52\text{ cm}^{-1}$  and  $1421.25\text{ cm}^{-1}$  respectively. The loss of one uptake within the carboxylic salt group and the guluronate fingerprint was due to a crosslinking reaction between the alginate and  $\text{CaCl}_2$  crosslinker involving ion exchange between the carboxylic group of guluronate acid and  $\text{Ca}^{2+}$  of crosslinkers.

The drug loading of glutathione in microspheres can be seen in Table 4 which also contains an analysis of entrapment efficiency and glutathione yield in microspheres. The determination of entrapment efficiency and drug loading used Na citrate  $0.5\text{ M}$  pH  $8.5$ . Na citrate solution was chosen as the medium within which the mechanism breaks down microspheres by replacing  $\text{Ca}^{2+}$  in crosslinked Ca carboxylic linkage with  $\text{Na}^+$ , thus rendering alginate soluble and causing glutathione to be re-dissolved. The data resulting from the examination of the glutathione content of microspheres produced was expressed as percentages. F1 was  $8.623\% \pm 0.12$  and F2 was  $7.81\% \pm 0.22$ .

**Table 4.** Drug loading, entrapment efficiency and yield of glutathione-Ca alginate microspheres.

Formula	Drug Loading $\pm\text{SD}(\%)$	EE $\pm\text{SD}(\%)$	Yield $\pm\text{SD}(\%)$
1	$8.623 \pm 0.12$	$54.46 \pm 2.4$	$94.03 \pm 3.07$
2	$7.81 \pm 0.22$	$48.49 \pm 2.37$	$93.34 \pm 3.65$

Formula I: Glutathione Microspheres + surfactant HLB 7

Formula II: Glutathione Microspheres - surfactant

The values are written as an average value mean  $\pm\text{SD}$  of drug loading, entrapment efficiency and yield of triplicates.

EE: Encapsulation efficiency; SD: Standard Deviation

The entrapment efficiency result for F1 was  $54.46\% \pm 2.40$ , while that for F2 was  $48.49\% \pm 2.37$ . From these

results, it could be seen that  $500\text{mg}$  of the drug could not be entrapped entirely by alginate at a concentration of  $2\%$  w/v. The resulting drug content remained relatively low. This was possibly due to the production of alginate microspheres using the current concentration of alginate and  $\text{CaCl}_2$  only being able to encapsulate this maximum capacity. Therefore, it is recommended to further optimize certain ratios of alginate polymer and crosslinker concentrations as a means of encapsulating a higher amount of glutathione. In this study, the  $2\%$  alginate concentration or  $1\text{M}$   $\text{CaCl}_2$  may need to increase so as to entrap and load larger amounts of the drug. Moreover, in order to produce optimum entrapment and drug loading efficiency, microspheres must consist of the necessary amount of both polymer and crosslinker to establish an optimal hydrogel composition which needs further experiments using several molar ratio composition.<sup>20,21</sup>

The ANOVA test results obtained did not differ significantly ( $p > 0.05$ ) because the concentration of alginate and  $\text{CaCl}_2$  used within both formulas was equal. Moreover, the addition of surfactant to glutathione did not affect microspheres entrapment efficiency. An analysis of F1 microspheres recovery confirmed yields as being  $94.03\% \pm 3.07$  (F1) and  $93.34\% \pm 3.65$  (F2). From these results, it could be seen that no difference in yield between F1 and F2 was existed because the amount of alginate and  $\text{CaCl}_2$  concentrations in both formulas was equal. The determination of yield recovery was aimed to quantify the extent of dry microspheres recovery of initial compounds added during the manufacture of microspheres (polymers and drug).<sup>22</sup> In future research, it is advisable to investigate the maximum capacity of the microspheres in order to obtain high drug loading, high entrapment efficiency and high yield by varying concentrations of alginate polymer and  $\text{CaCl}_2$ . Furthermore, the conducting of an in vitro release test is highly recommended.

Microspheres were subsequently subjected to stress tests which aimed at determining the stability of glutathione after microencapsulation (Figure 3, Table 5 and 6).

**Table 5.** Log Ct vs GSH remaining remaining after storage at temperatures of 50°C 0, 60°C and 80°C.

Days	Temperature 50°C				Temperature 60°C				Temperature 80°C			
	Log Ct	Log Ct Microsphere GSH <sub>-surf</sub> (F2)	Log GSH <sub>+surf</sub>	Log Microsphere Ct GSH <sub>+surf</sub> (F1)	Log Ct	Log Ct Microsphere GSH <sub>-surf</sub> (F2)	Log GSH <sub>+surf</sub>	Log Microsphere Ct GSH <sub>+surf</sub> (F1)	Log Ct	Log Ct Microsphere GSH <sub>-surf</sub> (F2)	Log GSH <sub>+surf</sub>	Log Microsphere Ct GSH <sub>+surf</sub> (F1)
0	2.356	2.343	2.406	2.410	2.358	2.344	2.406	2.411	2.378	2.410	2.340	2.410
1	2.355	2.342	2.405	2.409	2.357	2.341	2.406	2.410	2.357	2.406	2.339	2.402
3	2.348	2.337	2.404	2.407	2.343	2.333	2.404	2.407	2.274	2.343	2.26	2.36
5	2.337	2.295	2.402	2.405	2.33	2.289	2.400	2.404	2.175	2.293	2.17	2.32
<b>Regression curve (y=ax+b)</b>												
a	0.003	0.0091	0.0008	-0.0009	0.0005	0.0100	0.0012	0.0014	0.0414	-0.0248	0.0357	0.0186
b	2.357	2.35	2.406	2.41	2.36	2.3505	2.406	2.411	2.389	2.419	2.3575	2.415
r	-	-0.9878	-0.9796	0.9742	-	-0.9190	-0.9566	-0.9983	-0.993	-0.987	-0.979	-0.994
K	0.0085	0.0209	0.0018	0.0020	0.0133	0.023	0.0027	0.0032	-0.095	-0.057	-0.082	-0.043
Ln K	-4.767	-3.868	-6.319	-6.18	-4.319	-3.772	-5.914	-5.744	-2.35	-2.862	-2.501	-3.14

GSH: Glutathione; Surf: Surfactant

From the results of stress test, it was known that glutathione belonged to the first order because the plot of log Ct to t produced a straight line or linearity approaching 1. Based on the value of each compound, it was evident that the glutathione plus surfactant microspheres were more stable than glutathione with surfactant only. In addition, the glutathione with surfactant was more stable than the glutathione without surfactant (Figure 3). This was in accordance with the microencapsulation purpose of protecting glutathione from oxidation reactions.<sup>23</sup>

**Table 6.** Stability Linearity Curve of 0 order and 1<sup>st</sup> order at a temperature of 80°C.

Compounds	Linearity (r)	
	Zero Order	1 <sup>st</sup> Order
GSH <sub>-surf</sub>	-0.9972	-0.9930
Microspheres GSH <sub>-surf</sub> (F2)	-0.9871	-0.9897
GSH <sub>+surf</sub>	-0.9847	-0.9799
Microspheres GSH <sub>+surf</sub> (F1)	-0.9889	-0.9941

GSH: Glutathione; Surf: Surfactant

Based on the linearity of the stability value (r), it was apparent that the 1<sup>st</sup> order reaction was more linear (the value of r was close to 1), indicated that the reaction order of this microspheres system followed 1<sup>st</sup> order. Therefore, the determination of the constant value of glutathione degradation (k) used an equation formula of the 1<sup>st</sup> order. The averages produced by an MMP-1 test are presented in Table 7. Based on the statistical analysis of variance (ANOVA) within an MMP-1 expression test, a p-value (sig) of 0.000, less than 0.050, was obtained. The resulting glutathione-alginate microspheres, both those with additional surfactants and those without surfactants, were then mixed into the gel base for penetration evaluation. It was possible to evaluate the penetration test result from the MMP-1 level. The formula was able to penetrate when showing decreased levels of MMP-1 in mouse skin, having been exposed to ultra violet (UV) irradiation every two days, through the application of a dosage of 60 mJ/m<sup>2</sup> during each irradiation. The gel was applied to the skin twice a day, 20 minutes before irradiation (to give the topical absorption time into the skin) and four hours after irradiation (reactive oxygen species (ROS) initiated four hours after exposure). Topical application of the material was occurred on a day without irradiation. The gel base


was chosen because the microspheres were hydrophilic. Thus, it was appropriate to use gel as a carrier basis since it is elastic, easy to wash and has a cooling effect when applied to the skin on which it can readily be spread. In this study, the average MMP-1 expression in the treatment group smeared with glutathione microspheres gel with increased lipophilicity was lower than in either the glutathione gel without microspheres or control groups (Figure 4). The average level of MMP-1 expression in the control group consisting exclusively of gelling base was 72.03%, whereas the MMP-1 level of gel consisting of glutathione-alginate microspheres with increased lipophilicity was 15.44%. One-way ANOVA and post-hoc tests on the glutathione lipophilicity gel of the control group confirmed a significant increase.

**Table 7.** Study of MMP-1 expression of glutathione-Ca alginate microspheres.


Group	MMP-1 Expression (%)	SD
1	15.44	3.83
2	55.12	5.85
3	72.03	0.59

The values are presented as average mean values ± SD of percentage of MMP-1 expression. Each group consisted of triplicates. MMP-1: Matrix Metalloproteinase I; SD: Standard Deviation

Therefore, the results of the control group with glutathione lipophilicity gel increased significantly. It demonstrated that lipophilic glutathione microspheres were able to decrease MMP-1 expression in the dermis tissue of mice. This was due to formulas having increased their lipophilicity near the 2-3 Plog with the result that they penetrated the stratum corneum and entered the dermis network. Lipophilic glutathione increased on the gel microspheres. Increased MMP-1 expression was occurred after the skin of the mice was exposed to radiation for two weeks because the energy from UV radiation damages cell membranes and proteins. This, in turn, produces reactive oxygen species (ROS) which induce expression of proinflammatory cytokines binding to cell surface receptors including receptors of epidermal growth factor, interleukin (IL)-1, insulin keratinocyte growth factor and tumor necrosis factor (TNF).<sup>24-25</sup>



**Figure 3.** Correlation between Ln K and 1/T (temperature) of glutathione and glutathione microspheres. The data represented is Mean  $\pm$  SD. N average = three formulas in each group.



**Figure 4.** MMP-1 Expression of histology dermis networks of mice with immunohistochemical staining (A) Glutathione microspheres gel +surf, (B) Gel glutathione microspheres -surf, (C) Gel base (Magnification 400x). The yellow arrow indicates fibroblasts cells expressing MMP-1 (the surrounding cytoplasm is purplish to brownish), while the red arrows indicate fibroblast cells (cytoplasm around the bluish-colored) not expressing MMP-1

## Discussion

According to the results of this study, morphology changes were observed in the glutathione-alginate microspheres with the addition of surfactant compared to those without surfactant. Glutathione-alginate microspheres with the addition of surfactant with a HLB value equal to 7 possessed an almost spherical and smooth surface, whereas that of the glutathione microspheres without surfactant was not spherical and contained a number of holes. Almost spherical microspheres morphology was formed suggesting that non-spherical microspheres that may require a higher concentration of maltodextrin lyoprotectant which protects against mechanical stress and prevents aggregation during the freeze-drying process. This result was in accordance with those of a previous study using maltodextrin lyoprotectant to stabilize microsphere surfaces and improve dissolution properties.<sup>9, 26</sup>

In terms of resistance to oxidation, the glutathione plus surfactant microspheres were more resistance than either the glutathione with surfactant only or the glutathione without surfactant. This was in accordance with the microencapsulation's purpose of protecting glutathione from oxidation reactions.<sup>23</sup>

In addition, to increase entrapment efficiency and drug loading, higher concentrations of alginate polymer and CaCl<sub>2</sub> may be needed for future study of the optimum encapsulation process during the crosslinking process. Crosslinking of longer duration may need to be considered. Higher percentages of drug loading and entrapment efficiency of alginate microspheres have been shown to be necessary by other researchers employing a larger amount of alginate and CaCl<sub>2</sub> and crosslinking time in excess of one hour.<sup>27,28</sup>

Consequently, the optimized lipophilic glutathione-loaded alginate microspheres produced high in vivo effectiveness by decreasing MMP-1 expression in the dermis tissue of mice and penetrating the stratum corneum and dermis.

## Conclusion

Glutathione with surfactant loaded alginate microspheres has been successfully produced through aerosolization. The resulting small, spherical microspheres which were almost completely smooth were produced using alginate 2%. The enhanced lipophilicity of glutathione microspheres using surfactant with a HLB value equal to 7 was significantly more penetrative in nature than that of others, as indicated by the decreased levels of MMP-1.

## Acknowledgements

The author would like to thank the Faculty of Pharmacy, Universitas Airlangga and Airlangga Research and Innovative Institute for all support provided to the conduct of this research.

## Conflict of interests

The authors claim that there is no conflict of interest.

## References

1. Lushcak IV. Glutathione Homeostasis and Functions: Potential Targets for Medical Interventions. *J Amino Acids*. 2012;2012:1-26. doi:10.1155/2012/736837
2. Murray RK. Metabolism of xenobiotics. In: Murray RK, Bender DA, Botham KM, Kennelly PJ, Rodwell VW, Weil PA, editors. *Harper's Illustrated Biochemistry*. 28th ed. Michigan: McGraw-Hill; 2009. p. 612-3.
3. Dickinson DA, Forman HJ. Glutathione in defense and signaling: Lessons from a small thiol. *Ann NY Acad Sci*. 2002;973:488-504.
4. Nugrahaeni F, Melani Hariyadi D, Rosita N. Partition Coefficient and Glutathione Penetration of Topical Antiaging: Preformulation Study. *International Journal of Drug Delivery Technology*. 2018;8(2):39-43. doi:10.25258/ijddt.v8i2.13866
5. Choudhury PK, Kar M. Controlled release metformin hydrochloride microspheres of ethyl cellulose prepared by different methods and study on the polymer affected parameters. *J Microencapsul*. 2009;26(1):46-53. doi:10.1080/02652040802130503
6. Kim DG, Jeong YI, Choi C, Roh SH, Kang SK, Jang MK, et al. Retinol-encapsulated low molecular water-soluble chitosan nanoparticles. *Int J Pharm*. 2006;319(1-2):130-8. doi:10.1016/j.ijpharm.2006.03.040
7. Abdalla KF, Kamoun EA, Maghraby GM. Optimization of the entrapment efficiency and release of ambroxol hydrochloride alginate beads. *J Appl Pharm Sci*. 2015;5(4):13-9. doi:10.7324/japs.2015.50403
8. Lawrie G, Keen I, Drew B, Chandler-Temple A, Rintoul L, Fredericks P, et al. Interactions between alginate and chitosan biopolymers characterized using FTIR and XPS. *Biomacromolecules*. 2017;8(8):2533-41. doi:10.1021/bm070014y
9. Hariyadi DM, Purwanti T, Nirmala RN. Effect of Lactose and Maltodextrin on The Physical Characteristics of Ovalbumin-loaded Alginate Microspheres Produced by Aerosolization. 2nd Annual International Conference on Pharmacology and Pharmaceutical Sciences. 2014;5(2):26-9. doi:10.5176/2345-783x\_pharma14.29
10. Hariyadi DM, Lin SCY, Wang Y, Bostrom T, Turner MS, Bhandari B, et al. Diffusion loading and drug delivery characteristics of alginate gel microparticles produced by a novel impinging aerosols method. *J Drug Target*. 2010;18(10):831-41. doi:10.3109/1061186x.2010.525651
11. Mendes JBE, Riekes MK, de Oliveira VM, Michel MD, Stulzer HK, Khalil NM, et al. PHBV/PCL Microparticles for Controlled Release of Resveratrol: Physicochemical Characterization, Antioxidant Potential, and Effect on Hemolysis of Human Erythrocytes. *The Scientific World Journal*. 2012;2012:1-13. doi:10.1100/2012/542937

12. Griffith LG. Polymeric biomaterials. *Acta Mater.* 2000;48(1):263-77. doi:10.1016/s1359-6454(99)00299-2
13. Blessy M, Ruchi DP, Prajesh NP, Agrawal YK. Development of forced degradation and stability indicating studies of drugs - a review. *J Pharm Anal.* 2014;4(3):159-65. doi:10.1016/j.jpha.2013.09.003
14. Soni ML, Kumar M, Namdeo KP. Sodium alginate microspheres for extending drug release: formulation and in vitro evaluation. *International Journal of Drug Delivery.* 2011;2(1):64-8. doi:10.5138/ijdd.2010.0975.0215.02013
15. Khan H, Khan MF, Ali Khan B, Razaque G, Haque N, Akhter B, et al. Evaluation of the Interaction of Aluminium Metal with Glutathione in Human Blood Components. *Biomed Res.* 2012;23(2):237-40.
16. Singh MN, Hemant KSY, Ram M, Shivakumar HG. Microencapsulation: A promising technique for controlled drug delivery. *Res Pharm Sci.* 2010;5(2):65-77.
17. Alpert P, Oliver MJ. Drying without dying. In: *Desiccation and Survival in Plants: Drying Without Dying.* Black M and Pritchard HW, editors. Oxon, UK: CAB International/Wallingford; 2000. p. 3-43.
18. Hwang MY, Kim SG, Lee HS, Muller SJ. Generation and characterization of monodisperse deformable alginate and pNIPAM microparticles with a wide range of shear moduli. *Soft Matter.* 2017;13(34):5785-94. doi:10.1039/c7sm01079f
19. Dubey M, Shami TC, Rao KUB. Microencapsulation Technology and Applications. *Def Sci J.* 2009;59(1):82-95. doi:10.14429/dsj.59.1489
20. Patil P, Chavanke D, Wagh MA. A review on ionotropic gelation method: novel approach for controlled gastroretentive gelispheres. *Int J Pharm Pharm Sci.* 2012;4(4):27-32.
21. Kuo CK, Ma PX. Maintaining dimensions and mechanical properties of ionically crosslinked alginate hydrogel scaffolds in vitro. *J Biomed Mater Res A.* 2008;84A(4):899-907. doi:10.1002/jbm.a.31375
22. Zakir Hossain KM, Patel U, Ahmed I. Development of microspheres for biomedical applications: a review. *Prog Biomater.* 2015;4(1):1-19. doi:10.1007/s40204-014-0033-8
23. Bartolini D, Piroddi M, Tidei C, Giovagnoli S, Pietrella D, Manevich Y, et al. Reaction kinetics and targeting to cellular glutathione S-transferase of the glutathione peroxidase mimetic PhSeZnCl and its d,l-poly lactide microparticle formulation. *Free Radic Biol Med.* 2015;78:56-65. doi:10.1016/j.freeradbiomed.2014.10.008
24. Fagot D, Asselineau D, Bernerd F. Matrix metalloproteinase-1 production observed after solar simulated radiation exposure is assumed by dermal fibroblasts but involves a paracrine activation through epidermal keratinocytes. *Photochem Photobiol.* 2007;79(6):499-506. doi:10.1111/j.1751-1097.2004.tb01266.x
25. Brenneisen P, Sies H, Scharffetter-Kochanek K. Ultraviolet-B irradiation and matrix metalloproteinases: from induction via signaling to initial events. *Ann N Y Acad Sci.* 2002;973(1):31-43. doi:10.1111/j.1749-6632.2002.tb04602.x
26. Sansone F, Mencherini T, Picerno P, d'Amore M, Patrizia Aquino R, Rosaria Lauro M. Maltodextrin/pectin microparticles by spray drying as carrier for nutraceutical extracts. *J Food Eng.* 2011;105(3):468-76. doi:10.1016/j.jfoodeng.2011.03.004
27. Ullah Shah S, Socha M, Fries I, Gibaud S. Synthesis of S-nitrosoglutathione-alginate for prolonged delivery of nitric oxide in intestines. *Drug Deliv.* 2016;23(8):2927-35. doi:10.3109/10717544.2015.1122676
28. Khanna O, Larson JC, Moya ML, Opara EC, Brey EM. Generation of Alginate Microspheres for Biomedical Applications. *J Vis Exp.* 2012;66:3388. doi:10.3791/3388

# Pharmaceutical Sciences

(/)

ISSN: 1735-403X ([http://road.issn.org/issn\\_search?afs:query=1735-403X](http://road.issn.org/issn_search?afs:query=1735-403X)) eISSN: 2383-2886  
([http://road.issn.org/issn\\_search?afs:query=2383-2886](http://road.issn.org/issn_search?afs:query=2383-2886))

---

## Editors



**Editor-in-Chief**

Alireza Garjani, Pharm.D, Ph.D

Tabriz University of Medical Sciences, Iran

Email: garjania@tbzmed.ac.ir

**Associate Editor**

Ali Shayanfar, Pharm.D, Ph.D

Tabriz University of Medical Sciences, Iran

Email: shayanfara@tbzmed.ac.ir

<http://orcid.org/0000-0001-7507-9057>

**Associate Editor (Pharmacognosy)**

Sanaz Hamedeyazdan, Pharm.D, Ph.D

Tabriz University of Medical Sciences, Iran

Email: yazdans@tbzmed.ac.ir

<http://orcid.org/0000-0002-8950-9879>

---

**Advisory Board****Hadi Afrasiabi-Garekani**

Mashahd University of Medical Sciences, Iran

Email: hadiafrasiabi@yahoo.com

**Hossein Babaei**

Tabriz University of Medical Sciences, Iran

E-mail: babaeih@tbzmed.ac.ir



**Mohammad Barzegar-Jalali**

Tabriz University of Medical Sciences, Iran

Email: barzegar\_jalali@yahoo.com



**Siavoush Dastmalchi**

Tabriz University of Medical Sciences, Iran

Email: dastmalchi.s@tbzmed.ac.ir



**Robert J Dierksen**

University of Mississippi, USA

Email: rjd@olemiss.edu



**Alireza Ghanadi**

Isfahan University of Medical Sciences, Iran

Email: ghannadi@pharm.mui.ac.ir



**Taravat Ghafourian**

University of Sussex, UK

Email: T.Ghafourian@sussex.ac.uk



**Aliasgar Hemmati**  
Ahvaz University of Medical Sciences, Iran  
Email: hemmati\_aa@yahoo.com



**Nidal Jaradat**  
An-Najah National University, Palestine  
Email: nidaljaradat@najah.edu



**Fleming Martinez**  
Universidad Nacional de Colombia  
Email: fmartinezr@unal.edu.co



**Md. Moklesur Rahman Sarker**  
State University, Bangladesh  
Email: moklesur2002@yahoo.com



**Negar Motayaghani**  
David Geffen School of Medicine at UCLA, USA  
Email: Nmotayaghani@mednet.ucla.edu



**Seyed Alireza Mortazavi**

Shahid Beheshti University of Medical Sciences, Iran

Email: alirmortazavi@yahoo.com



**Khaled Nabih Zaki Rashed**

National Research Centre, Egypt

Email: khalednabih2015@yahoo.co.uk



**Hossein Nazemiyeh**

Tabriz University of Medical Sciences, Iran

Email: nazemih@tbzmed.ac.ir



**Ali Nokhodchi**

University of Sussex, UK

Email: A.Nokhodchi@sussex.ac.uk



**Ali Rajabi Siahboomi**

Colorcon Inc., Global Headquarters, Philadelphia, United States

Email: asiahboomi@colorcon.com



**Luca Rastrelli**

University of Salerno, Italy

Email: rastrelli@unisa.it



**Mahdi Rezayat**

Tehran University of Medical Sciences, Iran

Email: mhdirezayat43@gmail.com



**Omid Sabzevari**

Tehran University of Medical Sciences, Iran

Email: omid@tums.ac.ir



**Faiyaz Shakeel**

King Saud University, Saudi Arabia

Email: faiyazs@fastmail.fm



**Satya Sarker**

Liverpool John Moores University, UK

Email: S.Sarker@ljmu.ac.uk



**Afshin Zarghi**

Shahid Beheshti University of Medical Sciences, Iran

Email: zarghi@sbmu.ac.ir



**Mohammad Hossein Zarrintan**

Tabriz University of Medical Sciences, Iran

Email: zarrintan@tbzmed.ac.ir

[Submit Your Paper \(/Login\)](#)

[Editorial Board \(/EditorialBoard\)](#)

[Instructions for authors \(/InstructionsforAuthors\)](#)

[Contact editorial office \(/ContactUs\)](#)

[About \(/AimsandScope\)](#)

[Editorial policies \(/EditorialPolicies\)](#)

[Peer Review \(/PeerReview\)](#)

[Archive \(/Archive\)](#)

### Indexing & Abstracting:

**Scopus**<sup>®</sup> ([https://www.scopus.com/source/sourceInfo.uri?](https://www.scopus.com/source/sourceInfo.uri?sourceId=19700189400&origin=resultslist)

[sourceId=19700189400&origin=resultslist](https://www.scopus.com/source/sourceInfo.uri?sourceId=19700189400&origin=resultslist))

**CiteScore (2017): 1.26**

**SJR (2017): 0.30**

**SNIP (2017): 0.638**

([http://cassi.cas.org/publication.jsp?P=LgIBQf5Q2NQyz133K\\_II3zLPXfcr-WXfiqUFhuWc5h5Tugx6LjttiTLPXfcr-WXfimSBIkq8XcUjhmkOWtYxmzLPXfcr-WXfqJLb3NEY3CBvgLtWIG0ZUQ](http://cassi.cas.org/publication.jsp?P=LgIBQf5Q2NQyz133K_II3zLPXfcr-WXfiqUFhuWc5h5Tugx6LjttiTLPXfcr-WXfimSBIkq8XcUjhmkOWtYxmzLPXfcr-WXfqJLb3NEY3CBvgLtWIG0ZUQ)) (<http://jgateplus.com/search/footer-html/PublisherPartners.jsp>)

(<https://doaj.org/toc/2383-2886>)



([http://www.scimagojr.com/journalsearch.php?](http://www.scimagojr.com/journalsearch.php?q=19700189400&tip=sid&exact=no)

[q=19700189400&tip=sid&exact=no](http://www.scimagojr.com/journalsearch.php?q=19700189400&tip=sid&exact=no))

(<https://scholar.google.com/citations?user=58IjPBIAAAAJ&hl=en&authuser=1>)

([http://cassi.cas.org/publication.jsp?P=LgIBQf5Q2NQyz133K\\_II3zLPXfcr-WXfiqUFhuWc5h5Tugx6LjttiTLPXfcr-WXfimSBIkq8XcUjhmkOWtYxmzLPXfcr-](http://cassi.cas.org/publication.jsp?P=LgIBQf5Q2NQyz133K_II3zLPXfcr-WXfiqUFhuWc5h5Tugx6LjttiTLPXfcr-WXfimSBIkq8XcUjhmkOWtYxmzLPXfcr-)

WXfqJLb3NEY3CBvgLtWIG0ZUQ) **embase**<sup>®</sup>  
BIOMEDICAL ANSWERS


(<https://www.elsevier.com/solutions/embase-biomedical-research/embase-coverage-and-content>)



([http://cassi.cas.org/publication.jsp?P=LglBQf5Q2NQyz133K\\_II3zLPXfcr-](http://cassi.cas.org/publication.jsp?P=LglBQf5Q2NQyz133K_II3zLPXfcr-)

WXfiqUFHuWc5h5Tugx6LjttiTLPXfcr-WXfimSBIkq8XcUjhmkOWtYxmzLPXfcr-

WXfK8hX6TnCuU9w9UTjtlA)XtA)



(<https://doaj.org/toc/2383-2886>)

**Google scholar** (<https://scholar.google.dk/citations?user=58IjPBIAAAAJ&hl=en>)

Director-In-Charge:

Mohammad Reza Siahi (<http://pharmfac-en.tbzmed.ac.ir/?PageID=86&ID=31&BasesID=45>)

Editor-in-Chief:

Alireza Garjani (<http://pharmfac-en.tbzmed.ac.ir/?PageID=86&ID=15&BasesID=45>)



**No submission or publication fees**

(<http://ijisonline.org/ijpni/About>)







SJR

Scimago Journal &amp; Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

# Pharmaceutical Sciences

**Country** Iran - SIR Ranking of Iran

# 12

**Subject Area and Category** Pharmacology, Toxicology and Pharmaceutics  
Pharmacology, Toxicology and Pharmaceutics (miscellaneous)

H Index

**Publisher** Tabriz University of Medical Sciences**Publication type** Journals**ISSN** 23832886, 1735403X**Coverage** 2009-ongoing

Join the conversation about this journal

## Quartiles

Pharmacology, Toxicology and Pharmaceutics (miscellaneous)	2010	Q4
Pharmacology, Toxicology and Pharmaceutics (miscellaneous)	2011	Q4
Pharmacology, Toxicology and Pharmaceutics (miscellaneous)	2012	Q3
Pharmacology, Toxicology and Pharmaceutics (miscellaneous)	2013	Q4
Pharmacology, Toxicology and Pharmaceutics (miscellaneous)	2014	Q4
Pharmacology, Toxicology and Pharmaceutics (miscellaneous)	2015	Q3
Pharmacology, Toxicology and Pharmaceutics (miscellaneous)	2016	Q2
Pharmacology, Toxicology and Pharmaceutics (miscellaneous)	2017	Q2
Pharmacology, Toxicology and Pharmaceutics (miscellaneous)	2018	Q2



## SJR

Year	SJR
2010	0.102
2011	0.102
2012	0.133
2013	0.122
2014	0.133
2015	0.183
2016	0.293
2017	0.295
2018	0.252




## Citations per document



Total Cites

Self-Cites







External Cites per Doc

Cites per Doc



% International Collaboration





Citable documents

Non-citable documents



Cited documents

Uncited documents



Pharmaceutical Sciences

Show this widget in your own website

Q2  
Pharmacology, Toxicology and Pharmaceutics...  
best quartile

SJR 2018  
0.25

powered by scimagojr.com

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaq
```

F

Fadia Thamir Ahmed 9 months ago

Dear Mr

I am a pharmacist from Iraq and I have an article about the perceptions of pharmacy students about postgraduate studies, that I would like to publish in your journal. is the subject compatible with your journal system?

reply



**Elena Corera** 9 months ago

Dear Fadia,

thank you very much for your comment. Unfortunately, we cannot help you with your request, we suggest you to contact journal's editorial staff so they could inform you more deeply.

Best Regards,  
SCImago Team

### Leave a comment

Name

Email

(will not be published)




I'm not a robot

reCAPTCHA  
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2019. Data Source: Scopus®

EST MODUS IN REBUS  
Horatio (Satire 1,1,106)