

**International Journal of
Drug Delivery
Technology**

ISSN: 0975 4415
Peer Review Journal

International Journal of Drug Delivery Technology

ISSN: 0975 4415

Peer Review Journal

ARCHIVES ▾

HOME

EDITORIAL ▾

INSTRUCTIONS

CONTACT

SUBMIT MANUSCRIPT

Volume 7, Issue 1; January - March 2017

1. Development of An Antidiabetic Phytocomposite Loaded Phytoceutical Formulation, Its Quality Control and Pharmacokinetic Studies and Establishing *In Vitro- In Vivo* Correlation

De Baisakhki, Bhandari Koushik, Katakam Prakash, Adiki K Shanta, Mitra Analava

Abstract

2. Understanding the Impact of Polymer Ratio and Its Concentration on Omeprazole Release from Matrix Tablets: Response Optimization Study

Tiwari R, Tiwari G, Wal P, Wal A, Maurya P

Abstract

3. Development and Validation of a Simple HPLC-UV Method for The Quantification of Andrographolide In Rabbit Plasma

Syukri Y, Widarno I S, Adewiyah A, Wibowo A, Martien R, Lukitaningsih E, Nugroho A E

Abstract

4. Bosentan Monohydrate Vesicles Loaded Transdermal Drug Delivery System: *In Vitro In Vivo* Evaluation

Revathi M, Indira Muzib Y

Abstract

5. Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

Tristiana Erawati M, Retnowati, Amalia Wardatul F, Widji Soeratri

Abstract

6. Improved Solubility and Dissolution Rate of Ketoprofen by Beta Cyclodextrin Ternary Complexes Incorporating Hydrophilic Polymers

Mohammed Jafar, Sadath Ali, Hassan Mahmoud Ghonaim

Abstract

7. Acyclovir Loaded Solid Lipid Nanoparticle Based Cream: A Novel Drug Delivery System

EL- Assal M I A

Abstract

8. Anticancer Activity of Mixed Doxorubicin and Pravastatin in Nanoemulsions Against HCT 116 Colon Cancer Cells

Mayson H Alkhatib, Duaa K Zahim, Wadiyah S Backer


Abstract

9. Photostability Study on Character and Antioxidant Activity of Tomato Extract (*Solanum lycopersicum L.*) in Nanostructured Lipid Carrier (NLC) and Conventional Cream

Noorma Rosita, Dewi Melani Haryadi, Tristiana Erawati, Rossa Patria Nanda, Widji Soeratri

Abstract

Impact Factor: 1.529



UGC Approved Journal



This journal is present in UGC approved List of Journals for the purpose of Career Advancement Scheme (CAS) and Direct Recruitment of Teachers and other academic staff as required under the UGC (Minimum Qualifications for Appointment of Teachers and other Academic Staff in Universities and Colleges)

Other Journals published by International Society for Science and Nature



WWW.IJPQA.COM

International Journal of Pharmaceutical Quality Assurance



WWW.IJPCR.COM

International Journal of Pharmaceutical and Clinical Research



WWW.IJPPR.COM

International Journal of Pharmacognosy and Phytochemical Research



WWW.IJCPR.COM

International Journal of Current Pharmaceutical Review and Research



WWW.IJTPR.COM

International Journal of Toxicological and Pharmacological Research

The publication is licensed under Creative Commons License  View Legal Published by Dr. Yashwant Research Labs Pvt. Ltd. on behalf of International Society for Sci

Photostability Study on Character and Antioxidant Activity of Tomato Extract (*Solanum lycopersicum L.*) in Nanostructured Lipid Carrier (NLC) and Conventional Creame

Noorma Rosita*, Dewi Melani Haryadi, Tristiana Erawati, Rossa Patria Nanda, Widji Soeratri

Pharmaceutics Department of Faculty of Pharmacy, Universitas Airlangga, Surabaya, East Java, Indonesia.

Received: 18th Dec, 2016; Revised: 10th Feb, 2017; Accepted: 25th Feb, 2017; Available Online: 1st March, 2017

ABSTRACT

The aim of this study was to investigate the ability of NLC in increasing photostability of tomato extract in term of antioxidant activity. Photostability testing on antioxidant activity of samples were conducted by accelerating method using UV_B radiation 32.400 joule for 21 hours radiation. Antioxidant activity was measured by DPPH method. NLC was made by High Shear Homogenization (HPH) method at 24000 rpm for 4 cycles, while conventional creame was made by low speed at 400 rpm. The product were characterized include: pH, viscosity, and particle size. There were had difference characters and physical stability. NLC had smaller size, more homogenous and more stable than conventional creame. It was known that stability of antioxidant activity of tomato extract in NLC system higher than in conventional creame. That was showed with k value, as constanta of rate scavenging activity decreasing in antioxidant power between time (Sigma 2-tail < 0.005) of NLC and conventional creame were: $2.03 \times 10^{-2} \%/ \text{hour} \pm 0.08$ (3.94) and $4.71 \times 10^{-2} \%/ \text{hour} \pm 0.23$ (4.88) respectively.

Keywords: antioxidant, photostability, tomato extract, NLC, DPPH.

INTRODUCTION

The aging process of the skin consists of two processes, namely intrinsic aging and extrinsic aging. Intrinsic aging is the aging process that occurs naturally, caused by genetic factors, hormonal and racial. Extrinsic aging process is a process that is caused by factors outside the body such as sunlight (UV rays)¹. Photobiology effect of UV rays (UV_A and UV_B) produce free radicals that cause damage to the skin². It is therefore that the body needs antioxidants to protect the body from free radical attack³. The antioxidant protect the enzymes that repair DNA damages, thereby enhancing our body's ability to rejuvenate itself⁴. One antioxidant is derived from tomato extract (*Solanum lycopersicum L.*). Tomato extract is a natural source of antioxidants, especially lycopene as the highest content. Lycopene is one of about 600 types of carotenoids that act as antioxidants. This compound is a natural color forming pigment contained in tomatoes. Tomato extracts is unstable against light, heat, degradan and poorly soluble in water. As topical cosmetic preparations, a proper delivery system to improve the stability of the tomato extract was needed. One of delivery system was selected is *Solid Lipid Nanoparticle* (SLN). *Solid Lipid Nanoparticle* (SLN) is colloidal dispersion in nano size consists of a solid lipid stabilized by surfactants that work in the interface area⁵. Some advantages of SLN, for instance the ability to protect the active compounds from degradation, controlled release, the ability to load a

lipophilic or hydrophilic material, non-toxic and did not irritate.⁵ Limitations of SLN systems included loading capacity of the active ingredient was limited and the expulsion of active ingredient during storage. This was because of the high orderdness of the lattice crystal of solid lipid so that the active ingredient which has been entrapped easily squeezed out of the system. To overcome the limitations of the system SLN, a second-generation nanoparticle system that is *nanostructured lipid carrier* (NLC) was developed^{6,7}. NLC are formed with both liquid and solid lipid as a blend mixture in such a ratio that they are solid at room temperature. The active ingredient can be incorporated in particle matrix in a molecular dispersed form or it can be arranged as amorphous cluster⁶. SLN and NLC system were both ideal carrier cosmetic preparations because they contained lipid base provide emollient effect on the skin. The existence of liquid lipid decreased in the orderdness of solid lipid crystal lattice so that the active ingredient entrapped more stable and increased drug loading^{8,9}. The active ingredients that are entrapped in the system was expected to increase its stability and less contact with the free radicals exposed in the air. This research examined the influence of the NLC system against antioxidant stability of tomato extract which was compared to conventional creame. The parameters used to determine the stability of the antioxidant extract of tomato in the system, based on the value of a constant percent scavenging activity decreasing in antioxidant power

*Author for Correspondence: itanr@yahoo.com

between time (*k*). Stability test was done by acceleration test by 32,400 Joule UV_B rays exposure for 2, 5, 9, 15 and 21 hours.

MATERIALS AND METHODS

Materials

Tomato extract (the Asian Group CN Nutrition Lab), alcohol, distilled water and methanol, olive oil (PT Safarindo Internusa Jakarta), Kolliphor® P188 (Poloxamer 188) and tween 80 both were obtained from Sigma Aldrich.

Methods

Measurement of Tomato Extract Antioxidant activity (IC₅₀) by DPPH method^{4,10}. Each tomato extract solutions with various concentrations was added with 0.004% of DPPH solution and then allowed to stand for 30 minutes after the measurement of absorbance at a wavelength of maximum DPPH at three wavelengths, namely 505, 515, and 525 nm using the formula: Absorbance Calculate at 515 nm = A515 - ($\frac{A505+A525}{2}$).....(1)

Note: A515 nm = Absorbance of the sample solution at a wavelength 515 nm

A505 nm = Absorbance of the sample solution at a wavelength 505 nm

A525 nm = Absorbance of the sample solution at a wavelength 525 nm

A control = Absorbance of the DPPH solution as blanko

A sample = Absorbance of the sample solution

% free radical inhibition = 1 - ($\frac{\text{Absorbance Calculate of sample}}{\text{Absorbance Calculate of DPPH}}$) x 100%.....(2)

To determine IC₅₀ curves made between % inhibition data obtained as ordinate with concentration as absis. IC₅₀ value obtained by entering y = 50 into the regression equation y = bx + a wich was obtained from the curve of the relationship between free radical inhibition data to concentration, in order to get the value of x as the value IC₅₀.

NLC and conventional creame Preparation

NLC was made by diluting the tomato extract to molten lipid phase consisting of solid lipid and liquid lipid at 60 °C ± 5 °C (speed of 3,400 rpm for 2 minutes). The aqueous phase was added to the mixture was stirred for 5 minutes. Further stirring at a speed of 24,000 rpm for 8 minutes used Ultra Turax IKA® T-25 Digital High Shear Homogenizer. The next step was cooling process until it reached room temperature. Preparation of conventional cremes did like the NLC preparation but made with a hot plate Dragon Lab MS H-Pro with low speed (400 rpm). The composition of formula NLC and conventional cremes can be seen in table 1.

Sample Characterizations were done by measuring pH, viscosity, particle size, Polydispersity Index (PI)

Determination of pH

Determining pH of each test preparation was done using a pH meter.

Viscosity Measurement

Viscosity measurement was done using a Brookfield Cone and Plate Viscometry®. It was performed for each sample by diluting the sample 1: 9 with CO₂ free aquadest.

Table 1: Formula NLC and conventional creame (CC).

Ingredient	Function	Concentration (%v/w)	
		NLC	Conventional Creame (CC)
tomato extract (TE)	active ingredient	0.25	0.25
cetyl alcohol	solid lipid	18	18
olive oil	liquid lipid	2	2
Poloxamer 188	surfactant	5	5
Tween 80	surfactant	5	5
aquadest	water phase	ad 100	ad 100

Table 2: Characterisation of NLC and conventional creame.

Sample	Characterisation of NLC and conventional creame ± SD (CV)		
	pH	Viscosity	
NLC	5.59 ± 0.02 (0.27)	2,416.33 ± (7.72)	186.50
conventional creame	5.29 ± 0.03 (0.47)	1,563.33 ± (5.50)	85.82

Table 3: The constanta of rate scavenging activity decreasing in antioxidant power.

Sample	Mean constanta of rate scavenging activity decreasing in antioxidant power (%/hour) ± SD (CV)
NLC	2.03x10 ⁻² ± 0.08 (3.94)
conventional creame (CC)	4.71x 10 ⁻² ± 0.23 (4.88)
Tomato extract (TE)	4.41x10 ⁻² ± 0.2 (4.53)

Examination of the particle size and size distribution

Examination of the particle size consisting of two treatments before and after UV_B irradiated for 21 hours. Each sample was diluted to 200 ppm stirring for 3 minutes at 500 rpm. Dilution checked by Delsa Nano Particle Size on the angle of 65° and a temperature of 25°C. Data obtained were average particle diameter and Polydispersity Index (PI). All measurement was replicated 3 times.

Antioxidant Activity Photostability Test

Antioxidant Stability for each dosage was calculated by the percent inhibition using formula 1. Tomato extracts in conventional cremes and NLC were measured free radical activity inhibition percent after being exposed to UV_B rays produced from 32,400 Joule UV_B bulbs. The experimental conditions were controlled at 30 °C ± 5 °C. Free radical inhibition every 2, 5, 9, 15, and 21 hours after exposure to UV_B were determined. The results of calculations per cent inhibition between the time was made curve. Furthermore, *k* value as constanta of the rate of scavenging activity decreasing in antioxidant power were determined based on the appropriate reaction order.

RESULTS AND DISCUSSION




Figure 1: Curve of free radical inhibition of tomato extract.




Figure 2: Histogram of average particle size of NLC and Conventional cream (CC) without and after exposure to UV_B for 21 hours.




Figure 3: The curve of scavenging activity decreasing in antioxidant power.

Tomato Extract Antioxidant Activity

Antioxidants activity of tomato extracts were determined by DPPH method^{4,9}. DPPH free radical compounds are often used to test the antioxidant activity of several compounds or extracts of natural ingredients. DPPH compound is a free radical that has a purple color in solution. If there is a compound that can donate a proton, DPPH will be in a reduced form and its color intensity decreased. This study showed that the IC₅₀ of tomato extract was 0.0662% (Figure 1). This means that at 0.0662% tomato extract reduced 50% DPPH radical activity, or 0.14% tomato extract inhibited 100% free radicals. In this study, all samples used 0.25% tomato extract.

Physical Characteristic and Its Stability of NLC and Conventional Cream

The result of pH and viscosity of both samples were shown at Table 2. Both samples had an appropriate pH with skin (4.5 - 6.5). That was not irritable to the skin and comfortable to use. The viscosity of conventional cream was higher than NLC. It seemed associated with the particle size of the samples (Figure 2). The average particle size of NLC was 210 nm, while of conventional cream was about 662 nm approximately. It made NLC more viscous. The particle size of conventional cream is bigger than NLC because of the speed of mixing in conventional cream preparation was lower than NLC. Energy produced from mixing at low speed was not able to break the molten lipid droplets to be a small size. This was due to the energy produced was not capable to be distributed efficiently than when performed at high speed. By exposing UV_B radiation for 21 hours led to an increase in particle size. Greater increase occurred in the conventional

creames. The particle size of conventional creames was found to be 1,384 nm, while the NLC was 258 nm. It was caused by the energy of UV_B rays reduced the viscosity of the preparation, thus boosted the rate of flocculation of particles. The polydispersity Index (PI) NLC before and after irradiated <0.3. PI close to 0 indicated a homogenous dispersion, but when PI greater than 0.3 indicated a high heterogeneity¹⁰. It showed that the NLC had a homogeneous particle size. In conventional creames, UV_B exposure showed to degrade homogeneity of particle size, it showed with the change of PI value was 0.27 to be 0.42

Antioxidant of Tomato Extract Photostability Test in NLC and Conventional Creame

Antioxidant photostability test was conducted using accelerated method by exposure to UV_B radiation. The reduction of antioxidant activity over time can be seen in Figure 3. Stability antioxidants indicated by the value of k , as constant percent scavenging activity decreasing in antioxidant power between time. The greater the value of k indicated that the rate of scavenging activity decreasing in antioxidant power is high, it means that an antioxidant power was unstable. To determine the value of k , it had to know the order of the reaction previously. It was known that the rate of change percent reduction between the time progresses according to order 1. Order 1 means that the reaction rate was influenced one aspect, in this case was the content of active material¹¹. The following Table 3 contains the mean constant of rate scavenging activity decreasing in antioxidant power (%/hour) (k) samples. The value of k of tomato extract in conventional creame was not difference significantly with tomato extract itself, but greater than NLC. It showed that conventional creame could not be able to protect tomato extract from photo degradation. In NLC system, the active ingredient were entrapped in solid lipid, it would be protect the active ingredient from environmental influences.

CONCLUSIONS

It can be concluded that the NLC can improve the stability of the tomato extract antioxidant power than if only formulated in conventional creames.

REFERENCES

- Jadoon, S., Karim, S., Hassham, M., Bin, H., Akram, M. R., Khan, A. K., Murtaza, G., 2015, Anti-Aging Potential of Phytoextract Loaded-Pharmaceutical Creames for Human Skin Cell Longevity.
- Fisher, G. J., Kang, S., Varani, J., Bata-Csorgo, Z., Wan, Y., Datta, S., & Voorhees, J. J., 2002, Mechanisms of photoaging and chronological skin aging. *Archives of Dermatology*. 138(11), pp 1462-1470.
- Masaki, H. 2010. Role of antioxidants in the skin: Anti-aging effects. *Journal of Dermatological Science*, 58(2). pp 85-90.
- Vidhya, Suji, Dhatchayani, Priya, Rao Bhaskara, 2015, Evaluation of In-vitro Antioxidant, Antimicrobial Activities and GC-MS Analysis of *Alternanthera bettzickiana* Linn. Leaf Extracts, *International Journal of Pharmacognosy and Phytochemical Research*; 7(6); 1072-79.
- Ekambaram P., Shatali A. A. H., Priyanka K. 2011. *Solid Lipid Nanoparticle: A Review*, Scientific Review and Chemical Communication. Vol. 2. pp 80-102.
- Prithu Pathak, A.K. Pathak, 2015, Preparation and Evaluation of Nano-Structured Lipid Carriers of Azelaic Acid in Topical Formulation (Hydrogel), *International Journal of Current Pharmaceutical Review and Research*; 6(3); 188-194.
- Montenegro, Lucia, 2014, *Nanocarriers For Skin Delivery of Cosmetic Antioxidants*, *Journal of Pharmacy & Pharmacognosy Research*, Vol. 2, No. 4, pp 73-92.
- Loo CH, et al, 2013, *Effect of Composition in Nanostructured Lipid Carriers (NLC) on Skin Hydration and Occlusion*. *International Journal of Nanomedicine*, Vol.8, pp 13-22.
- Fidrianny Irda, Sefiany Esther, Ruslan Komar, 2015, In-vitro Antioxidant Activities from Three Organs of White Ambon Banana (*Musa AAA Group*) and Flavonoid, Phenolic, Carotenoid Content, *International Journal of Pharmacognosy and Phytochemical Research*; 7(3); 590-96.
- Ammar et al. Solid Acid Loaded Lipid Nanocarriers With Promotes Skin Antiaging And Antioxidant Efficacy. *Journal of Drug Delivery Science and Technology*. 2016, 31: 72-82.
- Sinko, P.J., 2012, *Martin's Physical Pharmacy and Pharmaceutical Sciences*. 6th Ed. New Jersey, pp 441-443.



INTERNATIONAL JOURNAL OF DRUG DELIVERY TECHNOLOGY

Home > About the Journal > Editorial Team

Editorial Team

Editor

Mr. Pradeep Kumar Tiwari, MRI Publication Pvt. Ltd.(OPC), India

ISSN: 0975-4415

USER

Username

Password

Remember me

LOGIN

NOTIFICATIONS

View

Subscribe

JOURNAL CONTENT

Search

International Journal of Drug Delivery Technology

ISSN: 0975 4415

Peer Review Journal

[ARCHIVES](#) ▾[HOME](#)[EDITORIAL](#) ▾[INSTRUCTIONS](#)[CONTACT](#)[SUBMIT MANUSCRIPT](#)

EDITOR IN CHIEF

Prof. Dina Nath Mishra
Professor and Head of Pharmaceutics, Department of Pharmaceutical Sciences,
Guru Jambheshwar University of Science and Technology, INDIA

Board Members

Dr. Shailendra K. Singh
Guru Jambheshwar University of Science and Technology, INDIA

Dr. Somnath Singh
Creighton University, Omaha, USA

Dr. Parshuram Roy
Himachal Institute of pharmaceutical & Research, HP, INDIA

Dr. Tathagata Dutta
University of Queensland, Brisbane, AUSTRALIA

Dr. Ashish Suttee
Lovely Professional University, Phagwara, INDIA

Dr. Kalpesh Gaur
Geetanjali College of Pharmaceutical Studies, Udaipur, INDIA

Dr. Vishal Gupta
Director, Research & Development Covidien, USA

Dr. Chandan M. Thomas
Department of Pharmaceutical Sciences, Lake Erie College of Osteopathic Medicine and School of Pharmacy 5000 Lakewood Ranch Blvd, Bradenton, Florida-34211

Prof. Kamla Pathak
Rajiv Academy of Pharmacy, Mathura, INDIA

Prof. V. R. Sinha
Panjab University, Chandigarh, INDIA

Prof. Pramil Tiwari
National Institute of Pharmaceutical Education and Research (NIPER), Mohali, INDIA

Prof. Arun Nanda
Faculty of Pharm. Sciences, Maharshi Dayanand University, Rohtak, INDIA

Prof. O.P. Katare
Panjab University, Chandigarh, INDIA

Dr. Amit Bhatia
Lovely Professional University, Punjab, INDIA

Dr. Anil Phillip
Rajiv Academy of Pharmacy, Mathura, INDIA

Dr. Dinesh Kaushik
Hindu College of Pharmacy, Sonapat, INDIA.

Dr. Munish Ahuja
Dept. of Pharm. Sciences, Guru Jambheshwar University of Science and Technology, Hisar, INDIA

Dr. Sanju Nanda
Dept. of Pharm. Sciences, M.D. University, Rohtak, INDIA


Dr. Rakesh P. Patel
S.K. Patel College of Pharm. Edu. & Res., Ganpat University, Gujarat, INDIA

Dr. Bhaskar Mazumder
Dept. of Pharmaceutical Sciences, Dibrugarh University, Dibrugarh, Assam, INDIA

Dr. Kalpana Nagpal
Apeejay Satya University, Sohna, Gurgaon, Haryana, INDIA

Submit Manuscript | Contact IJDDT | Join Editorial | Accepted Manuscripts | Home

Impact Factor: 1.529



UGC Approved Journal



This journal is present in UGC approved List of Journals for the purpose of Career Advancement Scheme (CAS) and Direct Recruitment of Teachers and other academic staff as required under the UGC (Minimum Qualifications for Appointment of Teachers and other Academic Staff in Universities and Colleges)

Other Journals published by International Society for Science and Nature



WWW.IJQA.COM

International Journal of Pharmaceutical Quality Assurance



WWW.IJPCR.COM

International Journal of Pharmaceutical and Clinical Research



WWW.IJPPR.COM

International Journal of Pharmacognosy and Phytochemical Research



WWW.IJCPR.COM

International Journal of Current Pharmaceutical Review and Research



WWW.IJTPR.COM

International Journal of Toxicological and Pharmacological Research

The publication is licensed under Creative Commons License  View Legal Published by Dr. Yashwant Research Labs Pvt. Ltd. on behalf of International Society for Sci

International Journal of Drug Delivery Technology

ISSN: 0975 4415

Peer Review Journal

ARCHIVES ▾	HOME	EDITORIAL ▾	INSTRUCTIONS	CONTACT	SUBMIT MANUSCRIPT
------------	------	-------------	--------------	---------	-------------------

INDIA

Published by: DYR Labs Pvt. Ltd.
1/14, Housing Board, devipura, Sikar, India 332001


Branch: Publishers, In-IJDDT
35, Printers Nagar II
Sitabari Tonk Road
Jaipur, Rajasthan

Emails

For Submission: ijddtjournal@gmail.com

[Submit Manuscript](#) | [Contact IJDDT](#) | [Join Editorial](#) | [Accepted Manuscripts](#) | [Home](#)

Impact Factor: 1.529



UGC Approved Journal



This journal is present in UGC approved List of Journals for the purpose of Career Advancement Scheme (CAS) and Direct Recruitment of Teachers and other academic staff as required under the UGC (Minimum Qualifications for Appointment of Teachers and other Academic Staff in Universities and Colleges)

Other Journals published by International Society for Science and Nature



WWW.IJPQA.COM

International Journal of Pharmaceutical Quality Assurance



WWW.IJPCR.COM

International Journal of Pharmaceutical and Clinical Research



WWW.IJPPR.COM

International Journal of Pharmacognosy and Phytochemical Research




WWW.IJPCR.COM

International Journal of Current Pharmaceutical Review and Research



WWW.IJTPR.COM

International Journal of Toxicological and Pharmacological Research

The publication is licensed under Creative Commons License  View Legal Published by Dr. Yashwant Research Labs Pvt. Ltd. on behalf of International Society for Sci

**SJR**

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

International Journal of Drug Delivery Technology

Country Australia - SIR Ranking of Australia**Subject Area and Category** Pharmacology, Toxicology and Pharmaceutics
Pharmaceutical Science**Publisher** International Journal of Drug Delivery Technology**Publication type** Journals**ISSN** 09754415**Coverage** 2011-ongoing**7**

H Index

Join the conversation about this journal

Quartiles

Pharmaceutical Science	2012	Q4
Pharmaceutical Science	2013	Q4
Pharmaceutical Science	2014	Q3
Pharmaceutical Science	2015	Q3
Pharmaceutical Science	2016	Q3
Pharmaceutical Science	2017	Q3

Pharmaceutical Science 2018 Q3







SJR

scientific discussion an average article of the journal is.

Year	SJR
2012	0.112
2013	0.132
2014	0.159
2015	0.220
2016	0.144
2017	0.123
2018	0.213

Citations per document

Total Cites Self-Cites



International Journal of Drug Delivery Technology

Q3 Pharmaceutical Science
best quartile

SJR 2018 0.21

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com/journalsearch.php?q=20500195212&tin=sid&clean=0" style="border: 1px solid black; padding: 2px 5px; display: inline-block;">https://www.scimagojr.com/journalsearch.php?q=20500195212&tin=sid&clean=0
```

D

Dr. Zainab Al-Sharifi 5 days ago

hi sir:

I would like to publish my paper in your journal, my name is dr. Zainab Al-Sharifi am from Iraq, Collage of medicine / University of Baghdad

reply

Leave a comment

Name

Email


(will not be published)

 I'm not a robot
reCAPTCHA
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2019. Data Source: Scopus®

EST MODUS IN REBUS
Horatio (Satire 1.1, 106)

