

FONT SIZE

HOME ABOUT LOGIN REGISTER SEARCH CURRENT
ARCHIVES ANNOUNCEMENTS ARTICLE TEMPLATE INDEXING

Home > About the Journal > Editorial Team

USER

Username
Password

Remember me

EDITORIAL TEAM

CHIEF EDITOR (VALID FROM JANUARY 1, 2019)

Dr. Heru Santoso Wahito NUGROHO, Poltekkes Kemenkes Surabaya (Health Polytechnic of Surabaya), Indonesia

EDITORS (VALID FROM JANUARY 1, 2019)

Joel Rey U. ACOB, Department of Nursing, Visayas State University, Philippines
Dr. Sahrir SILLEHU, Department of Public Health, STIKes (College of Health Science) "Maluku Husada", Indonesia
Tanko Titus AUTA, Department of Nursing, Niger State Ministry of Health, Minna, Nigeria
Wiwin MARTININGSIH, Poltekkes Kemenkes Malang (Health Polytechnic of Malang), Indonesia
Somsak THOJAMPA, School of Nursing, Naresuan University, Thailand
Dr. Hadi PRAYITNO, Department of Public Health, Universitas Jember, Indonesia
Dr. Mfuh Anita Y. LUKONG, Department of Nursing Science, Faculty of Allied Health Sciences, College of Health Sciences, Ahmadu Bello University, Nigeria
Victoria KALUSOPA, School of Nursing, University of Zambia, Zambia

ADMINISTRATOR (VALID FROM JANUARY 1, 2019)

Suparji SUPARJI, Poltekkes Kemenkes Surabaya, Indonesia

"HEALTH NOTIONS" ISSN: 2580-4936 (online version only), published by Humanistic Network for Science and Technology

Cemara street 25, 001/002, Dare, Ds./Kec. Sukorejo, Ponorogo, East Java, Indonesia, 63453

OPEN JOURNAL SYSTEMS

Journal Help

NOTIFICATIONS

View
Subscribe

JOURNAL CONTENT

Search
Search Scope
All

Browse


By Issue
By Author
By Title
Other Journals

INFORMATION

For Readers
For Authors
For Librarians

KEYWORDS

Anemia, Learning Achievement Index, Midwifery Students Anger Big five personality, Coronary heart disease, Personality Type Capacity Instruments, Disaster Mitigation, Campus Resilience To Disasters Child Health, Outbreak Child and Adolescent Psychiatry Childbirth assurance, people's habits, residence distance, travel time Macrosomia, Diabetes Mellitus, Junk Food, Parity, Macrosomia History Management Maternity nursing Metastatic bone disease, pathological fracture, clinical features Nursing Care Module, Pulmonary TB Nutritional deficiency, the Oral Cavity Hygiene, Oral Candidiasis PCOS Pregnant Women Social capital, Perception, Commitment, Behavior Socio-demographic characteristics, immunization, measles, rubella, tongkuno, indonesia adolescent health infertile maternity nursing multivessel disease



FONT SIZE

USER

Username

Password

Remember me

HOME ABOUT LOGIN REGISTER SEARCH CURRENT
 ARCHIVES ANNOUNCEMENTS ARTICLE TEMPLATE INDEXING

Home > Archives > Vol 2, No 9 (2018)

VOL 2, NO 9 (2018)

SEPTEMBER

TABLE OF CONTENTS

ARTICLES

Self-Management of Patients with Diabetes Mellitus: An Integrative Literature Review Chatarina Hatri Istiari	PDF 901-910
Correlation Between History of ISTC Training on Independent Practicing Doctor With Discovery of TB Suspected Children in Surabaya City Madu Puspita Nuansa Jatu, Fariani Syahrul	PDF 911-917
Navigating the Storm: Integrative Review of Attrition Factors Among Undergraduate Nursing Students Carmela Theresa T. De Leon	PDF 918-926
Knowledge, Attitudes and Application of Child Safety by Families in Karunrung Village, Rappocini, Makassar Baharuddin Baharuddin, Nasrullah Nasrullah, Muhasidah Muhasidah, Sri Angriani	PDF 927-929
Preparation and characterization of BSA-loaded Chitosan Microspheres Esti Hendradi, Retno Sari, Rifka Anggraini Anggai	PDF 930-934
The Efforts to Control and Prevent Cervical Cancer through Early Detection Using the VIA Test in the East Java Provincial Health Office Riza Muhammad Zulham, Chatarina Umbul Wahyuni, Muhammad Atoillah Isfandiari, Hasan Huda	PDF 935-942
Both Acute and Chronic Exercise Decrease Total Cholesterol Level in Human Blood Bambang Purwanto, Wahyu N. Pratiwi, Abdurachman Abdurachman	PDF 943-945
Optimization Ca-Alginate Beads as Drug Reservoir in the Patch System for Model Antinflammatory Drug Esti Hendradi, Isnaeni Isnaeni, Rifaatul Mahmudah	PDF 946-951
Psychological Factors Affecting the Incidence of Essential Hypertension in General Hospital of Makassar Yonathan Ramba, Suharto Suharto, M. Bagus Qomaruddin, Dewi Retno Suminar, Hendrik Hendrik	PDF 952-955
Risk Factors of HIV and AIDS Mortality At Ibnu Sina Hospital Silvanus Roga Misi, Chatarina Umbul Wahyuni, Hari Basuki Notobroto	PDF 956-960
Evaluation of POMP for Filariasis and Factors Associated with Its Problems in East Manggarai Regency in 2017 Fransiscus Salesius Onggang	PDF 961-968
Characterization of Indigenous Lactic Acid Bacteria (LAB) from the wild horse milk of Bima Narda Widiada	969-971
Effectiveness of A Combination of Ergonomic Exercise Therapy and Deep Breathing Relaxation to Reduce Muscle Pain (Myalgia) in Unskilled Workers (Dock Laborers) in Bima Port, Bima City in 2018 Kurniadi Kurniadi, Ana Triana Wulandari, Julhana Julhana	972-975
The Effectiveness of the Mother's Class in Efforts to Reduce Incidence of LBW and Efforts to Increase ASI Intake for LBW in Gunung Sari Health Center, West Lombok Siti Aisyah, Ni Nengah Arini Murni	976-980

OPEN JOURNAL SYSTEMS

Journal Help

NOTIFICATIONS

View
Subscribe

JOURNAL CONTENT

Search

Search Scope
All

Browse
By Issue
By Author
By Title
Other Journals

INFORMATION

For Readers
For Authors
For Librarians

KEYWORDS

Anemia, Learning Achievement Index, Midwifery Students **Anger** Big five personality, Coronary heart disease, Personality Type Capacity Instruments, Disaster Mitigation, Campus Resilience To Disasters Child Health, Outbreak Child and Adolescent Psychiatry Childbirth assurance, people's habits, residence distance, travel time Macrosomia, Diabetes Mellitus, Junk Food, Parity, Macrosomia History Management **Maternity nursing** Metastatic bone disease, pathological fracture, clinical features Nursing Care Module, Pulmonary TB Nutritional deficiency, the Oral Cavity Hygiene, Oral Candidiasis PCOS **Pregnant Women** Social capital, Percption, Commitment, Behavior Socio-demographic characteristics, immunization, measles, rubella, tongkuno, indonesia adolescent health infertile maternity nursing multivessel disease

CURRENT ISSUE

<http://heanoti.com/index.php/hn>

RESEARCH ARTICLE

URL of this article: <http://heanoti.com/index.php/hn/article/view/hn20905>

Preparation and characterization of BSA-loaded Chitosan Microspheres

Esti Hendradi^{1 (CA)}, Retno Sari², Rifka Anggraini Anggai³^{1(CA)}Department of Pharmaceutics, Faculty of Pharmacy, Airlangga University, Indonesia; esti-h@ff.unair.ac.id
(Corresponding Author)²Department of Pharmaceutics, Faculty of Pharmacy, Airlangga University, Indonesia³Department of Pharmaceutics, Faculty of Pharmacy, Airlangga University, Indonesia

ABSTRACT

Proteins are unstable molecules because of their complex structure and easily degraded by enzymatic systems. Biodegradable microspheres have been used in protein delivery system. Bovine serum albumin was chosen as a model protein. Chitosan/tripolyphosphate microparticles have already been used to encapsulated BSA. The aim of this work was to microencapsulate these protein-CS/TPP by ionic gelation and freeze-drying. The morphology, particle size, and encapsulation efficiency of the prepared microspheres were investigated. The results revealed that the microspheres exhibited good sphericity with ranged in size from 2.25 to 2.73 μm . The study found that the moisture content of microspheres was 2.8%, the encapsulation efficiency of BSA in CS-TPP microspheres ranged from 88.14 to 96.25%. The results suggested that ionotropic gelation method will be an effective method for fabricating chitosan microspheres for delivery of protein..

Keywords: Microspheres, BSA, Ionic Gelation

INTRODUCTION

Background

Inhalation therapy was development in decades with the specificity to delivery peptides, proteins and drugs with small molecules within uptake in the lung⁽¹⁾. Inhalation one of the treatment process by inhaling the drugs, that can directly enter in to the lungs as the target organ⁽²⁾.

Pharmacokinetic studies of inhaled macromolecular therapy showed very high bioavailability, with the large surface area of alveolar, low fluid volume on the lung surface, ultra thin diffusion layer, and paracellular transport ranged in pore size 1-5 nm⁽¹⁾. Delivery of protein/peptide in inhalation route is difficult because the process of denaturation proteins during manufacture, aerolization and storage, inefficient deliver. To overcome this problem, biocompatible polymer-based delivery system can be used to improve protein stability and increase absorption through the lungs, improve the aerodynamic properties of particles, and have a controlled release effect and free from natural phagocytosis clearance in the lungs⁽³⁾.

Polymeric nanoparticles (NPs) have been an attractive approach for lung delivery due to their ability to enter intracellular compartments and escape macrophages phagocytosis. Furthermore, they provide the possibility of achieving high drug loading capacity, control release, increased stability and high drug absorption and appropriate therapeutic target deposition⁽⁴⁾. Chitosan is the second longest polymer after cellulose and the only alkaline polysaccharide in nature, with a low immunity that makes chitosan possible as a carrier for drug delivery⁽⁵⁾. Chitosan-based polymers in nano- microparticles are mucoadhesive and able to open tight junctions between epithelial cells. The characteristics can stimulate the absorption of proteins/antigens by epithelial cells and macrophage cells⁽⁶⁾.

Bovine serum albumin (BSA) has the isoelectric value of 4.7 and was used as a protein model⁽⁷⁾. To delivering BSA as a protein model, chitosan has a major role affecting the shape of the microsphere, 2% chitosan in the formula produces a microspheres with a very spherical shape with a homogeneous microspheres size⁽⁸⁾. Several methods can be used to create chitosan particulate systems, such as ionic gelation, emulsion-droplet coalescence, co-preservation/precipitation, and reverse micellar⁽⁹⁾. Microspheres were prepared by ionotropic gelation methods with aerolization techniques has the advantage of maintaining the integrity of proteins trapped

in microspheres⁽¹⁰⁾. Chitosan microparticles were prepared by crosslinking with tripolyphosphate (TPP) through ionic interactions between the positive charge of the amino group and the negative ion content of the TPP⁽¹¹⁾. Enhanced crosslinker concentration in microspheres making increasing efficiency encapsulation, because the rigid and complex structure of particles are formed and can enhanced the drug content of the drug⁽¹²⁾.

Based on the previous consideration, we hypothesized that these TPP concentration can effect on physical the characteristics of microspheres and bovine serum albumin will be trapped in a microsphere system using ionic gelation method with freeze drying with an smaller particle size.

METHODS

Chemicals

Chitosan with a dacencylation degree of 87.7% was purchased from Biotech Surindo (Cirebon, Indonesia), Bovine Serum Albumin (BSA) was purchased from Sigma-Aldrich, Castle Hill, NSW, Australia. TPP was obtained from Sigma-Aldrich. Maltodextrin (Dextrose equivalence 12-15) was obtained from Qingdao Shengda Commercial & Trade Co.,Ltd, All other reagents were of analytical grade and were used without further purification.

Preparation of chitosan microspheres

Ionic gelation with freeze drying was used as a method in this study. Amount of BSA was dissolved in aqua dem. Acetic acid solutions was used to make the swelling properties of chitosan by stirred at 500 rpm. BSA solution was added into chitosan solution. TPP solution were prepared by dissolving TPP in aqua dem. Dispersion solution of BSA-chitosan was sprayed into an aqueous TPP cross-linking solution via peristaltic pump using speed controller. Microspheres were purified were centrifugated for 15 minutes at 2500 rpm to wash the crosslinker and acetic acid in solution. The sediment then was suspended in water. Freeze-dried BSA-loaded CS microspheres were prepared by adding maltodextrin as a lyoprotectant. Samples were initially frozen by storing at -80 °C overnight and freeze-dried.

Size and Surface Morphological

Microsphere particles were analyzed to provide data on particle size and size range using low angle laser light scattering (Mastersizer2000, Malvern Instruments, United Kingdom). The particle size was directly determined and was estimated as mean of 300 particle measurements (n=300) using optical microscopy.

Fourier transform infrared (FTIR)

FTIR spectra analysis was performed for the microsphere particles. Before analysis, 2 mg samples was mixed with 300 mg KBr powder to for a pellet. The KBr pellet was observe at 4000-450/cm using Jasco FT-IR 5300, Easton MD, USA.

Determination of protein loading capacity of microspheres

Particles microspheres (± 100 mg) was dissolved 30 ml aquadem and stirred at 1000 rpm for 4 hours and was centrifuged at 15,000 rpm for 20 minutes to precipitate particles. BSA concentration was measured using Bradford protein assay, measuring the absorbances by spectrophotometer UV/VIS V-630 (Jasco, jepang) at 586 nm. Each sample was assayed in triplicate (n=3). The encapsulation efficiency (E.E.) and loading capacity (L.C) of microspheres was determined as follows:

$$\text{E.E. (\%)} = \frac{\text{theoretical BSA amount} - \text{BSA in supernatant}}{\text{Theoretical BSA amount}} \times 100$$

$$\text{L.C. (\%)} = \frac{\text{theoretical BSA amount} - \text{BSA in supernatant}}{\text{Microspheres weight}} \times 100$$

Statistical analysis

Statistical analysis was performed by two-way analysis of variance (ANOVA) for repeated measurements. The post hoc (Bonferroni and Scheffe) test was used to perform multiple comparison analysis (SPSS program, USA, version 15) and differences were considered significant at a level of $p < 0.05$.

RESULTS

Table 1. Composition of BSA-loaded CS microspheres and the effect of concentration TPP on physical characteristic BSA-loaded CS microspheres

Formula	Ratio chitosan:TPP			Σ Particle Size ± SD (μm)	Σ Entrapment Efficiency ± SD (%)
	BSA	Chitosan	TPP		
1	0.01	1	2	2.40 ± 0.20	96.25 ± 3.91
2	0.01	1	2.6	2.41 ± 0.31	92.51 ± 0.14

Size and Surface Morphological

The chitosan and TPP composition of the microspheres has no influence on their morphology, were illustrated in figured 1 and figured 2.




Figure 1. Optical photograph of BSA (A), chitosan (B), tripolyphosphate (C), and microspheres with chitosan-tripolyphosphate ratio: 1:2 (F1), 1:2.6 (F2) (magnification x4.000)




Figure 2. Organoleptic of chitosan microspheres with chitosan-tripolyphosphate ratio 1:2 (F1), 1:2.6 (F2)

Fourier Transform Infrared (FT-IR)

FTIR spectra of BSA-loaded CS microspheres are shown in Figure 3.




Figure 3. Fourier transform infrared spectra of BSA (A), chitosan (B), tripolyphosphate (C), and microspheres with chitosan-tripolyphosphat ratio: 1:2 (F1), 1:2.6 (F2)

DISCUSSION

The characteristics of the BSA-loaded CS microspheres prepared with different concentration of chitosan were studied. The optical microscopy resulted BSA-loaded CS microspheres having spherical shape with mean sizes ranging from 2.38-2.41 μm (table 1 and figure 1). Concentration range in this study, as TPP increases microscale on particles are formed. BSA-loaded CS microspheres had smaller size, indicate that using gelation ionic and stirring speed was 1000 rpm in this study resulted the smaller particles size. The ionic gelation method has several advantages such as the use of aqueous solutions, the preparation of small size particles, the control of particle size by the variation of parameters such as chitosan and TPP concentrations, and the possibility of encapsulation of a large range of molecules⁽¹³⁾. Stirring speed of 1000 rpm was found to be optimal, yielding microspheres of uniform size and with a narrow size range and spherical shape⁽¹⁴⁾. The structure of chitosan-microspheres was showed that the particles have a uniform spherical shape and smooth surface.

Data of FTIR spectra BSA-loaded CS microspheres resulted that the absorption band at wavenumber 1640 cm^{-1} was higher, indicating that the amino groups on chitosan and TPP are involved in the formation microspheres. So we suppose that the phosphoric groups in tripolyphosphate are linkage with ammonium group of chitosan; inter- and intramolecular action are enhanced in BSA-loaded CS microspheres.

Table 1 shows the effects of concentration crosslinker on the efficiency encapsulation. To investigated the capability of BSA-loaded CS microspheres to encapsulated BSA, microspheres were prepared with various amounts of TPP. The BSA-loaded CS microspheres was resulted $>90\%$, indicated that the microspheres prepared with chitosan and TPP can trapped the BSA. The mechanical strength, physicochemical properties and particle absorption efficiency and extend drug release time were enhanced by crosslinking⁽¹⁵⁾.

CONCLUSION

Although the drug delivery system based on chitosan/TPP microparticles has its own advantages, the efficiency encapsulation of macromolecules with smaller particles size. It has been found in this study that an increase in the concentration of TPP to promote the formation of a compact and sufficient cross-linking structure of chitosan/TPP particles that can enhanced the efficiency encapsulation. The particles size of BSA-loaded CS microspheres were formed in smaller size indicate the microspheres will be an effective system for delivery the protein.

REFERENCES

1. Mahmud A, Discher ED. Lung vascular targeting through inhalation delivery: insight from filamentous viruses and other shapes. *International Union of Biochemistry and Molecular Biology Life*, 2011; 63: 612–607.
2. Milala AS. Technology of technology dry powder as a protein delivery system in to pulmonary. *Medicinus* 2013; 26: 46-39.
3. Sivadas N, O'Rourke D, Tobin A, Ramtoola Z., Kelly GJ. A Comparative study of a range of polymeric microspheres as potential carriers for the inhalation of proteins. *International Journal of Pharmaceutics*. 2008;358:159-67.
4. Al-Qadi S, Grenha A, Carion-recio D, Seijo B, Remmunan-Lopez C. Microencapsulated chitosan nanoparticles for pulmonary protein delivery: In vivo evaluation of insulin-loaded formulations. *Journal of Controlled Release*. 2012; 157: 390–383
5. Zhang HP, Luo XG, Lin XY, Lu X, Tang Y. the molecular understanding of interfacial interaction of functionalized graphene and chitosan. *Applied Surface Scientific*. 2016;360: 721-715.
6. Amidi, Maryam, Mastrobattista E, Jiskoot W, Wim EH. Chitosan-based delivery systems for protein therapeutics and antigens. *Advanced Drug Delivery Reviews*. 2010;62: 82–59.
7. Mattu C, Li R, Ciardelli G. Chitosan nanoparticles as therapeutic protein nanocarriers: the effect of ph on particle formation and encapsulation efficiency. *Polymer Composites* 2013.
8. Zou X, Zhao X, Ye L, Wang Q, Li H. Preparation and drug release behavior of pH-responsive bovine serum albumin-loaded chitosan microsphere. *Journal Of Industrial and Engineering Chemistry*. 2014:10-1.
9. Agnihotri, SA, Malikarjuna NN, Aminabhavi TM. recent advance on chitosan-based micro-and nanoparticles in drug delivery. *Journal Of Control Release*. 2004;100: 28-5.
10. Hariyadi, DM, Ma, T., Wang, Y, Bostrom, T, Turner MS, Bhandari B, Coombes AGA. The potential for production of freeze-dried oral vaccines using alginate hydrogel microspheres as protein carriers. *Journal of Drug delivery Science. Technology*, 201;24 :184-178.
11. Hejjaji EM, Smith AM, Morris GA. Designing chitosan-tripolyphosphate microparticles with desired size for specific pharmaceutical or forensic applications. *International Journal of Biological Macromolecules*. 2016;95:573-564.

12. Manjanna KM, Kumar TM, Pramod B. Shivakumar. calcium alginate cross-linked polymeric microbeads for oral sustained drug delivery in arthritis. *Drug Discoveries and Therapeutics*. 2010: 122-109.
13. Hassani S. Preparation of chitosan-TPP nanoparticles using microengineered membrans-effect of parameters and encapsulation of tacirone. *Col and Surf A: Physchem and Eng Asp*. 2015;482:43-34.
14. Nagda CD, Chotai NP, Nagda DC, Patel SB, Patel UL. Development and characterization of mucoadhesive microspheres for nasal delivery of ketorolac. *Pharmazie*. 2011;66:257-249.
15. Tsai HZ, Wang YZ. Properties of hydrophilice chitosan network membranes by introducing binary crosslink agent. *Polymer Bulletin* 2008;60:113-103.