

G-17

ISSN (cetak) : 2088-4559

ISSN (online): 2477-0256

Pharmaciiana

Accredited by Ministry of Research, Technology and Higher Education, Republic of Indonesia (Decree No. 32a/E/KPT/2017)

Jurnal Kefarmasian

Vol. 8 No. 2 Nov 2018



Search Scope

All ▼

Search

Browse

- By Issue
- By Author
- By Title
- Other Journals

HOME ABOUT LOGIN REGISTER SEARCH CURRENT ARCHIVES
ANNOUNCEMENTS SCOPE EDITORIAL BOARD AUTHOR GUIDELINE

Home > Vol 8, No 2 (2018)

Pharmaciana

Pharmaciana is a scientific journal published by the University of Ahmad Dahlan worked closely with Ikatan Apoteker Indonesia (IAI). Pharmaciana published two times a year, namely in May and November since 2011 with ISSN 2088-4559 and e-ISSN 2477-0256. The article published in the Journal Pharmaciana selected by editors and reviewed by the reviewer. Articles published in Pharmaciana must not be published in other journals or have been previously published.

Pharmaciana is indexed in google scholar, Garuda Scholar, Indonesian Publication Index (IPI), Sherpa Romeo, and BASE. Pharmaciana is planned to be indexed in Scopus. Since 2017 Pharmaciana is accredited by DIKTI (DGHE) of Indonesia No. [32a/E/KPT/2017](#) April 26, 2017

Announcements

IMPORTANT NOTIFICATION For May Issue 2018

Starting in May 2018 edition, the article will be published **in English**. Therefore, the author should submit articles **in English**. The author may also submit articles in Indonesian, but **will be charged** translations in accordance with terms and conditions.

For more information, don't hesitate to ask us if you have any questions. Please contact at pharmaciana@pharm.uad.ac.id

Posted: 2017-10-28

Pharmaciana Journal Call for Paper

Authors are invited to submit electronically no more than 12 pages full paper through journal website. The submitted paper should follow the format available on the [author guideline](#).

The scopes of accepted papers are: Pharmaceutics, Biopharmaceutics, Drug Delivery System, Physical Pharmacy, Chemical Pharmacy, Pharmaceutical Technology, Pharmaceutical Microbiology and Biotechnology, Pharmacology and Toxicology, Pharmacokinetics, , Pharmaceutical Chemistry, Pharmaceutical Biology, Community and Clinical Pharmacy, Regulatory Affairs and Pharmaceutical Marketing Research, and Alternative Medicines.

Posted: 2016-03-22

[More Announcements...](#)

Vol 8, No 2 (2018): Pharmaciana

Table of Contents

In silico study of the active compounds in bitter melon (<i>Momordica charantia</i> L) as antidiabetic medication	PDF
<i>Ruswanto Ruswanto, Richa Mardianingrum, Tresna Lestari, Tita Nofianti, Lilis Tuslinah, Dudi Nurmalik</i>	194
The potentiation effect of Bawang Dayak (<i>Sisyrinchium</i>	PDF

palmifolium L.) extract on T47D cell growth inhibition after 5-fluorouracil treatment <i>Azizah Yuniarti, Elza Sundhani, Nunuk Aries Nurulita</i>	204
In silico toxicity prediction of 1-phenyl-1-(quinazolin-4-yl) ethanol compounds by using Toxtree, pkCSM and preADMET <i>Yeni Yeni, Supandi Supandi, Fajar Merdekawati</i>	PDF 216
Neuropharmacological Activity of Nut Grass (Cyperus Rotundus L.) Rhizome Fraction <i>Lusi Putri Dwita, Ema Dewanti, Vera Ladeska, Sediarmo Sediarmo, Adnan Muntashir, Ulfa Safni, Rahma Wulan Sari</i>	PDF 224
Inhibition of HMG CoA Reductase and Lipid Peroxidation in The Rats Liver by Selected Zingiberaceae <i>Patonah Hasimun, Agus Sulaeman, Hendra Mahakam Putra, Heni Lindasari</i>	PDF 232
Anti Inflammatory And Anti Bacterial Activity Of Sumbawa Oil <i>Taufan Hari Sugara, Baiq Nurbaety</i>	PDF 240
REDUCED ULCEROGENECITY FROM SELF-NANO EMULSIFYING DRUG DELIVERY SYSTEM OF PIROXICAM <i>Iis Wahyuningsih, Wahyu Widyarningsih, Silviana Wulandari</i>	PDF 248
The validation of patient knowledge questionnaire for patients with diabetes mellitus and hyperlipidemia <i>Ginanjar Zukhruf Saputri, Haafizah Dania, Imaniar Noor Faridah, Faiza Chusnuni A</i>	PDF 256
Effect of the ethanolic extract of red roselle calyx (Hibiscus sabdariffa L.) on hematocrit, platelets, and erythrocytes in healthy volunteers <i>Siti Setianingsih, Laela Hayu Nurani, Abdul Rohman</i>	PDF 266
Evaluation of Adherence, Distress and Quality of Life For Type 2 Diabetes Melitus Patients In Puskesmas Wedarijaksa I and Puskesmas Trangkil Kabupaten Pati <i>Dyah Aryani Perwitasari, Siti Urbayatun, Oni Yulianta Willis</i>	PDF 274
Evaluasi terapi erythropoetin pada pasien hemodialisa di rumah sakit PKU Muhammadiyah Yogyakarta <i>Adnan Adnan, Haafizah Dania, Woro Supadmi</i>	PDF 282
The physical properties and stability of purple yam (Ipomoea batatas (L.) Lam) lipstick <i>susmiatun susmiatun, Anjar mahardian kusuma, Arif Budiman, Indri Hapsari</i>	PDF 290
Improving the in vitro penetration of niacinamide using span 60-based niosomal system in gel formulation <i>Rise Desnita, Sri Luliana, zainab kasim</i>	PDF 296
The Effect of Concentration Variation of Ethanolic Extract from Potato Peels (Solanum tuberosum L.) on the Physical Properties and Antibacterial Activity of Gels against Propionibacterium acnes <i>Dewi Rashati, Mikhania Cristiningtyas Eryani</i>	PDF 302
Formulation and evaluation of erythropoietin-alginate microspheres at different amount of drug <i>Dewi Hariyadi, Esti Hendradi, Khoirotul Lailiyah</i>	PDF 312
Antibacterial activity of dialkyl-arginate biosurfactant cream againsts Staphylococcus aureus an Pseudomonas aeruginosa <i>Kintoko Kintoko, Nining Sugihartini, Benazir Evita Rukaya</i>	PDF 320
Determination of Sun Protection Factor and Antioxidant Properties of Cream Formulation of Kencur (Kaempferia galanga L) and Temu Kunci (Boesenbergia pandurata (Roxb.) Schlecht) Rhizomes Extract <i>Shintia Lintang Charisma, Wiranti Sri Rahayu, Retno Wahyuingrum</i>	PDF 330
Isolation, identification, phytochemical screening, and antibacterial	PDF

activity of *Aspergillus* sp. MFD-01, an endophytic fungus derived from *Mesua ferrea*

Dwi Hartanti, Rida Yunita Andestia Sinaga, Asmiyenti Djaliasrin Djilil, Retno Wahyuningrum

338

Antifertility activities of srikaya leaves (*Annona squamosa* L.) ethanol extract against reproduction system : spermatogenesis and quality sperm study

[PDF](#)

Nur Laili Dwi Hidayati, Tita Nopianti, Yuliana Dewi, Yani Suryani

348

Pharmaciana indexed by:



MENDELEY



This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).

Search Scope
 All ▾
 Search

Browse

- By Issue
- By Author
- By Title
- Other Journals

Home > About the Journal > **Editorial Team**

Editorial Team

Editor in Chief

[Dr. Nurkhasanah Mahfudh](#), Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia

Editorial Board

- [Dr. Kenny Chitcholtan](#), Christchurch School of Medicine and Health Sciences, New Zealand
[Prof. Dr. Irwandi Jaswir](#), International Islamic University Malaysia (IIUM), Malaysia
[Dr. drh Supto Yuliani, MP](#), Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia
[Sukanya Dej-adisai Ph.D.](#), Faculty of Pharmaceutical Sciences, Prince of Songkla University, Hat-Yai, Songkhla 90110 Thailand, Thailand
[Dr. Fezah Binti Othman](#), Universiti Putra Malaysia, Malaysia
[Prof. Bob Wilffert](#), Department Pharmacotherapy and Pharmaceutical Care, University of Groningen, Netherlands
[Zullies Ikawati, Prof., Dr., Apt.](#), Fakultas Farmasi, Universitas Gadjah Mada, Yogyakarta, Indonesia
[Abdul Rohman, Prof., Dr., M.Si., Apt.](#), Fakultas Farmasi, Universitas Gadjah Mada Yogyakarta, Indonesia
[Akhmad Kharis Nugroho, Prof., Dr., M.Si., Apt.](#), Fakultas Farmasi, Universitas Gadjah Mada, Indonesia
[Dadan Hermawan, M.Si., Ph.D.](#), Fakultas MIPA, Universitas Jenderal Soedirman, Indonesia
[Dr. Moch. Saiful Bachri](#), Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia
[Prof. Dr. Achmad Mursyidi](#), Universitas Ahmad Dahlan, Indonesia
[Dr Isnaeni Yudi Haryanto](#), Fakultas Farmasi, Universitas Airlangga, Indonesia
[Keri Lestari Dandan, Dr., M.Si., Apt.](#), Fakultas Farmasi, Universitas Padjadjaran, Indonesia
[Phebe Hendra, M.Si., Apt., Ph.D.](#), Sanata Dharma University, Indonesia
[Dr Laela Hayu Nurani, M.Si., Apt.](#), Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia
[Anita Sukmawati, Ph.D., Apt.](#), Fakultas Farmasi, Universitas Muhammadiyah Surakarta, Indonesia
[Dr Tedjo Yuwono, Apt.](#), Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia
[Dr Nining Sugihartini, M.Si., Apt.](#), Faculty of Pharmacy, Ahmad Dahlan University, Indonesia
[Lannie Hadisoewignyo, Dr., M.Si., Apt.](#), Fakultas Farmasi, Universitas Katolik Widya Mandala Surabaya, Indonesia
[Iin Narwanti, M.Sc.](#), Fakultas Farmasi Universitas Ahmad Dahlan, Indonesia
[Nina Salamah, M.Sc., Apt.](#), Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia
[Faridah Baroroh, M.Sc., Apt.](#), Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia
[Susan Fitria Candradewi, M.Sc., Apt.](#), Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia
[Citra Ariani Edityaningrum, M.Si., Apt.](#), Fakultas Farmasi, Universitas Ahmad Dahlan

Pharmaciana indexed by:



This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).

Formulation and evaluation of erythropoietin-alginate microspheres at different amount of drug

Dewi Melani Hariyadi*, Esti Hendradi, Khoirotul Lailiyah

Department of Pharmaceutics, Faculty of Pharmacy

Universitas Airlangga, Surabaya, Indonesia

Submitted: 08-03-2018

Reviewed: 04-07-2018

Accepted: 02-11-2018

ABSTRACT

This research formulate erythropoietin-alginate microspheres and to evaluate characteristics of erythropoietin-alginate microspheres at different amount of drug using aerosolization. Amount of erythropoietin are 10,000 IU (F1); 20,000 IU (F2); 60,000 IU (F3). The mixture of erythropoietin-alginate was sprayed into CaCl_2 and was stirred at 1000 rpm for 30 minutes. Formulas resulted spherical shape of microspheres. The size of microspheres was 2.77 μm for F1; 3.89 μm for F2; and 4.42 μm for F3. The results of swelling index showed that swelling index of microspheres increased by increasing the concentration of erythropoietin. The results were in accordance with the size of the microspheres that increased with increasing concentration of drug. The yields of microspheres obtained were respectively 91.92%; 87.53%; 86.50% for F1, F2 and F3. It can be concluded that the particle size of microspheres, swelling index increased by increasing concentration of erythropoietin. In contrast, yield of microspheres decreased by increasing drug concentration. In conclusion, formulas of microspheres were potential in terms of characteristics and may recommend for further in vivo study.

Keywords: erythropoietin, alginate microspheres, drug concentration, aerosolization

***Corresponding Author:**

Dewi Melani Hariyadi

Faculty of Pharmacy, Universitas Airlangga, Surabaya

Jl. Dharmawangsa Dalam Surabaya

E-mail address: dewi-m-h@ff.unair.ac.id

INTRODUCTION

Erythropoietin is a 30.4 kDa glycoprotein hormone which is the main regulator of red blood cell production (Prenggono, 2015). Erythropoietin has a protective role against tissue ischaemia (Pachos, 2008). Erythropoietin, which is a protein, has several weaknesses, such as unstable in biological fluids because of easily digested by enzymes, cannot be fully absorbed from the digestive tract due to a high molecular weight resulting to low bioavailability, and has short half-life (Wang *et al.*, 2013). Erythropoietin is stable at pH 4-9 and has the highest activity at pH 7-8 (Chang *et al.*, 2013). Erythropoietin is also stable at -20°C or -70°C (Chang *et al.*, 2013). At pH 9 and 40 °C, erythropoietin will undergo dimerization to form large molecules that can induce the body's immune response, which can harm patients (Chang *et al.*, 2013). Some alternative is by formulatemicrospheressystem to release the drug in a sustained release manner (Chaudhary *et al.*, 2010).

Microspheres are a drug delivery system in which the drug is diluted or dispersed homogeneously in the polymer matrix (Kumar *et al.*, 2011). The advantage of a microsphere delivery system is that the drug's release properties in the body occur gradually, making it suitable for carrying the necessary drugs into the body with a fixed amount. In addition, the microspheres can also protect peptide/protein drugs from hydrolytic enzyme attacks (Takada, 2008).

The manufacture of erythropoietin microspheres by ionic gelation method and aerosolization technique requires polymers and crosslinking solutions. One of the most commonly used natural polymers is alginate.

Alginate is a polysaccharide which is a salt of alginic acid, comprising a monomer (1-4)- β -D-manuronic acid (unit M) and α -L-guluronic acid (unit G) that varies in quantity and distribution along its polymer chain (Pal *et al.*, 2013). Na alginate is widely applied for biomedical purposes as a drug delivery system due to its biocompatibility, biodegradability, low toxicity, relatively low cost, and lighter gelation by the addition of divalent cations such as Ca^{2+} (Hariyadi *et al.*, 2014).

The frequently used crosslinking cations are divalent and trivalent cations (Ahirrao *et al.*, 2014). The addition of monovalent metal ions to the alginate will form a soluble salt (Lee and Money, 2012). Pb, Cu, Cd cations are rarely used because they have some toxicity (Zhai, 2012). The use of Ba and Sr cations should be in low concentration as they are slightly toxic (Zhai, 2012). Ca^{2+} cations are often used because of their low toxicity and lack of high affinity for alginates (Zhai, 2012).

The ionic gelation method is based on the ability of the polyelectrolyte to be crosslinked in the presence of crosslinking ions forming a hydrogel. This method is prepared by dripping the drug-polymer solution into a cation polyvalent solution. The cation diffuses into the droplets of a polymer solution, forming a three-dimensional crosslinked formation (Patil *et al.*, 2012). The ionic gelation method is suitable for the active ingredients of proteins because of the simple and light process, which is greatly helpful for maintaining the bioactivity of the proteins (Koppolu *et al.*, 2014). The advantage of the ionic gelation method is that all polyelectrolytes are diluted in the water so that the protein can be encapsulated without the use of organic solvents and the ionic gelation method does not use high temperatures so that protein integrity is maintained. In addition, this method is considered simple, fast, cost-effective, and has a small particle size diameter (Hariyadi *et al.*, 2014).

Microspheres that have been formed will undergo the drying process. The drying process of microspheres in this research was the freeze drying technique, because it is able to maintain protein integrity during the drying process (Abdelwahed *et al.*, 2006). The freeze drying process requires pressure during the freezing and drying processes, requiring a lyoprotectant (Abdelwahed *et al.*, 2006). A lyoprotectant can stabilize the microsphere through the mechanism of hydrogen bond formation with polar groups on the surface of the microsphere at the end of the drying process in order to maintain the microsphere structure and replace the water's position on the microsphere surface (Abdelwahed *et al.*, 2006). One of the usable lyoprotectants is maltodextrin, because it protects protein stability during the freezing and drying (Hariyadi *et al.*, 2014).

Factors that may affect the characteristics of the resulting microsphere are the number of polymers, the number of crosslinkers, the amount of drug ingredients used, and the crosslinking time

(Jin *et al.*, 2009). The higher the concentration of the drug, the higher the microspheres size that is obtained (Wang *et al.*, 2013). In a study conducted by Balasubramaniam *et al.* (2007) on the effect of drug concentration on microsphere size using the emulsification technique, it was found that the greater the concentration of drug used, the greater the microsphere size that is obtained. This was because the higher the concentration of the drug, the greater the droplet size produced, resulting in the larger size of the formed microsphere (Balasubramaniam *et al.*, 2007). The microspheres size range generated from the study was 47-50 μm (Balasubramaniam *et al.*, 2007). In addition, increasing the concentrations of medicinal ingredients may also increase the efficiency of encapsulation (Balasubramaniam *et al.*, 2007). The differences of the research conducted by Balasubramaniam *et al.* (2007) with this research is on aerosolization technique used in method of produced microspheres, therefore it is expected that the microsphere particle size formed is $<5 \mu\text{m}$. The determination of target size of microspheres of $<5 \mu\text{m}$ is adjusted to the particle size requirement for suspensions (Patel, 2010).

This study was conducted to determine the effect of erythropoietin amount (10,000, 20,000, 60,000 IU) on erythropoietin-alginate microspheres characteristics using ionotropic gelation method and aerosolization technique.

MATERIALS AND METHODS

Materials

Pharmaceutical grade Erythropoietin (Daewoong Inc.), Sodium alginate (Sigma-Aldrich Inc.), $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (Solvay Chemicals International), demineralized water, pro-analysis Na_2HPO_4 (Merck), KH_2PO_4 (Merck), NaCl (Merck), NaOH (Merck), and pharmaceutical grade Maltodextrin (PT Bratachem).

Research Method

Microspheres formula

Sodium alginate (2 gram) were dissolved into 100 mL of demineralized water. Erythropoietin was dispersed into 100 mL of alginate solution based on drug concentration. After that, CaCl_2 solution 1M was made in 200 ml aqua demineralized by stirring homogeneously. The formed erythropoietin-alginate solution was then sprayed into CaCl_2 solution using aerosol spray with a 35 μm hole size and a constant pressure of 40 psi while being continuously stirred using a magnetic stirrer at 1000 rpm for 30 minutes. The microspheres suspension that had been formed was centrifuged at 4000 rpm for 10 minutes, then washed with demineralized water 3 times to separate the microspheres from the CaCl_2 solution. The washed microspheres were collected and dried using the freeze drying technique at -26°C for 38 hours.

Table I. Erythropoietin-alginate microspheres formulas

Materials	Function	F1	F2	F3
Erythropoietin	Active agent	10.000 IU	20.000 IU	60.000 IU
Na Alginate	Polymer	2% (in 100ml)	2% (in 100ml)	2% (in 100ml)
CaCl_2	Crosslinker	1 M (in 200 ml) 5%	1 M (in 200 ml) 5%	1 M (in 200ml) 5%
Maltodextrin	Lyoprotectant	(from wet microspheres weight)	(from wet microspheres weight)	(from wet microspheres weight)

Evaluation of erythropoietin-alginate microspheres

Particle size distribution

The evaluation of erythropoietin-alginate microsphere size distribution was performed using 400x optical microscope magnification. The diameter of erythropoietin-alginate microspheres was observed with a 400x magnification of 300 particles. Then, the mean diameter was determined and the microsphere size distribution curve was made. The average diameter was calculated using the formula:

$$d_{vs} = \frac{nd^3}{nd^2}$$

Microspheres shape and surface morphology

The morphological evaluation of erythropoietin-alginate microspheres and surface forms was performed using a Scanning Electron Microscope (SEM). This evaluation was performed by placing a sample on the handle of the preparation with an adhesive material containing metal grains, such as the metal Pt. The gold in the Chamber was evaporated so that the gold steam coated the entire surface of the microparticles. The surface of the gold-coated microparticles was observed using SEM to observe the morphology of the shape and surface of the microspheres.

Moisture content determination

The moisture content determination on erythropoietin alginate microspheres can be determined using the moisture balance. This evaluation can be performed by weighing 0.5 grams of microspheres and inserting them into the moisture balance. The tool would work for 10 minutes. The moisture content (MC) was calculated by the equation: $MC = \frac{\text{Initial weight} - \text{Final weight}}{\text{Final weight}} \times 100\%$

Determination of Swelling Index (SI)

The determination of swelling index was performed by weighing 50 mg microspheres and then adding 5 ml PBS pH 7.4 in a vial. The swelling index determination was performed at 24 and 30 hours. After the observation time had been reached, the wet microspheres were filtered using filter paper. After no PBS was dripped from the filter paper, the wet microspheres were transferred to a dry filter pad and then the microspheres were flipped continuously to remove the adsorbed water on the microsphere surface until the filter paper became not too wet. Afterwards, the wet microspheres were stirred at 37°C for 2 hours or until the microspheres weight became constant. The microspheres was weighed as the swelling weight. The swelling index (SI) value was calculated using the formula:

$$SI = \frac{\text{Swelling weight} - \text{Dry weight}}{\text{Dry weight}}$$

The swelling index observation was also performed based on the microsphere size. A little amount of microsphere were taken to be observed for microspheres size during swelling using an optical microscope. The swelling index (SI) value was calculated using the formula:

$$\frac{\text{Swelling microsphere size} - \text{Dry microsphere size}}{\text{Dry microsphere size}}$$

Yield determination

The yield value was determined by the ratio of the total weight of the dry microspheres obtained to the amount of weight of erythropoietin, sodium alginate, and maltodextrin. The yield value can be calculated by the formula:

$$\text{Yield}(\%) = \frac{\text{total microspheres weight (mg)}}{\text{alginate + erythropoietin + maltodextrin weight (mg)}} \times 100\%$$

The data from each examination were compared with each of the formulas. The data analysis was performed using SPSS 23 statistical program using the One Way ANOVA method.

RESULTS AND DISCUSSION

The mean diameter for blank microspheres was 2.31 μm and the mean diameters of F1 microspheres, F2, F3 using 400 times magnification were $2.77 \pm 0.08 \mu\text{m}$; $3.89 \pm 0.12 \mu\text{m}$; and $4.42 \pm 0.06 \mu\text{m}$ respectively (Figure 1). These results indicated that blank microspheres were smaller than the three formulas microspheres with erythropoietin. The observation also showed an increase in the size of the microsphere as the levels concentration of erythropoietin from F1 to F3 increased with the values of 10,000, 20,000, and 60,000 IU. The average microspheres diameter result met the specification of $<5 \mu\text{m}$, where the size is considered good for a parenteral suspension (Patel, 2010).




Figure 1. Histogram of diameters of microspheres sizes of blank microspheres and microspheres of F1, F2 and F3

The microspheres size increased by addition of erythropoietin levels was due to two factors. First, the higher the amount of erythropoietin, the more erythropoietin can filled the remaining empty egg-box structures. Second, there was a repulsive force between erythropoietin and alginate. Alginate as a bioelectrolyte can caused repulsive force with erythropoietin. The repulsive force occurred between the COO^- group of alginate with the COO^- group of erythropoietin. The larger the erythropoietin, the greater the repulsive force that occurs with alginate. The existence of such repulsive forces may increase the viscosity, resulting in increased microsphere size (Chaudhari *et al.*, 2015).

In the analysis of the effect of erythropoietin levels on microspheres, sig values of $0.000 < 0.05$ for F1:F2; $0.000 < 0.05$ for F1:F3; and $0.001 < 0.05$ for F2:F3 were obtained. The three comparisons of the formula showed sig values of < 0.05 , so we can say there is a significant difference in microsphere size between F1, F2, and F3.

The morphological observations of the shape and surface of the microspheres using the Scanning Electron Microscope (SEM) of the F1, F2, F3 resulted in spherical and smooth shapes (Figure 2). These results can be obtained due to the addition of maltodextrin as a lyoprotectant which made the form of the microsphere to be spherical with a smooth and flat surface. The maltodextrin would filled the cavity on the surface of the microsphere and form the hydrogen bond with the polar group on the microsphere surface during the freeze drying process (Abdelwahed *et al.*, 2006).




Figure 2. The shape and surface morphology of the erythropoietin microspheres (A) F1, (B) F2, (C) F3 observed using the Scanning Electron Microscopy at 5000x magnification

The result of MC determination showed that the average of water content in the three formulas were $5.85 \pm 0.10\%$; $5.08 \pm 0.13\%$; and $8.08 \pm 0.21\%$ as seen in Table I. The MC of three formulas corresponded to specifications of MC of $<10\%$. The high moisture content can decrease the stability of the active ingredients in the microspheres due to the formation of particle agglomeration (Shan *et al.*, 2016).

Table II. Moisture content of microspheres formula

Microspheres Formula	MC (%) \pm SD	CV (%)
F1	5.85 ± 0.10	1.71
F2	5.08 ± 0.13	2.56
F3	8.08 ± 0.21	2.60

From the determination of swelling index, the average swelling index at 24 hours based on the weights of microspheres in formula F1, F2 and F3 were 1.300 ± 0.071 ; 1.463 ± 0.046 ; 1.749 ± 0.044 , while during the observation at 30 hours, the average swelling index based on the weight of the microsphere in the formula F1, F2, and F3 were 1.867 ± 0.056 ; 2.272 ± 0.154 ; 2.475 ± 0.189 . The average swelling indices at 24 hours based on microspheres sizes in the formula F1, F2, and F3 were 1.13 ± 0.10 ; 1.37 ± 0.05 ; 1.67 ± 0.04 , respectively. Meanwhile, during the observation at 30 hours, the

average swelling indices based on the microspheres size in formulas F1, F2, and F3 were 1.85 ± 0.04 ; 2.07 ± 0.12 ; 2.28 ± 0.15 .




Figure 3. The swelling index histogram based on weight of microspheres F1, F2, F3




Figure 4. The swelling index histogram based on the size of microspheres of F1, F2, F3

The average swelling indices based on microsphere weight at 24 hours and 30 hours yielded gave greater result than swelling indices based on microspheres sizes. However, if it was tested statistically using the independent t-test, the difference of swelling indices between microsphere weight and microsphere size was not significant due to the 2-tailed sig value > 0.05 . The increase of swelling index values at 24 to 30 hours based on the weight and size of microspheres can be said to differ significantly because the independent t-test results showed that the 2-tailed sig value < 0.05 .

When we looked at the average swelling index results, both based on microsphere weight and microsphere size, it seems that both yielded the same resulted that the swelling index value increased with the increased of the ingredients added in the formula. This corresponded to the average yield of microspheres in which microsphere sizes increased with the ingredients added in the formula. The swelling index results based on microsphere weight and microspheres size were also in accordance

with the theory, where the higher the observation time, the higher the swelling index value was obtained (Navneet *et al.*, 2011).

The swelling index evaluation was used to ensure the polymer can expand, then it can released drug. Alginate, in releasing drug substances, had diffusion, swelling, and erosion mechanisms (Fernanda *et al.*, 2014). The results of the evaluation have been found to be in accordance with the theory in which alginate can expand, as evidenced by the addition of weight and size of microspheres after being put aside for 24 and 30 hours.

From the result of yield determination, the average yields obtained in formula F1, F2, and F3 were $91.92 \pm 3.33\%$; $87.53 \pm 6.01\%$; $86.50 \pm 4.59\%$. From these results, it is known that the value of the yield fallen along with the increase of the ingredient content of the drug added in the formula. This was because microspheres have entrapment capacities in entrapping drug ingredients (Rastogi *et al.*, 2007). The increasing yield values indicated that erythropoietin with a level of $>10,000$ IU is not entirely entrapped in the microspheres and was dissolved when washed with water since erythropoietin was also soluble in water. The results of statistical analysis showed that the effect of erythropoietin content on yield was determined by sig values of $0.534 > 0.05$ for F1: F2; $0.403 > 0.05$ for F2: F3; and $0.963 > 0.05$ for F2: F3. The three comparisons of the formula showed sig values of < 0.05 , so we can say there is a significant difference in microspheres size between F1, F2, and F3.




Figure 5. Histogram of % yield of microspheres of F1, F2, F3

CONCLUSION

The formulation of erythropoietin-alginate microspheres by the ionic gelation method and aerosolization technique with different erythropoietin amount (10,000, 20,000, 60,000 IU) resulted in microspheres which were spherical and smooth. Increasing levels of erythropoietin increased the size of the microspheres and swelling index significantly, but did not have a significant effect on yield.

ACKNOWLEDGMENTS

The author would like to thank the Faculty of Pharmacy, Universitas Airlangga for all supports in this research.

REFERENCES

- Abdelwahed, W., Degobert, G., Stainmesse, S., and Fessi, H., 2006. Freeze-drying of nanoparticles: Formulation, process and storage considerations, *Advanced Drug Delivery Reviews*, 58: 1693-1696.

- Ahirrao, S. P., Gide, P. S., Shrivastav, B., and Sharma, P., 2014. Iontropic Gelation : A promising cross linking technique for hydrogels, *Journal of Pharmaceutic and Nanotechnology*, 2(1): 1–6.
- Balasubramaniam, Vinay U. Rao, M. Vasudha, Jeevan Babu and P.S. Rajinikanth., 2007. Sodium alginate microspheres of metformin HCl: formulation and in vitro evaluation. *Current Drug Delivery*, 4(3): 249-256.
- Koppolu, B.P., Sean G. Smith, Sruthi Ravindranathan, Srinivas Jayanthi, Thallapuram K.S. Kumar, David A. Zaharoff, 2014. Controlling chitosan-based encapsulation for protein and vaccine delivery, *Biomaterials*, 35(14): 4382–4389.
- Chang, S.H. , Kim, H.J., and Kim, C. W., 2013. Analysis of the structure and stability of erythropoietin by pH and temperature changes using various LC/MS. *Bull, Korean Chemical Society*, 34(9): 2663-2670.
- Chaudhari, S., R. Kar, J. and S. Singhal, R., 2015. Immobilization of proteins in alginate: functional properties and applications, *Current Organic Chemistry*, 19(17): 1732-1754.
- Chaudhary, K., Haddadin, S., Nistala, R. and Papageorgio, C., 2010. Intraperitoneal Drug Therapy: An Advantage, *Current Clinical Pharmacology*, 5(2): 82-88.
- Fernanda M. Carbinatto, Ana Dórisde, Castro Raul C., Evangelista Beatriz S.F.Cury, 2014. Insights into the swelling process and drug release mechanisms from cross-linked pectin/high amylose starch matrices, *Asian Journal of Pharmaceutical Sciences*, 9(1) : 27-34.
- Hariyadi D.M., HendradiEsti, Purwanti Tutiek, Fadil F.D.G.P., Ramdani, C.N., 2014. Effect of crosslinking agent and polymer on the characteristic of ovalbumin loaded alginate microspheres, *International Journal Of Pharmacy And Pharmaceutical Sciences*, 6(4): 1-6.
- Inanlou, F., Pejvak, K., Naser, M., Soheila, M.B., Hossein, Z., Chang, Y.F., Bazrafshan, M.R., and Saraei, F., 2015. Preparation of microspheres containing leptospiral antigen using biodegradable alginate polymers, *Veterinarski Arhiv* 85(4): 395-406.
- Jin, M., Zheng, Y., Hu, Q., 2009. Preparation and Characterization of Bovine Serum Albumin Alginate/Chitosan Microspheres for Oral Administration, *Asian Journal of Pharmaceutical Sciences*: 215-221.
- Kumar, B.T., Chandiran, I.S., Bhavya, B, and Sindhuri, M., 2011. Microparticulate Drug Delivery System : a Review, *Indian Journal of Pharmaceutical Science and Research*, 1: 19-31
- Pal, K., Allan T.Paulson, Dérick Rousseau, 2013. 14 - Biopolymers in Controlled-Release Delivery Systems in *Handbook of Biopolymers and Biodegradable Plastics*, 329-363.
- Lee, K. and Mooney, D., 2012. Alginate: Properties and biomedical applications, *Progress in Polymer Science*, 37(1): 106-126.
- Navneet, G., Akanksha, G. and Neetesh, J., 2011. Formulation design and in-vitro evaluation of metformin microspheres using ionotropic gelation technique, *Journal of Pharmacy Research*, 4(7): 2103-2106.
- Paschos, N., Lykissas, M.G., Beris, A.E., 2008. The Role of Erythropoietin as an Inhibitor of Tissue Ischemia, *International Journal of Biological Sciences* :161-168.
- Patel, R., 2010, Parenteral Suspension: An Overview, *International Journal of Current Pharmaceutical Research*, 2(3): 1-10.
- Patil, P., Chavanke, D., Wagh, M., 2012. A Review on Iontropic Gelation Method: Novel Approach for Controlled Gastroretentive Gelspheres, *International Journal of Pharmacy and Pharmaceutical Sciences*, 4(4): 27-32.
- Prenggono, M.D., 2015. Eritropoetindan penggunaan eritropoetin pada pasien kanker dengan anemia, *CDK-224*, 42(1): 20-26.
- Rastogi, R., Y. Sultana, M., Aqil, S. Kumar, K., Chuttani, A.K. Mishar, 2007. Alginate microspheres of isoniazid for oral sustained drug delivery, *International Journal of Pharmaceutics*: 334 (1-2): 71-77.
- Shan, L., Tao, E. X., Meng, Q. H., Hou, W. X., Liu, K., Shang, H. C., Tang, J. B., and Zhang, W. F., 2016. Formulation, optimization, and pharmacodynamic evaluation of

- chitosan/phospholipid/ β -cyclodextrin microspheres, *Drug, Design, Development and Therapy*, 10 : 417-429.
- Takada, K., 2008. Microfabrication-derived DDS: From batch to individual production, *Drug Discoveries and Therapeutics*, 2(3) :140-155.
- Yang, J., Xie, Y. and He, W. 2011. Research progress on chemical modification of alginate: A review. *Carbohydrate Polymers*, 84(1): 33-39.
- Ying Wang, Xiaoping Yang, Wentao Liu, Feng Zhang, Qing Cai, Xuliang Deng, 2013. Controlled release behaviour of protein-loaded microparticles prepared via coaxial or emulsion electrospray, *J Microencapsul*, 30(5): 490–497.
- Wang, L., Liu, Y., Zhang, W., Chen, X., Yang, T. and Ma, G., 2013. Microspheres and Microcapsules for Protein Delivery: Strategies of Drug Activity Retention, *Current Pharmaceutical Design*, 19(35): 6340-6352.
- Zhai, P., 2012. Alginate microspheres for protein delivery in tissue engineering, *A Thesis Division of Biomedical Engineering University of Saskatchewan*: 1-72.