

DAFTAR PUSTAKA

- Abikusno, N. (2007). Older population in Indonesia: trends, issues and policy responses. Bangkok: YNFPA, p:23.
- American Thoracic Society. (2002). ATS Statement: Guidelines for the Six-Minute Walk Test. *American Journal of Respiratory Critical Care Medicine*, 166(111–117). DOI: 10.1164/rccm.166/1/111
- Askandar, T., Poernomo, BS., Chairul, E., Djoko, S., Gatot, S. (2015). *Buku Ajar Ilmu Penyakit Dalam Fakultas Kedokteran Universitas Airlangga Rumah Sakit Dr Soetomo*, 2nd edn, Airlangga University Press, Surabaya.
- Basmajian. (1990). *Therapeutic Exercise*, 5th edn, Lippincot William and Wilkins, Baltimore.
- Bernstein, ML., Despars, JA., Singh, NP., Avalos, K., Stansbury, DW., Light RW. (1994). Reanalysis of the 12 minute walk in patients with chronic obstructive pulmonary disease. *Chest*, 105:163–7.
- Bodine, SC., Stitt, TN., Gonzalez, M., Kline, WO., Stover, GL., Bauerlein, R., Zlotchenko, E., Scrimgeour, A., Lawrence, JC., Glass, DJ., Yancopoulos, GD. (2001). Akt/mTOR pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy in vivo. *Nat Cell Biol*, 3:1014–19.
- Bohannon, RW., Bubela, D., Magasi, S., McCreath, H., Wang, YC., Reuben, D., Gershon, R. (2014). Comparison of walking performance over the first 2 minutes and the full 6 minutes of the Six-Minute Walk Test. *BMC Research Notes*, 7, 269. <http://doi.org/10.1186/1756-0500-7-269>
- British Geriatric Society. (2004). Fit For Frailty.
- Brooks, Dina. (2002). Reliability of the two-minute walk test in individuals with transtibial amputation. *Archives of Physical Medicine and Rehabilitation*, 83(11), 1562–1565
- Brooks, D., Parsons, J., Hunter, JP., Devlin, M., Walker, J. (2001). The 2-minute walk test as a measure of functional improvement in persons with lower limb amputation. *Arch Phys Med Rehabil*, 82:1478–83
- Chung, HY., Cesari, M., Anton, S., Marzetti, E., Giovannini, S., Seo, AY., Carter, C., Yu, B.P., Leeuwenburgh, C. (2009) Molecular inflammation: Underpinnings of aging and age-related diseases. *Ageing Res*, 8, 18-30.
- Clegg, A., Young, J., Iliffe, S., Rikkert, MO., Rockwood, K. (2013). Frailty in elderly people. *Lancet*, 381(868):752-762.
- Connelly, D. M., Thomas, B. K., Cliffe, S. J., Perry, W. M., & Smith, R. E. (2009). Clinical Utility of the 2-Minute Walk Test for Older Adults Living in Long-Term Care. *Physiotherapy Canada*, 61(2), 78–87. <http://doi.org/10.3138/physio.61.2.78>
- Davies, GA., Bolton. (2010). Age-Related Changes in the Respiratory Sistem. In Fillit, H.M., Rockwood,K, Woodhouse,K eds. Brokleshurst's textbook geriatri. 7th ed., Saunders Elsevier, Philadelphia, pp. 101-162.
- Eiser N, Willsher D, Dore CJ. (2003). Reliability, repeatability and sensitivity to change of externally and self-paced walking tests in COPD patients. *Respir Med*, 97(4):407–414.

- Enright, PL. (2009). Aging of The respiratory System. In WR Hazzard (eds), *Hazzard's Geriatric Medicine and Gerontology*, 6th edn, The McGraw-Hill Companies, Inc, USA, pp. 751-66
- Fielding, RA., Vellas, B., Evans, WJ., Bhasin, S., Morley, JE., Newman, AB., Abellan, van Kan G., Andrieu, S., Bauer, J., Breuille, D., Cederholm, T., Chandler, J., De Meynard, C. (2011). Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. International Working Group on Sarcopenia. *J Am Med Dir Assoc*, 12: 249-56.
- Finch, E., Brooks, D., Stratford, PW., Mayo, NE. (2002). Physical rehabilitation outcome measures: a guide to enhanced clinical decision making. 2nd ed. Hamilton, ON: BC Decker.
- Fried, LP. (2001). Frailty in older adults, evidence for a phenotype. *Gerontology Med Sci*; pp 146.
- Gomes, MJ., Martinez, PF., Pagan, LU., Damatto, RL., Mariano Cezar, MD., Ruiz Lima, AR., Okoshi, MP. (2017). Skeletal muscle aging: influence of oxidative stress and physical exercise. *Oncotarget*, 8(12), 20428–20440. <http://doi.org/10.18632/oncotarget.14670>
- Goroll, AH., Mulley, AG. (2009). *Primary Care Medicine, Office Evaluation and Management of the Adult Pain*, 6th Ed. Philadelphia. Lippincott William, pp. 239
- Gremiaux, V., Iskandar, M., Kervio, G., Deley, G., Pérénou, D., Casillas, JM. (2008). Comparative analysis of oxygen uptake in elderly subjects performing two walk tests: the six-minute walk test and the 200-m fast walk test. *Clin Rehabil*, 22(2):162-8.
- Guccione, AA., Wong, RA., Avers, D. (2012). *Geriatric Physical Therapy*, 3rd edn, Elsevier Mosby, Missouri.
- Guyatt, GH., Sullivan, MJ., Thompson, PJ., Fallen, EL., Pugsley, SO., Taylor, DW. (1985). The 6-minute walk: a new measure of exercise capacity in patients with chronic heart failure. *Can Med Assoc J*, 132(8):919-23
- Hamid, A. (2007). Penduduk Lanjut Usia di Indonesia dan Masalah Kesejahteraannya, viewed 5 January 2017, <http://www.kemsos.go.id/>
- Harahap, H., Widodo, Y., Mulyati Sri., 2005. Penggunaan berbagai cut-off indeks massa tubuh sebagai indikator obesitas terkait penyakit degeneratif di Indonesia. *Gizi Indon*, 31, pp. 1-12.
- Hiengkaew, V., Jitaree, K., Chaiyawat, P. (2012). Minimal detectable changes of the Berg balance scale, Fugl-Meyer Assessment Scale, Timed “Up & Go” Test, gait speeds, and 2-minute walk test in individuals with chronic stroke with different degrees of ankle plantarflexor tone. *Archive of Physical Medicine Rehabilitation*, 93:1201–1208.
- Johnston, K. N., Potter, A. J., & Phillips, A. C. (2017). Minimal important difference and responsiveness of 2-minute walk test performance in people with COPD undergoing pulmonary rehabilitation. *International Journal of Chronic Obstructive Pulmonary Disease*, 12, 2849–2857. <http://doi.org/10.2147/COPD.S143179>
- Katch, VL., McArdle, WD., Katch, F. I. (2011). Measuring and Evaluating Human Energy-Expenditure Capacities During Exercise. In: *Essential of*

- Exercise Physiology, 5th edition.* Philadelphia: Lippincott William & Wilkins, pp. 203-233.
- Karnik, K., Mazzatti, DJ. (2009). Review of Tools and Technologies to Assess Multi-System Functional Impairment and Frailty. Sharnbrook, U.K. *Clinical Medicine: Geriatrics*, 3.
- Kirkwood, TBL. (2010). *Evolution Theory and the Mechanisms of Aging*. In: Fillit, H.M, Rockwood, K, Woodhouse, K eds. Brokleshurst's textbook geriatri., Saunders Elsevier, Philadelphia, pp. 18-25.
- Leung AS, Chan KK, Sykes K, Chan KS. (2006). Reliability, validity, and responsiveness of a 2-min walk test to assess exercise capacity of COPD patients. *Chest*, 130(1):119–125.
- Lewis, CB., Bottomley, JM. (2008). Geriatric Rehabilitation: A Clinical Approach, 3rd ed. New Jersey: Pearson Pentice Hall, pp: 210-2.
- Masoro, EJ. (2010). Physiology of Aging. In : Fillit,H.M., Rockwood,K, & Woodhouse,K. Eds. *Brokleshurst's Textbook of Geriatric Medicine and Gerontology*. 7th edition. Philadelphia, Saunders Elsevier. pp.51
- Manton, KG., Land, KC. (2000). Multidimensional Disability/Mortality Trajectories at Ages 65 and Over: The Impact of State Dependence. *Social Indicators Research*, 51: 193. <https://doi.org/10.1023/A:1006966525964>
- Meng, SJ., Yu, LJ. (2010). Oxidative Stress, Molecular Inflammation and Sarcopenia. *International Journal of Molecular Sciences*, 11(4), 1509–1526. <http://doi.org/10.3390/ijms11041509>
- Motl, RW., Suh, Y., Balantrapu, S. (2012). Evidence for the different physiological significance of the 6- and 2-minute walk tests in multiple sclerosis. *BMC Neurology*, 12:6.
- Nusdwinuringtyas N, Laksmi W, Bachtiar A, 2011. Healthy adults maximum oxygen uptake prediction from a six minute walking test. *Med J Indones* 20(3): 195-9
- Noonan, V., Dean, E. (2000). Submaximal Exercise Testing. *Physical Therapy*, 80:782-807.
- Papathanasiou, JV., Ilieva, E., Marinov, B. (2013). Six-minute walk test: an effective and necessary tool in modern cardiac rehabilitation. *Hellenic J Cardiol*, 54(2):126-30.
- Pin, TW., Lewis, J. (2012). Pilot normative study on 2- and 6-minute walk tests on typically developing children in Australia. Australasian Academy of Cerebral Palsy and Developmental Medicine Conference. *Developmental Medicine Children Neurology*, 54:85.
- Poerwadi, T. (2015). Dignity and Happiness for the Golden Age. Zifatama Publisher. Sidoarjo, pp:51-81.
- Prentice, W., 2011. Maintaining Aerobic Capacity and Endurance during Rehabilitation. In W. Prentice, *Rehabilitation Techniques for Sports Medicine and Athletic Training*. New York: McGraw-Hill, pp. 214-25.
- Reid, MB., Li, YP. (2001). Tumor necrosis factor- α and muscle wasting: A cellular perspective. *Respir*, 2, 269-272.
- Reid, L., Thomson, P., Besemann, M., Dudek, N. (2015). Going Places: Does The Two-Minute WALK Test Predict The Six-Minute WALK Test In Lower Extremity Amputees? *J Rehabil Med*, 47: 256–261

- Rockwood., Song, X., Bergman, H. (2005.) A global clinical measure of fitness and frailty in elderly people. *CMAJ*, p: 489.
- Roman, D., Mahoney, K., Mohamadi, A. (2013) Sarcopenia: what's in a name? *J Am Med Dir Assoc*, 14: 80-2.
- Satyawati, R., Pratiwi, N. (2017). 15 Feet walking test as a Measure of functional Mobility In Frailty Elderly
- Setiati, S., Alwi, I., Sudoyo, AW., Simadibrata, M., Setiyohadi, B., Syam, AF. (2014). Proses Menua dan Implikasi Kliniknya, in Setiati S, Harimurti K, dan Govinda A (eds.), *Ilmu Penyakit Dalam*, 6th edn, Interna Publishing, Jakarta, pp. 3669-79.
- Setiati, S., Alwi, I., Sudoyo, AW., Simadibrata, M., Setiyohadi, B., Syam, AF. (2014). Kerapuhan dan Sindrom Gagal Pulih, in Setiati S, Rizka A, (eds.), *Ilmu Penyakit Dalam*, 6th edn, Interna Publishing, Jakarta, pp. 3725-29.
- Setiati, S., Alwi, I., Sudoyo, AW., Simadibrata, M., Setiyohadi, B., Syam. AF. (2014). Sarkopenia, in Setiati S dan Dwimartutie N (eds.), *Ilmu Penyakit Dalam*, 6th edn, Interna Publishing, Jakarta, pp. 3717-29.
- Strandberg, TE., Pitkala, KH., Tilvis, RS. (2011). Frailty in Older People, *European Geriatric Medicine Journal*, 2:344-55.
- Solway, S., Brooks, D., Lacasse, Y., Thomas, S. (2001). A qualitative systematic overview of the measurement properties of functional walk tests used in the cardiorespiratory domain. *Chest*, 119:256–270.
- Syddall, H. (2010). Prevalence and correlates of frailty among community-dwelling older men and women. *BMC Geriatrics*.
- Sykes, K., Roberts, A. (2004). The Chester Step Test - a Simple yet Effective Tool for The Prediction of Aerobic Capacity. *Physiotherapy*, 90, pp. 183-188.
- Tamis, WP., Choi, HL. (2017): Reliability, validity, and norms of the 2- min walk test in children with and without neuromuscular disorders aged 6–12, *Disability and Rehabilitation*. <http://dx.doi.org/10.1080/09638288.2017.1294208>
- Thompson, WR., Gordon, NF., Pescatello, (2010). *ACSM's Guidelines for Exercise Testing and Prescription*, 8 edn, Wolters Kluwer, Philadelphia.
- Timiras. PS. (2007). *Physiological Basis of Aging and Geriatrics*, 4th edn, Informa Healthcare, New York.
- Tumbelaka, A. R., Riono, P., Sastroasmoro, S., Wirjodiarjo, M., Pudjiastuti, P., Firman, K., 2014. Pemilihan Uji Hipotesis. In: S. Sastroasmoro & S. Ismael, eds. Dasar-dasar Metodologi Penelitian Klinis. Jakarta : CV. Sagung Seto, pp. 328-350.
- Upton CJ, Tyrrell JC, Hiller EJ. Two minute walking distance in cystic fibrosis. *Arch Dis Child* 1988; 63:1444–1448
- Xue, Q. (2011). The Frailty Syndrome: Definition and Natural History. *Cinical Geriatric Medicine*. Elsivier, pp: 6.
- Vermeulen. (2011). Predicting ADL disability in community-dwelling elderly people using physical frailty indicators: a systematic review BMC Geriatrics.