

p-ISSN 2085-191X (print)  
e-ISSN 2338-7610 (online)

# Biosaintifika

Journal of Biology & Biology Education



Biosaintifika

Volume 7

Number 2

Pages 79-147

Semarang  
September 2015

p-ISSN 2085-191X (print)  
e-ISSN 2338-7610 (online)

A  
G

SALINAN

LAMPIRAN  
KEPUTUSAN MENTERI RISET, TEKNOLOGI,  
DAN PENDIDIKAN TINGGI REPUBLIK  
INDONESIA  
NOMOR 2/E/KPT/2015  
TENTANG  
HASIL AKREDITASI TERBITAN BERKALA  
ILMIAH PERIODE II TAHUN 2015

<b>Bidang Ilmu</b>	<b>No</b>	<b>Nama Jurnal</b>	<b>ISSN</b>	<b>Penerbit</b>	<b>Peringkat</b>
Agama	1	Mutawatir	2088-7523	Jurusan Al-Qur'an dan Hadis Fakultas Ushuluddin dan Filsafat UIN Sunan Ampel Surabaya	Terakreditasi B
	2	ADDIN	0854-0594	Pusat Penelitian dan Pengabdian Pada Masyarakat (P3M) STAIN Kudus	Terakreditasi B
	3	Teologia: Jurnal Ilmu- ilmu Ushuluddin	0853-3857	Fakultas Ushuluddin UIN Walisongo Semarang	Terakreditasi B
	4	Jurnal Review Politik	2088-6241	Konsorium Dosen Ilmu Politik dan Politik Islam bekerjasama dengan Prodi Filsafat Politik Islam UIN Sunan Ampel Surabaya	Terakreditasi B
	5	Al-Daulah	2089-0109	Prodi Siyasa Jinayah Fakultas Syariah dan Hukum UIN Sunan Ampel Surabaya	Terakreditasi B
	6	Musawa, Jurnal Studi Gender dan Islam	1412-3460	PSW UIN Sunan Kalijaga	Terakreditasi B
Ekonomi	1	Jurnal Ekonomi dan Kebijakan (JEJAK)	1979-715X	Jurusan Ekonomi Pembangunan, Fakultas Ekonomi Universitas Negeri Semarang	Terakreditasi B

Bidang Ilmu	No	Nama Jurnal	ISSN	Penerbit	Peringkat
	2	Asia Pacific Management and Business Application (APMBA)	2252-8997	Jurusan Manajemen Fakultas Ekonomi dan Bisnis Universitas Brawijaya	Terakreditasi B
	3	Jurnal Bisnis dan Manajemen	1412-3681	Departemen Manajemen & Bisnis, Fakultas Ekonomi Universitas Padjdjaran	Terakreditasi B
Kesehatan	1	Global Medical and Health Communication	2301-9123	Pusat Penerbitan Universitas - Lembaga Penelitian dan Pengabdian Kepada Masyarakat (P2P-LPPM) Fakultas Kedokteran Universitas Islam Bandung	Terakreditasi B
	2	Jurnal Respiriologi Indonesia (JRI)	0853-7704	Perhimpunan Dokter Paru Indonesia (PDPI)	Terakreditasi A
	3	Folia Medica Indonesia	2355-8393	Graha Masyarakat Ilmiah Kedokteran (Gramik) Fakultas Kedokteran Universitas Airlangga	Terakreditasi B
MIPA	1	Biosaintifika	2085-191X	Jurusan Biologi, Fakultas Matematika dan Ilmu Pengetahuan Alam, Universitas Negeri Semarang bekerjasama dengan Perhimpunan Biologi Indonesia dan Konsorsium Biologi	Terakreditasi B
	2	Jurnal Ilmiah Kimia Molekul	1907-9761	Jurusan Kimia FMIPA Unsoed	Terakreditasi B
Pendidikan	1	Jurnal Pendidikan IPA Indonesia	2339-1286	Program Studi Pendidikan IPA, Fakultas Matematika dan Ilmu Pengetahuan Alam, Universitas Negeri Semarang bekerjasama dengan Perkumpulan Pendidikan IPA Indonesia (JPPII)	Terakreditasi B

# Biosaintifika

Journal of Biology & Biology Education

[HOME](#) [ABOUT](#) [LOGIN](#) [REGISTER](#) [SEARCH](#) [CURRENT](#) [ARCHIVES](#) [CONFERENCE PUBLICATION COLLABORATION](#)

Home > Vol 11, No 2 (2019)

## Biosaintifika: Journal of Biology & Biology Education

Nationally Accredited by Ministry of Research, Technology and Higher Education, Number 2/E/KPT/2015

Biosaintifika: Journal of Biology & Biology Education [p-ISSN 2085-191X| e-ISSN 2338-7610] published scientific papers on the results of biology and biology education research (see Focus and Scope). Editor accepts the article has not been published in other media with the writing format as listed on page manuscript writing guidelines (see Author Guidelines). The journal has been indexed in Google Scholar, Microsoft Academic, DOAJ, Dimension, DOI Crossref, EBSCO, CABI, SINTA 2 and published three times a year, on April, August & December and published by Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Semarang Perhimpunan Biologi Indonesia in collaboration with Perhimpunan Biologi Indonesia Tingkat Pusat.

Home > About the Journal > [Editorial Team](#)

### Editorial Team

#### Editor-in-Chief

1. Dr. Ir. Dyah Rini Indriyanti, [SCOPUS ID: 57194044439] Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Semarang, Indonesia

#### International Editorial Board

1. Prof. Dr. Fahrul Zaman Huyop, [SCOPUS ID: 10240240100] Department of Biotechnology and Medical Engineering, Universiti Teknologi Malaysia, Skudai, Malaysia
2. Dr. Mulyoto Pangestu, (Scopus ID= 35488897500) Education Program in Reproductive Biology Dept. Obstetric and Gynaecology, Monash University, Australia
3. Assoc. Prof. Dr. Kok Boon Neoh, (Scopus ID= 34873230600) Laboratory of Innovative Humano-habitability, Kyoto University, Japan
4. Dr. Rosarin Rujananon, [SCOPUS ID: 37122824900] Technical University of Hamburg, Germany
5. Dr. Tanuji Kudre, [SCOPUS ID: 55243377400] CSIR-CFTRI & NCL, Pune, India
6. Prof. Donald Stewart, (Scopus ID= 7403643344) Griffith Graduate Research School, Griffith University, Brisbane, Australia
7. Dr. Mehradj Ahmad, (Scopus ID= 35730106500) Mahidol University, Thailand
8. Dr. Mukh Syaifudin, (Scopus ID= 6507831675) Radiation Biology, Pusat Teknologi Keselamatan dan Metrologi Radiasi (PTKMR), Badan Tenaga Nuklir Nasional (BATAN), Indonesia
9. Prof. Dr. Ir. Yohanes Budi Widanarko, (Scopus ID= 6507055401) Department of Food Technology, Soegijapranata Catholic University, Semarang, Indonesia
10. Dr. Amir Hamidy, (Scopus ID= 24341129500) Research Center For Biology - Cibinong Science Center (CSC), Bogor, Indonesia
11. Prof. Dr. Ir. S. Sudarsono, (Scopus ID= 16450098400) Department of Agronomy and Horticulture, Institut Pertanian Bogor, Indonesia
12. Dr. Iin Kurnia, [SCOPUS ID 57193537585] Biomedika PTKMR BATAN Jakarta, Indonesia
13. Mohammad Irham M.Sc, (Scopus ID= 55312221800) Museum Zoologicum Bogoriense, Research Center for Biology-Indonesian Institute of Sciences, Bogor, Indonesia, Indonesia

#### Associate Editors

1. Prof. Dr. Enni Suwarsi Rahayu, [SCOPUS ID: 56644635000] Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Semarang, Indonesia
2. Talitha Widiatningrum, Ph.D, [SCOPUS ID: 56798544200] Department of Biology, Faculty of Mathematics and Sciences, Semarang State University, Indonesia
3. Prof. Dr. Ir. Amin Retnoningsih, [SCOPUS ID: 51864647000] Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Semarang, Indonesia
4. Dr. Margareta Rahayuningsih, (SCOPUS ID: 57094066700) Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Semarang, Indonesia
5. Dr. Niken Subekti, [SCOPUS ID: 57195315756] Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Semarang, Indonesia



#### ABOUT THE JOURNAL

[Aims and Scope](#)

[Publication Ethics](#)

[Indexing & Abstracting](#)

[Editorial Team](#)



#### ABOUT THE JOURNAL

[Aims and Scope](#)

[Publication Ethics](#)

[Indexing & Abstracting](#)

[Editorial Team](#)

[Reviewer Team](#)

[Contact](#)

#### COLLABORATE WITH



perhimpunan biologi indonesia

#### FOR AUTHORS

[Guidelines for Author](#)

[Peer Review Process](#)

[Author Fees](#)

[Online Submission](#)

[Google Citation Analysis](#)

Vol 8, No 1 (2016)



































March 2016

DOI: <https://doi.org/10.15294/biosaintifika.v8i1>

Available online since 9th March 2016

Table of Contents

Articles

<p><b>Knowledge Understanding and Utilization of Medicinal Plants by Local Community Tompu District of Kaili, Sigi Biromaru, Central Sulawesi</b>  <i>Slamet Ifandi, Jumari Jumari, Sri Widodo Agung Suedy</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.4529">10.15294/biosaintifika.v8i1.4529</a>  Views of Abstract: 766   PDF: 584</p>	PDF 1-11
<p><b>Influence of Polysaccharide Krestin from <i>Coriolus versicolor</i> Extract on Nitrite and Malondialdehyde Concentrations of <i>Mus musculus</i> Serum Exposed by <i>Mycobacterium tuberculosis</i></b>  <i>Sri Puji Astuti Wahyuningsih, Manikya Pramudya, Sugiharto Sugiharto</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.4969">10.15294/biosaintifika.v8i1.4969</a>  Views of Abstract: 508   PDF: 384</p>	PDF 12-17
<p><b>Effect of Turmeric Powder Supplementation to The Age of Sexual Maturity, Physical, and Chemical Quality of The First Japanese Quails (<i>Coturnix japonica</i>) Egg</b>  <i>Tyas Rini Saraswati, Silvana Tana</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.4982">10.15294/biosaintifika.v8i1.4982</a>  Views of Abstract: 806   PDF: 482</p>	PDF 18-24
<p><b>Bird Communities In Seblat Nature Recreation Park (SNRP) North Bengkulu, Bengkulu</b>  <i>Eki Susanto, Yeni Aryati Mulyani, Bambang Suryobroto</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.4948">10.15294/biosaintifika.v8i1.4948</a>  Views of Abstract: 412   PDF: 371</p>	PDF 25-32
<p><b>Plant Species Composition and Their Conspecific Association in Natural Tropical Rainforest, South Papua</b>  <i>Agustinus Murdjoko, Djoko Marsono, Ronggo Sadono, Suwarno Hadisusanto</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5217">10.15294/biosaintifika.v8i1.5217</a>  Views of Abstract: 546   PDF: 504</p>	PDF 33-46
<p><b>Characteristics of superior soybean breeding lines tolerance to rust (<i>Phakopsora pachyrhizi</i> Syd.)</b>  <i>Alfi Inayati, Eriyanto Yusnawan</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5081">10.15294/biosaintifika.v8i1.5081</a>  Views of Abstract: 390   PDF: 315</p>	PDF 47-55
<p><b>Analysis on the Absolute Growth Rate of <i>Rhizophora mucronata</i> Seedling in Silvicultural Pond Canals by the Influence of Initial Condition and Changes of Environment Quality</b>  <i>Endah Dwi Hastuti, Rini Budihastuti</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5358">10.15294/biosaintifika.v8i1.5358</a>  Views of Abstract: 441   PDF: 328</p>	PDF 56-63
<p><b>Ethnobotanical Study of Rattans on Suku Anak Dalam Community in Bukit Duabelas Nasional Park</b>  <i>Dwi Mairida, Muhadiono Muhadiono, Iwan Hilwan</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5164">10.15294/biosaintifika.v8i1.5164</a>  Views of Abstract: 595   PDF: 455</p>	PDF 64-72
<p><b>Growth Response of <i>Sorghum bicolor</i> (L.) Moench. Cultivars to Trivalent Chromium Stress</b>  <i>Sri Kasmiyati, Santosa Santosa, Irfan Dwidja Priyambada, Kumala Dewi, Sucahyo Sucahyo, Rintawati Sandradewi</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5178">10.15294/biosaintifika.v8i1.5178</a>  Views of Abstract: 627   PDF: 379</p>	PDF 73-86
<p><b>Optimization of In Vitro Conservation Protocol of <i>Carica pubescens</i> Lenne &amp; K. Koch through Medium Concentration, Temperature and Irradiation Duration Decrease</b>  <i>Enni Suwarsi Rahayu, Noor Aini Habibah</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5371">10.15294/biosaintifika.v8i1.5371</a>  Views of Abstract: 315   PDF: 298</p>	PDF 87-95
<p><b>Exploration of Potential Actinomycetes from CIFOR Forest Origin as Antimicrobial, Antifungus, and Producing Extracellular Xylanase</b>  <i>Sipriyadi Sipriyadi, Yulin Lestari, Aris Tri Wahyudi, Anja Meryandini, Maggy Thenawidjaja Suhartono</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5052">10.15294/biosaintifika.v8i1.5052</a>  Views of Abstract: 564   PDF: 441</p>	PDF 96-104
<p><b>Optimization of In Vitro Conservation Protocol of <i>Carica pubescens</i> Lenne &amp; K. Koch through Medium Concentration, Temperature and Irradiation Duration Decrease</b>  <i>Enni Suwarsi Rahayu, Noor Aini Habibah</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5371">10.15294/biosaintifika.v8i1.5371</a>  Views of Abstract: 315   PDF: 298</p>	PDF 87-95
<p><b>Exploration of Potential Actinomycetes from CIFOR Forest Origin as Antimicrobial, Antifungus, and Producing Extracellular Xylanase</b>  <i>Sipriyadi Sipriyadi, Yulin Lestari, Aris Tri Wahyudi, Anja Meryandini, Maggy Thenawidjaja Suhartono</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5052">10.15294/biosaintifika.v8i1.5052</a>  Views of Abstract: 564   PDF: 441</p>	PDF 96-104
<p><b>An Analysis of Partial DNA Sequence of Meisa1 Gene on Sweet and Bitter Cassavas (<i>Manihot esculenta</i> Crantz.)</b>  <i>Dewi Indriyani Roslmi, Fitriyatun Nisa, Herman Herman</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5365">10.15294/biosaintifika.v8i1.5365</a>  Views of Abstract: 353   PDF: 275</p>	PDF 105-112
<p><b>Mass Cultivation of Entomopathogenic Nematode in Artificial Media</b>  <i>Dyah Rini Indriyanti, Nur Lailatul Muharromah</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5579">10.15294/biosaintifika.v8i1.5579</a>  Views of Abstract: 437   PDF: 387</p>	PDF 113-120
<p><b>Characterization of Carambola (<i>Averrhoa carambola</i> L.) Plant Collection of Cibirong Plant Germplasm Garden Based on Phenotypic and Genetic Characters</b>  <i>Dody Priadi, Ambar Yuswi Perdani, Yuli Sulistyowati, Fiqolbi Nuro Pohan, Enung Sri Mulyaningsih</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.5199">10.15294/biosaintifika.v8i1.5199</a>  Views of Abstract: 529   PDF: 327</p>	PDF 121-128
<p><b>Antioxidant Activity from Various Tomato Processing</b>  <i>Retno Sri Iswari, R. Susanti</i>   <a href="https://doi.org/10.15294/biosaintifika.v8i1.4722">10.15294/biosaintifika.v8i1.4722</a>  Views of Abstract: 622   PDF: 473</p>	PDF 129-134



**ABOUT THE JOURNAL**

Aims and Scope
Publication Ethics
Indexing & Abstracting
Editorial Team
Reviewer Team
Contact



**FOR AUTHORS**

Guidelines for Author
Peer Review Process
Author Fees
Online Submission
Google Citation Analysis
Scopus Citation Analysis



**Readers**

ID 111,522	TR 304
US 12,462	AU 283
IN 1,329	NL 241
MY 1,126	IR 233
PH 581	DE 184
CN 519	PK 173
TH 489	FR 172
BR 419	CA 171
SG 390	TM 170
JP 390	MX 167
GB 373	RU 167

Pageviews: 415,993  
Flags Collected: 157

**FLAG counter**  
00968877 View My Stats

**USER**

Username

Password

Remember me

**JOURNAL CONTENT**

Search

Search Scope

**USER**

Username

Password

Remember me

**JOURNAL CONTENT**

Search

Search Scope

- Browse**
- >> By Issue
  - >> By Author
  - >> By Title
  - >> Other Journals



Biosaintifika 8 (1) (2016)

# Biosaintifika

Journal of Biology & Biology Education

<http://journal.unnes.ac.id/nju/index.php/biosaintifika>



## Influence of Polysaccharide Krestin from *Coriolus versicolor* Extract on Nitrite and Malondialdehyde Concentrations of *Mus musculus* Serum Exposed by *Mycobacterium tuberculosis*

✉ Sri Puji Astuti Wahyuningsih, Manikya Pramudya, Sugiharto

DOI: 10.15294/biosaintifika.v7i2.3955

Departement of Biology, Faculty of Science and Technology, Universitas Airlangga, Indonesia.

### History Article

Received 7 January 2016  
Approved 23 February 2016  
Published 29 March 2016

### Keywords:

Malondialdehyde; nitrite; polysaccharide krestin; tuberculosis

### Abstract

*Mycobacterium tuberculosis* is a major infection agent of tuberculosis that is controlled by the response of cell-mediated immunity. It is macrophages and cytolytic T lymphocytes. Activated macrophages will produce free radicals. Excessive free radicals cause tissue damage. Polysaccharide krestin contains  $\beta$ -glucan. It is a scavenger of free radicals. This research aimed to identify the influence of polysaccharide krestin from *C. versicolor* on nitrite and malondialdehyde concentrations of mice serum exposed by *M. tuberculosis*. Nitrite concentration was determined by nitrite assay. Malondialdehyde concentration was determined by TBARS assay. The result showed that adding polysaccharide krestin before exposure (P1) and adding polysaccharide krestin before-after exposure (P3) had the best potential to decrease nitrite concentration. Nitrite concentrations of P1 and P3 were  $1.364 \pm 0.523$  M and  $1.456 \pm 0.712$  M respectively. Meanwhile, P1 group and adding polysaccharide krestin after exposure (P2) had the best potential to decrease malondialdehyde concentration. Malondialdehyde concentrations of P1 and P2 were  $1125.86 \pm 97.96$   $\mu$ M and  $953.86 \pm 328.16$   $\mu$ M respectively. Their nitrite and malondialdehyde concentrations decreased, compared to K and K- groups. The research conclusion was that adding polysaccharide krestin before exposure could decrease both nitrite and malondialdehyde concentrations.

### How to Cite

Wahyuningsih, S., Pramudya, M., & Sugiharto, S. (2016). Influence of Polysaccharide Krestin from *Coriolus versicolor* Extract on Nitrite and Malondialdehyde Concentrations of *Mus musculus* Serum Exposed by *Mycobacterium tuberculosis*. *Biosaintifika: Journal of Biology & Biology Education*, 8(1).

© 2016 Semarang State University

✉ Correspondence Author:  
Mulyorejo C Campus Surabaya, 60115  
E-mail: [sri-p-a-w@fst.unair.ac.id](mailto:sri-p-a-w@fst.unair.ac.id)

p-ISSN 2085-191X  
e-ISSN 2338-7610

## INTRODUCTION

Tuberculosis is infectious disease caused by *Mycobacterium tuberculosis*. In 2014, tuberculosis killed 1.5 million people worldwide. Tuberculosis ranks alongside HIV as leading cause of death worldwide (WHO, 2015). Individual who is infected by infectious disease, virulent, and pathogen will give immune responses, as both specific immune response and nonspecific immune response. Macrophage is one of the nonspecific immune system components. Macrophage will engulf and kill *Mycobacterium tuberculosis* by phagocytosis. During phagocytosis, reactive oxygen intermediates and nitric oxide in the form of peroxynitrite radicals has able to kill pathogen (Abbas et al., 2012). Reactive oxygen intermediates are needed to kill pathogen, however excessive reactive oxygen intermediates will cause DNA damage and induce lipid peroxidation. Excessive nitric oxide will increase amount of reactive nitrogen species. It is produced from reaction between nitric oxide and superoxide radical. Reactive nitrogen species are strong pro-inflammation molecule. Reactive nitrogen species will cause tissues damage. High concentration of nitric oxide kill pathogens, immune cells, and other cells in the vicinity (Zhang et al., 2011). Free radical is difficult to measure directly, but it can be measured effectively using lipid peroxidation concentration that measured by malondialdehyde concentration (Dalle-Donne et al., 2006). Nitric oxide can be measured by nitrite concentration (Jose et al., 2003). In normal condition, antioxidant system will neutralize free radical. Low endogenous antioxidant will form more free radicals. Body needs exogenous antioxidant. *Coriolus versicolor* is medicinal mushroom used in Japan, China, Korea and other Asian countries. Polysaccharide krestin is extraction product of *Coriolus versicolor*. Polysaccharide krestin imitates activity of superoxide dismutase. Superoxide dismutase will change superoxide into hydrogen peroxide (Ooi and Liu, 2000). Powdered polysaccharide krestin contain 34 – 35 soluble carbohydrate (91 – 93%  $\beta$ -glucan), 28 – 35% protein, 7% moisture, 6 – 7% ash, and the rest are free sugars and amino acids (Cui and Chisti, 2003).  $\beta$ -Glucan can be used as antioxidant and scavenger for free radicals (Chen and Seviour, 2007). The objective of this research was to know the influence of polysaccharide krestin from *Coriolus versicolor* extract on nitrite and malondialdehyde concentrations of mice serum exposed by *Mycobacterium tuberculosis*.

## METHODS

### Research site

The research was conducted in the Molecular Genetics Laboratory, Departement of Biology, Faculty of Science and Technology, Universitas Airlangga, Jalan Mulyorejo Kampus C, Surabaya, East Java, Indonesia.

### Material

Polysaccharide krestin was obtained from *Coriolus versicolor* extract method, adapted from Wahyuningsih (2014), *Mycobacterium tuberculosis* (0.25 Mc. Farland) was got from Balai Besar Laboratorium Kesehatan, Surabaya, East Java, Indonesia. Malondialdehyde assay kits were purchased from Northwest Life Science (Washington DC). Nitrite assay used Griess reagent. 10mL Griess reagent I was made from 0.3% sulfanilamide in 2.5% phosphoric acid. 10mL Griess reagent II was made from 0.3% naphthylendiamine in 2.5% phosphoric acid.

### Animal and experimental design

This research used 30 adult female mice (*Mus musculus*) strain Balb/C, age between 8 – 10 weeks, weight between 30 – 40 g. Mice were obtained from Faculty of Pharmacy, Airlangga University (Surabaya, Indonesia). Mice went to acclimatization for a week. Completely randomized design was used with six groups and five repetition, namely control (K), positive control (K+), negative control (K-). It is treated by polysaccharide krestin before being exposed by *M. tuberculosis* (P1), by polysaccharide krestin after being exposed by *M. tuberculosis* (P2) and and by polysaccharide krestin before-after being exposed by *M. tuberculosis* (P3). Mice were exposed to *M. tuberculosis* ( $10^6$  cell/mL) twice through intraperitoneally with two weeks gap from the first to the second exposure. 50 mg/kg BW of polysaccharide krestin was added by gavage in seven days.

### Serum isolation

One week after polysaccharide krestin adding, 1mL intra-cardiac blood was taken using 23G injection needle and collected to 1.5mL micro tube. Blood was incubated for 2 hours. Blood was centrifuged at 3000 rpm for 10 min. Supernatant layer was taken for nitrite and malondialdehyde assays.

### Nitrite assay

After creating the dilution (0 M, 0.2 M, 0.4 M, 0.6 M, 0.8 M, 1.0 M) from nitrite standard (0.69 mg NaNO<sub>2</sub> in 100mL aquadest), standard

nitrite regression equation  $y = ax + b$  was made.  $y$  is optical density value, and  $b$  is nitrite concentration. 50  $\mu\text{L}$  serum was prepared in 1.5mL micro tube. 100  $\mu\text{L}$  Griess reagent I and 100  $\mu\text{L}$  Griess Reagent II were added to serum. After 10 – 15 min incubation, optical density values were read at 540 nm. Nitrite concentrations (M) were known after putting optical density values in standard nitrite regression equation.

#### Thiobarbituric Acid Reactive Substance (TBARS) assay

For serum preparation, 200  $\mu\text{L}$  TCA 10% was added to 100  $\mu\text{L}$  of each serum. After 5 min incubation on ice, serum was centrifuged at 14.000 rpm for 5 min. 200  $\mu\text{L}$  of each serum was transferred to 1.5mL micro tubes. To each of the serum, 200  $\mu\text{L}$  TBA was added. Serum was homogenized and incubated at 100°C for 60 min. After cooling down, serum was homogenized and centrifuged briefly. Optical density values were read at 540 nm. Malondialdehyde concentrations ( $\mu\text{M}$ ) were known after put optical density values in standard malondialdehyde regression equation  $Y = 0.0007X - 0.0003$ ;  $y$  is optical density value and  $b$  is malondialdehyde concentration.

#### Data Analysis


All values were analyzed statistically by one way analysis of variances (ANOVA) to determine whether there were differences in the average of variable among treatments. If there were significant differences, Duncan test ( $\alpha = 0.05$ ) was taken to determine the best treatment.

## RESULT AND DISCUSSION

### Nitrite Concentration

Free radical is difficult to be directly measured. Free radical is measured by other substance indirectly like nitrite ( $\text{NO}_2^-$ ). Nitrite is product of nitric oxide,  $\text{O}_2$  and  $\text{H}_2\text{O}$  reaction (Gharavi and El-Kadi, 2003). According to the result we obtained (Figure 1), the highest nitrite concentration was shown by treatment K- (2.938 M) and the lowest nitrite concentration was shown by P1 group (1.364 M). ANOVA Statistical analysis showed that there was significant difference in the average of variable among treatments ( $0.036 < 0.05$ ). According to Duncan test resulted in Figure 1 (different characters showed significant difference), the best treatment was identified. No significant difference was observed in groups of K, K+ and P2. Groups of P1 and P3 showed significant differences compared to K- group. Figure 1 showed that the present of polysaccharide kres-

tin before exposure (P1) and before-after exposure (P3) decreased nitrite concentration significantly.



**Figure 1.** Nitrite Concentration among Research Groups. Note: different letters in each section show significant difference.

Reactive nitrogen intermediates are free radicals that able to kill pathogen. In macrophage, reactive nitrogen intermediates are formed by inducible nitric oxide synthase (iNOS). iNOS catalyzes arginine conversion to citrullin and produced nitric oxide. In phagolysosome, nitric oxide, hydrogen peroxide and superoxide form peroxynitrite radicals to kill pathogen (Abbas et al., 2012). In cells/tissues, nitric oxide is not stable and has very short half life (Zhang et al., 2011). Nitric oxide can make reaction with  $\text{O}_2$  and  $\text{H}_2\text{O}$  to form stable metabolites including nitrite and nitrate. Generally, Nitric oxide is neutralized by endogenous antioxidant but exogenous antioxidant sometimes is needed to utilize body system against excessive free radicals. Polysaccharide krestin are exogenous that potential antioxidant in term of combating the free radicals. Chen and Seviour (2007) stated that polysaccharide krestin containing  $\beta$ -glucan could be used as free radicals scavenger.

In this research, adding polysaccharide krestin showed significantly different nitrite concentration among treatment groups. K- group that was exposed by *Mycobacterium tuberculosis* without adding polysaccharide krestin showed highest nitrite concentration. Excessive free radicals cannot be neutralized when free radicals were increased and amount of endogen antioxidant was constant. It is important to remember that nitric oxide made by iNOS is beneficial to the host defense reaction by contributing to microbial killing. In this condition, nitric oxide has cyto-protective characteristic. When large amounts of nitric oxide formed by iNOS surpass the physiological amounts of nitric oxide, nitric oxide has cytotoxic characteristic. Nitric oxide action in immune regulation depends on con-



centrations, cellular environment and whether cells have undergone prior activation (Guzik and Guzik, 2003).


According to Figure 1, group of K+ added by polysaccharide krestin without *M. tuberculosis* exposure had similar average nitrite concentration with K group although the concentration was lower than K- group. The presence of nitrite concentration of K group showed that body still produced free radical although it did not exposed by pathogen. As stated by Caramori and Papi (2004), oxidant generation was part of the normal metabolism of many types of cells and is critical for cell homeostasis.

Significant decline of nitrite concentration was showed by P1 and P3 groups. In P1 group, adding polysaccharide krestin before exposure could stimulate enzyme formation related to suppression of free radical. Cui and Chisti (2007) stated that polysaccharide of *C. versicolor* might give benefit in general healthy by inducing enzymes that mop up free radicals and mitigate oxidative damage. Adding polysaccharide krestin before exposure can prevent free radical formation. This evidence was supported by Pham-Huy et al. (2008) that classify antioxidant based on their mode on action including preventive antioxidant. Modes of preventive antioxidant actions are non-radical decomposition of LOOH and H<sub>2</sub>O<sub>2</sub>, sequestration of metal by chelating and quenching of active O<sub>2</sub>. Polysaccharide krestin induces antioxidant formation that breaks free radical chain including superoxide dismutase, catalase and glutathione peroxidase. As stated by Ooi and liu (2000), polysaccharide krestin has SOD-mimicking activity. Polysaccharide krestin induces glutathione peroxidase in mouse peritoneal macrophage (Pang et al., 2000). Enzyme that related to nitric oxide is superoxide dismutase that changed into superoxide radical from reduction of univalent oxygen molecule into H<sub>2</sub>O<sub>2</sub>. It was changed to H<sub>2</sub>O and O<sub>2</sub> by catalase. Reactive nitrogen species would not be formed because nitric oxide did not react with superoxide. There was **no structural change** of polysaccharide krestin which added before exposure because  $\beta$ -glucan of polysaccharide krestin was resistant from enzymatic proteolysis of gastric. In P3 group, polysaccharide krestin which was added before exposure induced antioxidant forming so that antioxidant was formed and could bind with free radical.

According to Figure 2, P2 group that added polysaccharide krestin showed no significant difference. In this treatment, polysaccharide krestin did not actively work because polysaccharide krestin that was used had variation in molecular

weight and branches. As stated by Chan et al. (2009), most of the  $\beta$ -glucan based on extract had variation in molecular weight and branches.  $\beta$ -Glucan with different molecular weight and branches possibly have variation in immune potency. Chan et al. (2009) also stated that variation in molecular size, branching frequency and solution conformation had an impact on the elimination half life, volume of distribution and clearance.

### Malondialdehyde Concentration



**Figure 2.** Malondialdehyde Concentration among Research Groups. Note: different letters in each section show significant difference.

Reactive oxygen species or nitrogen oxygen species are generally too reactive and have short half life to allow direct measurement in cells/tissues or body fluid. Molecular products that were formed from the reaction of reactive oxygen species or reactive nitrogen species with bio-molecules are more stable than reactive oxygen species or reactive nitrogen species themselves. Commonly, reactive oxygen species or reactive nitrogen species have been tracked by measuring stable metabolites and/or concentrations of their oxidation target products including malondialdehyde, lipid peroxidation left product (Dalle-Donne et al., 2006). According to the result in Figure 2, the highest malondialdehyde concentration was shown by K- group (1665.86  $\mu$ M) and the lowest malondialdehyde concentration was shown by P2 group (953.86  $\mu$ M). ANOVA statistical analysis showed that there was significant difference in the average of variable among treatments (0.028 < 0.05). Analysis was continued to Duncan test to determine the best treatment. According to Duncan test result in Figure 2 (different characters

showed significant difference), no significant difference was shown by K+ and P3 groups. Group of K showed significant difference compared to P2 group. Significant difference was also shown by P1 and P2 groups compared to K- group. Figure 2 showed that the presence of polysaccharide krestin before exposure (P1) and after exposure (P2) decreased malondialdehyde concentration significantly.

Reactive oxygen intermediates are formed by activated macrophage. Reactive oxygen intermediates such as superoxide radical, hydroxyl radical, and single oxygen are needed to kill pathogen along with nitric oxide (Abbas et al., 2012) but excessive hydroxyl radical can make reaction with poly unsaturated fatty acid to form malondialdehyde. In this research, adding polysaccharide krestin showed difference on nitrite concentration among groups. According to Figure 2, K- group that was exposed by *M. tuberculosis* without adding polysaccharide krestin showed higher malondialdehyde concentration than K group. Excessive free radical without sufficient antioxidant increased malondialdehyde production. This evidence is supported by Dagli et al. (2003), who stated that excess malondialdehyde, as a by-product of free radicals can be related to decreased blood and tissue antioxidant levels.

Group of K+, added by polysaccharide krestin without *Mycobacterium tuberculosis* exposure showed similar average of malondialdehyde concentration with K group, although the concentration was lower than K- group. Under normal circumstances, the major sources of oxidants and free radicals produced in the body occur by mitochondrial activity, electron transport chains, phagocyte cells and the endogenous enzyme system (Dagli et al., 2004).

Significant decrease of malondialdehyde concentrations were observed in P1 and P2 groups. In P1 group, that was added polysaccharide krestin before *M. tuberculosis* exposure, could stimulate enzyme formation to bind with free radical. Polysaccharide krestin induced glutathione forming.

In P2 group that was added polysaccharide krestin after *M. tuberculosis* exposure, stimulated forming of macrophage as phagocytes cell, therefore, the immune system was raised. This evidence was supported by Cui and Chisti (2003) who stated that main effects of using polysaccharide krestin are inducing production of interleukin-6, interferon, immunoglobulin-G, macrophages and T-lymphocytes. The increasing of activated macrophages helped raising a better condition of mice, therefore, production of endogenous

antioxidant increased indirectly. Polysaccharide krestin could bind with receptors on immune cell membrane, particularly macrophage because it had  $\beta$ -glucan. Multicellular organisms possess receptors, called pattern recognition receptors (PRRS), to detect innately non-self structures (including pathogen-associated molecular patterns or PAMPs) (Brown and Gordon, 2005). Chen and Seviour (2007) reported that  $\beta$ -glucan probably act as PAMPs and are recognized by appropriate cell surface receptors, initiating immune responses. Specific receptor to recognize  $\beta$ -glucan is Dectin-1. Malondialdehyde forming could be prevented by glutathione. Based on Pham-Huy et al. (2008), glutathione is classified in preventive antioxidant. Polysaccharide krestin stimulated glutathione forming. Polysaccharide krestin induced glutathione peroxides activity in mouse peritoneal macrophages (Pang et al., 2000). Ozkan et al. (2010) reported that significant increased of glutathione peroxides and decreased of malondialdehyde were observed in acetylsalicylic acid and  $\beta$ -glucan treatment.  $\beta$ -Glucan appeared to attenuate the gastric damage. Glutathione peroxidase in erythrocytes oxidized will reduce glutathione into glutathione sulphide. Glutathione sulphide protected lipid membrane and haemoglobin from oxidation of peroxide. Glutathione is formed by glutamic acid, serine and glycine (Eide, 2011). Polysaccharide krestin contain large amounts of aspartic acid and glutamic acid (Ooi and Liu, 2000). Acidic and neutral amino acids such as aspartic acid, threonine, serine, glutamic acid, glycine, alanine, valine, and leucine (Cui and Chisti, 2003). Polysaccharide krestin could induce glutathione forming because it contained the same amino acid as glutathione.

Group of P3 showed lower malondialdehyde concentration than K- group but did not show significant difference. In P3 group, polysaccharide krestin was added twice, before and after *M. tuberculosis* exposure. Therefore immunostimulator capability of polysaccharide krestin increased. This evidence was supported by research conducted by Nurdini (2015) that reported lowest interleukin-10 concentration found in mice that were added polysaccharide krestin before and after *M. tuberculosis* exposure. While interleukin-10 was in low concentration, production of Th1 increased. Raising amount of Th1 increased amount of activated macrophages so macrophages could do phagocytosis. Phagocytosis produced reactive oxygen intermediates and nitric oxide. While more of reactive oxygen intermediates and nitric oxide were produced, possibilities of nitrite and malondialdehyde forming were higher. The capa-

bility of polysaccharide krestin to mop up free radical would decrease because the amount of free radicals exceeds the capacity of polysaccharide krestin.

There was different result in P3 nitrite group and P3 malondialdehyde group. P3 nitrite group showed significant difference compared to K- group but P3 malondialdehyde group did not. The difference could happen because immune response of each individual could be different toward nitrite and malondialdehyde formation. Fungal  $\beta$ -glucan-induced immune responses are different in their actions based on individual immune system (Chen and Seviour, 2007).

Research results useful to give the immune system so as to prevent the transmission of pulmonary TB disease due to infection with *M. tuberculosis*. It is given that TB is a tropical disease that desperately need attention to be tackled. *Mycobacterium tuberculosis* is easily transmitted through the air when the patient coughs or sneezes TB, even when spitting and talking. One patient can spread the TB bacteria in 10-15 people in one year. In general, the reaction of the immune system can stop the progression of TB germs. If the immune system decreases the bacteria will grow, and consequently in a few months in question would be tuberculosis.

## CONCLUSION

It was concluded that adding polysaccharide krestin could influence nitrite and malondialdehyde concentration of mice serum exposed by *Mycobacterium tuberculosis*. Adding polysaccharide krestin before exposure (P1) and before-after exposure (P3) could decrease nitrite concentration. Adding polysaccharide krestin before exposure (P1) and after exposure (P2) could decrease malondialdehyde concentration. Based on the evidences, adding polysaccharide krestin before *Mycobacterium tuberculosis* exposure could decrease both of nitrite and malondialdehyde concentrations.

## ACKNOWLEDGEMENT

The authors are grateful for the financial support provided by the Center through Annual Research Project DIPA PUPT 2014, Airlangga University.

## REFERENCES

Abbas, A.K., Litchmann, A.H.H. & Pillai, S. (2012). *Cellular and Molecular Immunology*, 7<sup>th</sup> Edition. Philadelphia: WB Saunders Company.

- Brown, G.D., Taylor, P.R., Reid, D.M., Willment, J.A., Williams, D.L., Martinez-Pomares, L., Wong, S.Y., & Gordon, S. (2002). Dectin-1 is a Major  $\beta$ -Glucan Receptor on Macrophages. *J Exp Med*, 196(3), 407-412.
- Caramori, G. & Papi, A. (2004). Oxidant and Asthma. *Thorax*, 59, 170-173.
- Chan, G.C., Chan, W.K., & Sze, D.M. (2009). The Effect of  $\beta$ -Glucan on Human Immune and Cancer Cells. *Bio Med Central*, 2 (25), 1-11.
- Chen, J. & Seviour, R. (2007). Medicinal Important of Fungal Beta-(1-3), (1-6)-Glucans. *Mycol Res*, 111(Pt 6), 635-652.
- Cui, J. & Chisti, Y. (2003). Polysaccharopeptides of *Coriolus versicolor*: Physiological Activity, Uses, and Production. *Biotechnol Adv*, 21(2), 109-122.
- Dagli, M., Eryilmaz, A., Besler, T., Akmansu, H., Acar, A. & Hakan, K. (2011). Role of Free Radical and Antioxidant in Nasal Polyps. *Laryngoscope*, 114, 1200-1204.
- Dalle-Donne, I., Rossi, R., Colombo, R., Giustarini, D., & Milzani, A. (2006). A Biomarkers of Oxidative Damage in Human Disease. *Clin Chem*, 52(4), 601-623.
- Eide, D.J. (2001). Functional Genomics and Metal Metabolism. *Genome Biol*, 2(10), 1028.
- Gharavi, N. & El-Kadi, E.O. (2003). Measurement of Nitric Oxide in Murine Hepatoma Hepalclc7 Cells by Reserved Phase Hple with Fluorescence Detection. *J Pharmaceut Sci*, 6(2), 302-307.
- Guzik, T.J., Korb, R., & Guzik, T.A. (2003). Nitric Oxide and Superoxide in Inflammation and Immune Regulation. *J Physiol Pharmacol*, 54(4), 469-487.
- Jose, A., Gonzalves, N., Benayas, C.G. & Arenas, J. (2003). Semiautomated Measurement of Nitrate in Biology Fluids. *Europ J Clin Chem*, 17, 7-9.
- Nurdini, R.F. (2015). Pengaruh Polisakarida Krestin *Coriolus versicolor* Terhadap Konsentrasi IL-10 dan Histologis Paru Pada *Mus musculus* yang Terpapar *Mycobacterium tuberculosis*. *Skripsi*. Surabaya: Fakultas Sains dan Teknologi, Universitas Airlangga.
- Ooi, V.E.C & Liu, F. (2000). Immunomodulation and Anti-Cancer Activity of Polysaccharide-Protein Complexes. *Curr Med Chem*, 7, 715-729.
- Ozkan, O.V., Ozturk, O.H., & Aydin, M. 2010. Effects of  $\beta$ -glucan Pretreatment on Acetylsalicylic Acid-Induced Gastric Damage: An Experimental Study in Rats. *Curr Ther Res Clin Exp*, 71(6), 369-383.
- Pang, Z.J., Chen, Y., Zhou, M., & Wan, J. (2000). Effect of Polysaccharide krestin on Glutathione Peroxidase Gene Expression in Mouse Peritoneal Macrophages. *Br J Biomed Sci*, 57(2), 130-136.
- Pham-Huy, L.A., He, H. and Pham-Huy, C. (2008). Free Radicals, Antioxidants in Disease and Health. *Int J Biomed Sci*, 4(2): 89-96.
- WHO [World Health Organization]. (2015). *Global Tuberculosis Report*. Switzerland Geneva: World Health Organization.
- Zhang, W., Liu, H., Rojas, M., Caldwell, R.W. & Caldwell, R. B. (2011). Anti Inflammatory Therapy for Diabetic Retinopathy. *Immunotherapy*, 3(5), 609-628.