

The Correlation of Bone Mineral Density (BMD), Body Mass Index (BMI) and Osteocalcin in Postmenopausal Women

by Eha Renwi Astuti

Submission date: 31-Mar-2022 01:22PM (UTC+0800)

Submission ID: 1797624441

File name: ody_Mass_Index_BMI_and_Osteocalcin_in_Postmenopauseal_Women..pdf (346.23K)

Word count: 3693

Character count: 20419



The Correlation of Bone Mineral Density (BMD), Body Mass Index (BMI) and Osteocalcin in Postmenopausal Women

Nike Hendrijantini^{1*}, Rostiny Alie¹, Rosy Setiawati², Eha Renwi Astuti³ and Manggala Pasca Wardhana⁴

¹Department of Prosthodontics, Faculty of Dental Medicine, Airlangga University, Indonesia

²Department of Radiology, Faculty of Medicine, Airlangga University, Indonesia

³Department of Radiology, Faculty of Dental Medicine, Airlangga University, Indonesia

⁴Department of Obstetrics and Gynecology, Faculty of Medicine, Airlangga University, Indonesia

*Corresponding author: Nike Hendrijantini, Department of Prosthodontics, Faculty of Dentistry, Airlangga University, Surabaya, Indonesia, Tel: 62-81-134-1610; E-mail: hendrijantininike@yahoo.com

Received date: June 15, 2016; Accepted date: July 25, 2016; Published date: August 01, 2016

Copyright: © 2016 Hendrijantini N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Osteoporosis is related to the decrease of bone density, osteoporosis requires attention from dentists because it can also occurs in the jaw bone. Osteoporosis is calculated with a quantitative assessment of bone density, which is called Bone Mineral Density (BMD). The best imaging modalities to assess bone density are the Dual Energy examination methods X-ray Absorptiometry (DEXA). In addition to DEXA, osteoporosis examination can be performed with measurement of Body Mass Index (BMI), which is a ratio of weight and height. Process of bone formation by osteoblasts can be examined by bone marker such as Osteocalcin, as a parameter (alone or in combination with BMD) to determine metabolic bone disorders during bone formation and bone remodeling (bone turnover).

Aim: To analyze the correlation among BMD, BMI and Osteocalcin in Postmenopausal women, to investigate the possibility of using osteocalcin examination to predict the mandibular bone osteoporosis.

Method: Fifty four women above 51 years old who are at least 1 year postmenopausal, underwent BMD (using DEXA examinations), BMI, and Osteocalcin. The results of each examination were counted, and the correlations among each examinations were evaluated using Spearman's correlation test.

Result: Means of BMD, BMI, and Osteocalcin in postmenopausal women were 1.606, 25.189, and 30.566 respectively. BMD were significantly correlated with BMI (Spearman's rank correlation coefficient $r=0.414$, $p<0.05$). While, BMD were significantly correlated with Osteocalcin ($r=-0.343$, $p<0.05$). Moreover, BMI were significantly correlated with Osteocalcin ($r=-0.274$, $p<0.05$).

Conclusion: There were significant correlations among BMD, BMI, and Osteocalcin results in postmenopausal women. It is concluded that each examination of BMD, BMI, and Osteocalcin can be used to identify the risk of osteoporosis in postmenopausal women. Therefore simple examination of osteocalcin can be used to predict mandibular bone loss.

Keywords: BMD; BMI; Osteocalcin; Osteoporosis; Postmenopausal women

Introduction

Osteoporosis is a disease which requires more attention from the dentists, osteoporosis is related to the decrease of bone density, and it can also occurs in the jaw bone. The osteoporotic jaw bone will experience the decrease of jaw bone density, the increase alveolar jaw bone resorption, and also the increase risk of periodontal disease and tooth loss. It is crucial for dentists and especially prosthodontist to diagnose jaw bone density, especially in postmenopausal women before prosthodontic treatment. There are numerous failure in prosthodontic treatment due to osteoporosis. Osteoporosis will cause mandible resorption, which in turn cause the failure of osteointegration between the implant and the jaw bone [1].

Problem of osteoporosis in postmenopausal women is associated with the decrease of estrogen level. This condition will accelerate bone fragility [2]. World Health Organisation (WHO) defines osteoporosis as a disease that is characterized by low bone mass and microarchitectural destruction of bone tissue, leading to bone fragility and increased risk of fracture. Diagnosis of osteoporosis according to WHO is obtained from any of the three regions (total hip, femoral neck and lumbar spine). Study Group Meeting of WHO assessed fracture risks and applied it to screen postmenopausal osteoporosis, and recorded that more than 75 million people in the United States, Europe and Japan were suffering from osteoporosis on 2006. Osteoporosis caused more than 8.9 million fractures per year that occurred in the United States and Europe [3]. While, research of Health Department showed that two out of five people in Indonesia have the risk of osteoporosis [4].

Diagnosis of osteoporosis is calculated with a quantitative assessment of bone density, which is called Bone Mineral Density (BMD) [5]. The best imaging modalities to assess bone density are the Dual Energy examination methods x-ray absorptiometry (DEXA) and Quantitative Computerized Tomography (CT). CT is more sensitive but with the added risk of more radiation exposure, several experts prefer DEXA examination due to the less radiation exposure. BMD in postmenopausal women is determined by peak bone mass and the amount of bone lost since menopause [6].

In addition to DEXA, osteoporosis examination can be done with Body Mass Index (BMI). BMI is a ratio of weight and height with units (kg/m²) [7]. Regarding to BMD and BMI, there are four important factors to assess the quality of bone, these are bone density; bone turnover (remodeling); bone size; and bone geometry (microarchitecture) [6].

Osteocalcin is known as bone Gla protein, a marker of bone formation, vitamin K and vitamin D-dependent protein produced by osteoblasts and most widely studied of the non-collagen protein in bone [8]. Osteocalcin plays an important role in the process of mineralization and calcium ion homeostasis process. Osteocalcin is synthesized during the bone formation and it exhibits a compact, calcium dependent, alpha helical formation, in which the Gamma Carboxyglutamic Acid (GLA) residues bind and promote absorption to hydroxyapatite in the bone matrix, thus leading to bone mineralization. Calcium and phosphorus deficient osteoporotic woman may have a decrease rate of bone mineralisation due to a reduction in hydroxyapatite crystal formation [9].

The examination of osteocalcin is a good parameter for determining bone metabolism disorders during bone formation and bone turnover. This marker is often used as an early biomarker of bone-forming therapy and to assess the effectiveness of treatment outcome [6].

Relationship between BMD and BMI is still not clearly defined, some authors found out a positive relationship, while others showed that BMI is a risk factor for osteoporosis. There are some researches showed positive correlation between systemic osteoporosis and the decrease of mandibular bone density [10].

There is increasing evidence that a combination of a biochemical marker and BMD may be a better predictor of fracture than BMD alone, yet biochemical marker such as osteocalcin may also predict fracture risk independently of BMD. High bone turnover can disrupt the trabecular architecture by increasing the incidence of trabecular perforation and buckling, thus reducing bone strength, without necessarily appreciably affecting BMD [9].

In this study we analyzed the correlation among BMD (calculated from lumbar spine), BMI and Osteocalcin in postmenopausal women. While BMD is still the gold standart for osteoporosis diagnosis, the device is unfortunately limited in our country, osteocalcin examination in the other hand is easier, cheaper and more simple. Researcher wants to investigate the possibility of using osteocalcin examination to predict the mandibular bone osteoporosis.

Materials and Methods

Study design

This is a cross sectional observation study, conducted at Pramita and Prodia Laboratory. This study is using fifty four postmenopausal women who recieved prosthodontic treatment from January until March 2014 in prosthodontic clinic Airlangga University as subjects, and all of them were given informed consent to be examined. Characteristics of subjects were at least 1 year postmenopausal women, above 51 years old, no history of systemic diseases (Hyperparathyroidism, Paget's Disease, Osteomalacia, Renal Osteodystrophy, Osteogenesis Imperfecta), no history of cancer or kidney diseases, no history of hormonal therapy and smoking.

BMD measurement

BMD is the main criteria used to diagnose and monitor osteoporosis. DEXA is the current gold standard examination of osteoporosis in male and female, which is used to examine the bone mass of total hip, femoral neck, lumbar spine, or whole body. The purpose of bone mass measurement are to determine diagnosis of osteoporosis, to predict the occurrence of fractures, to assess changes in bone density after treatment [11]. In this study, position of patients were in lumbar spine with center of laser beam in lumbal. Lumbar spine is measured in this research, due to the faster bone loss nature compared with the hip in post-menopausal women [12]. Measurement of bone density usually expressed in T-scores, where the standard deviations number of bone density varies from the average bone density in normal subjects with the same sex. WHO define osteoporosis as a condition in which bone mineral density is below -2.5 SD, osteopenia in which bone mineral density between -1 to -2.5 SD, while normal in which bone mineral density is above -1 SD [13].

BMI measurement

BMI is a person's weight in kilograms divided by the square of height in meters. Height was measured using a 0.1 microtoise cm precision. Body weight was measured using a digital weight scales Seca brands with 0.1 kg precision. BMI figure obtained by using the weight formula (kg/m²).

Osteocalcin measurement

The examination of osteocalcin is a good parameter to determine bone metabolism disorders during bone formation and bone turnover. Osteocalcin level were examined using enzyme-linked immunosorbent assay (ELISA). Normal values were: 10.1 ± 9.4 ng/ml [6]. *In vivo* quantitative measurement of N-MID osteocalcin levels were performed in serum or plasma. Examination of the N-Mid Osteocalcin with postmenopausal scale 15-46 ng/ml. Stability samples 2-80C (immediately after sampling), 200C (4 hours after sampling, 2 hours if the sample correction carried out in room temperature), -800°C (more than 1 month). This examination is used as a therapeutic control antiresorp. Principles of inspection: The principle of this test is a sandwich ELISA.

In the first incubation, the sample will react with the monoclonal antibody specific N-MID osteocalcin and biotin-labeled monoclonal antibodies specific N-MID osteocalcin labeled ruthenium complex to form a sandwich. Furthermore, the addition of streptavidin-coated

microparticles that cause complex bound to the solid phase via interaction of biotin and streptavidin. The reaction mixture is put into the measuring cell where the microparticles were captured with a magnet on the electrode surface, the compounds are not bound disposed pro cell. Voltage provision to the electrodes induces chemiluminescent emission. Results are determined via a calibration curve made by means of a two-point calibration curves and masks available on the reagent barcode.

Ethical Clearance

This study has received ethical clearance from the Commission on Ethics Health Research (KKPEK).

10 Statistical Analysis

Data analysis was performed using SPSS software (version 21.0; SPSS, Inc., Chicago, IL). Subject characteristics are presented as mean ± SD unless otherwise stated. Agreement among BMD and BMI, and osteocalcin determined using Spearman's rank correlation coefficient. Significant correlation were found if p<0.05.

Results

Characteristics of participants: There were fifty four postmenopausal women, with average age of 63, 78 years (range 51-83 years old), with 83.33% subjects were in the range of 51-65 years old and 16.67% subjects were in range above 66 years old.

Difference mean of BMD, BMI, and osteocalcin are listed in Table 1. Mean of BMD, BMI, and osteocalcin were 1.606 ± 1.213, 25.189 ± 4.076, 30.566 ± 9.144. Based on BMD value, 27.8% of subjects were osteoporosis, 44.4% of subjects were osteopenia, and 27.8% were normal.

Agreement among BMD, BMI, and osteocalcin with non-parametric method are listed in Table 2, with Spearman rank correlation. Assessments of BMD and BMI were significantly related (p<0.05, r=0.414, positive relation). BMD and osteocalcin were significantly related (p<0.05, r=-0.343, negative relation), as well as BMI and osteocalcin were also significantly related (p<0.05, r=-0.274, negative relation). BMD and BMI had a stronger relation compared with BMI and osteocalcin.

Correlation types are listed in Figure 1. Correlation between BMD and BMI showed linear positive, because dots in graph above were placed in positive axis. Higher BMD linear also has higher BMI. Correlation between BMD and osteocalcin were negative because dots in graph above were placed in negative axis. Higher BMD has lower Osteocalcin, as well as correlation between BMI and osteocalcin.

Category	Mean	Standard Deviation
BMD	1.606	1.213
BMI	25.189	4.076
Osteocalcin	30.566	9.144

Table 1: Difference Means of BMD, BMI, and osteocalcin with standard deviation.

Correlations					
			BMD	BMI	Osteocalcin
Spearman's rho	BMD	Correlation Coefficient	1	.414**	-.343*
		Significance tailed)	(2-	.002	0.011
		N	54	54	54
	BMI	Correlation Coefficient	.414**	1	-.274*
		Significance tailed)	(2-	0.002	.045
		N	54	54	54
Osteocalcin	Correlation Coefficient	-.343*	-.274*	1	
	Significance tailed)	(2-	0.011	0.045	
	N	54	54	54	

** Cor. is significant at .01 level 2-tail...
* Cor. is significant at .05 level 2-tail...

Table 2: Correlations among BMD, BMI, and osteocalcin with Spearman correlation statistical test.




Figure 1: Scatterplot correlations from BMD and BMI, BMD and osteocalcin, BMI and osteocalcin.

Discussion

In this study, bone mineral density was evaluated using a DEXA of the spine. Density of the spine is most responsive to drugs intervention and monitoring for patient. DEXA is a technology to determine the relationship between bone density and fracture risk, and has been validated in a longitudinal cohort study. DEXA obtained some information about bone density in a given area in grams/cm², comparison between mean level of bone density compared to the mean level of bone density with the same ethnic adults called T score in percent (%) [11].

This study showed that BMD and BMI had significant correlation (positive correlation) showed in Table 2. This result was consistent with those reported by Hsu et al. [14] and Ahmad et al. [15] which reported a significant positive relationship among weight, BMI and BMD. BMI is a good indicator and often used for measuring BMD [16], this is similar with the research of Iqbal which suggested that BMI is a good indicator for referring patients to do DEXA scanning on Bone Mineral Density (BMD) [17]. The results of the bivariate analysis of Herkulina study showed positive and significant correlation between Body Mass Index (BMI) and Bone Mineral Density (BMD) [18].

The study of Kim et al. also stated that low body weight is more likely associated with osteoporosis [19]. All of height, weight and BMI positively correlated with the dependent variable BMD T-score. As well as research of Hubungan [20] showed that low BMI and decreased bone strength associated with reduced bone mass in all parts of the body. Study of Mendez et al. showed relationship of BMI and BMD was significantly positive at the lumbar spine, pelvis and femoral neck [21].

Correlation between BMD and osteocalcin had significant relationship (negative correlation) showed in Table 1, this is well according to the theory that as a specific marker of osteoblast function, level of osteocalcin have been shown to correlate with bone formation rates, osteoporotic post-menopausal women will have an increase of serum OC concentration. The same statement was made by Kalaiselvi et al. [22] which reported a negative correlation found between the osteocalcin level and the BMD in post-menopausal women (Pearson's correlation $p < 0.05$ and $r = -0.595$). That study in 60 post-menopausal women at Sree Balaji Medical College and Hospital and Bharat Scan Limited at Chennai, India, as well as study of Susanto [9] reported a significant inverse correlation between serum osteocalcin and femoral neck BMD ($p < 0.05$, and $r = -0.29$).

BMI and osteocalcin have significant correlation (negative correlation) showed in Table 2. The same statement is made by Kalaiselvi et al. [22] which reported that a significant correlation was found between osteocalcin level and BMD in post-menopausal women (Pearson's correlation $p < 0.05$, and $r = 0.339$).

Despite BMI usage as a measurement to predict whether a person is overweight or not, it has some drawbacks and may not be an accurate tool for assessing weight status and height. This study is similar with that of body weight or BMI indicates the value inversely with markers of bone turnover, where people with low body weight has higher bone turnover, while overweight and obese have lower serum osteocalcin compared to normal weight. Effect of obesity on the risk of fracture depends on the definition, if defined by BMI or body weight, obesity may protect against bone mineral loss or vertebral fractures. However, if obesity is based on the percentage of body fat, obesity may be a risk factor for osteoporosis [9].

Osteoporosis is not only occurred in lumbar, femur and radius bone, but also on jaw bone, researches have concluded that density of mandible bone related with density of other bones significantly [1]. Based on this research, there was a correlation among BMD, BMI and osteocalcin. Those examinations therefore can be used to predict osteoporosis of the mandible bone.

Conclusion

There were significant correlations among BMD, BMI, and Osteocalcin results in postmenopausal women. BMD and BMI had

positive correlation, while BMD and osteocalcin had negative correlation, and BMI and osteocalcin had also negative correlation. It is concluded that examination of BMD, BMI, and Osteocalcin can be used to identify the risk of osteoporosis in postmenopausal women. Therefore simple examination of osteocalcin can be used to predict bone loss in mandibular bone.

Acknowledgement

To all of respondents whom involved in this study, and also Prodia laboratorium for examining osteocalcin.

References

1. Lindawati SK (2012) Dampak Osteoporosis Tulang Rahang Terhadap Perawatan Prostodontik dan Kualitas Hidup Lansia. *Pidato Pengukuhan Guru Besar FKG UI*, pp. 3-15.
2. Abas BJ dan, Sri P (2007) Risiko Osteoporosis di Indonesia. *Publitsbang Gizi dan Makanan Depkes RI Bogor* 30: 1-11.
3. World Health Organization (2004) "WHO Scientific Group on The Assessment of Osteoporosis at Primary Health Care Level. Summary Meeting report Brussel, Belgium, pp.2
4. Ichramshah AR, dan Bambang S (2010) Penyakit Osteoporosis.
5. Compston J, Cooper A, Cooper C, Francis R, Kanis JA, et al. (2013) Guideline for the diagnosis and management of osteoporosis in postmenopausal women and men from the age of 50 years in the UK. *Maturitas* 62: 105-8.
6. Adi P (2007) Peran Pertanda Tulang Dalam Serum Pada Tata Laksana Osteoporosis. *Universa Medicina* 26: 152-9.
7. Centers For Disease Control (2009) Body Mass Index: "Considerations for Practitioners. Department of Health and Human Services, USA. pp. 1.
8. Lee AJ, Hodges S, Eatell R (2000) Measurement of Osteocalcin. *Association of Clinical Biochemist* 37: 432-446.
9. Susanto LTM (2011) Serum Osteocalcin and bone mineral density in postmenopausal women. *Universa Medicina* 30: 155-161.
10. Ishii K, Taguchi A, Nakamoto T, Ohtsuka M, Sutthiprapaporn P, et al. (2007) Diagnostic efficacy of alveolar bone loss of the mandible for identifying postmenopausal women with femoral osteoporosis. *Dentomaxillofac Radiol* 36: 28-33.
11. Kawiayana IKS (2009) Osteoporosis, Patogenesis Diagnosis dan Penanganan Terkini. *J Penyakit Dalam* 10: 158-162.
12. Bonnick SL, Shulman L (2006) "Monitoring Osteoporosis Therapy: Bone Mineral Density, Bone Turnover Markers, or Both?". *The American Journal of Medicine* 119: 255-315.
13. Margo U, Wulandari M, Zilfa KP (2010) Faktor-Faktor Yang Berhubungan Dengan Kepadatan Tulang Pada Wanita Postmenopause. *Jurnal Unimus* 6: 1-2;
14. Hsu YH, Venners SA, Terwedow HA, Feng Y, Niu T, et al. (2006) Relation of body composition, fat mass, and serum lipids to osteoporotic fractures and bone mineral density in Chinese men and women. *Am J Clin Nutr* 83: 146-54.
15. Ahmad EA, Abbas M, Amir R, Mohammad S (2012) The Relationship between Body Weight, Body Mass Index (BMI) and Bone Mineral Density (BMD) of the Lumbar Spine and Femoral neck in Professional Cyclist of Iran and Tour de France. *Iranian Journal of Health and Physical Activity* 4: 67-71.
16. Budi S, Sri P, Wasilah R, dan Retno P (2011) Hubungan Indeks Massa Tubuh Dengan Densitas Mineral Tulang Pada Perempuan Dewasa Muda. *Pusat Penelitian dan Pengembangan Gizi dan Makanan, Badan Penelitian dan Pengembangan Kesehatan. Kemenkes RI* 34: 93-103.
17. Iqbal SI, Morch LS, Rosenzweig M, Dela F (2005) The Outcome of Bone Mineral Density Measurements on Patients Referred From General Practise. *Journal of Clinical Densitometry* 8: 178-182.

Citation: Hendrijantini N, Alie R, Setiawati R, Astuti ER, Wardhana MP (2016) The Correlation of Bone Mineral Density (BMD), Body Mass Index (BMI) and Osteocalcin in Postmenopausal Women. *Biol Med (Aligarh)* 8: 319. doi:[10.4172/0974-8369.1000319](https://doi.org/10.4172/0974-8369.1000319)

Page 5 of 5

-
18. Herkulina F (2012) Hubungan Indeks Massa Tubuh dan Komposisi Tubuh dengan Densitas Massa Tulang Wanita Dewasa di Perwatusi Kota Pontianak. Electronic Tesis dan disertasi Universitas Gajah Mada.
 19. Kim SJ, Yang WG, Cho E, Park E (2012) Relationship between Weight, Body Mass Index and Bone Mineral Density of Lumbar Spine in Women. *J Bone Metab* 19: 95-102.
 20. Hubungan NA (2004) Tingkat Konsumsi Kalsium, Protein dan Status Gizi dengan Derajat Osteoporosis Pada lansia.
 21. Mendez JP, Mejia DR, Pedraza J, Varquez RMC, Soriano R, et al. (2013) Bone Mineral Density in Postmenopausal Mexican-Mestizo Women with Normal Body Mass Index, Overweight, or Obesity. *Menopause* 20: 568-72.
 22. Kalaselvi, Prabhu K, Ramesh M, Venkatesan V (2013) The Association of Serum Osteocalcin with the Bone Mineral Density in Post-Menopausal Women. *J Clin Diagn Res* 7: 814-816.

The Correlation of Bone Mineral Density (BMD), Body Mass Index (BMI) and Osteocalcin in Postmenopausal Women

ORIGINALITY REPORT

19%

SIMILARITY INDEX

7%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Richard Eastell, Rosemary A. Hannon. "Biomarkers of bone health and osteoporosis risk", Proceedings of the Nutrition Society, 2008
Publication 3%
- 2** www.liebertpub.com
Internet Source 2%
- 3** Bone Densitometry in Clinical Practice, 2010.
Publication 1%
- 4** Harald Herkner, Ulla Waldenhofer, Anton N. Laggner, Marcus Müllner et al. "Clinical application of rapid quantitative determination of cardiac troponin-T in an emergency department setting", Resuscitation, 2001
Publication 1%
- 5** Merve Nur Ceylan, Sevginur Akdas, Nuray Yazihan. "Is Zinc an Important Trace Element on Bone-Related Diseases and Complications? A Meta-analysis and Systematic Review from

Serum Level, Dietary Intake, and Supplementation Aspects", Biological Trace Element Research, 2020

Publication

6

Lee, A. J., S. Hodges, and R. Eastell. "Measurement of Osteocalcin", *Annals of Clinical Biochemistry An international journal of biochemistry and laboratory medicine*

Publication

1 %

7

www.ncbi.nlm.nih.gov

Internet Source

1 %

8

Harry Freitag Luglio Muhammad, Erri Larene Safika, Fani Cahya Wahyuni, Aviria Ermamilia, Emy Huriyati. "The Effect of Zumba Training on Body Composition, Dietary Intake, Sleep Quality, and Duration in Adult Sedentary Women With Overweight", *Topics in Clinical Nutrition*, 2019

Publication

1 %

9

aatmabharat.blogspot.com

Internet Source

1 %

10

Andrew A. Almeida, Emily H. Reeve, Rachel L. Dickinson, Megan Carty, Julia Gilpin, Deborah L. Fearheller. "Civilians Have Higher Adherence and More Improvements in Health with a Mediterranean Diet and Circuit Training Program Compared to Firefighters", *Journal of Occupational & Environmental Medicine*, 2022

<1 %

11

univmed.org

Internet Source

<1 %

12

Bin Chen, Gongwen Liu, Xiao Wang, Yan Gao, Aifei Wang, Guangfei Li, Peng Jia, Jun Dai, Hui Zhang, Youjia Xu. "The Influence of Body Fat Content and Distribution on Bone Mass in Healthy Chinese Adults", Research Square Platform LLC, 2022

Publication

<1 %

13

O. Ström, F. Borgström, John A. Kanis, Juliet Compston, Cyrus Cooper, Eugene V. McCloskey, Bengt Jönsson. "Osteoporosis: burden, health care provision and opportunities in the EU", Archives of Osteoporosis, 2011

Publication

<1 %

14

Phui-Ly Liew, Weu Wang, Yi-Chih Lee, Ming-Te Huang, Yang-Chu Lin, Wei-Jei Lee. "Gallbladder Disease among Obese Patients in Taiwan", Obesity Surgery, 2007

Publication

<1 %

15

digitalcommons.library.umaine.edu

Internet Source

<1 %

16

primarycare.imedpub.com

Internet Source

<1 %

17

Annie W. C. Kung, Andrew Y. Y. Ho, Philip D. Ross, Jean-Yves Reginster. "Development of a clinical assessment tool in identifying Asian men with low bone mineral density and comparison of its usefulness to quantitative bone ultrasound", *Osteoporosis International*, 2004

Publication

<1 %

18

K V Haderslev. "Vitamin D status and measurements of markers of bone metabolism in patients with small intestinal resection", *Gut*, 5/1/2003

Publication

<1 %

19

repository-tnmgrmu.ac.in

Internet Source

<1 %

20

A R Hermus, A G Smals, L M Swinkels, D A Huysmans, G F Pieters, C F Sweep, F H Corstens, P W Kloppenborg. "Bone mineral density and bone turnover before and after surgical cure of Cushing's syndrome.", *The Journal of Clinical Endocrinology & Metabolism*, 1995

Publication

<1 %

21

Dini Setiarsih, Bambang Wirjatmadi, Merryana Adriani. "Bone Density Status and Vitamin D and Calcium Concentrations in Pregnant and Non-Pregnant Women", *Makara Journal of Health Research*, 2016

<1 %

22

"Poster Presentations - Clinical abstracts (ICOBR 2014)", IBMS BoneKEy, 2014.

Publication

<1 %

23

"Preventing cardiogenic shock protects patients with acute MI", Postgraduate Medical Journal, 2003

Publication

<1 %

24

Hanlon, Phil, Carlisle, Sandra, Lyon, Andrew. "EBOOK: The Future Public Health", EBOOK: The Future Public Health, 2012

Publication

<1 %

25

Li Zhu, Jing Zhang, Jing wang, Xiaohong Lv, Dan Pu, Yanhua Wang, Qian Men, Lan He. "Uncoupled bone remodeling is characteristic of bone damage in premenopausal women with new-onset systemic lupus erythematosus", Lupus, 2021

Publication

<1 %

26

Neeta G. Nayak, Laura D. Carbone. "Management of osteoporosis (an update on diagnosis and management, with an emphasis on non-hormonal treatments)", Primary Care Update for OB/GYNS, 1997

Publication

<1 %

27

Shun-zhi Liu, Li-fang Tian, Peng Xu, Gui-hua Zhuang, Fang Zheng, Juan Tian, Qi-Lan Ning, Bo-Feng Zhu, She-Min Lu, Hong Yan. "Analysis

<1 %

of correlation between blood biochemical indicators and bone mineral density of post-menopausal women", Molecular Biology Reports, 2010

Publication

28

link.springer.com

Internet Source

<1 %

29

www.netnest.com.au

Internet Source

<1 %

30

"Abstracts of the 35th European Symposium on Calcified Tissues", Calcified Tissue International, 2008

Publication

<1 %

31

"Oesophagus posters 442-464", Gut, 2002

Publication

<1 %

32

Jos H. H. Thijssen. "Gene polymorphisms involved in the regulation of bone quality", Gynecological Endocrinology, 2009

Publication

<1 %

33

Shahzad, K.. "Analytic evaluation of the @b-human chorionic gonadotropin assay on the Abbott IMx and Elecsys2010 for its use in doping control", Clinical Biochemistry, 200711

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

The Correlation of Bone Mineral Density (BMD), Body Mass Index (BMI) and Osteocalcin in Postmenopausal Women

[GRADEMARK REPORT](#)

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
