

ANTIMICROBIAL RESISTANCE DETECTION OF CITROBACTERFREUNDII ISOLATED FROM RECTAL SWAB IN EAST JAVA

by Wiwiek Tyasningsih

Submission date: 04-Nov-2021 05:08PM (UTC+0800)

Submission ID: 1692804560

File name: Antimicrobial_Resistance_Detection_of.pdf (1M)

Word count: 2443

Character count: 14407



ISSN: 2230-9926

Available online at <http://www.journalijdr.com>

IJDR

International Journal of Development Research
Vol. 07, Issue, 11, pp.17119-17122, November, 2017



ORIGINAL RESEARCH ARTICLE

OPEN ACCESS

ANTIMICROBIAL RESISTANCE DETECTION OF *CITROBACTERFREUNDII* ISOLATED FROM RECTAL SWAB IN EAST JAVA

^{1*}Mustofa Helmi Effendi, ¹Intan Permatasari Hermawan, ²Wiwiek Tyasningsih, ³Suwarno, ¹Intan Galuh Bintari, ¹Risi Cicilia and ¹Rinda Dewi Safitri

¹Department of Veterinary Public Health, College of Veterinary Medicine, Airlangga University

²Laboratory of Bacteriology, Department of Microbiology and Mycology, College of Veterinary Medicine, Airlangga University

³Laboratory of Virology, Department of Microbiology and Mycology, College of Veterinary Medicine, Airlangga University, Surabaya, Indonesia

ARTICLE INFO

Article History:

Received 05th August 2017
Received in revised form 29th September, 2017
Accepted 18th October, 2017
Published online 30th November, 2017

Key Words:

Antimicrobial resistance, *Citrobacterfreundii*, Rectal swab, bla_{TEM} gene, ESBL.

ABSTRACT

The aim of this study was to detect bla_{TEM} gene which the agent of antibiotic resistance in *Citrobacterfreundii* isolated from rectal swab in East Java. A total of 275 rectal swab samples were collected from 12 district of East Java. Then, samples were isolated with selective media and biochemical tests. Antibiotic Resistance test was using the disc diffusion against amoxycillin, amoxycillin-clavulanic acid, ampicillin-sulbactam, cefotaxime, ceftazidime, sulfametaxole-trimethoprim, and tetracycline and then detection of bla_{TEM} gene used PCR. The number of *Citrobacterfreundii* was 7 (2,55 %). Antimicrobial resistance showed 2 broilers and 1 dairy cattle from Surabaya resistant to ampicillin-sulbactam and tetracycline, 4 milkfish from Sidoarjo resistant to amoxicillin and ampicillin-sulbactam. All seven samples were positive bla_{TEM} gene with 445 bp size used PCR.

Copyright ©2017, Mustofa Helmi Effendi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Mustofa Helmi Effendi, Intan Permatasari Hermawan, Wiwiek Tyasningsih, Suwarno, Intan Galuh Bintari, Risi Cicilia and Rinda Dewi Safitri. 2017. "Antimicrobial resistance detection of citrobacterfreundii isolated from rectal swab in east java", International Journal of Development Research, 7, (11), 17119-17122.

INTRODUCTION

Citrobacter freundii is a group of Enterobacteriaceae bacteria that can be isolated from stool specimens or intestinal tract of humans and animals (Kao et al., 2010). Farm products became the biggest source of fecal contamination that can cause human pathogens. Fecal contamination of water or feces usually present in the environment or migrate through the water. Still little is known about the effects of antibiotic resistance in animal faeces. Antibiotic resistance in the particles containing the gene resistant genetically and can spread through animal feces in the soil and aquatic ecosystems (Adelowo and Fagade, 2010). Excessive use of antibiotics as well as non-compliance with medication causing downtime

residues in poultry meat, eggs and milk or other dairy products (Maynard et al., 2003). National Institutes of Health reported that the death rate in the United States reached 90,000 people per year due to antibiotic resistance. Antimicrobial Resistance Bank CDC in 2013 reported That *Citrobacterfreundii* resistant to the antibiotic Amoxicillin, Ampicillin, cefazolin and Cefoxitin (Guilfoile et al., 2007). Antibiotic resistance to beta-lactam antibiotics caused by the production of beta-lactamase, an enzyme that can hydrolyze beta-lactam ring so that the bacterial cell wall synthesis process is not disrupted. One of the enzymes produced by bacteria is the extended-spectrum beta-lactamase, or ESBL which is the result of mutations of beta-lactamase. Most ESBL derived from TEM-type enzyme encoded by the gene bla_{TEM}. bla_{TEM} genes are genes that cause antibiotic resistance in the plasmid of the most frequently detected in clinical populations of gram-negative microorganisms (Wilopo et al., 2015). Based on this background it is necessary to detect the gene encoding bla_{TEM}

*Corresponding author: Mustofa Helmi Effendi,

Department of Veterinary Public Health, College of Veterinary Medicine, Airlangga University.

¹⁸ as antimicrobial resistance in bacteria *Citrobacterfreundii* isolated from rectal swab of animal that endanger public health.

³ MATERIALS AND METHODS

Samples

A total of 275 rectal swab samples were collected from 12 district. Rectal swab samples were taken aseptically. The sampling technique is purposive sampling. Rectal swab taken with a sterile cotton swab that has been prepared and then inserted into the sample pot containing physiological saline and has been labelled the previous location. Rectal swabs taken directly from the rectum or spontaneous defecation (rectal toucher). Taken stool samples must be free of urine to prevent bacterial contamination of urine. All samples were stored in the cool box or directly checked exceed 30-40 minutes.

Table 1. Samples

No	District	Sample	Number of sample
1	Surabaya	Catfish, Tilapia fish, Parrot fish, Shark catfish, Shrimp, Dairy cattle, Broiler	33
2	Sidoarjo	Tilapia fish, Milkfish, Catfish, Shark catfish, Beef cattle	31
3	Gresik	Tilapia fish, ¹⁹ mp, Milkfish	8
4	Bojonegoro	Tilapia fish, Catfish, Gurame fish	12
5	Blitar	Tilapia fish, Catfish, Gurame fish	2
6	Probolinggo	Tilapia fish, Milkfish, Shrimp, Grouper fish, Mullet fish	11
7	Mojokerto	Tilapia fish, Catfish, Gurame fish, Shark catfish	22
8	Malang	¹⁶ pia fish, Milkfish, Shrimp	8
9	Nongkojajar	Dairy cattle	39
10	Grati	Dairy cattle	48
11	Senduro	Dairy cattle	30
12	Batu	Dairy cattle	31
Total sample			275

Table 2. Results of resistance phenotype

No	Code	Sample	Location	Resistance phenotype
1	AW-1	Broiler	Wonokromo, Surabaya	SAM, TET
2	AK-5	Broiler	Keputran, Surabaya	SAM, TET
3	SB-3	Dairy cattle	BendulMerisi, Surabaya	SAM, TET
4	SPT.K-1	Beef cattle	Krian, Sidoarjo	AML, SAM
5	IB.S-1	Milkfish	Sidoarjo	AML, SAM
6	IB.S-2	Milkfish	Sidoarjo	AML, SAM
7	IB.S-4	Milkfish	Sidoarjo	AML, SAM

AML= amoxicillin, SAM= ampicillin-sulbactam, TET= Tetracycline

Isolation of *Citrobacterfreundii*

All sa⁹ples were diluted and then isolated on Eosin Methylene Blue Agar and Mac Conkey's Agar. The inoculated plates were then incubated at 37°C for 24 hours and then observed macroscopically. Characteristics of bacterial colonies *Citrobacterfreundii* on EMBA appeared diameter of 2-4 mm, smooth, low, convex, moist, metallic green, on MCA appear pink colonies (Holt *et al.*, 1994). Furthermore, all bacterial isolated and identified by magnification microscopic examination with negative Gram staining. Negative gram stain showed short stems (bacilli) and then all bacterial isolated and identified by biochemical test (Lay, 1994). Identification test using Triple Sugar Iron Agar, glucose, lactose, sucrose,

mannitol, maltose, Simon's Indol Motility, MR- Vogen Prekauer (VP), Simon's Citrate Agar and Urea Agar.

⁴ Antibiotic Susceptibility Testing

The antimicrobial sensitivity phenotypes of bacteria isolates were determined using a Kirby-Bauer disk diffusion assay according to the standards and interpretive criteria described by the Clinical and Laboratory Standards Institute. And used stand art Mac Farland number 0.5 so that the concentration of bacteria to be equivalent to 3.0×10^8 CFU / ml (Mcfarland, 1907). Inspection began with a cotton stick sterile dipped into the bacterial suspension then ¹⁵ssed on the walls of the tube until the cotton is not too wet, then applied to the surface of the Muller Hinton Agar until blended, wait for about 15 minutes, then put the disc containing antibiotics on it and incubated at 37°C for 19-24 hours. Antibiotic resistance tests against Amoxycillin 20 mg, Amoxicillin-clavulanate 20/10 mg, ampicillin-sulbactam 10µg / 10µg, Cefotaxime 30 g, Ceftazidime 30 g, Sulfametaxole-Trimethoprim 23.75 / 1,25ug, Tetracycline 30 ug. Observations of each antibiotic susceptibility test results based on the diameter of that zone was measured using calipers (Ahmed and Shimamoto, 2011).

Detection of bla_{TEM} Using PCR

DNA extraction

DNA bacteria extraction were grown on agar overnight at a ⁵ mperature of 36°C and then taken colonies using 1µl sterile plastic loop and transferred into 184 mL of sterile distilled water and added to 5 µl lysozyme (5 mg / ml) followed by incubation at 56°C for 30 minutes.

Furthermore, the DNA was diluted to 100µl with buffer kit (QIAamp DNA mini kit 50). A total of 1µl DNA solution used for PCR amplification (Moenstein *et al.*, 2007).

Amplification

Primers used for detection of gene bla_{TEM} shown in Table 3. In the process anne⁵; (attachment) used 1µl DNA solution Hot Star Taq Qiagen Master Mix (Qiagen Nr. 203 445) and 10 pmol primer specific genes reach a final volume of ⁶ 5µl. Amplification followed denaturation process at 95°C for 15 min, 30 cycles of denaturation at 94°C for 30 seconds, annealing at 50°C for 30 seconds, extension at 72°C for 2 minutes followed by a final extension at of 72°C for 10 minutes (Moenstein *et al*¹⁰, 2007). The next step was reacting the PCR products by electrophoresis in 1% (w / v) agarose gel, then stained with ethidium-bromide and visualized using UV light. primers and DNA sequence was shown in Table 2. (Ahmed and Shimamoto, 2011). In genedetection bla_{TEM} positive results when there are 445 base paresized DNA band (Moenstein *et al.*, 2009).

RESULTS

Isolation of *Citrobacterfreundii* ³

The results of this study from the isolation and identification of bacteria from a total of 275 samples found positive *Citrobacterfreundii* were seven samples (2,55%) that the samples of rectal swab of Boriler from Wonokromo Surabaya (AW-1), Broiler from Keputran Surabaya (AK-5), Dairy Cattle from Bendul Merisi Surabaya (SB-3), Beef cattle from KrianSidoarjo (SPT.K-1), Milkfishes from Sidoarjo (IBS-1), (IBS-2), (IBS-4).

Table 3. Primer used for PCR (Moenstein *et al.*, 2007)

Primer	Sekunes (5 to 3)	Gene
TEM-F	TCGCCGCATACACTATTCTCAGAATGA	Bla _{TEM}
TEM-R	ACGCTCACCGCTCCAGATTTAT	Bla _{TEM}

Antibiotic Susceptibility Testing

Interpretation of the results obtained diameter of antibiotic zone that Broiler from Keputran (AK-5) and Wonokromo (AW-1) and Dairy cattle (SB-3) are resistant to ampicillin-sulbactam and tetracycline, but Beef cattle from Krian (SK-1), Milkfishes from Sidoarjo (IBS-1), (IBS-2), (IBS-4) are resistant to Amoxicillin and Ampicillin-Sulbactam. But on the other antibiotics are still visible intermediete and sensitive.

Detection of bla_{TEM} Using PCR

PCR result identified of bla_{TEM} in *Citrobacterfreundii* showed on seven samples of broiler from Keputran (AK-5), broilers from Wonokromo (AW-1), Dairy cattle from BendulMerisi (SB-3), Beef cattle from Krian (SK-1), Milkfishes from Sidoarjo (IBS-1), (IBS-2), (IBS-4). The results showed the band in 445 basepare (bp).




Figure 1. PCR assay to detect bla_{TEM} Lane 1= AK-5; Lane 2= AW-1; Lane 3= SB-3; M= Marker

DISCUSSION

The increasing prevalence of ESBL-producing Enterobacteriaceae very harmful to people's health, because the transmission can be derived from food products of animal origin or can be derived from animal faeces. In this study found 7 (2,55%) samples positive *Citrobacterfreundii* from the total 275 samples of rectal swab. Seven samples were broiler from Keputran (AK-5), broiler from Wonokromo (AW-1), Dairy cattle from BendulMerisi (SB-3), Beef cattle from Krian (SPT.K-1), Milkfishes from Sidoarjo (IBS-1), (IBS-2), (IBS-4).

Three samples were AK-5, AW-1, SB-3 found to be resistant to the antibiotic ampicillin-sulbactam, and tetracycline. And SPT.K-1, IBS-1, IBS-2, IBS-4 found to be resistant to the amoxicillin and ampicillin-sulbactam (Table 4). Then followed the PCR test to detect gene bla_{TEM} as agents ESBL and found positive for the band DNA of seven samples in 445 basepare (Figure 1).

In research done by Ahmed and Shimamoto (2011) that the gene coding for antimicrobial resistance bla_{TEM} as found as many as 23 isolates (20.5%) of 34 isolates of gram-negative bacteria in the case of mastitis in Egypt and in *Citrobacterfreundii* found bla_{TEM} genes as much as 3 isolates. Results of research conducted by Kao *et al.* (2010) found a gene in *Citrobacterfreundii* showed bla_{TEM} much as 2 isolates of 30 isolates sampled. *Extended Spectrum Beta Lactamase* (ESBL) derived from TEM-type enzyme encoded by the gene bla_{TEM}. The Bla_{TEM} genes are genes that cause antibiotic resistance in the plasmid of the most frequently detected in clinical populations (14) gram-negative microorganisms (Wilopo *et al.*, 2015). *Extended Spectrum Beta Lactamase* (ESBL) have the ability to transfer genes from one organism into another organism, so that the spread of resistance very easily occur between strains and even between species (Wilopo *et al.*, 2015). Plasmids are also responsible for coding genes that carry resistance genes to other drug classes (eg, classes (aminoglycoside). This makes selection of antibiotics against ESBL-producing organisms are very limited (Jacob and Munoz, 2005). Members of the family Enterobacteriaceae including *Citrobacterfreundii* often expressing plasmid-encoded β lactamase. Generally ESBL genes derived from TEM-1, TEM-2 or SHV-1 experienced mutase and change the configuration of amino acids around the active site of beta-lactamase. This situation makes the spectrum beta-lactam antibiotics susceptible to hydrolysis by this enzyme (Bush and Jacoby, 2010).

Conclusion

The conclusion of this study shows that seven samples positive *Citrobacterfreundii* found in broiler chickens, dairy cattle, beef cattle and milkfishes in East Java resistance to antibiotics. And *Citrobacterfreundii* showed as ESBL agent to detect bla_{TEM} gene using PCR with 445 basepare size.

Acknowledgement

11 Supported by Ministry of Research, Technology and Higher Education Indonesia on research of *Penelitian Terapan Unggulan Terpadu* fiscal year 2017.

REFERENCES

- Adelowo, O.O and Fagade, O.E 2010. Phylogenetic characterization, Antimicrobial Susceptibilities, And Mechanisms Of Resistance In Bacteriaisolates From A PoultryWaste-Polluted River, Southwestern Nigeria. *Turk J Biol*, 36. 37-45.
- Ahmed, A.M and Shimamoto, T 2011. Molecular characterization Of Antimicrobial Resistance In Gram-Negative Bacteria Isolated From Bovine Mastitis In Egypt. *MicrobiolImmunol*. 55 : 318-327.
- Bush, K., Jacoby, G, A 2010. Updated Functional Classification of Beta-lactamases. *Antimicrob. Agents Chemother*. 54 (3): 969-976.
- Clinicaland Laboratory Standarts Institute 2016. Performance Standards for Antimicrobial Susceptibility Testing. 26th edition. USA.
- Guilfoile, P and Alcamo, I.E. 2007 *Antibiotic-Resistant Bacteria*. E-book. Chelsea House Publisher, USA. 38
- Holt, J. G., Krieg, N. R., Sneath, P. H, A., Staley, JT and Williams, ST 1994. *Bergey's Manual of Determinative Bacteriology*. 9th Edn. Baltimore: Williams & Wilkins.

- Jacob, G, A and Munoz-Price LS 2005. The New Beta-lactamases. *N. Engl. J. Med.* 352 (4): 380-291.
- Kao, CC Liu, MF, Lin, CF, Huang YC, Liu PY, Chang, CW and Shi, ZY 2010. Antimicrobial Susceptibility and Multiplex PCR Screening of Genes From Amp C Isolates of *Enterobacter cloacae*, *Citrobacter freundii*, and *Serratia marcescens*. *Journal of Microbiology, Immunology and Infection.* 43 (3): 180-197.
- Lay, B.W 1994. Analisis Mikroba di Laboratorium. Jakarta: Rajawali.
- Mcfarland. 1907. Nephelometer: an instrument for media used for estimating the number of bacteria in suspensions used for calculating the opsonic index and for vaccines. *J Am Med Assoc.*, 1907; 14:1176-8.
- Maynard, C., Fairbrother, JM, Bekal, S., Sanschargin, F., Levesque, RC and Brousseau, R 2003. Antimicrobial Resistance Genes In enterotoxigenic *Escherichia coli* O149: K91 isolates Obtained Over a 23 Year Period From Pigs. *Antimicrob. Agents Chemother.* 47: 3124-3221.
- Monstein, HJ, Ostholm-Bulkhed, A., Nilsson, MV, M, Dombusch., K & Nilsson, LE 2007. Multiplex PCR amplification assay for the detection of blaSHV, blaTEM, and blaCTX-M genes in Enterobacteriaceae. *APMIS.* 115: 1400-1408
- Wilopo, BAP, Sudigdoadi, S., Sahiratmadja, E and Goddess, I, M, W 2015. Loop-Mediated Isothermal Amplification for Detection Gene Encoding bla TEM For Extended-Spectrum Beta-Lactamase in Enterobacteriaceae isolates. *MKB.* 47 (4).

ANTIMICROBIAL RESISTANCE DETECTION OF CITROBACTERFREUNDII ISOLATED FROM RECTAL SWAB IN EAST JAVA

ORIGINALITY REPORT

17%

SIMILARITY INDEX

15%

INTERNET SOURCES

10%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|----|
| 1 | www.fkh.unair.ac.id
Internet Source | 3% |
| 2 | mafiadoc.com
Internet Source | 2% |
| 3 | academicjournals.org
Internet Source | 2% |
| 4 | Ahmed, Ashraf M., Toshi Shimamoto, and Tadashi Shimamoto. "Characterization of integrons and resistance genes in multidrug-resistant <i>Salmonella enterica</i> isolated from meat and dairy products in Egypt", <i>International Journal of Food Microbiology</i> , 2014.
Publication | 1% |
| 5 | H.-J. MONSTEIN. "Multiplex PCR amplification assay for the detection of <i>bla</i> ^{SHV} , <i>bla</i> ^{TEM} and <i>bla</i> ^{CTX-M} genes in <i>Enterobacteriaceae</i> ", <i>Apmis</i> , 12/2007
Publication | 1% |

6	studentsrepo.um.edu.my Internet Source	1 %
7	journalijdr.com Internet Source	1 %
8	pubmed.ncbi.nlm.nih.gov Internet Source	1 %
9	revistainfectio.org Internet Source	1 %
10	www.nature.com Internet Source	1 %
11	phcogj.com Internet Source	1 %
12	dare.uvu.vu.nl Internet Source	1 %
13	ADELOWO, Olawale Olufemi and FAGADE, Obasola Ezekiel. "Phylogenetic characterization, antimicrobial susceptibilities, and mechanisms of resistance in bacteria isolates from a poultry waste-polluted river, southwestern Nigeria", TUBITAK, 2012. Publication	<1 %
14	ir.uiowa.edu Internet Source	<1 %
15	biomedres.us Internet Source	<1 %

16 dairy-cattle.extension.org <1 %
Internet Source

17 krishikosh.egranth.ac.in <1 %
Internet Source

18 Christy Manyi-Loh, Sampson Mamphweli, Edson Meyer, Anthony Okoh. "Antibiotic Use in Agriculture and Its Consequential Resistance in Environmental Sources: Potential Public Health Implications", *Molecules*, 2018 <1 %
Publication

19 Prind Triajeng Pungkasanti, Saifur Rohman Cholil, B. Very Christioko. "PERANCANGAN SISTEM PEMILIHAN BUDIDAYA IKAN AIR TAWAR BERBASIS WEB", *Jurnal Pengembangan Rekayasa dan Teknologi*, 2019 <1 %
Publication

20 www.ncbi.nlm.nih.gov <1 %
Internet Source

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

ANTIMICROBIAL RESISTANCE DETECTION OF CITROBACTERFREUNDII ISOLATED FROM RECTAL SWAB IN EAST JAVA

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
