

Lampiran c1. Bukti korespondensi dengan editor Karil JIB No 1 pada usulan pertama

No	Permasalahan	Komentar Tim PAK DIKTI	Klarifikasi
1	<p>Karil no 1</p> <p>Judul Artikel: Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion</p> <p>(Penulis ke 1 dari 4 penulis) dan sebagai Corresponding Author</p> <p>International Journal of Drug Delivery Technology; Vol.7, Issues 1; Januari-Maret 2017; Hal: 42-44 ISSN: 0975 4415 doi:10.25258/ijddt.v7i1.8915 Impact Factor: 1.529 SJR 2018: 0,21; Q3: Coverage: 2011-ongoing H index: 7 Similarity Index (Turnitin): 3%</p>	<p>Tidak ada masalah. Dapat diterima sebagai syarat khusus</p>	<p>Tambahan Kronologi dan bukti korespondensi dengan editor jurnal terlampir</p>

Kronologi korespondensi dengan editor jurnal untuk Karil JIB No 1

No	Tanggal	Keterangan
1	16 November 2016 (email 1: editor)	Pengiriman artikel melalui web jurnal; <i>Manuscript Submission Successful</i>
2	26 November 2016 (email 2: editor)	Naskah dengan ID 11262016DDTB <i>accepted after minor modifications</i>
3	26 November 2016 (email 3: author)	Pengiriman artikel yang sudah direvisi melalui web jurnal ada bukti artikel yang telah direvisi
4	26 November 2016 (email 4: editor)	Pemberitahuan tidak ada biaya publikasi, dan diminta mengirimkan <i>Copyright transfer agreement</i>
5	26 November 2016 (email 5: author)	Pengiriman <i>Copyright transfer agreement</i> melalui web jurnal ada bukti Copyright transfer agreement yang dikirim
6	26 Januari 2017 (email 6: editor)	Konfirmasi dari editor artikel akan segera di publish
7	21 Maret 2017 (email 7: editor)	Konfirmasi naskah telah terbit pada IJDDT, Vol7, Issue1, 2017

Manuscript Submission Successful

Dari: **IJDDT Submission System** (ijddtjournal@gmail.com)

Kepada: **era_ffua@yahoo.co.id**

Tanggal: **Jumat, 18 November 2016 15.44 GMT+7**

Dear Author,

Kindly find the details of your manuscript submission at the journal website.

1. Name of Corresponding Author

Tristiana Erawati

2. Email of Corresponding Author

era_ffua@yahoo.co.id

Institute of Affiliation of Corresponding Author

UNIVERSITAS AIRLANGGA

3. Phone/Mobile Number

+628121671607

4. Address

Pharmaceutics Department of Faculty of Pharmacy, Airlangga University, Kampus B UNAIR,
Jl Dharmawangsa Dalam
Surabaya 60286
Indonesia
[Map It](#)

5. Title of Manuscript

Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

6. Abstract

This study was aimed to determine the characteristic and tretinoin release in nanoemulsion using virgin coconut oil (VCO) as oil phase compared with emulsion. The characteristics of the tretinoin nanoemulsion (TN) were observed in terms of droplet morphology by Transmission Electron Microscopy (TEM) and droplet size by particle analyzer and light microscope and the pH value by pH meter. The release rate of tretinoin in nanoemulsion and emulsion was measured by Franz diffusion cell using cellophane membrane. Result of this research showed the droplet morphology of tretinoin nanoemulsion and emulsion were spherical. The droplet size of tretinoin nanoemulsion (72.57 ± 18.16 nm) was smaller than tretinoin emulsion (10.54 ± 0.61 m). The pH value of tretinoin nanoemulsion and tretinoin emulsion was 6.24 ± 0.01 and 6.21 ± 0.02 . In interval times 5 – 60 minutes the tretinoin release rate (flux) in nanoemulsion was 0.158 ± 0.016 $\mu\text{g}/\text{cm}^2/\text{minute}$ higher than in emulsion which was 0.048 ± 0.016 $\text{g}/\text{cm}^2/\text{minute}$. In interval times 60 – 180 minutes, tretinoin release rate (flux) in nanoemulsion was 0.046 ± 0.005 $\text{g}/\text{cm}^2/\text{minute}$ lower than in emulsion which was 0.090 ± 0.016 $\text{g}/\text{cm}^2/\text{minute}$. In interval times 180 – 720 minutes the tretinoin release rate (flux) in nanoemulsion was 0.025 ± 0.001 $\text{g}/\text{cm}^2/\text{minute}$ which had no significant different compared to in emulsion which was 0.022 ± 0.002 $\text{g}/\text{cm}^2/\text{minute}$. The statistical analysis of the tretinoin release rate value using independent T-test result was known that there were significant different between nanoemulsion (TN) and emulsion (TE). Conclusion: The droplet size of tretinoin nanoemulsion was below 100 nm, pH value 6.24 ± 0.01 and the tretinoin release rate in nanoemulsion using VCO was higher than in emulsion.

7. Keywords (2-10)

Characterization, Nanoemulsion, Release rate (Flux), Virgi Coconut Oil (VCO), Tretinoin.

8. Complete manuscript in one file (word)

- [TRISTIANA-ERAWATI_UNIVERSITAS-AIRLANGGA_INDONESIA.doc](#)

Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

Dari: IJDDT Journal (ijddtjournal@gmail.com)

Kepada: era_ffua@yahoo.co.id

Tanggal: Sabtu, 26 November 2016 15:09 GMT+7

Dear author

Your manuscript number "I1262016DDTB" bearing with title "Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion" **has been provisionally accepted after minor modifications**. It may be published in the upcoming issue of the journal subject to deposition of Open Access Fee USD67.50.

The authors are required to use the **paypal link** available at the website under instructions area. Other modes are not acceptable.

Important: The Fee will be revised to USD 100 for manuscripts submitted after 1-Dec-16.

Note: The authors may **reply this mail** with modified manuscript containing minor typography errors if left coincidentally during submission. The errors must be highlighted clearly in **red color font**. The modifications are subject to approval of editorial board. Any typological/coincidental error left after this stage will not be considered. In case the modifications are not minor, then authors may be asked to withdraw and resubmit manuscript.

Important: After making a transaction, please confirm transaction by providing transaction details as a reply to this mail only.

1. Name of Corresponding Author	Tristiana Erawati
2. Email of Corresponding Author	era_ffua@yahoo.co.id
Institute of Affiliation of Corresponding Author	UNIVERSITAS AIRLANGGA
3. Phone/Mobile Number	+628121671607
4. Address	Pharmaceutics Department of Faculty of Pharmacy, Airlangga University, Kampus B UNAIR, Jl Dharmawangsa Dalam Surabaya 60286 Indonesia Map It
5. Title of Manuscript	Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion
6. Abstract	This study was aimed to determine the characteristic and tretinoin release in nanoemulsion using virgin coconut oil (VCO) as oil phase compared with emulsion. The characteristics of the tretinoin nanoemulsion (TN) were observed in terms of droplet morphology by Transmission Electron Microscopy (TEM) and droplet size by particle analyzer and light microscope and the pH value by pH meter. The release rate of tretinoin in nanoemulsion and emulsion was measured by Franz diffusion cell using cellophane membrane. Result of this research showed the droplet morphology of tretinoin nanoemulsion and emulsion were spherical. The droplet size of tretinoin nanoemulsion (72.57 ± 18.16 nm) was smaller than tretinoin emulsion (10.54 ± 0.61 μ m). The pH value of tretinoin nanoemulsion and tretinoin emulsion was 6.24 ± 0.01 and 6.21 ± 0.02 . In interval times 5 – 60 minutes the tretinoin release rate (flux) in nanoemulsion was 0.158 ± 0.016 μ g/cm ² /minute higher than in emulsion which was 0.048 ± 0.016 g/cm ² /minute. In interval times 60 – 180 minutes, tretinoin release rate (flux) in nanoemulsion was 0.046 ± 0.005 g/cm ² /minute lower than in emulsion which was 0.090 ± 0.016 g/cm ² /minute. In interval times 180 – 720 minutes the tretinoin release rate (flux) in nanoemulsion was 0.025 ± 0.001 g/cm ² /minute which had no significant different compared to in emulsion which was 0.022 ± 0.002 g/cm ² /minute. The statistical analysis of the tretinoin release rate value using independent T-test result was known that there were significant different between nanoemulsion (TN) and emulsion (TE). Conclusion: The droplet size of tretinoin nanoemulsion was below 100 nm, pH value 6.24 ± 0.01 and the tretinoin release rate in nanoemulsion using VCO was higher than in emulsion.
7. Keywords (2-10)	Characterization, Nanoemulsion, Release rate (Flux), Virgi Coconut Oil (VCO), Tretinoin.
8. Complete manuscript in one file (word)	TRISTIANA-ERAWATI_UNIVERSITAS-AIRLANGGA_INDONESIA.doc

--
Editor
International Journal of Drug Delivery Technology
ISSN: 0975-4415
website: www.ijddt.com

List of correction:

No	Name Of Paragraph	Line	Uncorrected text	Correction
1	Abstract	8 m μm
2	Key words	1, Virgi, Virgin
3	RESULT AND DISCUSSION	5 emulsion nanoemulsion ...
4	RESULT AND DISCUSSION	16 emulsion nanoemulsion ...

Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

Tristiana Erawati M.*, Retnowati, Amalia Wardatul F., Widji Soeratri

Pharmaceutics Department of Faculty of Pharmacy, Airlangga University,

Kampus B UNAIR, Jl Dharmawangsa Dalam Surabaya, 60286,

*Corresponding Author: era_ffua@yahoo.co.id or tristiana-e-m@ff.unair.ac.id

ABSTRACT

This study was aimed to determine the characteristic and tretinoin release in nanoemulsion using virgin coconut oil (VCO) as oil phase compared with emulsion. The characteristics of the tretinoin nanoemulsion (TN) were observed in terms of droplet morphology by Transmission Electron Microscopy (TEM) and droplet size by particle analyzer and light microscope and the pH value by pH meter. The release rate of tretinoin in nanoemulsion and emulsion was measured by Franz diffusion cell using cellophane membrane. Result of this research showed the droplet morphology of tretinoin nanoemulsion and emulsion were spherical. The droplet size of tretinoin nanoemulsion (72.57 ± 18.16 nm) was smaller than tretinoin emulsion (10.54 ± 0.61 μm). The pH value of tretinoin nanoemulsion and tretinoin emulsion was 6.24 ± 0.01 and 6.21 ± 0.02 . In interval times 5 – 60 minutes the tretinoin release rate (flux) in nanoemulsion was 0.158 ± 0.016 $\mu\text{g}/\text{cm}^2/\text{minute}$ higher than in emulsion which was 0.048 ± 0.016 $\text{g}/\text{cm}^2/\text{minute}$. In interval times 60 – 180 minutes, tretinoin release rate (flux) in nanoemulsion was 0.046 ± 0.005 $\text{g}/\text{cm}^2/\text{minute}$ lower than in emulsion which was 0.090 ± 0.016 $\text{g}/\text{cm}^2/\text{minute}$. In interval times 180 – 720 minutes the tretinoin release rate (flux) in nanoemulsion was 0.025 ± 0.001 $\text{g}/\text{cm}^2/\text{minute}$ which had no significant different compared to in emulsion which was 0.022 ± 0.002 $\text{g}/\text{cm}^2/\text{minute}$.

The statistical analysis of the tretinoin release rate value using independent T-test result was known that there were significant different between nanoemulsion (TN) and emulsion (TE). Conclusion: The droplet size of tretinoin nanoemulsion was below 100 nm, pH value 6.24 ± 0.01 and the tretinoin release rate in nanoemulsion using VCO was higher than in emulsion.

Key words: Characterization, Nanoemulsion, Release rate (Flux), **Virgin** Coconut Oil (VCO), Tretinoin.

INTRODUCTION

Virgin coconut oil (VCO) is one of vegetables oil that used as oil phase in nanoemulsion. In the previous study nanoemulsion using VCO produced droplet size smaller than nanoemulsion using corn oil and soybean oil.¹ Smaller droplet size caused rapid drug release that can increase penetration into the skin. Nanoemulsion also known can increase solubility of drug. In this research VCO used in nanoemulsion for delivery system of Tretinoin an antiaging,^{2, 3} it was practically insoluble in water. Tretinoin nanoemulsion using VCO was than characterized and tested on the tretinoin release compared with tretinoin in emulsion.

MATERIAL AND METHODS

Research Materials

Tretinoin (PT. Cortico Mulia), virgin coconut oil (VCO), Tween 80 (Sigma Aldrich), Span 80 (Sigma Aldrich), ethanol 96 % (E-Merck), NaH₂PO₄ (E-Merck) and Na₂HPO₄ (E-Merck), aquademineralisata (PT Brataco), Methanol (E-Merck)

Research Instruments

Stirrer plate (Dragon Lab MS-Pro), ultrasonic (Branson 3510), shaker machine (Wine shake), pH meter (Eutech Instruments pH 700), particle analyzer (Delsa Nano C), Franz diffusion cell with cellophane membranes, spectrophotometer (Shimadzu UV-1800), Transmission Electron Microscopy (TEM-type JOEL JEM-1400), light microscope.

Methods

Nanoemulsion and emulsion preparation:

Nanoemulsion and emulsion preparation was done by Erawati and Soeratri method.⁴ Formulas tretinoin nanoemulsion (TN) and tretinoin emulsion (TE) was presented in Table 1. The tretinoin nanoemulsion was characterized included; pH, droplet morphology by TEM-type JOEL JEM-1400, droplet size and polydispersity index by particle analyzer Delsa Nano C. The tretinoin emulsion characterized included; pH, droplet morphology and droplet size by light microscope.

Table 1. Formula of Tretinoin in Nanoemulsion (TN) and Emulsion (TE) using VCO as oil phase

Materials	Concentration (%)	
	TN	TE
Tretinoin	0.1	0.1
VCO	2.66	1,66
Span 80	1.92	14,17
Tween 80	18.66	6,40
Ethanol 96%	3.42	-
Phosphate buffer solution pH 6.0 ± 0.5	ad 100	ad 100

Release Study

A cellophane membrane was immersed in water for ± 12 hours. It was drained until no water is dripping before used, and then was mounted on the surface of the receptor compartment of Franz diffusion cell. Franz diffusion cell Receptor compartment was filled with phosphate buffer

medium pH 6.0 ± 0.2 up to full (temperature $32 \pm 2^\circ\text{C}$, at 100 rpm). Then, 2 ml of tretinoin nanoemulsion or tretinoin emulsion was inserted into the donor compartment. Samples (1 ml) were taken within a certain time interval, i.e. at 0, 5, 10, 15, 30, 45 minutes, and then 1, 1.5, 2, 3, 4, 6, 8, 10, 12 hours. Immediately after sampling medium was replaced with phosphate buffer pH 6.0 ± 0.2 with a volume of samples taken. Subsequently, samples were taken and were observed with UV-Vis spectrophotometer. Tretinoin concentration in the sample was calculated using the standard curve regression equation, then correction to the measured concentration using the Wurster equation.

RESULT AND DISCUSSION

The characteristics of tretinoin nanoemulsion (TN) and tretinoin emulsion (TE) included pH, droplet size and polydispersity index was presented in Table 2. The pH value of both nanoemulsion and emulsion tretinoin were in the range of pH skin, it was expected that would not cause irritation when used.

Table 2. Characteristics of Tretinoin Nanoemulsion (TN) and Tretinoin Emulsion (TE)

Characteristic	TN	TE
pH	6.29 ± 0.01	6.21 ± 0.015
Droplet size	85.53 ± 6.28 nm	11.70 ± 2.51 μm
Polydispersity Index	0.676 ± 0.05	-

Tretinoin **nanoemulsion** droplet size was about 85.53 ± 6.28 nm; it was smaller than tretinoin emulsion droplet size of 11.70 ± 2.51 μm (Table 2). Droplet morphology of tretinoin nanoemulsion by TEM and tretinoin emulsion by light microscope both was appeared spherical. The release profiles of tretinoin in nanoemulsion (TN) and in emulsion (TE) were presented in Figure 1 and the tretinoin release-rate (Flux) in nanoemulsion and emulsion was presented in

Table 3 and Figure 2. From the release profile it was known that there were three interval times that gave different release rate (Flux). In first five minutes, the amount of tretinoin release from nanoemulsion (TN) and emulsion (TE) was equal; it may be caused by tretinoin existing in the water phase. From 5 to 60 minutes tretinoin release rate (Flux) from nanoemulsion (TN) was $0.158 \pm 0.016 \mu\text{g}/\text{cm}^2/\text{minute}$, higher than in emulsion (TE) is $0.048 \pm 0.016 \text{ g}/\text{cm}^2/\text{minute}$. It can be caused by droplet size of tretinoin nanoemulsion was smaller than tretinoin emulsion so it more rapid to release. In the second interval times (from 60 to 180 minutes) tretinoin release rate (flux) from nanoemulsion (TN) is $0.046 \pm 0.005 \text{ g}/\text{cm}^2/\text{minute}$ lower than in emulsion (TE) is $0.090 \pm 0.016 \text{ g}/\text{cm}^2/\text{minute}$, it can be caused by concentration tretinoin in nanoemulsion became lower than in emulsion. In the third interval times (from 180 to 720 minutes) tretinoin release rate (flux) in nanoemulsion is $0.025 \pm 0.001 \text{ g}/\text{cm}^2/\text{minute}$ had no significant difference with in emulsion which was $0.022 \pm 0.002 \text{ g}/\text{cm}^2/\text{minute}$. The result of statistical analysis by independent T-test with degree of confidence 95% ($\alpha = 0.05$), known that release-rate (Flux) tretinoin in nanoemulsion was higher than in emulsion.




Figure 1. Release profile of tretinoin in nanoemulsion (TN) and in emulsion (TE)




Figure 2. Histogram release flux of tretinoin in nanoemulsion (TN) and in emulsion (TE) in interval times 5 – 60, 60 – 180 and 180 – 720 minutes

Table 3. Release Flux of Tretinoin Nanoemulsion (TN) and Tretinoin Emulsion (TE) in interval times 5 – 60, 60 – 180 and 180 – 720 minutes

Sample	Release Flux in interval times (minutes)		
	mean \pm SD ($\mu\text{g}/\text{cm}^2 \cdot \text{minute}$)		
	5 – 60	60 – 180	180 – 720
TN	0.158 \pm 0.016	0.046 \pm 0.005	0.025 \pm 0.001
TE	0.048 \pm 0.016	0.090 \pm 0.016	0.022 \pm 0.002

CONCLUSION:

The conclusion of this study is tretinoin nanoemulsion using virgin coconut oil (VCO) produced droplet sizes below 100 nm, the pH value 6.24 ± 0.01 and the value of tretinoin release rate in nanoemulsion was higher than in the emulsion.

ACKNOWLEDGEMENT:

The authors want to thank Indonesia Government who gives research fund through Airlangga University and Faculty of Pharmacy Airlangga University.

REFERENCES:

1. Erawati T. Hendradi E., Soeratri W., Praformulation Study Of *p*-Methoxycinnamic Acid (APMS) Nanoemulsion Using Vegetable Oils (Soybean Oil, Corn Oil, VCO), *Int. J Pharm. Pharm. Sci.*, Vol. 6, Issue 2, 2014, p 99-101
2. Kligman AM, Grove GL, Hirose R, *et al.*, Topical tretinoin for photo aged skin, *J Am Acad. Dermatology*; 15: 1986, p 836–859
3. Stefanaki C, Stratigos A, Katsambas A., Topical retinoids in the treatment of photoaging, *Journal of Cosmetic Dermatology*, July 2005, 4, 2005, p 130–134
4. Erawati T. and Soeratri W., Characterization and the Release Test of Anti-Aging Tretinoin in Nanoemulsion Using Olive Oil, *International conference on Medicines and Health Sciences*, 2016, Jember University, Indonesia

Re: Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

Dari: IJDDT Journal (ijddtjournal@gmail.com)

Kepada: era_ffua@yahoo.co.id

Tanggal: Sabtu, 26 November 2016 15.10 GMT+7

Dear Author,

There are no publication charges upto dec 31, 2016. You are required to submit copyright transfer agreement only

On Sat, Nov 26, 2016 at 1:39 PM, IJDDT Journal <ijddtjournal@gmail.com> wrote:

Dear author

Your manuscript number "11262016DDTB" bearing with title "Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion" has been provisionally accepted after minor modifications. It may be published in the upcoming issue of the journal subject to deposition of Open Access Fee USD67.50.

The authors are required to use the **paypal link** available at the website under instructions area. Other modes are not acceptable.

Important: The Fee will be revised to USD 100 for manuscripts submitted after 1-Dec-16.

Note: The authors may **reply this mail** with modified manuscript containing minor typographic errors if left coincidentally during submission. The errors must be highlighted clearly in **red color font**. The modifications are subject to approval of editorial board. Any typographical/coincidental error left after this stage will not be considered. In case the modifications are not minor, then authors may be asked to withdraw and resubmit manuscript.

Important: After making a transaction, please confirm transaction by providing transaction details **as a reply to this mail only**.

1. Name of Corresponding Author

Tristiana Erawati

2. Email of Corresponding Author

era_ffua@yahoo.co.id

Institute of Affiliation of Corresponding Author

UNIVERSITAS AIRLANGGA

3. Phone/Mobile Number

+628121671607

4. Address

Pharmaceutics Department of Faculty of Pharmacy, Airlangga University, Kampus B UNAIR,

Jl Dharmawangsa Dalam

Surabaya 60286

Indonesia

[Map It](#)

5. Title of Manuscript

Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

6. Abstract

This study was aimed to determine the characteristic and tretinoin release in nanoemulsion using virgin coconut oil (VCO) as oil phase compared with emulsion. The characteristics of the tretinoin nanoemulsion (TN) were observed in terms of droplet morphology by Transmission Electron Microscopy (TEM) and droplet size by particle analyzer and light microscope and the pH value by pH meter. The release rate of tretinoin in nanoemulsion and emulsion was measured by Franz diffusion cell using cellophane membrane. Result of this research showed the droplet morphology of tretinoin nanoemulsion and emulsion were spherical. The droplet size of tretinoin nanoemulsion (72.57 ± 18.16 nm) was smaller than tretinoin emulsion (10.54 ± 0.61 m). The pH value of tretinoin nanoemulsion and tretinoin emulsion was 6.24 ± 0.01 and 6.21 ± 0.02 . In interval times 5 – 60 minutes the tretinoin release rate (flux) in nanoemulsion was 0.158 ± 0.016 $\mu\text{g}/\text{cm}^2/\text{minute}$ higher than in emulsion which was 0.048 ± 0.016 $\text{g}/\text{cm}^2/\text{minute}$. In interval times 60 – 180 minutes, tretinoin release rate (flux) in nanoemulsion was 0.046 ± 0.005 $\text{g}/\text{cm}^2/\text{minute}$ lower than in emulsion which was 0.090 ± 0.016 $\text{g}/\text{cm}^2/\text{minute}$. In interval times 180 – 720 minutes the tretinoin release rate (flux) in nanoemulsion was 0.025 ± 0.001 $\text{g}/\text{cm}^2/\text{minute}$ which had no significant different compared to in emulsion which was 0.022 ± 0.002 $\text{g}/\text{cm}^2/\text{minute}$. The statistical analysis of the tretinoin release rate value using independent T-test result was known that there were significant different between nanoemulsion (TN) and emulsion (TE). Conclusion: The droplet size of tretinoin nanoemulsion was below 100 nm, pH value 6.24 ± 0.01 and the tretinoin release rate in nanoemulsion using VCO was higher than in emulsion.

7. Keywords (2-10)

Characterization, Nanoemulsion, Release rate (Flux), Virgi Coconut Oil (VCO), Tretinoin.

8. Complete manuscript in one file (word)

[TRISTIANA-ERAWATI_UNIVERSITAS-AIRLANGGA_INDONESIA.doc](#)

–
Editor
International Journal of Drug Delivery Technology
ISSN: 0975-4415
website: www.ijddt.com

--
Editor
International Journal of Drug Delivery Technology
ISSN: 0975-4415
website: www.ijddt.com

TRANSFER OF COPYRIGHT AGREEMENT

INTERNATIONAL JOURNAL OF DRUG DELIVERY TECHNOLOGY

Title: Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

Author(s) name(s): Tristiana Erawati M.*, Retnowati, Amalia Wardatul F., Widji Soeratri

Corresponding authors name address, affiliation and email: Tristiana Erawati M.,
Pharmaceutics Department of Faculty of Pharmacy, Airlangga University, Kampus B UNAIR,
Jl. Dharmawangsa Dalam Surabaya, 60286, era_ffua@yahoo.co.id or tristiana-e-m@ff.unair.ac.id

I warrant that the material / manuscript submitted to IJDDT is original and has not been and will be published, whole or in part, in print or electronic format, in any other journal.

Each author has participated sufficiently in the work to take public responsibility for appropriate portions of the content. The order of authorships on the byline is a joint decision of all the coauthors. In case the manuscript should be accepted for publication, I have the consent of each author to transfer and assign any and all right, title and interest, including copyright of the article referenced above.

I hereby assign and transfer copyright and all rights under it to the International Journal of Drug Delivery Technology.

Corresponding author: Tristiana Erawati M.

Title: Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion



Signature: -----

Re: Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

Dari: IJDDT Journal (ijddtjournal@gmail.com)
 Kepada: era_ffua@yahoo.co.id
 Tanggal: **Kamis, 26 Januari 2017 16:57 GMT+7**

Dear Author,
The manuscript will be published in upcoming edition.

On Sat, Nov 26, 2016 at 1:40 PM, IJDDT Journal <ijddtjournal@gmail.com> wrote:

Dear Author,
 There are no publication charges upto dec 31, 2016. You are required to submit copyright transfer agreement only

On Sat, Nov 26, 2016 at 1:39 PM, IJDDT Journal <ijddtjournal@gmail.com> wrote:

Dear author

Your manuscript number "11262016DDTB" bearing with title "Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion" has been provisionally accepted after minor modifications. It may be published in the upcoming issue of the journal subject to deposition of Open Access Fee USD67.50.

The authors are required to use the **paypal link** available at the website under instructions area. Other modes are not acceptable.

Important: The Fee will be revised to USD 100 for manuscripts submitted after 1-Dec-16.

Note: The authors may **reply this mail** with modified manuscript containing minor typographic errors if left coincidentally during submission. The errors must be highlighted clearly in **red color font**. The modifications are subject to approval of editorial board. Any typographical/coincidental error left after this stage will not be considered. In case the modifications are not minor, then authors may be asked to withdraw and resubmit manuscript.

Important: After making a transaction, please confirm transaction by providing transaction details as a reply to this mail only.

1. Name of Corresponding Author

Tristiana Erawati

2. Email of Corresponding Author

era_ffua@yahoo.co.id

Institute of Affiliation of Corresponding Author

UNIVERSITAS AIRLANGGA

3. Phone/Mobile Number

+628121671607

4. Address

Pharmaceutics Department of Faculty of Pharmacy, Airlangga University, Kampus B UNAIR,
 Jl Dharmawangsa Dalam
 Surabaya 60286
 Indonesia
[Map It](#)

5. Title of Manuscript

Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

6. Abstract

This study was aimed to determine the characteristic and tretinoin release in nanoemulsion using virgin coconut oil (VCO) as oil phase compared with emulsion. The characteristics of the tretinoin nanoemulsion (TN) were observed in terms of droplet morphology by Transmission Electron Microscopy (TEM) and droplet size by particle analyzer and light microscope and the pH value by pH meter. The release rate of tretinoin in nanoemulsion and emulsion was measured by Franz diffusion cell using cellophane membrane. Result of this research showed the droplet morphology of tretinoin nanoemulsion and emulsion were spherical. The droplet size of tretinoin nanoemulsion (72.57 ± 18.16 nm) was smaller than tretinoin emulsion (10.54 ± 0.61 m). The pH value of tretinoin nanoemulsion and tretinoin emulsion was 6.24 ± 0.01 and 6.21 ± 0.02 . In interval times 5 – 60 minutes the tretinoin release rate (flux) in nanoemulsion was 0.158 ± 0.016 µg/cm²/minute higher than in emulsion which was 0.048 ± 0.016 g/cm²/minute. In interval times 60 – 180 minutes, tretinoin release rate (flux) in nanoemulsion was 0.046 ± 0.005 g/cm²/minute lower than in emulsion which was 0.090 ± 0.016 g/cm²/minute. In interval times 180 – 720 minutes the tretinoin release rate (flux) in nanoemulsion was 0.025 ± 0.001 g/cm²/minute which had no significant different compared to in emulsion which was 0.022 ± 0.002 g/cm²/minute. The statistical analysis of the tretinoin release rate value using independent T-test result was known that there were significant different between nanoemulsion (TN) and emulsion (TE). Conclusion: The droplet size of tretinoin nanoemulsion was below 100 nm, pH value 6.24 ± 0.01 and the tretinoin release rate in nanoemulsion using VCO was higher than in emulsion.

7. Keywords (2-10)

Characterization, Nanoemulsion, Release rate (Flux), Virgi Coconut Oil (VCO), Tretinoin.

8. Complete manuscript in one file (word)

[TRISTIANA-ERAWATI_UNIVERSITAS-AIRLANGGA_INDONESIA.doc](#)

--
 Editor
 International Journal of Drug Delivery Technology
 ISSN: 0975-4415
 website: www.ijddt.com

--
 Editor
 International Journal of Drug Delivery Technology
 ISSN: 0975-4415
 website: www.ijddt.com

--
 Editor
 International Journal of Drug Delivery Technology
 ISSN: 0975-4415
 website: www.ijddt.com

 11262016DDTB_TRISTIANA-ERAWATI_UNIVERSITAS-AIRLANGGA_INDONESIA_revised.doc
 116.5kB

Re: Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

Dari: IJDDT Journal (ijddtjournal@gmail.com)
 Kepada: era_ffua@yahoo.co.id
 Tanggal: Selasa, 21 Maret 2017 13.29 GMT+7

Please find the attachment:

On Thu, Jan 26, 2017 at 3:27 PM, IJDDT Journal <ijddtjournal@gmail.com> wrote:

Dear Author,
 The manuscript will be published in upcoming edition.

On Sat, Nov 26, 2016 at 1:40 PM, IJDDT Journal <ijddtjournal@gmail.com> wrote:

Dear Author,
 There are no publication charges upto dec 31, 2016. You are required to submit copyright transfer agreement only

On Sat, Nov 26, 2016 at 1:39 PM, IJDDT Journal <ijddtjournal@gmail.com> wrote:

Dear author

Your manuscript number "11262016DDTB" bearing with title "Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion" has been provisionally accepted after minor modifications. It may be published in the upcoming issue of the journal subject to deposition of Open Access Fee USD67.50.

The authors are required to use the **paypal link** available at the website under instructions area. Other modes are not acceptable.

Important: The Fee will be revised to USD 100 for manuscripts submitted after 1-Dec-16.

Note: The authors may **reply this mail** with modified manuscript containing minor typography errors if left coincidentally during submission. The errors must be highlighted clearly in **red color font**. The modifications are subject to approval of editorial board. Any typographical/coincidental error left after this stage will not be considered. In case the modifications are not minor, then authors may be asked to withdraw and resubmit manuscript.

Important: After making a transaction, please confirm transaction by providing transaction details as a reply to this mail only.

1. Name of Corresponding Author

Tristiana Erawati

2. Email of Corresponding Author

era_ffua@yahoo.co.id

Institute of Affiliation of Corresponding Author

UNIVERSITAS AIRLANGGA

3. Phone/Mobile Number

+628121671607

4. Address

Pharmaceutics Department of Faculty of Pharmacy, Airlangga University, Kampus B UNAIR,
 Jl Dharmawangsa Dalam
 Surabaya 60286
 Indonesia
[Map It](#)

5. Title of Manuscript

Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

6. Abstract

This study was aimed to determine the characteristic and tretinoin release in nanoemulsion using virgin coconut oil (VCO) as oil phase compared with emulsion. The characteristics of the tretinoin nanoemulsion (TN) were observed in terms of droplet morphology by Transmission Electron Microscopy (TEM) and droplet size by particle analyzer and light microscope and the pH value by pH meter. The release rate of tretinoin in nanoemulsion and emulsion was measured by Franz diffusion cell using cellophane membrane. Result of this research showed the droplet morphology of tretinoin nanoemulsion and emulsion were spherical. The droplet size of tretinoin nanoemulsion (72.57 ± 18.16 nm) was smaller than tretinoin emulsion (10.54 ± 0.61 m). The pH value of tretinoin nanoemulsion and tretinoin emulsion was 6.24 ± 0.01 and 6.21 ± 0.02 . In interval times 5 – 60 minutes the tretinoin release rate (flux) in nanoemulsion was 0.158 ± 0.016 µg/cm²/minute higher than in emulsion which was 0.048 ± 0.016 g/cm²/minute. In interval times 60 – 180 minutes, tretinoin release rate (flux) in nanoemulsion was 0.046 ± 0.005 g/cm²/minute lower than in emulsion which was 0.090 ± 0.016 g/cm²/minute. In interval times 180 – 720 minutes the tretinoin release rate (flux) in nanoemulsion was 0.025 ± 0.001 g/cm²/minute which had no significant different compared to in emulsion which was 0.022 ± 0.002 g/cm²/minute. The statistical analysis of the tretinoin release rate value using independent T-test result was known that there were significant different between nanoemulsion (TN) and emulsion (TE). Conclusion: The droplet size of tretinoin nanoemulsion was below 100 nm, pH value 6.24 ± 0.01 and the tretinoin release rate in nanoemulsion using VCO was higher than in emulsion.

7. Keywords (2-10)

Characterization, Nanoemulsion, Release rate (Flux), Virgi Coconut Oil (VCO), Tretinoin.

8. Complete manuscript in one file (word)


[TRISTIANA-ERAWATI_UNIVERSITAS-AIRLANGGA_INDONESIA.doc](#)

--
 Editor
 International Journal of Drug Delivery Technology
 ISSN: 0975-4415
 website: www.ijddt.com

--
 Editor
 International Journal of Drug Delivery Technology
 ISSN: 0975-4415
 website: www.ijddt.com

--
 Editor
 International Journal of Drug Delivery Technology
 ISSN: 0975-4415
 website: www.ijddt.com

--
Editor
International Journal of Drug Delivery Technology
ISSN: 0975-4415
website: www.ijddt.com

 IJDDT, Vol7, Issue1, Article5.pdf
293.4kB

Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion

Tristiana Erawati M^{*}, Retnowati, Amalia Wardatul F, Widji Soeratri

Pharmaceutics Department of Faculty of Pharmacy, Airlangga University, Kampus B UNAIR, Jl Dharmawangsa Dalam Surabaya, 60286.

Received: 15th Nov, 2016; Revised: 21st Jan, 2017; Accepted: 3rd Feb, 2017; Available Online: 1st March, 2017

ABSTRACT

This study was aimed to determine the characteristic and tretinoin release in nanoemulsion using virgin coconut oil (VCO) as oil phase compared with emulsion. The characteristics of the tretinoin nanoemulsion (TN) were observed in terms of droplet morphology by Transmission Electron Microscopy (TEM) and droplet size by particle analyzer and light microscope and the pH value by pH meter. The release rate of tretinoin in nanoemulsion and emulsion was measured by Franz diffusion cell using cellophane membrane. Result of this research showed the droplet morphology of tretinoin nanoemulsion and emulsion were spherical. The droplet size of tretinoin nanoemulsion (72.57 ± 18.16 nm) was smaller than tretinoin emulsion (10.54 ± 0.61 μ m). The pH value of tretinoin nanoemulsion and tretinoin emulsion was 6.24 ± 0.01 and 6.21 ± 0.02 . In interval times 5 – 60 minutes the tretinoin release rate (flux) in nanoemulsion was 0.158 ± 0.016 μ g/cm²/minute higher than in emulsion which was 0.048 ± 0.016 g/cm²/minute. In interval times 60 – 180 minutes, tretinoin release rate (flux) in nanoemulsion was 0.046 ± 0.005 g/cm²/minute lower than in emulsion which was 0.090 ± 0.016 g/cm²/minute. In interval times 180 – 720 minutes the tretinoin release rate (flux) in nanoemulsion was 0.025 ± 0.001 g/cm²/minute which had no significant different compared to in emulsion which was 0.022 ± 0.002 g/cm²/minute. The statistical analysis of the tretinoin release rate value using independent T-test result was known that there were significant different between nanoemulsion (TN) and emulsion (TE). Conclusion: The droplet size of tretinoin nanoemulsion was below 100 nm, pH value 6.24 ± 0.01 and the tretinoin release rate in nanoemulsion using VCO was higher than in emulsion.

Keywords: Characterization, Nanoemulsion, Release rate (Flux), Virgin Coconut Oil (VCO), Tretinoin.

INTRODUCTION

Virgin coconut oil (VCO) is one of vegetables oil that used as oil phase in nanoemulsion. In the previous study nanoemulsion using VCO produced droplet size smaller than nanoemulsion using corn oil and soybean oil.¹ Smaller droplet size caused rapid drug release that can increase penetration into the skin. Nanoemulsion also known can increase solubility of drug. In this research VCO used in nanoemulsion for delivery system of Tretinoin an antiaging,^{2, 3} it was practically insoluble in water. Tretinoin nanoemulsion using VCO was than characterized and tested on the tretinoin release compared with tretinoin in emulsion.

MATERIAL AND METHODS

Research Materials

Tretinoin (PT. Cortico Mulia), virgin coconut oil (VCO), Tween 80 (Sigma Aldrich), Span 80 (Sigma Aldrich), ethanol 96 % (E-Merck), NaH₂PO₄ (E-Merck) and Na₂HPO₄ (E-Merck), aquademineralisata (PT Brataco), Methanol (E-Merck)

Research Instruments

Stirrer plate (Dragon Lab MS-Pro), ultrasonic (Branson 3510), shaker machine (Wine shake), pH meter (Eutech Instruments pH 700), particle analyzer (Delsa Nano C), Franz diffusion cell with cellophane membranes,

spectrophotometer (Shimadzu UV-1800), Transmission Electron Microscopy (TEM-type JOEL JEM-1400), light microscope.

Methods

Nanoemulsion and emulsion preparation:

Nanoemulsion and emulsion preparation was done by Erawati and Soeratri method.⁴ Formulas tretinoin nanoemulsion (TN) and tretinoin emulsion (TE) was presented in Table 1. The tretinoin nanoemulsion was characterized included; pH, droplet morphology by TEM-type JOEL JEM-1400, droplet size and polydispersity index by particle analyzer Delsa Nano C. The tretinoin emulsion characterized included; pH, droplet morphology and droplet size by light microscope.

Release Study

A cellophane membrane was immersed in water for ± 12 hours. It was drained until no water is dripping before used, and then was mounted on the surface of the receptor compartment of Franz diffusion cell. Franz diffusion cell Receptor compartment was filled with phosphate buffer medium pH 6.0 ± 0.2 up to full (temperature $32 \pm 2^\circ\text{C}$, at 100 rpm). Then, 2 ml of tretinoin nanoemulsion or tretinoin emulsion was inserted into the donor compartment. Samples (1 ml) were taken within a certain time interval, i.e. at 0, 5, 10, 15, 30, 45 minutes, and then 1, 1.5, 2, 3, 4, 6, 8, 10, 12 hours. Immediately after

*Author for Correspondence: era_ffua@yahoo.co.id




Figure 1: Release profile of tretinoin in nanoemulsion (TN) and in emulsion (TE).




Figure 2: Histogram release flux of tretinoin in nanoemulsion (TN) and in emulsion (TE) in interval times 5 – 60, 60 – 180 and 180 – 720 minutes.

Table 1: Formula of Tretinoin in Nanoemulsion (TN) and Emulsion (TE) using VCO as oil phase.

Materials	Concentration (%)	
	TN	TE
Tretinoin	0.1	0.1
VCO	2.66	1,66
Span 80	1.92	14,17
Tween 80	18.66	6,40
Ethanol 96%	3.42	-
Phosphate buffer solution pH 6.0 ± 0.5	ad 100	ad 100

sampling medium was replaced with phosphate buffer pH 6.0 ± 0.2 with a volume of samples taken. Subsequently, samples were taken and were observed with UV-Vis spectrophotometer. Tretinoin concentration in the sample was calculated using the standard curve regression equation, then correction to the measured concentration using the Wurster equation.

Table 2: Characteristics of Tretinoin Nanoemulsion (TN) and Tretinoin Emulsion (TE).

Characteristic	TN	TE
pH	6.29 ± 0.01	6.21 ± 0.015
Droplet size	85.53 ± 6.28 nm	11.70 ± 2.51 µm
Polidispersity Index	0.676 ± 0.05	-

RESULT AND DISCUSSION

The characteristics of tretinoin nanoemulsion (TN) and tretinoin emulsion (TE) included pH, droplet size and polidispersity index was presented in Table 2. The pH value of both nanoemulsion and emulsion tretinoin were in the range of pH skin, it was expected that would not cause irritation when used.

Tretinoin nanoemulsion droplet size was about 85.53 ± 6.28 nm; it was smaller than tretioin emulsion droplet size of 11.70 ± 2.51 µm (Table 2). Droplet morphology of

Table 3: Release Flux of Tretinoin Nanoemulsion (TN) and Tretinoin Emulsion (TE) in interval times 5 – 60, 60 – 180 and 180 – 720 minutes.

Sample	Release Flux in interval times (minutes)		
	mean \pm SD ($\mu\text{g}/\text{cm}^2\cdot\text{minute}$)		
	5 – 60	60 – 180	180 – 720
TN	0.158 0.016	\pm 0.046 \pm 0.005	0.025 \pm 0.001
TE	0.048 0.016	\pm 0.090 0.016	\pm 0.022 \pm 0.002

tretinoin nanoemulsion by TEM and tretinoin emulsion by light microscope both was appeared spherical. The release profiles of tretinoin in nanoemulsion (TN) and in emulsion (TE) were presented in Figure 1 and the tretinoin release-rate (Flux) in nanoemulsion and emulsion was presented in Table 3 and Figure 2. From the release profile it was known that there were three interval times that gave different release rate (Flux). In first five minutes, the amount of tretinoin release from nanoemulsion (TN) and emulsion (TE) was equal; it may be caused by tretinoin existing in the water phase. From 5 to 60 minutes tretinoin release rate (Flux) from nanoemulsion (TN) was $0.158 \pm 0.016 \mu\text{g}/\text{cm}^2/\text{minute}$, higher than in emulsion (TE) is $0.048 \pm 0.016 \text{g}/\text{cm}^2/\text{minute}$. It can caused by droplet size of tretinoin nanoemulsion was smaller than tretinoin emulsion so it more rapid to release. In the second interval times (from 60 to 180 minutes) tretinoin release rate (flux) from nanoemulsion (TN) is $0.046 \pm 0.005 \text{g}/\text{cm}^2/\text{minute}$ lower than in emulsion (TE) is $0.090 \pm 0.016 \text{g}/\text{cm}^2/\text{minute}$, it can caused by concentration tretinoin in nanoemulsion became lower than in emulsion. In the third interval times (from 180 to 720 minutes) tretinoin release rate (flux) in nanoemulsion is $0.025 \pm 0.001 \text{g}/\text{cm}^2/\text{minute}$ had no

significant different with in emulsion which was $0.022 \pm 0.002 \text{g}/\text{cm}^2/\text{minute}$. The result of statistical analysis by independent T-test with degree of confident 95% ($\alpha = 0.05$), known that release-rate (Flux) tretinoin in nanoemulsion was higher than in emulsion.

CONCLUSION

The conclusion of this study is tretinoin nanoemulsion using virgin coconut oil (VCO) produced droplet sizes below 100 nm, the pH value 6.24 ± 0.01 and the value of tretinoin release rate in nanoemulsion was higher than in the emulsion.

ACKNOWLEDGEMENT

The authors want to thank Indonesia Government who gives research fund through Airlangga University and Faculty of Pharmacy Airlangga University.

REFERENCES

1. Erawati T. Hendradi E., Soeratri W., Praformulation Study Of *p*-Methoxycinnamic Acid (APMS) Nanoemulsion Using Vegetable Oils (Soybean Oil, Corn Oil, VCO), *Int. J Pharm. Pharm. Sci.*, Vol. 6, Issue 2, 2014, p 99-101.
2. Kligman AM, Grove GL, Hirose R, *et al.*, Topical tretinoin for photo aged skin, *J Am Acad. Dermatology*; 15: 1986, p 836–859.
3. Stefanaki C, Stratigos A, Katsambas A., Topical retinoids in the treatment of photoaging, *Journal of Cosmetic Dermatology*, July 2005, 4, 2005, p 130–134.
4. Erawati T. and Soeratri W., Characterization and the Release Test of Anti-Aging Tretinoin in Nanoemulsion Using Olive Oil, *International conference on Medicines and Health Sciences*, 2016, Jember University, Indonesia.