


Journal of Applied Environmental and Biological Sciences (JAEBS)



An International Peer-reviewed journal

Number of issues per year: 12

ISSN (Print): 2090-4274

ISSN (Online): 2090-4215

[Home](#)[Journals](#)[Contact Us](#)

TEXTROAD JOURNALS

Journal of Applied Environmental and Biological Sciences



J. Appl. Environ. Biol. Sci., Vol.8 No. 1: pp. 1-191, Year 2018

Journal of Applied Environmental and Biological Sciences (JAEBS) Monthly Publication



Number of issues per year: 12

ISSN: 2090-4274 (Print)

ISSN: 2090-4215 (Online)

Journal of Applied Environmental and Biological Sciences (JAEBS) is a peer-reviewed, open access international scientific journal dedicated for rapid publication of high quality original research articles as well as review articles in the all areas of Applied Environmental and Biological Sciences.

Scope

Journal of Applied Environmental and Biological Sciences (JAEBS) is devoted to the monthly publication of research papers of outstanding significance in the all fields of environmental sciences, environmental engineering, environmental Pollution, green chemistry, environmentally friendly synthetic pathways, alternatively fuels, environmental analytical chemistry, biomolecular tools and tracers, water and soil, environmental [management, economics, humanities], Mathematics, multidisciplinary aspects such as Business Management, Organizational Behavior, all areas of biological sciences, including cell biology, developmental biology, structural biology, microbiology, molecular biology & genetics, biochemistry, biotechnology, biodiversity, ecology, marine biology, plant biology, bioinformatics, toxicology, developmental biology, structural biology, microbiology, molecular biology & genetics, biotechnology, biodiversity and related fields. The journal presents the latest developments in the fields of environmental social marketing, environmental journalism, environmental education, sustainability education, environmental interpretation, and environmental health communication.

Editorial Board

Editor -in-Chief

William Ebomoyi

Ph.D., Professor, Department of Health Studies, College of Health Sciences, Chicago State University, **USA**.

E-mail: editor@textroad.com

Associate Editors

Prof. Dr. Sanaa T. El-Sayed

Ex Head of Biochemistry Department, Professor of Biochemistry, Genetic Engineering & Biotechnology Division, National Research Centre, **Egypt**

Saeid Chekani Azar

PhD of Veterinary Physiology; Faculty of Veterinary, Department of Physiology, Ataturk University, Erzurum 25010, **Turkey**.

Prof. Dr. Sarwoko Mangkoedihardjo

Professor, Professional Engineer of Indonesian Society of Sanitary and Environmental Engineers, **Indonesia**

Prof. Dr. Ashraf Latif Tadross

Head of Astronomy Department, Professor of Star Clusters and Galactic Structure, National Research Institute of Astronomy & Geophysics (NRIAG), 11421 Helwan, Cairo, **Egypt**.

Dr. Chandrasekar Raman

Research Associate, Department of Biochemistry & Molecular Biophysics, Biotechnology Core Facility, 238, Burt Hall, Kansas University, Manhattan 66506, KS, **USA**.

Dr. YUBAO CUI

Associate Professor, Department of Laboratory Medicine, Yancheng Health Vocational & Technical College, Jiangsu Province, P. R. **China**

Dr. Muhammad Altaf Khan

Department of Mathematics, Abdul Wali Khan University Mardan **Pakistan**

Dr. Fahrettin Tilki

Assoc. Professor, Artvin Coruh University, Faculty of Forestry, Department of Forest Science, Artvin, **TURKEY**.

Dr. Ibtisam abd el ghany hammad

Associate Professor of Genetics, Faculty of Science, Helwan University. **Egypt**.

Dr. Charalambos Tsekeris

Department of Psychology, Panteion University of Social and Political Sciences, Athens, **Greece**.

Dr. Elsayed E. Hafez

Associate Professor, Molecular Biology, Plant Molecular Pathology & Arid Lands Institute, **Egypt**.

Dr. Naushad Mamode Khan

University of Mauritius, Reduit, **Mauritius**.

Mirza Hasanuzzaman

Department of Agronomy, Faculty of Agriculture, Sher-e-Bangla Agricultural University, Dhaka-1207, **Bangladesh**.

Dr. Hala Ahmed Hafez Kandil

Professor Researcher, National Research Centre, Plant Nutrition Dept. El-Bhouth St. Dokki, Giza, **Egypt**.

Dr. Yule Yue Wang

Biotechnology and Medicinal Biochemistry, Division of Life Science, The Hong Kong University of Science & Technology, **Hong Kong**.

Dr. Aziza Sharaby

Professor of Entomology. Plant Protection Department, National Research Center. Cairo, **Egypt**.

Dr. Sulaiman

Assistant Professor, Department of Biochemistry, Abdul wali Khan University Mardan, Khyber Pakhtunkhwa, **Pakistan**.

Editors

Maulin P Shah

PhD-Microbiology, Chief Scientist & Head Industrial Waste Water Research Laboratory, Division of Applied & Environmental Microbiology, Enviro Technology Limited, Ankleshwar-393002, Gujarat, **India**

Dr. Josphert N. Kimatu

Department of Biological Sciences. South Eastern University College, **Kenya**.

Dr. Mukesh Kumar Meena

Assistant Professor (Crop Physiology), Department of Crop Physiology, University of Agricultural Sciences, Raichur-584104, Karnataka , **India**

Jehngir Khan

Lecturer in Zoology Department, Abdul Wali Khan University Mardan (AWKUM), Buner Campus, Buner, Khyber Pakhtunkhwa, **Pakistan**.

Syed Muhammad Nurulain

Medical Research Specialist, FMHS, UAE University, **Emirates**

Dr. Ayman Batisha

Environment and Climate Research Institute, National Water Research Center, Cairo, **Egypt**.

Dr. Hakeem Ullah

Assistant Professor, Department of Mathematics Abdul Wali Khan University Mardan **Pakistan**.

DR. DATTA ASARAM DHALE

Assistant Professor, Post Graduate Department of Botany, Ghogrey Science College, Dhule, Maharashtra State, **India**.

Dr. Muhammad Ismail Mohmand

Tutor/Administrator in the Excellence Training Den College in Newcastle, **United Kingdom**

Prof. Dr. Valdenir José Belinelo

Department of Health Sciences and Postgraduate Program in Tropical Agriculture, Federal University of Espirito Santo (UFES), São Mateus, ES, **Brazil**.

Siva Sankar. R

Department of Ecology and Environmental Sciences, School of Life Sciences, Pondicherry University, **India**.

Table of Contents, January 2018

Siti Farida Noorlayla

Family Support with Coping Mechanism of Elderly Hypertension in Elderly Integrated Health Service Post in Kediri

J. Appl. Environ. Biol. Sci. 2018 8(1): 1-5. [[Abstract](#)] [[Full-Text PDF](#)] [[Full-Text XML](#)]

Nahal Boudierba Nora, Belmekki Anouar, Ghassel Aicha, Kadi Hamid

Antioxidant Activities and Antifungal Screening of *Salvia officinalis* Flavonoids

J. Appl. Environ. Biol. Sci. 2018 8(1): 6-13. [[Abstract](#)] [[Full-Text PDF](#)] [[Full-Text XML](#)]

Sana Yasin, Tariq Ali, Umar Draz, Low Tan Jung, M. Ayaz

Formal Analysis of Malicious Attacks of Ad Hoc Distance Vector Protocol in Mobile Ad-hoc and Sensor Networks (MAHSNs)

J. Appl. Environ. Biol. Sci. 2018 8(1): 14-22. [[Abstract](#)] [[Full-Text PDF](#)] [[Full-Text XML](#)]

Nurdina

Analysis of Lawrence Green Theory Concerning Pregnancy Sexuality Behavior to Sexual Behavior Change in Primigravida in Kediri

J. Appl. Environ. Biol. Sci. 2018 8(1): 23-28. [[Abstract](#)] [[Full-Text PDF](#)] [[Full-Text XML](#)]

Sidra Arooj, Syed Asif Ali Naqvi, Muhammad Sohail Amjad Makhdum

Impact of Environmental Degradation on Wheat Production in Pakistan

J. Appl. Environ. Biol. Sci. 2018 8(1): 29-34. [[Abstract](#)] [[Full-Text PDF](#)] [[Full-Text XML](#)]

Muhammad Naeem Shahid, Dr. Muhammad Ayub Siddiqui, Muhammad Haris Qureshi, Faraz Ahmad

Corporate Governance and its Impact on Firm's Performance: Evidence from Cement Industry of Pakistan

J. Appl. Environ. Biol. Sci. 2018 8(1): 35-41. [[Abstract](#)] [[Full-Text PDF](#)] [[Full-Text XML](#)]

Sentot Imam Suprpto

Post Accreditation Performance on Service Quality of Outpatient in Gambiran Hospital

J. Appl. Environ. Biol. Sci. 2018 8(1): 42-47. [[Abstract](#)] [[Full-Text PDF](#)] [[Full-Text XML](#)]

Sumaira Aslam, Muhammad Naeem Shahid, Muhammad Haris Qureshi, Dr Amina Muzaffar Qureshi

Investigating Innovativeness and Emotional Intelligence as Mediator to Explore Entrepreneurial Marketing Strategy Focused on Firm Performance: A Case in Pakistan

J. Appl. Environ. Biol. Sci. 2018 8(1): 48-60. [[Abstract](#)] [[Full-Text PDF](#)] [[Full-Text XML](#)]

Ghulam Murtaza Bodla, Muhammad Tahir khaliy

Treatment of Mental Health Issues through Positive Psychology Interventions

J. Appl. Environ. Biol. Sci. 2018 8(1): 61-64. [[Abstract](#)] [[Full-Text PDF](#)] [[Full-Text XML](#)]

Widya Paramita Lokapinasari, Sri Hidanah, Suharsono, Anisah Fathinah, Arrifah Ratna Dewi, Anita Dwi Andriani, Balqis Karimah, Tri Nurhajati, Koesnoto Soepranianondo, Mirni Lamid

Potency of Probiotics on HDL, LDL, Cholesterol and Total Protein of Egg's Quail (*Coturnix coturnix japonica*)

J. Appl. Environ. Biol. Sci. 2018 8(1): 65-69. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Muhammad Nisar, Mussawar Shah, Asadullah, Dilkash Sapna, Anas Baryal, Zia Ur Rahman

Land Ownership Disputes Due to Land Commercialization and Its Effect on Social Structure in Tehsil Dargai District Malakand

J. Appl. Environ. Biol. Sci. 2018 8(1): 70-74. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Dr. Muhammad Shabbir Ch., Muhammad Arqam Mushtaq, Muhammad Idrees, Muhaammad Atif

Socio-Economic and Cultural Determinants of Criminal Youth Behavior: A Case of Faisalabad City in Punjab, Pakistan

J. Appl. Environ. Biol. Sci. 2018 8(1): 75-82. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Humaira Gul, Seema Gul, Husna, Aqib Sayyed, Naeema Begum, Sheheryar and Zahida Parveen

Interactive Role of Potassium Nitrate and Sodium Chloride on Growth and Biochemical Constituents of Brassica Junceal.

J. Appl. Environ. Biol. Sci. 2018 8(1): 83-95. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Muhammad Nisar, Dilkash Sapna, Anas Baryal, Ejaz Ahmad, Zia Ur Rahman

Effects of Mothers Employment on Children Socialization (A Case Study of District Charsadda)

J. Appl. Environ. Biol. Sci. 2018 8(1): 96-101. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Yung Savitri and Sarwoko Mangkoedihardjo

Technical Evaluation of Residential Wastewater Treatment Plant in West Lombok Regency

J. Appl. Environ. Biol. Sci. 2018 8(1): 102-105. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Uzma Niaz, Zunera Jamil, Dr. Muhammad Shabbir Ch., Shazia Nasim

Socio-Economic and Cultural Problems of Brick Kiln's Working Families: (A Case Study of District Faisalabad)

J. Appl. Environ. Biol. Sci. 2018 8(1): 106-111. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Itbar Khan, Azhar Mehmood, Nabi Bux Jumani

The relationship of Instructional Beliefs of Teacher Educators with their Classroom Practices in Khyber Pakhtunkhwa

J. Appl. Environ. Biol. Sci. 2018 8(1): 112-119. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Erma Retnaningtyas

The Role of Midwife on the Use of Poedji Rochjati Score Card with Knowledge of Cadres in Early Detection of Pregnant Women Risk in Working Area Sidomulyo Health Center

J. Appl. Environ. Biol. Sci. 2018 8(1): 120-124. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Lailan Ni'mah, Fidelis Boy Manurung, Eka Pramita

Lightweight Concrete Production by Gypsum from Waste Materials of Clamshell and Eggshell

J. Appl. Environ. Biol. Sci. 2018 8(1): 125-133. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Fayyaz Ahmed Anjum, Dr Iffat Batool, Muhammad Umar Fayyaz

Internal Consistency Reliability of Modified Hand Test

J. Appl. Environ. Biol. Sci. 2018 8(1): 134-143. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Rabia Safdar, M. Imran, N. Sadiq, Farooq Ahmad

Unsteady Rotational Flow of a Burgers' Fluid through a Pipe with Non-Integer Order Fractional Derivatives

J. Appl. Environ. Biol. Sci. 2018 8(1): 144-156. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

N. Sadiq, M. Imran, Rabia Safdar, Hassan Waqas

Rotational Flow of Second Grade Fluid with Caputo-Fabrizio Fractional Derivative in an Annulus

J. Appl. Environ. Biol. Sci. 2018 8(1): 157-168. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Danial Habib, M. Anwar Kamal, Farooq Ahmad, Sajjad Hussain

Partial Slip and Buoyancy Effects on MHD Flow Through Porous Medium Adjacent to Porous Accelerated Sheet in Presence of Heat Radiation

J. Appl. Environ. Biol. Sci. 2018 8(1): 169-174. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Muhammad Nisar, Dilkash Sapna, Anas Baryal, Saima Gul, Zia Ur Rahman

Effects of Disabled Children on Their Families (A Case Study of District Charsadda)

J. Appl. Environ. Biol. Sci. 2018 8(1): 175-179. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Muhammad Nisar, Anas Baryal, Dilkash Sapna, Zia Ur Rahman

People's Perception Regarding Jirga in Pakhtun Society

J. Appl. Environ. Biol. Sci. 2018 8(1): 180-183. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Muhammad Amjid, Shazia Zamir

Comparison of Leadership Attributes Among Students at Cadet and Public Sectors Colleges

J. Appl. Environ. Biol. Sci. 2018 8(1): 184-191. [\[Abstract\]](#) [\[Full-Text PDF\]](#) [\[Full-Text XML\]](#)

Potency of Probiotics on HDL, LDL, Cholesterol and Total Protein of Egg's Quail (*Coturnix coturnix japonica*)

Widya Paramita Lokapirnasari^{1*}, Sri Hidanah¹, Suharsono², Anisah Fathinah³, Arrifah Ratna Dewi³, Anita Dwi Andriani², Balqis Karimah³, Tri Nurhajati¹, Koesnoto Soepranianondo¹, Mirni Lamid^{1,4,5}

¹Department of Animal Husbandry Faculty of Veterinary Medicine, Airlangga University, Mulyorejo Street, Surabaya 60115, Indonesia,

²Department of Anatomy Veteriner, Faculty of Veterinary Medicine, Airlangga University, Mulyorejo Street, Surabaya 60115, Indonesia

³Agribusiness Veteriner Faculty of Veterinary Medicine, Airlangga University, Mulyorejo Street, Surabaya 60115, Indonesia

⁴Faculty of Fisheries and Marine, Airlangga University, Surabaya, Indonesia

⁵Proteomics Laboratory, Institute of Tropical Disease, Airlangga University, Surabaya-Indonesia

Received: September 12, 2017

Accepted: November 28, 2017

ABSTRACT

The purpose of this research was to know the potency of probiotic supplementation to content of HDL, LDL, Cholesterol and total protein of egg's quail *Coturnix coturnix japonica*. This research consist of five treatments, each treatment consisted of four replications and each replication consisted by ten quail. The treatment were T0: control (without probiotic), T1: 0.05 gram probiotic/ kg feed, T2: 0.1 gram probiotic/kg feed, T3:0.025 gram probiotic/L drinking water and T4: 0.05 gram probiotic/L drinking water. The probiotic consisted of combine *Lactobacillus casei* and *Lactobacillus rhamnosus* of 1.2×10^8 CFU/gram. The results showed that the probiotic supplementation both in feed and water gave a significant impact to reduce of Low Density Lipoprotein (LDL) and cholesterol, improve of High Density Lipoprotein (HDL) level. Moreover, the quails feed with free antibiotic feed with free antibiotic feed + 0.025 gram probiotic/L drinking water and 0.1 gram probiotic/kg feed could reduce ($P<0.05$) egg cholesterol, LDL and could improve HDL and the total proteins. It can be concluded that using probiotic contain *Lactobacillus casei* and *Lactobacillus rhamnosus* could improve egg quality by reduced cholesterol, LDL and increase HDL and total protein of egg's quail *Coturnix coturnix japonica*.

KEYWORDS: Probiotic, HDL, LDL, Cholesterol, total protein

INTRODUCTION

Quail has advantageous attributes such as small body size, rapid growth rate and early sexual maturity [1]. Quail also has cheaper production cost, and don't need a large land, easy to handle by human. Therefore they used for commercial rearing, egg and meat production under intensive management[2].

Probiotics are defined as beneficial live microorganisms, when administered in adequate amount may support a beneficial health on people with specific illnesses[3]. Probiotics, including live yeasts and bacteria, as feed additive or supplements and known give beneficial effect to the host by improving intestinal microbial balance, producing several substances that act as antimicrobial, stimulating the immune response, increase the intestinal microflora balance, modulating the immune response, and lowering cholesterol levels, enhance nutrient bioavailability, carcass yield and quality of Japanese quail[4, 5].

*Lactobacillus sp*is one of the isolates of bacteria that has a potential as a probiotic from the group of lactic acid bacteria. Lactic acid bacteria are Gram-positive, coccus or rod, non spore, acid tolerant, catalase negative, facultative an aerob and the major end product was lactic acid. The use of lactic acid bacteria provides benefits for animals and humans against pathogenic bacteria. Prolonged and continuous use of antibiotics in animal feed may result in antibiotic resistance. Probiotics such as lactic acid bacteria can be used as a substitute for antibiotic growth promoters that have a positive impact on livestock growth and disease prevention. Several studies have shown that the use of probiotics can decrease triglyceride and cholesterol content in serum and egg yolks [6-9]. The purpose of this research to evaluate the effect of probiotics supplementation containing *Lactobacillus casei* and *Lactobacillus rhamnosus* as alternative *Antibiotic Growth Promoter* to HDL, LDL, cholesterol and total protein of egg's *Coturnix coturnix japonica*.

*Corresponding author: Widya Paramita Lokapirnasari, Department of Animal Husbandry Faculty of Veterinary Medicine, Airlangga University, Mulyorejo Street, Surabaya 60115, Indonesia, email: widyaparamitalokapirnasari@gmail.com

MATERIALS AND METHODS

About 200 of 14weekold quails were randomized into five treatments with four repetitions, each replication consisted of ten quails, with this following treatment: T0: control (without probiotic), T1: 0.05 gram probiotic/ kg feed, T2: 0.1 gram probiotic/kg feed, T3: 0.025 gram probiotic/L drinking water and T4: 0.05 gram probiotic/L drinking water. Egg's quail were collected every day. The data collection of this research by taking five eggs randomly on each repetition in the last week of treatment.

a. **Cholesterol**, samples and reagents are prepared at room temperature after that pipette into tubes as below:

Tubes	Blank	Sample	CAL. Standard
R1.Monoreagent	1.0 mL	1.0 mL	1.0 mL
Sample	-	10 µL	-
CAL Standard	-	-	10 mµ

Mixing samples and reagents until homogen and incubate 10 minutes or 5 minutes at room temperature (37°C). After incubate, record the absorbance (A) of the standard and samples at λ500 nm against the reagent blank.

The color will be stable for 30 minutes and protected it from sun/light.

Calculations of total cholesterol was total cholesterol (mg/dL) = A sample/A standard × C standard

If using samples with concentrations higher than 600 mg/ dL, it should be diluted with saline to be 1:2 and then assayed again. Multiply the result by 2.

If results are to be expressed as SI units apply: mmol/L = mg/ dL × 0.0259

Cholesterol assay procedure,

Temperature at 25°C- 37°C, Optical path 1 cm, λ500 nm and Hg 546 nm

Measurement Against reagent blank

	Sample	Standard
Supernatant	100 µL	-
Standard	-	100 µL
Reagent	1000 µL	1000 µL

Calculation of cholesterol

Cholesterol in supernatant (mg/dL) = Δ E sample/ Δ E standard × concentration standard (mg/dL)

The concentration standard is the concentration of the total cholesterol in the cholesterol standard solution.

- I. **HDL assay procedure:** samples and reagents prepare in room temperature and pipette into centrifuge tubes 0.2 mL standard or sample and 0.4 mL precipitating reagent with ratio sample/reagent = ½ and dilution factor = 3. Vortex and allow to stand at room's temperature for 10 minutes after that centrifuge at 4000 rpm for 10 minutes, or at 12000 rpm for 2 minutes. Separate the clear supernatant within 2 hours. If supernatant triglycerides (>350 g/ dL) the sample should be diluted with saline (1:2) and then assayed again. Multiply the result by 2.

II.

II. Colorimetry : the cholesterol standard (50 mg/ dL) and the cholesterol MR Monoreagent of the kit prepare in room's temperature and pipette into tubes as below:

TUBES	Blank	Standard Supernatant	Sample Supernatant
Monoreagent	1.0 mL	1.0 mL	1.0 mL
Supernatant	-	-	50 µL
Standard	-	50 µL	-

Mixing and let the tubes at room temperature for 10 minutes or at 37°C for 5 minutes and read the absorbance (A) of the supernatant and the standard at λ500 nm against the reagent blank. The color will be stable for 30 minutes and protected it from sun/light.

Calculations of HDL:

HDL - cholesterol (mg/dL) = A supernatant/A standard × C standard

The results are expressed as SI units: mmol/L = mg/dL × 0.0259

b. LDL

Assay procedure

Precipitation

Sample 100µL and precipitating reagent 1000 µL

Mix and incubate at room temperature for 15 minutes, then centrifuge for 20 minutes at 2500 g. Within one hour after centrifugation, transfer 100 µL of the clear supernatant to the reaction solution for the determination of cholesterol.

The cholesterol standard has to be diluted 1 + 10 with NaCl (9 g/L). After dilution the standard is treated like the supernatant.

c. Protein assay procedure

Pipette into tubes:

TUBES	Blank	Sample	CAL Standard
R1.Biuret	1.0 mL	1.0 mL	1.0 mL
Sample	-	20 µL	-
CAL. Standard	-	-	20 µL

Mixed and incubated the tubes for 10 minutes at 37°C and read the absorbance (A) of the samples and the standard at λ540 nm against the reagent blank. The color will be stable for at least 1 hour

Calculations of total protein:

Total protein (g/dL) = A sample/A standard × C standard

Samples with concentrations higher than 12 g/ dL should be diluted 1:2 with saline and assayed again. Multiply the results by 2.

If results are to be expressed as SI units apply: g/ dL x 10= g/L

Statistical Analysis

Data analysis use Analysis of Variant (ANOVA). If the result show different or significantly different then continue with Duncan Multiple Range Test[10]. Statistical analysis using SPSS for Windows 21.0.

RESULTS

Cholesterol, HDL, LDL of egg’s quail

The results of *Analysis of Variance* (ANOVA) showed that the addition of probiotics through feed and drinking water have an effect on egg’s HDL, LDL and cholesterol (p<0.05) compare with control (Table 1).

Table 1. HDL, LDL, and Cholesterol Levels of egg quail in 4 week of treatment

Treatment	Variable		
	HDL(mg/g)	LDL (mg/g)	Cholesterol (mg/g)
T0	4.15 ^b ± 0.01	1.38 ^b ± 0.01	19.02 ^d ± 0.02
T1	4.68 ^d ± 0.02	1.49 ^c ± 0.01	16.74 ^b ± 0.01
T2	7.15 ^e ± 0.02	1.27 ^a ± 0.01	22.97 ^e ± 0.02
T3	3.88 ^a ± 0.01	1.74 ^d ± 0.02	10.41 ^a ± 0.01
T4	4.64 ^c ± 0.02	1.38 ^b ± 0.01	17.23 ^c ± 0.02

Total Protein of egg’s quail

The results of *Analysis of Variance* (ANOVA) showed that the addition of probiotics through feed and drinking water have an effect on egg’s total protein content (p<0.05) compare with control (Table 2).

Table 2. Total Protein of egg quail in 4 week of treatment

Treatment	Variable
	Total Protein(mg/g)
T0	136.43 ^a ± 0.02
T1	215.48 ^c ± 0.01
T2	168.81 ^b ± 0.02
T3	212.62 ^d ± 0.02
T4	194.05 ^c ± 0.02

DISCUSSION

The results of the quail egg cholesterol showed a differences compared to the controls (Table 1). In T3 treatment (0.025 gram probiotic/L drinking water) showed the highest significantly different effect which means T3 (10.41 mg/g) has the lowest cholesterol content compared with T0 (19.02 mg/g) and the others treatment groups (Table 1). Treatment T3 was the best result that can reduce the yolk cholesterol. To know the most influence treatment continued with Duncan's Multiple Range Test. The lowering of cholesterol levels in the egg yolk from the treatment caused by the amount of probiotics administration through drinking water which more effective to produce the amount of microbes in the digestive tract. This result in line with the research[11] that probiotics *Lactobacillus Plus* supplementation with treatment 0, 0.2%, 0.4% and 0.6% consisted by 2.56×10⁸ CFU showed significantly different in egg cholesterol. Increasing the number of microbes inhibit Hydroxi Metyl Glutaryil-KoA reductase (HMG-KoA reductase) enzyme work which plays a role in the formation of mevalonate in the process of cholesterol synthesis so can't change into cholesterol.

The research on improving egg characteristics is essential, not only to produce better quality eggs but also to provide food safety in a balanced diet for poultry industries [12]. Cholesterol is important for health because it is used as a constituent of hormones and for the production of bile acids [13], but excessive cholesterol consumption will endanger the health because it can cause atherosclerosis (blockage of arteries).

The results of Analysis of Variance (ANOVA) in HDL levels of quail eggs showed significance different ($P < 0.05$) among treatments. The results of analysis by Duncan's Multiple Range test showed differences in HDL levels among egg yolk T0 with T1, T2, T3 and T4. The highest HDL showed in 0.1 gram probiotic/kg feed (T2) treatment. This is in line with the study [14] that the fermented synbiotics supplementation affect to the HDL enhancement or good fats in laying hens. The fermentation synbiotics supplementation to 30% level in feed can raise blood serum HDL in laying hens. Supplementation of protexin into the laying hens diet has been shown to reduce egg yolk and serum cholesterol level. The mean of serum cholesterol level in the study was decrease from 188.50 mg/dL in the control group to be 186.35 mg/dL in the probiotic supplementation group. It is possible that microorganisms in the probiotics can assimilate the cholesterol in the tractus digestivus for cell metabolism, thus reducing the total absorption [15]. Another reason to reduced the cholesterol in host which fed by probiotics, suggested by [16] that probiotics could inhibit enzyme hydroxymethyl-glutarylco enzyme A, which it involved in the gastrointestinal tract. Another mechanism through probiotics can cause hypercholesterolemia action through bile acids (deoxycholic bile acids and cholic) which produced by hepatocytes and conjugated with taurine and glycine amino acid. In the small intestine, this acid absorbed and directed to the liver, and lower bile acid recycling will eventually lower serum cholesterol level. Probiotics can affect the binding of cholesterol and the inhibition of micelle formation combined with the fermentation effect on short chain fatty acid (SCFA) production is a proposed mechanism for explaining potential cholesterol effects [17]. Probiotics may change the enzyme, which is related in the process.

The enhancement of HDL levels usually followed by reduction levels of LDL and cholesterol. The results of LDL levels showed significant differences ($P < 0.05$) among treatments. The lowest LDL (1.27 mg/g) and high HDL (7.15 mg/g) showed by 0.1 gram probiotic/kg feed (T2) treatment. The reduction in LDL caused by the balances between HDL and LDL levels. The statement in line with [18] which states that the feed supplement can increase HDL levels and directly affect the reduction in LDL levels. LDL that has been formed in the liver will be sent to the blood the over production of LDL will be bound by Lecithin - Cholesterol Acyl Transferase (LCAT) enzyme in the process of Reverse Cholesterol Transport (RCT). LCAT has a function to bind lipoproteins or free fat in the blood plasma and send it to the liver. HDL as a lipoprotein that plays a role in the RCT process will increase the transport of cholesterol from the tissues to be returned to the liver and excreted through bile [19].

Total Protein

Based on the results of statistical analysis (Table 2) showed that the treatment had significantly different effect ($P < 0.05$) on quail egg yolk protein level. The result showed that the addition of probiotics in feed and drink can increase protein levels. The high average of egg yolk protein showed in T1 (215.48), T3 (212.62), T1 (207.62), T4 (194.05) and T2 (168.81 mg/g). The lowest total protein content of egg showed in control treatment (T0) without addition of probiotic. Protein content in feed affects the protein composition in eggs. Protein in feed mostly used for egg production, only a small portion for basic living higher production level then need higher protein [20]. The addition of probiotics in feed can increase microbial activity, enzyme activity and protein digestibility as well as metabolic energy of feed in the quail's digestive tract, so can accelerate the rate of feed movement, so the absorption of feed substances become larger and impact the enhancement of feed efficiency and quail production [21]. So that the protein content in probiotics can affect the protein content in quail eggs.

CONCLUSION

The results showed that probiotics 0.1 gram /kg feed and 0.025 gram /L drinking water could reduce cholesterol level, LDL level, increase HDL level and total protein of egg's quail *Coturnix coturnix japonica*.

ACKNOWLEDGMENTS

We thanks to Rector of Universitas Airlangga, Dean of Veterinary Medicine, Universitas Airlangga and Kemenristekdikti who have provided the fund for this research.

REFERENCES

1. Siyadati S.A., A.Mirzaei-Aghsaghali and K.H Ghazvinian,2011. Effect of varying ratio of energy and protein on live performance and visceral organs of male Japanese Quail, *Annals of Biological research*, 2 (2): 137-144.
2. Agboola, A.F., B.R.O.Omidwura., D.Y.Ologbosere and E.A.Iyayi, 2016. Determination of Crude Protein and Metabolisable Energy of Japanese Quail (*Coturnix coturnix japonica*) during Laying Period.
3. FAO, 2001. Health and Nutritional Properties of Probiotics in Food including Powder Milk with Live Lactic Acid Bacteria.
4. Hume, M.E, 2011. Historic perspective: prebiotics, probiotics, and other alternatives to antibiotics *Poult. Sci.*, 90 (2011): 2663-2669.
5. Siadati, S.A., Y. Ebrahimnezhad., G.Salehi Jouzani and J.Shayegh, 2017. Evaluation of Probiotic Potential of Some Native Lactobacillus Strains on the Growth Performance and Serum Biochemical Parameters of Japanese Quails (*Coturnix Coturnix Japonica*) during Rearing Period. *Revista Brasileira de Ciência Avícola*, 19 (3): 399-408.
6. Mohan B., R. Kadirvel., M. Bhaskaran and M. Natarajan, 1995. Effect of probiotic supplementation on serum/yolk cholesterol and on egg shell thickness in layers. *Br Poult Sci.*36:799–803.
7. Kalavathy R., N. Abdullah., S.Jalaludin., M. Wong, and Y.W Ho, 2009.Effects of Lactobacillus cultures on performance of layinghens, and total cholesterol, lipid and fatty acid composition of egg yolk. *J Sci Food Agric.* 89:482–486.
8. Saleh A.A., K. Hayashi., A. Ohtsuka, 2013. Synergistic effect of feeding *Aspergillus Awamori* and *Saccharomyces Cerevisiae* on growth performance in broiler chickens; promotion of protein metabolism and modification of fatty acid profile in the muscle. *J Poult Sci.* 50:242–250.
9. Zaidi, N.A., A.A.A.Hamid and T.H.T.A.Hamid, 2017. Lactic acid bacteria with antimicrobial properties isolated from the intestines of Japanese quail (*Coturnix Coturnix Japonica*). *G. War. Sains* 1(1) (2017): 10–12
10. Kusningrum, R.S, 2008. *Experiment Design*. Airlangga University Press. Surabaya.
11. Adi P., Heriyanto., Osfar Sjoifan., M. Halim Natsir, 2015. Effect of Addition of Lactobacillus Plus Probiotic Powder as Feed Additive on Quail Egg Quality (*Coturnix coturnix japonica*). Faculty Animal Husbandry University of Brawijaya Malang.
12. Duru, M., A. A.Duru., K.Karadaş., E.Eyduran.,H.Cinli and M. M. Tariq,2017. Effect of Carrot (*Daucus carota*) Leaf Powder on External and Internal Egg Characteristics of Hy-Line White Laying Hens. *Pakistan Journal of Zoology*, 49(1).
13. Aviati, V., S.M. Mardiaty., T.R. Saraswati, 2014. Cholesterol levels of quail eggs after giving of turmeric powder in feed. *Anatomy and Physiology Bulletin Volume XXII*, Nomor 1 March.
14. Yusrizal., Noverdiman., M. Fahmida and Yatno, 2012. The influence of the combination of probiotic and prebiotic (symbiotic) cement palm kernel (BIS) for fermentation ammonia emission reduction, health status and laying hens performance. In *Proceeding in SINas 2012*.
15. Khan, S. H., M. Atif., N.Mukhtar., A.Rehman and G.Fareed, 2011. Effects of supplementation of multi-enzyme and multi-species probiotic on production performance, egg quality, cholesterol level and immune system in laying hens. *Journal of Applied Animal Research*, 39(4), 386-398.
16. Fukushima M., M Nakano, 1995. The effect of probiotic on faecal and liver lipid classes in rats. *British Journal of Nutrition.*73:701–710.
17. St-Onge, B., J. Dufort and R. Lepage, 2000: Measuring tree height using scanning laser altimetry. In *Proceedings of the 22nd Annual Canadian Remote Sensing Symposium*. Victoria, B.C., 21–25 August, 425–32.
18. Rosadi, I, 2013. Levels HDL (High Density Lipoprotein) and LDL (Low Density Lipoprotein) Blood on a variety of local female ducks whose feeds are supplemented with probiotics. *JurnalIlmiah Animal Husbandry* 1(2): 597-605, July 2013.
19. Pusparini, 2006. Low density lipoprotein small solid as a risk factor at atherosclerosis. *Universa Medicina.*; 25(1):22-23.
20. Lamid, M., N.N.T. Puspaningsih, S. Mangkoedihardjo, 2013.Addition of lignocellulolytic enzymes into rice straw improves in vitro rumen fermentation products. *J. Appl. Environ. Biol. Sci.*, 3(9): 166-171.
21. Suherman, A.F., M.H. Natsir and O. Sjoifan, 2015.Effect of addition of probiotic lactobacillus plus flour form as feed additive to production appearance of quail. 1-8.

INSTRUCTION TO AUTHORS

Manuscript Submission:

Send your manuscript with attachment by mailing it to info@textroad.com, textroadjournals@gmail.com along with [covering letter](#).

Manuscript Preparation:

- * Title
- * Author names and addresses
- * Abstracts (Not more than 300 words)
- * Key words
- * Introduction
- * Materials and Methods
- * Results and Discussions
- * References (Use numbering in the text instead of full references).
Give full references at the end of the file
- * Photographs should be of high quality (Minimum 300-600 dpi)
- * Graphs should be in clearly visible form so that it may become easy to redraw
- * The manuscript must be submitted in MS-WORD file format.

INSTRUCTIONS TO AUTHORS

Submission

Submit manuscripts as e-mail attachment to the Editorial Office at:

textroadjournals@gmail.com or info@textroad.com along with [covering letter](#). A manuscript number will be mailed to the corresponding author same day or within 48 hours. The authors may also suggest two to four reviewers for the manuscript (JBASR may designate other reviewers). There is no page limit. The submitting author takes responsibility for the paper during submission and peer review.

Terms of Submission

Papers must be submitted on the understanding that they have not been published elsewhere (except in the form of an abstract or as part of a published lecture, review, or thesis) and are not currently under consideration by another journal. The submitting author is responsible for ensuring that the article's publication has been approved by all the other coauthors. All enquiries concerning the publication of accepted papers should be addressed to editor@textroad.com.

Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors within one or two weeks. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the JBASR to publish manuscripts within 4 weeks after submission.

Style of Manuscripts

Manuscripts should be written in clear, concise and grammatically correct English (with 10 font size and Times New Roman font style) so that they are intelligible to the professional reader who is not a specialist in any particular field. Manuscripts that do not conform to these requirements and the following manuscript format may be returned to the author prior to review for correction. The entire manuscript, including references, should be typed single spaced on one side of the paper. All pages should be numbered consecutively in the bottom centre starting from the title page. The manuscript should be presented in the following order.

Title and Authorship Information

The title should be a brief phrase (capitalize first letter of each word in the title) describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

Abstract

All manuscripts should not exceed 250-300 words and should describe the scope, hypothesis or rationale for the work and the main findings. Complete sentences, active verbs, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Keywords

Key words (5-7 words) should be provided below the Abstract to assist with indexing of the article. These should not duplicate key words from the title.

Introduction

This section should include sufficient background information, provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. The aims of the manuscript should be clearly stated. The introduction should not contain either findings or conclusions. It should be understandable to colleagues from a broad range of scientific disciplines.

Materials and Methods

This should be complete enough to provide sufficient detail to allow the work to be repeated by others. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

Results

Results should be presented in a logical sequence in the text, tables and figures; repetitive presentation of the same data in different forms should be avoided. The results should not contain material appropriate to the Discussion. It should be written in the past tense when describing findings in the authors' experiments. Results should be explained, but largely without referring to the literature.

Discussion

The discussion should consider the results in relation to any hypotheses advanced in the Introduction and place the study in the context of other work. Results and Discussion sections can be combined.

Conclusions

If an optional conclusion section is used, its content should not substantially duplicate the abstract.

Acknowledgment

The acknowledgments of people, grants, funds, etc should be brief.

References

Bibliographic references in the text appear like [1, 2, 5, 6], using square brace in superscript. References should be numbered consecutively, with style:

Journal paper:

1. Hadjibabaie, M., N. Rastkari, A.Rezaie and M. Abdollahi, 2005. The Adverse Drug Reaction in the Gastrointestinal Tract: An Overview. Intl. J. Pharmacol., 1 (1): 1-8.

Books:

1. Daniel A. Potter, 2002. Destructive turfgrass insects: Biology, diagnosis and control. Wiley Canada Publishers, pp: 24-67.

Chapters in Book:

1. Bray R.A., 1994. The leucaena psyllid. In: Forage Tree Legumes in Tropical Agriculture (eds R.C. Gutteridge and H.M. Shelton) pp. 283-291. CAB International, Oxford.

Titles of journals should be given in full. 'In press' can only be used to cite manuscripts actually accepted for publication in a journal. Citations such as 'manuscript in preparation' or 'manuscript submitted' are not permitted. Data from such manuscripts can only be mentioned in the text as 'unpublished data'.

A Report:

1. Makarewicz, J.C., T. Lewis and P. Bertram, 1995. Epilimnetic phytoplankton and zooplankton biomass and species composition in Lake Michigan, 1983-1992. U.S. EPA Great Lakes National Program, Chicago, IL. EPA 905-R-95-009.

Conference Proceedings:

1. Stock, A., 2004. Signal Transduction in Bacteria. In the Proceedings of the 2004 Markey Scholars Conference, pp: 80-89.

A Thesis:

1. Strunk, J.L., 1991. The extraction of mercury from sediment and the geochemical partitioning of mercury in sediments from Lake Superior, M. S. thesis, Michigan State Univ., East Lansing, MI.

Tables and Equations

Tables and equations should not be submitted in a format exceeding the A4 page size (in portrait form). **All tables should be embedded within the manuscript, and must be captioned and numbered sequentially.** Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text.

Figures / Illustrations / Photographs

Graphics should be supplied as high resolution (at least 300-600 dp.i.) electronic files. Digital images supplied only as low-resolution print-outs cannot be used. Graphs, diagrams, chromatograms, photos, etc. should be prepared as clear, original positives, suitable for reproduction. **All figures should be embedded within the manuscript, and must be captioned and numbered sequentially.**

Proofs

Proofs will be sent via e-mail as an Acrobat PDF file (e-mail attachment) and should be returned within 3 days of receipt. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

Check List

We recommend that you ask a colleague to read over your paper prior to submission to ensure it is of a high standard and conforms to a high level of scientific writing.

Before submission of your manuscript, please check that:

- All references cited in the text are included in the reference section.
- All figures and tables are cited in the text.
- Figures are at least 300 d.p.i.
- The pages are numbered.



Home Journals Instructions to Authors **Manuscript Submission** Join Us Contact Us

	Journal of Basic and Applied Scientific Research		Journal of Social Sciences and Humanity Studies
	Journal of Basic and Applied Chemistry		Journal of Basic Sciences and Applied Research
	Journal of Applied Environmental and Biological Sciences		Journal of Computer Sciences and Communication
	Journal of Pharmaceutical and Biomedical Sciences		Journal of Engineering and Higher Technology
	Journal of Agriculture and Food Technology		Current Economics and Management Research

