

Volume - 11

No. 8

August-2018

ISSN: 0972-8988

EISSN: 2231-0916



NAAS Rating : 5.71

Indexed in ESCI-Thomson Reuters, PubMed,
PubMed Central, DOAJ, Scopus, CABI, CAS etc.

Veterinary World

Open access and peer reviewed journal



Editorial office

Veterinary World,
Star, Gulshan Park,
NH-8A, Chandrapur Road,
Wankaner - 363621,
Dist. Morbi, Gujarat, India
Website: www.veterinaryworld.org
E-mail: editorveterinaryworld@gmail.com

Veterinary World

Editorial Office: Veterinary World, Star, Gulshan Park, NH-8A, Chandrapur Road, Wankaner - 363621, Dist. Morbi, Gujarat, India

ISSN: 0972-8988, EISSN: 2231-0916, www.veterinaryworld.org

Editor-in-Chief

Anjum V. Sherasiya - Ex-Veterinary Officer, Department of Animal Husbandry, Gujarat State, India

Associate Editors

Shambhunath Choudhary - Department of Biomedical & Diagnostic Sciences, College of Veterinary Medicine, The University of Tennessee, 2407 River Drive, Room A 201, Knoxville, TN 37996, U.S.A.

Suresh H. Basagoudanavar - FMD Vaccine Research Laboratory, IVRI, Bangalore, Bangalore- 560024, Karnataka, India

Editorial board

R. G. Jani - Ex-Coordinator Wildlife Health, Western Region Centre, Indo-US Project, Department of Veterinary Medicine, Veterinary College, Anand - 388001, Gujarat, India

G. N. Gongal - Technical Officer, WHO South-East Asia Regional Office, New Delhi - 110002, India

Ranganath Mamidi - Dr. Julian E Stelzer's Lab, Department of Physiology & Biophysics, Medical School, Case Western Reserve University, Cleveland, OH - 44106, U.S.A.

Md. Tanvir Rahman - Department of Microbiology and Hygiene, Faculty of Veterinary Science, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh

Deepmala Agarwal - Cancer Prevention Laboratory, Pennington Biomedical Research Center, Baton Rouge, LA, U.S.A.

Foud Kasim Mohammad - Professor, Department of Pharmacology & Toxicology, Vice President for Administrative & Financial Affairs, University of Mosul, P.O. Box 11136, Mosul, Iraq

Abdel-Baset Nasr Sayed Ahmed - Professor and Head, Department of Animal Nutrition and Clinical Nutrition, Faculty of Veterinary Medicine, Assiut University, Assiut, Egypt

Nicole Borel - Department of Pathology, Vetsuisse Faculty, University of Zurich, CH-8057 Zurich, Switzerland

B. A. Lubisi - Virology, MED Programme, ARC - Onderstepoort Veterinary Institute, No. 100 Old Soutpan Road, Onderstepoort, Tshwane, 0110, South Africa

Kumar Venkitanarayan - Associate Professor, Graduate Programs Chair, Honors and Pre-Vet Programs Advisor, Department of Animal Science, University of Connecticut, Storrs, CT 06269, U.S.A.

Kemin Xu - Department of Veterinary Medicine, University of Maryland, College Park College Park, MD, 20742, U.S.A.

Vassilis Papatsiros - Faculty of Veterinary Medicine, Department of Medicine (Porcine Medicine), University of Thessaly, Thessaly, Greece

Mathias Devreese - Laboratory of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Ghent University, Belgium Sumeet Sharma - Edmonton North Animal Hospital, Edmonton, Alberta, T5X 3Y7, Canada

K. P. Singh - School of Medicine and Dentistry, University of Rochester, Department of Environmental Medicine, Room: 4- 6820, 601 Elmwood Avenue, Box-EHSC, Rochester, New York-14620, U.S.A.

Raj Mohan Raja Muthiah - Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts, U.S.A. Ashok

K. Chockalingam - Division of Applied Regulatory Science, U.S. Food and Drug Administration, 10903, New Hampshire Avenue, Silver Spring, Maryland 20993, U.S.A.

Ashutosh Wadhwa - Poxvirus and Rabies Branch, Division of High-Consequence Pathogens and Pathology, National Center for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention, 1600 Clifton Road, NE, Mail Stop G33, Atlanta, GA 30 333, U.S.A.

Luiz Otavio Pereira Carvalho - Laboratory of Immunomodulation and Protozoology, Oswaldo Cruz Institute, Ministry of Health (Brazil), Pavilhão "108" - Sala: 09, Av. Brasil, 4365 - Manguinhos, Rio de Janeiro - RJ, CEP: 21040-360, Brazil.

Mallikarjun Bidarimath - Cornell Stem Cell Program, Department of Biomedical Sciences, T2-012 Veterinary Research Tower, Cornell University, College of Veterinary Medicine, Ithaca, NY 14853-6401, USA.

Ratings of Veterinary World: NAAS - National Academy of Agricultural Sciences - 5.71, Scimago Journal Rank -0.284, Citescore - 0.57, SNIP - Source Normalized Impact per Paper - 0.570

Indexing and abstracting

Academic Journals Database, AGORA, AGRICOLA, AGRIS, CABI, CAS, DOAJ, EBSCO, EMBASE, ESCI - Thomson Reuters, Gale, Google Scholar, HINARI, Indian Animal Science Abstracts, Indian Science Abstracts, JournalSeek, Open J-gate, ProQuest, PubMed, PubMed Central, SCOPUS, TEEAL

Publisher: Veterinary World

Veterinary World is an open access journal, each issue available free of cost at www.veterinaryworld.org.

We accept online submission only. For more information regarding submission and publication charges, please visit www.veterinaryworld.org

Rs.7000 for Indian/USD 175 for Abroad per copy

Printed and Published by Dr. Anjum V. Sherasiya on behalf of Veterinary World. Printed and Published at Star, Gulshan Park, N.H. 8A, Chandrapur Road, Wankaner-363621, Dist. Morbi, Gujarat, India.
Editor: Dr. Anjum V. Sherasiya

Veterinary World

ISSN: 0972-8988, EISSN: 2231-0916, www.veterinaryworld.org

Volume-11

No.8

August-2018

The articles in Veterinary World are open access articles licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Research (Published online: 01-08-2018)

1. Genetic and phylogenetic analysis of the outer capsid protein genes of Indian isolates of bluetongue virus serotype-16
Arpit Saxena, Sanchay K. Biswas, Karam Chand, Jishnu Naskar, Ankita Chauhan, Gulam Mohd, Neha Tewari, Kurat-ul-Ain, Muthannan A. Ramakrishnan and Awadh Bihari Pandey
Veterinary World, 11(8): 1025-1029

Research (Published online: 01-08-2018)

2. Seroprevalence of brucellosis in small ruminants in organized and unorganized sectors of Gujarat state, India
A. Kanani, S. Dabhi, Y. Patel, V. Chandra, O. R. Vinodh Kumar and R. Shome
Veterinary World, 11(8): 1030-1036

Research (Published online: 02-08-2018)

3. Pathogens isolated from clinical cases of urinary tract infection in dogs and their antibiogram
Manisha Punia, Ashok Kumar, Gaurav Charaya and Tarun Kumar
Veterinary World, 11(8): 1037-1042

Review (Published online: 02-08-2018)

4. Red flour beetle (*Tribolium castaneum*): From population genetics to functional genomics
Harshit Kumar, Manjit Panigrahi, Supriya Chhotaray, V. Bhanuprakash, Rahul Shandilya, Arvind Sonwane and Bharat Bhushan
Veterinary World, 11(8): 1043-1046

Research (Published online: 03-08-2018)

5. The effect of cashew leaf extract on small intestine morphology and growth performance of Jawa Super chicken
H. Setiawan, M. E. Jingga and H. T. Saragih
Veterinary World, 11(8): 1047-1054

Research (Published online: 04-08-2018)

6. Investigation of haptoglobin, serum amyloid A, and some biochemical parameters in calves with omphalitis
K. Bozukluhan, O. Merhan, M. Ogun, B. Kurt, M. Cihan, E. E. Erkilic, G. Gokce, U. Aydin and A. Ozcan
Veterinary World, 11(8): 1055-1058

Research (Published online: 04-08-2018)

7. Post-epidemic awareness and knowledge of Lassa fever among residents in affected community in Ibadan, Oyo State, Nigeria
E. J. Awosanya
Veterinary World, 11(8): 1059-1063

Research (Published online: 06-08-2018)

8. Effectiveness of poultry litter amendments on bacterial survival and *Eimeria* oocyst sporulation
Essam S. Soliman, Nahla H. Sallam and Eman M. Abouelhassan
Veterinary World, 11(8): 1064-1073

Research (Published online: 07-08-2018)

9. Sources of contamination, prevalence, and antimicrobial resistance of thermophilic *Campylobacter* isolated from turkeys
Radia Bouhamed, Leila Bouayad, Sara Messad, Safia Zenia, Malek Naim and Taha-Mossadak Hamdi
Veterinary World, 11(8): 1074-1081

Research (Published online: 09-08-2018)

10. Antimicrobial resistance genes in pathogenic *Escherichia coli* isolated from diseased broiler chickens in Egypt and their relationship with the phenotypic resistance characteristics
Mohamed M. Amer, Hoda M. Mekky, Aziza M. Amer and Hanaa S. Fedawy
Veterinary World, 11(8): 1082-1088

Research (Published online: 09-08-2018)

11. Determination of hematological and serum biochemical reference values for indigenous sheep (*Ovis aries*) in Dhaka and Chittagong Districts of Bangladesh
Md. Kaisar Rahman, Shariful Islam, Jinnat Ferdous, Md. Helal Uddin, Muhammad Belal Hossain, Mohammad Mahmudul Hassan and Ariful Islam
Veterinary World, 11(8): 1089-1093

Research (Published online: 10-08-2018)

12. A cross-sectional study of the welfare of calves raised in smallholder dairy farms in Meru, Kenya, 2017
Emily K. Kathambi, John A. Van Leeuwen, George K. Gitau and Shawn L. McKenna
Veterinary World, 11(8): 1094-1101

Research (Published online: 10-08-2018)

13. Prevalence and risk factors for *Salmonella* spp. contamination in broiler chicken farms and slaughterhouses in the northeast of Algeria
Samia Djeflal, Bakir Mamache, Rachid Elgroud, Sana Hireche and Omar Bouaziz
Veterinary World, 11(8): 1102-1108

Research (Published online: 12-08-2018)

14. Molecular and immunological characterization of *Hyalomma dromedarii* and *Hyalomma excavatum* (Acari: Ixodidae) vectors of Q fever in camels
Hend H. A. M. Abdullah, Eman E. El-Shanawany, Sobhy Abdel-Shafy, Hala A. A. Abou-Zeina and Eman H. Abdel-Rahman
Veterinary World, 11(8): 1109-1119

Research (Published online: 14-08-2018)

15. Genetic and phenotypic characterization of the native rabbits in Middle Egypt
El-Sayed Mahfouz Abdel-Kafy, Sahar Saad El-Din Ahmed, Amira El-keredy, Neama Ibrahim Ali, Sherif Ramadan and Ahmed Farid
Veterinary World, 11(8): 1120-1126

Research (Published online: 16-08-2018)

16. Potency of lactic acid bacteria isolated from balinese bovine (*Bos sondaicus*) intestinal waste from slaughterhouse to improve nutrient content of wheat pollard as animal feedstuff by fermentation process
Widya Paramita Lokapirnasari, Adriana Monica Sahidu, Koesnoto Soepranianondo, Agus Supriyanto, Andreas Berny Yulianto and Anam Al Arif
Veterinary World, 11(8): 1127-1134

Research (Published online: 16-08-2018)

17. Seasonal changes of rumen and intestine morphology of the Qinghai yak (*Bos grunniens*)
Bao A. Ding, Shuang Q. Ma, Zong R. Li, Xi L. Li and Stephen R. Madigosky
Veterinary World, 11(8): 1135-1138

Research (Published online: 17-08-2018)

18. Slaughter of pregnant goats for meat at Nsukka slaughterhouse and its economic implications: A public health concern
Onyinye Josephine Okorie-Kanu, Ekene Vivienne Ezenduka, Christian Onwuchokwe Okorie-Kanu, Chidiebere Ohazurike Anyaoha, Chukwuebuka Anselm Attah, Toochukwu Eleazar Ejiofor and S. Onyinye Onwumere-Idolor
Veterinary World, 11(8): 1139-1144

Research (Published online: 23-08-2018)

19. Isolation and identification of bacteria from fresh guava (*Psidium guajava*) sold at local markets in Mymensingh and their antibiogram profile
Md. Atiqur Rahman Sarker, Md. Mazedul Haque, Rafia Afroze Rifa, Fateha Akther Ema, Md. Ariful Islam and Mst. Minara Khatun
Veterinary World, 11(8): 1145-1149

Research (Published online: 23-08-2018)

20. Sporadic cases of lumpy skin disease among cattle in Sharkia province, Egypt: Genetic characterization of lumpy skin disease virus isolates and pathological findings
Fatma M. Abdallah, Hend M. El Damaty and Gamilat F. Kotb
Veterinary World, 11(8): 1150-1158

Research (Published online: 25-08-2018)

21. Morphologic and morphometric characteristics of ascaroid worm, *Ophidascaris piscatoris* in *Xenochrophis piscator* snake in Sidoarjo, Indonesia
Lucia Tri Suwanti, Inggarsetya Syah Audini, Setiawan Koesdarto and Emmanuel Djoko Poetranto
Veterinary World, 11(8): 1159-1163

Research (Published online: 25-08-2018)

22. An epidemiological investigation on occurrence of enterohemorrhagic *Escherichia coli* in raw milk
H. D. Vanitha, C. Sethulekshmi and C. Latha
Veterinary World, 11(8): 1164-1170

Research (Published online: 27-08-2018)

23. Serosurveillance of Brucella antibody in food animals and role of slaughterhouse workers in spread of Brucella infection in Southeast Nigeria

Samuel Okezie Ekere, Emmanuel Okechukwu Njoga, Joseph Ikechukwu Onunkwo and Ugochinyere Juliet Njoga
Veterinary World, 11(8): 1171-1178

Research (Published online: 28-08-2018)

24. Intestinal parasites among migrant barn swallows (*Hirundo rustica*) in the central region of Mazandaran Province, Northern Iran

Mahdi Fakhari, Tooran Nayeri Chegeni, Reza Bastani, Zahra Hosseini, Reza Saberi and Saber Armat
Veterinary World, 11(8): 1179-1182

Research (Published online: 28-08-2018)

25. Assessment of antibody assay methods in determination of prevalence of infectious bursal disease among local chickens and guinea fowls in Kwara state, North Central Nigeria

Oluwafemi Babatunde Daodu, Oladapo Oyedele, Julius Olaniyi Aiyedun, Hauwa Motunrayo Ambali, Rafiu Adebisi Kadir, Oluwakemi Christiana Daodu, Isaac Dayo Olorunshola and Arimie Deborah Adah
Veterinary World, 11(8): 1183-1187

Research (Published online: 29-08-2018)

26. Mixing two different propolis samples potentiates their antimicrobial activity and wound healing property: A novel approach in wound healing and infection

Noori Al-Waili
Veterinary World, 11(8): 1188-1195

Research (Published online: 30-08-2018)

27. Amino acid sequence based on Cytochrome b gene in Kejobong goat and its genetic relationships among several local goats in Asia

Dela Ayu Lestari, Endang Purbowati, Sutopo Sutopo and Edy Kurnianto
Veterinary World, 11(8): 1196-1202

Potency of lactic acid bacteria isolated from balinese bovine (*Bos sondaicus*) intestinal waste from slaughterhouse to improve nutrient content of wheat pollard as animal feedstuff by fermentation process

Widya Paramita Lokapirnasari¹, Adriana Monica Sahidu², Koesnoto Soepranianondo¹, Agus Supriyanto³,
Andreas Berny Yulianto⁴ and Anam Al Arif¹

1. Department of Animal Husbandry, Faculty of Veterinary Medicine, Jl. Mulyorejo, Kampus C, Universitas Airlangga, Surabaya, Indonesia; 2. Department of Marine, Faculty of Fisheries and Marine, Jl. Mulyorejo, Kampus C, Universitas Airlangga, Surabaya, Indonesia; 3. Department of Biology, Faculty of Science and Technology, Jl. Mulyorejo, Kampus C, Universitas Airlangga, Surabaya, Indonesia; 4. Doctoral of Veterinary Science, Faculty of Veterinary Medicine, Jl. Mulyorejo, Kampus C, Universitas Airlangga, Surabaya, Indonesia.

Corresponding author: Widya Paramita Lokapirnasari, e-mail: widyaparamitalokapirnasari@gmail.com

Co-authors: AMS: adriana_monica16@yahoo.co.id, KS: koesnotosp@yahoo.com, AS: bio_asp@yahoo.co.id, ABY: bernyjuliantomiroen@gmail.com, AA: a_alarif@yahoo.com

Received: 24-04-2018, **Accepted:** 06-07-2018, **Published online:** 16-08-2018

doi: 10.14202/vetworld.2018.1127-1134 **How to cite this article:** Lokapirnasari WP, Sahidu AM, Soepranianondo K, Supriyanto A, Yulianto AB, Al Arif A (2018) Potency of lactic acid bacteria isolated from Balinese bovine (*Bos sondaicus*) intestinal waste from slaughterhouse to improve nutrient content of wheat pollard as animal feedstuff by fermentation process, *Veterinary World*, 11(8): 1127-1134.

Abstract

Aim: The purpose of this study was to know the genetic and biochemical identification of isolated lactic acid bacteria (LAB) from Balinese bovine (*Bos sondaicus*) intestinal waste, acidity, and ox bile salts and to inhibit the growth pathogen of *Staphylococcus aureus* and *Escherichia coli* and the potential of those isolated to improve nutrient value of wheat pollard as animal feed ingredient by fermentation process.

Materials and Methods: This research was divided into three stages. The first stage, isolated LAB were obtained from the bovine intestines at a slaughterhouse in Indonesia. Small intestinal samples were collected from 10 healthy Balinese beef cattle (*B. sondaicus*). The isolated LAB were identified by VITEK 2, polymerase chain reaction, and 16S rDNA. The basic local alignment search tool (BLAST) was performed to determine the phylogenetic tree. The second stage, the LAB were screened for their tolerance at pH 2, 3, and 4; bile salt, and antagonistic to enteric pathogen. In the third stage, to determine the potency of this isolate to increase nutrient content of wheat pollard by facultative anaerobe fermentation for 3 and 5 days.

Results: The result of the first stage showed that the isolate could be identified as *Lactobacillus casei* WPL 315. The result of the second stage showed that the isolate tolerance to low pH (pH 2, pH 3, and pH4) for 90 min and 24 h, and this isolate had viability tolerance in 0.3% bile salt. The isolate can inhibit *S. aureus* and *E. coli*. The result of the third stage by proximate analysis showed that crude protein increased by 23.08% after fermentation, while crude fiber decreased by 61.24% on the level 0.5% *L. casei* subsp. WPL 315 in the 3-day fermentation.

Conclusion: Based on the results, it showed that *L. casei* WPL 315 derived from indigenous intestinal Balinese beef cattle (*B. sondaicus*) has tolerant characteristic on acidity and ox bile salts, has antagonistic effect against *E. coli* and *S. aureus*, and has the ability to increase crude protein and decrease crude fiber content of wheat pollard. It would be interesting to determine whether the strain has a probiotic candidate.

Keywords: *Escherichia coli*, *Lactobacillus casei*, probiotics, *Staphylococcus aureus*, wheat pollard.

Introduction

Feeding cost is the biggest component in the production cost of the poultry industry. To decrease feeding costs, some efforts have been taken by poultry farmers such as the addition of feed additive. The addition of various feed additives to the poultries has an important role in stimulating growth and decreasing number of feed conversion that can give positive effect on chicken growth [1]. Probiotic is one of the

feed additives that have been recently developed in the poultry industries, non-pathogen living organism that has mechanism to preserve microbiota balance in the digestive tract by influencing gastric microbiota as well as eliminating microorganism of host-pathogen by creating an inconvenient atmosphere for pathogenic bacterial growth [2]. The most common microorganism species used as probiotics are *Lactobacillus*, *Bifidobacterium* [3,4] *Lactococcus*, *Leuconostoc*, *Enterococcus*, and *Carnobacterium* [5], *Lactobacillus acidophilus*, *Lactobacillus sporogenes*, *Lactobacillus plantarum*, *Lactobacillus rhamnosus*, *Lactobacillus delbrueckii*, *Lactobacillus reuteri*, *Lactobacillus fermentum*, *Lactococcus lactis*, *Lactobacillus cellobiosus*, *Lactobacillus brevis*, and *Lactobacillus casei* [6].

Probiotic ability can be explained through various mechanisms. The microorganism can produce

Copyright: Lokapirnasari, et al. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

an antimicrobial substance, compete, and colonize in the gastrointestinal tract [7,8]. Probiotic can modulate immune cells. Probiotic is directly taken up through transcytosis by microfold epithelial cells and engulfed by macrophages or dendritic cells, which eventually triggers an immune response. Cytokines modulate the immune functions of dendritic, T and B cells [9]. Probiotic has a role in increased feed consumption for most livestock. The condition is caused by the increasing of feed digestibility in an animal that causes digestive tract that can be emptied soon so feed efficiency can be achieved. Probiotic not only increases feed consumption but also promotes growth so that it can enhance the feed conversion [10].

The use of other alternative feed ingredient from agricultural by-products (i.e., wheat pollard, rice bran, and maize bran) or agricultural wastes (i.e., rice straws, maize straws, maize leaf, and sugarcane leaves) was needed to maintain the availability of feed supply. Agricultural by-products or agricultural wastes that are available all years have low crude protein and high crude fiber content [11], such as rice bran, wheat pollard, cotton, and tofu wastes. Wheat pollard is agricultural by-products that are mostly used in livestock feeding because it is easy to get and costs lower. The limitation of wheat pollard utilization as the mixture in the livestock feed because of its low protein content, high crude fiber, and low digestibility.

To increase the feed quality based on exploration of indigenous lactic acid bacteria (LAB) from bovine intestine in the slaughterhouse, we isolated and identified LAB on acidity survival, ox bile salts survival, and to inhibit the growth of pathogen *Staphylococcus aureus* and *Escherichia coli* and the potential of the isolate, and to improve nutrient value of wheat pollard as animal feed ingredient by fermentation process.

Materials and Methods

Ethical approval

The research does not need ethical approval. However, samples were collected as per standard collection methods without any harm or stress to the animals.

Research procedure

This research was divided into three stage. In the first stage, LAB isolate was obtained from the bovine intestines which were identified by VITEK 2, polymerase chain reaction (PCR), and nucleotide sequencing of 16S rDNA by comparing them to the GenBank database. The basic local alignment search tool (BLAST) was performed to determine the kinship arrangement based on the phylogenetic tree. This research only determined one species of *Lactobacillus*. In the second stage, the LAB isolate was screened further for their tolerance to low pH, at pH 2, 3, and 4 as well as ox bile salts tolerance. In the third stage, to know further the ability of this isolate to animal feed ingredient (wheat pollard) conducted fermentation

process was conducted on facultative anaerobe condition for 3 days and 5 days.

First stage

Genotypic identification

DNA amplification with PCR and identifying coding genes based on nucleotide sequence of 16S rDNA genomes.

Isolation of strain from the small intestine of bovine

Small intestinal samples were collected from 10 healthy Balinese beef cattle from a slaughterhouse in Indonesia. All samples were cultivated using a modified de Man Rogosa and Sharp (MRS) broth and agar. Bacterial colonies which showed clear zone surrounding their colonies were selected to biochemical identification by VITEK 2, PCR, and 16S rDNA, and a further test of basic probiotic properties including acid and ox bile salts tolerance assay, and antagonistic to enteric pathogen.

DNA isolation

Ingredients used in the DNA isolation process were as follows: Lysozyme 10 mg/mL, buffer TE 50 mM (50 mM Tris Cl [pH 8.0]; 50 mM EDTA), buffer STEP (sodium dodecyl sulfate 0.5%, 50 mM Tris Cl [pH 8.0], 0.4 M EDTA, and proteinase K), Na-acetate 3M, Phenol: chloroform:isoamyl alcohol (25:24:1), ethanol 70%, cold absolute ethanol, and distilled water.

Ingredients used in the 16S rDNA gene amplification were buffer 2.5 µl, dNTP 2.0 µl, MgSO₄ 1.0 µl, DNA template 2.0 µl, primer forward PB 36 (10 pmol) 1.0 µl, primer reverse PB 38 (µmol) 1.0 µl, distilled water 10.3 µl, enzyme high fidelity Taq polymerase 0.2 µl, and PCR product detection with electrophoresis: Buffer TBE (tris base/boric acid/EDTA) 0.5×, agarose, and ethidium bromide. DNA isolation was performed using Ausubel methods [12].

DNA Amplification with PCR

High fidelity platinum Taq DNA polymerase (Invitrogen™ Platinum™ Taq DNA Polymerase High Fidelity, US) kit with primer forward PB36 5'-AGR GTT TGA TCM TGG CTC AG-3' (Invitrogen) and primer reverse PB38 5'-GMT ACC TTG TTA CGA CTT-3' (Invitrogen) that produced ± 1400pb were used for PCR.

Master mix of used amplification reaction was 10× high fidelity PCR buffer 2.5 ml, 10 mM dNTP mix 2 ml, 50 mM MgSO₄ 1 ml, primer forward 1 ml (10 pmol/µl), primer reverse 1 ml (10 pmol/µl), template cDNA 2 ml, platinum tag high fidelity 0.2 ml, and distilled water until it reached total volume of 20 ml. The used PCR condition was pre-denaturation at 95°C for 5 min, denaturation at 95°C for 1 min, annealing at 50°C for 1 min, extension at 72°C for 1 min, 30 cycles, and a final extension at 72°C for 10 min. PCR result was analyzed by electrophoresis gel on 2%

of an agarose gel that contains ethidium bromide. 5 µl DNA added with 2 µl loading dye was added into agarose holes, and then run in the 100-volt tension for more or less 30 min.

Analysis of DNA sequence coding 16S rDNA

DNA sequencing coding 16S rDNA was performed by 1st Base Serdang, Malaysia. Analysis of sequencing result was performed through BLAST nucleotide sequencing from 16S rDNA sequencing result with the available database on www.ncbi.nlm.nih.gov.

Biochemical identification

Biochemical identification by VITEK 2 microbial identification system version: 05.01 (BioMérieux) was applied in examining WPL 315 isolates. The VITEK 2 system (bioMérieux) is an integrated modular system that consists of a filling-sealer unit, a reader-incubator, a computer control module, a data terminal, and a multicopy printer. The system detects bacterial growth and metabolic changes in the microwells of thin plastic cards using a fluorescence-based technology. Different microwell cards contain biochemical substrates [13].

Second stage

LAB survival test on acidity and survival test on ox bile salts, antagonistic test, on enteric pathogen bacteria

Ingredients used in this research included an antagonistic test on enteric pathogen microbe used in MRSB/de MRS Broth (Oxoid) media, nutrient agar media (NA and Oxoid), and nutrient broth (NB and Oxoid). Media used in the survival test on acidity were MRSB (Oxoid), MRSA (Oxoid), 0.85% of sterile NaCl, as well as HCl. Media used in the bile salts test were MRSB (Oxoid), MRSA (Oxoid), 0.85% of sterile NaCl, as well as ox gall 0.3% (Oxoid). Media used to test crude protein proximate analysis were Tablet Kjeldhal (Merck), H₂SO₄ (Merck), NaOH 40% (Merck), boric acid (Merck), methyl red (Merck) indicator, Brom cresol green (Merck), H₂SO₄ 0.01 N (Merck), and Aquadest.

Selection of LAB as a probiotic candidate

The isolate assumed to have the ability as probiotic was selected through various tests, so superior isolate of LAB was chosen to be tested *in vitro*. The tests were as follows.

LAB survival test on acidity and survival test on ox bile salts

Acid tolerance was assayed as reported by Succi *et al.* with modification [14], in 10 mL of MRS broth adjusted to pH values of 2.0, 3.0, and 4.0 with 3.0 M HCl. MRS broth at pH 7 served as control. All tests were carried out in duplicate.

The modification method of Gilliland and Kim [15] was employed in this study to know the effects of ox bile salts 0.3% (w/v) (Sigma, Milan,

Italy) in MRS broth. All the samples were incubated at 37°C, 24 h. The aliquots were 10-fold diluted and viable bacteria (CFU/mL) were enumerated by spot plating on MRS agar (48 h, 37°C, and anaerobic conditions) [16].

Antagonistic test on enteric pathogen bacteria

The antagonistic test was assayed as reported by Jin *et al.* [17] with modification. Antagonistic test on enteric pathogen was performed with an agar diffusion method with modification in the pouring of pathogenic bacteria culture. LAB culture was grown on MRSB medium at 37°C for 18-20 h. After that, pathogenic bacteria were inoculated as much as 1 ose in the NB media, to be incubated for 24 h at 37°C. After incubation ended, 0.2 mL of the incubated bacteria was taken and placed into 100 mL NA media (0.2%) to be mixed well (homogeneous), and then placed into Petri dish with 1-20 mL for each dish until solid. After agar media became solid, a hole was created in the agar media with 6 mm diameter. Five holes were created for each Petri dish.

LAB culture from MRSB was spotted into the hole as much as 50 µl and then incubated for 24 h at 37°C. MRSB medium without LAB was used as the control. The observation was performed by measuring the clear zone around the hole using Vernier calipers. LAB antagonistic activity on enteric pathogen was shown as the diameter of created clear zone.

Third Stage: Potency of *L. casei* WPL 315 on fermented wheat pollard

*Inoculation of *L. casei* WPL 315 on fermented wheat pollard*

To know isolate ability on the nutritional content changes of crude protein and crude fiber, fermentation process was performed through following treatment P0: 100 g of wheat pollard without *L. casei* WPL 315 + molasses 4% addition (as control), treatment P1: 100 g of wheat pollard with addition of 0.5% *L. casei* WPL 315 + molasses 4% (3-day fermentation), treatment P2: 100 g of wheat pollard with addition of 1.0% *L. casei* WPL 315 + molasses 4% (3-day fermentation), treatment P3: 100 g of wheat pollard with addition of 0.5% *L. casei* WPL 315 + molasses 4% (5-day fermentation), and treatment P4: 100 g of wheat pollard with addition of 1.0% *L. casei* WPL 315 + molasses 4% (5-day fermentation). The fermentation process was done in anaerobe condition. The experimental design used in this research was a completely randomized design in triplicate for each treatment. The molasses was mixed with Aquadest as much as 20% from sample weight, and the isolate based on treatment level was poured in the mixture (molasses + Aquadest) and then sprayed evenly on the wheat pollard. The mixture was then fermented in anaerobe condition in the plastic bag for 3 days and 5 days. The isolate concentration of *L. casei* WPL 315 used in this research was 1.2×10^8 CFU/mL. After incubation ended, it was dried and continued into the proximate

analysis of crude fiber and crude protein according to AOAC [18].

Statistical analysis

Data were analyzed using one-way analysis of variance (ANOVA). If the significant differences were found, the test would be continued using Duncan's multiple range test on 5% significance level.

Results

Genotypic and biochemical identification

DNA amplification with PCR and identifying coding genes based on nucleotide sequence of 16S rDNA genomes

In this research, a colony of WPL 315 LAB was capable of growing on MRSA medium. Based on Gram staining, this LAB isolate was Gram-positive (GP), rod-shaped, and positive motility.

An advanced test was conducted on code WPL 315 LAB isolate with 16S rDNA and phylogenetic tree structure with 91-98% similarity. The majority of bacteria resembling WPL 315 isolate originated from *Lactobacillus* genus. Based on the degree of similarity of nucleotide structure, the closeness in position with *L. casei* ATCC 334 (accession NC_008526.1; 98% identity, Table-1), and inherited traits in congruence with microbe identification system, the isolated strain was identified as *L. casei* WPL 315.

Biochemical assay of LAB isolate WPL 315 was investigated using the VITEK 2 Compact system. The GP card of the VITEK 2 system includes biochemical tests to determine carbohydrate usage, enzyme activity, and resistance to certain compounds that can be used to identify GP, non-spore-forming bacteria [19]. The result of phenotypic identification of LAB isolates WPL 315 by VITEK 2 microbial identification system version: 05.01 (BioMérieux) as shown in Table-2.

LAB survival test on acidity

In the digestive tract, bactericidal effect from acid happened at pH under 2.5 [20]. The result of survival test on acidity showed that *L. casei* WPL 315 tolerance to low pH (Table-3).

Survival test on ox bile salts

The results of this research showed that *L. casei* WPL 315 has viability tolerance in ox bile salts 0.3% and the concentration of *L. casei* WPL 315 in ox bile salts 0.3% (Table-4).

Antagonistic test on enteric pathogen bacteria

The result of the antagonistic test on enteric bacteria shows that *L. casei* WPL 315 has an antagonistic effect against *E. coli* and *S. aureus*. The index antibacterial is shown in Table-5.

Inoculation of *L. casei* WPL 315 on fermented wheat pollard

The result of statistical analysis using one-way ANOVA showed that the use of *L. casei* WPL 315 on wheat pollard fermentation had a significant effect in the pH, crude protein, and crude fiber content of wheat pollard ($p < 0.05$). The result of wheat pollard fermentation showed the decreasing of crude fiber content and the increasing of crude protein content at 0.5% *L. casei* WPL 315 isolate within 3-5-day fermentation as shown in Table-6.

Discussion

Genotypic and phenotypic identification

DNA amplification with PCR and identifying coding genes based on nucleotide sequence of 16S rDNA genomes

To identify and determine the taxonomy of bacteria from several environment sources and identify the phylogenetic characterization, 16S rDNA gene sequencing can be applied since this molecule exists in every organism with identical function in all organisms [21-23]. The BLAST nucleotide (BLASTn) program (available at <http://blast.ncbi.nlm.nih.gov>) was used to screen candidate genes based on sequence similarity [24]. An advanced assay was conducted on code WPL 315 LAB isolate with 16S rDNA and phylogenetic tree structure with 91-98% similarity. The majority of bacteria resembling WPL 315 isolate originated from *Lactobacillus* genus. Based on the degree of similarity of nucleotide structure, the closeness in position

Table-1: Similarity identity

Description	Identities	Accession (sequence ID):
<i>Lactobacillus casei</i> ATCC 334 chromosome, complete genome	98%	ref NC_008526.1
<i>Lactobacillus rhamnosus</i> GG whole genome sequence, strain GG (ATCC 53103)	98%	ref NC_013198.1
<i>Lactobacillus sakei</i> strain 23K complete genome	94%	ref NC_007576.1
<i>Pediococcus clausenii</i> ATCC BAA-344, complete genome	94%	ref NC_016605.1
<i>Pediococcus pentosaceus</i> ATCC 25745, complete genome	94%	ref NC_008525.1
<i>Lactobacillus buchneri</i> CD034, complete genome	92%	ref NC_018610.1
<i>Lactobacillus reuteri</i> DSM 20016, complete genome	92%	ref NC_009513.1
<i>Lactobacillus plantarum</i> WCFS1, complete genome	92%	ref NC_004567.2
<i>Lactobacillus fermentum</i> IFO 3956 DNA, complete genome	92%	ref NC_010610.1
<i>Lactobacillus brevis</i> ATCC 367, complete genome	91%	ref NC_008497.1

Table-2: Biochemical test of LAB isolate WPL 315 by VITEK 2

Biochemical test	Reaction	Biochemical test	Reaction
LAC (Lactose)	+	dRIB (d-Ribose)	+
SAC (Saccharose/Sucrose)	+	dGAL (D-Galactose) Galactose	+
Sucrose			
Gluconate	+	Celobiose	+
dRIB (D-Rybose) (Ribose)	+	dRAF (D-Raffinose) Raffinose	-
dXYL (D-Xylose) Xylose	-	Mannitol	+
ARG (Arginin)	-	Ramnose	-
Arabinose	-	Esculin	+
BXYL (beta-xylosidase)	+	LeU (leucine arylamidase)	+
BGAL (beta-galactosidase)	+	AlaA (alanine arylamidase)	+
APPA (Ala-Phe-Pro Arylamidase)	+	GLYG (Glycogene)	-
ELLM (Ellman)	-	MTE (Maltotriose)	-
dMNE (D-Mannose)	+	PLE (Palatinose)	-
BMAN (Beta-mannosidase)	-	AGLU (alpha-glucosidase)	+
INU (Inulin)	-	PSCNa (putrescine assimilation)	-
OLD (oleandomycin resistance)	+	POLYB_R (Polymixin_B resistance)	+
LysA (L-lysine Arylamidase)	+	PheA (phenylalanine arylamidase)	+
PyrA (L-pyrrolidonyl arylamidase)	+	TyrA (Tyrosine Arylamidase)	+
CDEX (cyclodextrin)	-	INO (Inositol)	-
MdX (Methyl-d-xyloside)	-	GlyA (glycine arylamidase)	-
dMLZ (D-melezitose)	-	IRHA (L-rhamnose)	+
PHC (phosphoryl choline)	-	dTAG (d-Tagatose)	+
dGLU (D-glucose)	+	NaCl 6.5% (growth in 6.5% NaCl)	+
ESC (esculin hydrolyze)	+	ProA (L-proline arylamidase)	+
AspA (L-aspartate arylamidase)	-	BNAG (beta-N-acetyl-glucosaminidase)	+
AGAL (alpha-galactosidase)	+	MdG (methyl-A-D Glucopyranoside acidification)	-
dGAL (D-Galactose)	+	dMAN (D-Mannitol)	+
AMAN (alpha-mannosidase)	-	BGLU (beta-glucosidase)	-
NAG (N-acetyl-D-glucosamine)	+	dTRE (D-Trehalose)	+
PVATE (pyruvate)	+	KAN (kanamycin resistance)	+
TTZ (tetrazolium red)	+		

LAB=Lactic acid bacteria

Table-3: LAB survival test on acidity**Survival test on acidity of *L. casei* WPL 315**

Time	MRS agar (control) (CFU/mL)	MRS agar pH 2 (CFU/mL)	MRS agar pH 3 (CFU/mL)	MRS agar pH 4 (CFU/mL)
90 (min)	2.90×10 ⁸	6.20×10 ⁷	1.50×10 ⁸	2.60×10 ⁸
Duplicate	3.00×10 ⁸	6.20×10 ⁷	2.40×10 ⁸	2.90×10 ⁸
24 (h)	1.10×10 ⁹	1.00×10 ⁷	2.00×10 ⁷	2.25×10 ⁸
Duplicate	1.20×10 ⁹	1.00×10 ⁷	2.00×10 ⁷	2.40×10 ⁸

LAB=Lactic acid bacteria, *L. casei*=*Lactobacillus casei***Table-4:** LAB survival test on oxbile salts after 24 h, starting inoculums 2.90×10⁸

Lactide acid bacteria viability isolate (ox bile tolerance 0.3%)	Isolate
9.6×10 ⁷ CFU/ml	<i>L. casei</i> WPL 315

LAB=Lactic acid bacteria, *L. casei*=*Lactobacillus casei*

with *L. casei* ATCC 334 (accession NC_008526.1; 98% identity, Table-1), and inherited traits in congruence with microbe identification system, the isolated strain was identified as *L. casei* WPL 315.

VITEK2 Compact (bioMerieux, France) is an automated system able to identify microorganisms by testing 59 biochemical properties and also handle many samples in one reaction. VITEK2 compact was used for this study to differentiate isolates at a strain level by analyzing and comparing the phenotypes. Strains

were individually grown on MRS agar. Colonies were picked and mixed in a 0.45% NaCl solution until the McFarland standard measured 0.50-0.63 on the VITEK 2 DensiCheck instrument (bioMerieux). GP colorimetric identification cards (bioMerieux) and the tubes containing the bacteria were assembled in a cassette and assayed using the VITEK2 compact system. Data were analyzed using the VITEK 2 software version VT2-R03.1.

LAB isolated from the intestine of local beef cattle produced several enzymatic activities: Beta-xylosidase, beta-galactosidase, Ala-Phe-Pro Arylamidase, l-lysine arylamidase, l-pyrrolidonyl arylamidase, alpha-galactosidase, leucine arylamidase, alanine arylamidase, alpha-glucosidase, phenylalanine arylamidase, tyrosine arylamidase, and beta-n-acetylglucosaminidase. The proteolytic system of LAB is composed of a cell

Table-5: LAB survival test on *E. coli* and *S. aureus*

Antagonistic test on enteric bacteria	Diameter inhibition (mm)
<i>E. coli</i>	2.0
<i>S. aureus</i>	1.5

LAB=Lactic acid bacteria, *L. casei*=*Lactobacillus casei*, *E. coli*=*Escherichia coli*, *S. aureus*=*Staphylococcus aureus*

Table-6: Analysis result on nutrient content changes in wheat pollard fermentation using *L. casei* WPL 315 isolate

Treatment	pH	Crude protein	Crude fiber
P0 (control)	7 ^a ±0.11	13.0 ^a ±0.21	12.9 ^a ±0.21
P1 (0.5%, 3 days)	5 ^b ±0.13	16.3 ^c ±0.18	5.0 ^c ±0.18
P2 (1.0%, 3 days)	5 ^b ±0.10	14.0 ^{ab} ±0.45	8.2 ^{ab} ±0.45
P3 (0.5%, 5 days)	5 ^b ±0.10	15.0 ^b ±0.19	6.0 ^b ±0.19
P4 (1.0%, 5 days)	5 ^b ±0.13	14.5 ^{ab} ±0.47	8.0 ^{ab} ±0.47

^{a,b,c}Means in the same column with the different superscript are significantly different at ($p < 0.05$).

L. casei=*Lactobacillus casei*

envelope-associated proteinase, peptide transport systems, and intracellular peptidases. It can hydrolyze proteins to small peptides and amino acids which are essential for rapid microbial growth [25].

β -glucosidases enzymes are responsible for the catalyze of β -1,4-glycosidic bonds of various oligosaccharides, disaccharides, and alkyl- and aryl- β -d-glucosides [26], responsible for the hydrolysis of ce-lo-oligosaccharides and cellobiose, an important fiber source in cereal feeds. In addition, these enzymes hydrolyze toxic and/or bitter glucosides, release aromatic compounds, and synthesize various oligosaccharides, glycoconjugates, and alkyl- and amino-glucosides [27].

LAB survival test on acidity

The result of survival test on acidity showed that *L. casei* WPL 315 tolerance to low pH (Table-3). This was comparable with *L. casei* IS-7257 has viability as much as 5.22 ± 0.31 log CFU/mL. The survival on acid tolerance indicated the ability of the isolate to survive in stomach that has extreme pH (pH 2) and could survive in the gastrointestinal tract process where hydrolytic and gastric juice are secreted [28].

These results are in agreement with those obtained from previous similar studies, where *Lactobacillus* strains were able to survive when exposed by pH 2.5-4.0 but displayed loss of viability at lower pH values [29,30]. Lactic acid produced by *Lactobacillus* creates an acid environment that can inhibit the growth of pathogenic bacteria [31]. Other research showed that some LAB strains have function as competitive inhibitors on pathogenic organism [32], the strains include *L. casei* 99p, *L. rhamnosus* GG, *L. casei* Shirota, *Bifidobacterium breve* Yacult, and *L. acidophilus* [33].

Survival test on bile salts

Bile tolerance and acid tolerance are required for bacterial growth in the small intestine and survive passage through the stomach. The result of this research

shows that *L. casei* WPL 315 has viability tolerance in 0.3% bile salts. The results showed that the concentration of *L. casei* WPL 315 in 0.3% bile salts was 9.6×10^7 CFU/mL in MRS agar (Table-4). Similar observations were also reported by Srinu *et al.* [34] and Balasingham *et al.* [35] that LAB strains survived and tolerated at 0.3-2.0% bile salts (Oxgall). The viability tolerance in the bile salts condition is one of the main criteria for *in vitro* selection of potentially probiotic bacteria and microbes [36]. Because the bacterial cell wall is comprised mainly of phospholipids, bile salts which are an emulsifier and solubilizes the lipid that can damage the bacterial cells [37].

Antagonistic test on enteric pathogen bacteria

Inhibition of pathogens by the intestinal microbiota has been called bacterial antagonism, bacterial interference, barrier effect, colonization resistance, and competitive exclusion. Mechanisms by which the indigenous intestinal bacteria inhibit pathogens include competition for colonization sites, competition for nutrients, production of toxic compounds, or stimulation of the immune system [38]. LAB strains have potency in creating bactericidal bioactive peptides. Bacteriocins are also produced by species from *Lactobacillus*, *L. acidophilus* produces lactacin B or F, and *L. casei* B80 produces casein 80 [39,40]. Antimicrobial activity produced by LAB strain is not correlated with the acidity level in the medium. It has been reported that LAB strain has a strong inhibitory effect on *S. aureus* growth in milk. The inhibition ability is correlated with the existence of bacteriocins production, hydrogen peroxide production, and organic acids production such as lactic acid and acetic acid [41,42].

Inoculation of *L. casei* WPL 315 on fermented wheat pollard

The result of wheat pollard fermentation showed the increase of nutrient that was shown by the decreasing of crude fiber content and the increasing of crude protein content at 0.5% level within 3-5-day fermentation as shown in Table-6. The result of the statistical analysis showed that the use of *L. casei* WPL 315 on wheat pollard fermentation had a significant effect in the content of pH wheat pollard ($p < 0.05$). The result of pH level analysis showed the decrease of pH within the incubation process for all treatment groups compared to that in the control group (P0). *L. casei* WPL 315 treatment showed that the lowest pH was achieved in the treatment that used 0.5% isolate addition in the fermentation process because it was caused by LAB activity in recasting activity on water-soluble carbohydrate contained in the wheat pollard in the form of lactic acid. The decrease in pH level was followed by the decrease in carbohydrate level.

The result of the statistical analysis showed that the use of *L. casei* WPL 315 on wheat pollard fermentation had a significant effect on the content of crude fiber in wheat pollard ($p < 0.05$). The result of the analysis

showed a decreasing level of crude fiber content for all treatment compared to that in the control group (P0). The lowest crude fiber content was achieved in the treatment group that used 0.5% isolate within 3 days' fermentation process. The decrease of crude fiber content was correlated with the isolate ability to degrade organic matter derived from complex molecules becoming simplest molecules: Cellulose was degraded into cello-oligosaccharide to be then degraded into cellobiose; in the end, cellobiose was degraded into glucose [6]. Probiotics also stimulate activities of cellulolytic bacteria to degrade crude fiber [10].

The result of the statistical analysis showed that the use of *L. casei* WPL 315 on wheat pollard fermentation had a significant effect on crude protein content ($p < 0.05$). The result showed an increasing level of crude protein content for all treatment compared to the control group (P0). The highest crude protein content was achieved by adding 0.5% isolate within 3-day fermentation. This was caused by increased activity of *L. casei* WPL 315 in binding N as the basic matter to synthesize protein. Thus, the increase of nitrogen level allowed bacteria to grow and perform activity optimally that made crude protein level in wheat pollard increased higher compared to that in other treatment groups because bacteria are a single cell protein. The increase of crude protein content was also caused by the decrease of other compounds including nitrogen-free extract produced by fermented crude fiber [4]. Enzyme β galactosidase, glycols, and lactate dehydrogenase could be produced by LAB. It has a role in decreasing pH in the gastrointestinal tract, so it will inhibit *E. coli* growth and other pathogenic bacteria that need pH 6-7 [20].

Conclusion

The result of the research showed that *L. casei* WPL 315 derived from indigenous intestinal Balinese beef cattle (*Bos sondaicus*) has tolerant characteristic on acidity and ox bile salts and has antagonistic effect against *E. coli* and *S. aureus*.

Authors' Contributions

The work was done by WPL who designed the research and WPL, AS, as well as ABY who conducted the experimental work. WPL, AMS, and AA analyzed and interpreted the data and drafted the manuscript. WPL, KS, and AA participated in doing data collection, data analysis, data interpretation, and writing the manuscript. All authors read and approved the final manuscript.

Acknowledgments

The authors would like to thank the Ministry of Research, Technology, and Higher Education, Indonesia, Rector and Director of Research and Innovation Department, Airlangga University, that have funded the Excellent Research Universities (PTUPT) No. 004/ADD/SP2H/LT/DRPM/VIII/2017

Competing Interests

The authors declare that they have no competing interests.

References

1. Waldroup, P.W., Oviedo-Rondon, E.O. and Fritts, C.A. (2003) Comparison of bio-mos and antibiotic feeding programs in broiler diets containing copper sulfate. *Int. J. Poult. Sci.*, 2: 28-31.
2. Astini, W. (2014) Potential of Commercial Probiotic on Body Weight Gain, Feed Consumption, Feed Conversion Ratio. Universitas Airlangga, Surabaya.
3. Dongarrà, M.L., Rizzello, V., Muccio, L., Fries, W., Cascio, A., Bonaccorsi, I. and Ferlazzo, G. (2013) Mucosal immunology and probiotics. *Curr. Allergy Asthma Rep.*, 13(1): 19-26.
4. Besseling-van der Vaart, I., Heath, M.D., Guagnini, F. and Kramer, M.F. (2016) *In vitro* evidence for efficacy in food intolerance for the multispecies probiotic formulation ecologic® tolerance (Syngut™). *Benef. Microbes*, 7(1): 111-118.
5. Muñoz-Atienza, E., Gómez-Sala, B., Araújo, C., Campanero, C., Del Campo, R., Hernández, P.E., Herranz, C. and Cintas, L.M. (2013) Antimicrobial activity, antibiotic susceptibility and virulence factors of lactic acid bacteria of aquatic origin intended for use as probiotics in aquaculture. *BMC Microbiol.*, 13(1): 1.
6. Amara, A.A. and Shibl, A. (2015) Role of probiotics in health improvement, infection control and disease treatment and management. *Saudi Pharm. J.*, 23(2): 107-114.
7. Buffie, C.G. and Pamer, E.G. (2013) Microbiota-mediated colonization resistance against intestinal pathogens. *Nat. Rev. Immunol.*, 13(11): 790-801.
8. Vuong, C.N., Chou, W.K., Hargis, B.M., Berghman, L.R. and Bielke, L.R. (2016) Role of probiotics on immune function and their relationship to antibiotic growth promoters in poultry, a brief review. *Int. J. Probiotics Prebiotics*, 11(1): 1-6.
9. Dicks, L.M.T. and Botes, M. (2010) Probiotic lactic acid bacteria in the gastrointestinal tract: Health benefits, safety and mode of action. *Benef. Microbes*, 1(1): 11-29.
10. Bidarkar, V.K., Swain, P.S., Ray, S. and Dominic, G. (2014) Probiotics: Potential alternative to antibiotics in ruminant feeding. *Trends Vet. Anim. Sci.*, 1: 1-4.
11. Gabriel, A., Victor, N. and du Preez James, C. (2014) Cactus pear biomass, a potential lignocellulose raw material for single-cell protein production (SCP): A review. *Int. J. Curr. Microbiol. Appl. Sci.*, 3(7): 171-197.
12. Ausubel, F.M., Brent, R., Kingston, R.E., Mppre, D.D., Seldman, J.G., Smith, J.A. and Struhl, K. (1992) Short Protocols in Molecular Biology. 2nd ed. John Wiley & Sons, New York, Chichester, Brisbane, Toronto, Singapore.
13. Crowley, E., Bird, P., Fisher, K., Goetz, K., Boyle, M., Benzinger, M.J. Jr., Juenger, M., Agin, J., Goins, D. and Johnson, R. (2012) Evaluation of the VITEK 2 Gram-negative (GN) microbial identification test card: Collaborative study. *J AOAC Int.*, 95(3): 778-785.
14. Succi, M., Tremonte, P., Reale, A., Sorrentino, E., Grazia, L., Pacifico, S. and Coppola, R. (2005) Bile salt and acid tolerance of *Lactobacillus rhamnosus* strains isolated from parmigiano reggiano cheese. *FEMS Microbiol. Lett.*, 244(1): 129-137.
15. Gilliland, S.E. and Kim, H.S. (1984) Effect of viable starter culture bacteria in yogurt on lactose utilization in humans. *J. Dairy Sci.*, 67(1): 1-6.
16. Solieri, L., Bianchi, A., Mottolose, G., Lemmetti, F. and Giudici, P. (2014) Tailoring the probiotic potential of non-starter *Lactobacillus* strains from ripened parmigiano reggiano cheese by *in vitro* screening and principal component analysis. *Food Microbiol.*, 38: 240-249.
17. Jin, L.Z., Ho, Y.W., Abdullah, N., Ali, M.A. and Jalaludin, S. (1996) Antagonistic effects of intestinal *Lactobacillus*

- isolates on pathogens of chicken. *Lett. Appl. Microbiol.*, 23(2): 67-71.
18. AOAC. (1990) Official Methods of Analysis. 15th ed. Association of Official Analytical Chemists, Washington, DC, USA.
 19. Pincus, D.H. (2006) Microbial Identification using the BioMérieux Vitek 2 System. Encyclopedia of Rapid Microbiological Methods. Parenteral Drug Association, Bethesda, MD.
 20. Surono, I.S. (2003) *In vitro* probiotic properties of indigenous dadih lactic acid bacteria. *Asian Aust. J. Anim. Sci.*, 16(5): 726-731.
 21. Fan, C., Li, S., Li, C., Ma, S., Zou, L. and Wu, Q. (2012) Isolation, identification and cellulase production of a cellulolytic bacterium from intestines of giant panda. *Acta Microbiol. Sin.*, 52(9): 1113-11121.
 22. Huang, S., Sheng, P. and Zhang, H. (2012) Isolation and identification of cellulolytic bacteria from the gut of *Holotrichia parallela* larvae (*Coleoptera: Scarabaeidae*). *Int. J. Mol. Sci.*, 13(3): 2563-2577.
 23. Torres, A.R., Araujo, W.L., Cursino, L., Hungria, M., Plotegher, F., Mostasso, F.L. and Azevedo, J.L. (2008) Diversity of endophytic enterobacteria associated with different host plants. *J. Microbiol.*, 46(4): 373-379.
 24. Lee, H., Baek, H., Lim, S.B., Hur, J.S., Shim, S., Shin, S.Y., Han, N.S. and Seo, J.H. (2015) Development of species-specific PCR primers and polyphasic characterization of *Lactobacillus sanfranciscensis* isolated from Korean sourdough. *Int. J. Food Microbiol.*, 200: 80-86.
 25. Zotta, T., Ricciardi, A. and Parente, E. (2007) Enzymatic activities of lactic acid bacteria isolated from Cornetto di Matera sourdoughs. *Int. J. Food Microbiol.*, 115(2): 165-172.
 26. Yan, T.R., Lin, Y.H. and Lin, C.L. (1998) Purification and characterization of an extracellular β -glucosidase II with high hydrolysis and transglucosylation activities from *Aspergillus niger*. *J. Agr. Food Chem.*, 46(2): 431-437.
 27. Bhatia, Y., Mishra, S. and Bisaria, V.S. (2002) Microbial β -glucosidases: Cloning, properties, and applications. *Crit. Rev. Biotechnol.*, 22(4): 375-407.
 28. Surono, I.S. (2004) Probiotik Susu Fermentasi dan Kesehatan. YAPMMI, Jakarta. p149-151.
 29. Jacobsen, C.N., Nielsen, V.R., Hayford, A.E., Møller, P.L., Michaelsen, K.F., Paerregaard, A., Sandström, B., Tvede, M. and Jakobsen, M. (1999) Screening of probiotic activities of forty-seven strains of *Lactobacillus* spp. by *in vitro* techniques and evaluation of the colonization ability of five selected strains in humans. *Appl. Environ. Microb.*, 65(11): 4949-4956.
 30. Dunne, C., O'Mahony, L., Murphy, L., Thornton, G., Morrissey, D., O'Halloran, S., Feeney, M., Flynn, S., Fitzgerald, G., Daly, C. and Kiely, B. (2001) *In vitro* selection criteria for probiotic bacteria of human origin: Correlation with *in vivo* findings. *Am. J. Clin. Nutr.*, 73(2): 386s-392s.
 31. Bernardeau, M., Vernoux, J.P., Henri-Dubernet, S. and Gueguen, M. (2008) Safety assessment of dairy microorganisms: The *Lactobacillus* genus. *Int. J. Food Microb.*, 126: 278-285.
 32. Massi, M.B.V.F., Federici, D. and Matteuzzi, P. (2004) Identification e based on PCR combined with automated ribotyping for tracking probiotic *Lactobacillus* strains colonizing the human gut and vagina. *J. Appl. Microbiol.*, 96(4): 777-786.
 33. Nomoto, K. (2005) Prevention of infection by probiotics. *J. Biosci. Bioeng.*, 100: 583-592.
 34. Srinu, B., Rao, T.M., Reddy, P.V.M. and Reddy, K.K. (2013) Evaluation of different lactic acid bacterial strains for probiotic characteristics. *Vet. World*, 6(10): 785-788.
 35. Balasingham, K., Valli, C., Radhakrishnan, L. and Balasuramanyam, D. (2017) Probiotic characterization of lactic acid bacteria isolated from swine intestine. *Vet. World*, 10(7): 825-829.
 36. Hawaz, E. (2014) Isolation and identification of probiotic lactic acid bacteria from curd and *in vitro* evaluation of its growth inhibition activities against pathogenic bacteria. *Afr. J. Microbiol. Res.*, 8: 1419-1425.
 37. Musikasang, H., Tani, A., H-kittikun, A. and Maneerat, S. (2009) Probiotic potential of lactic acid bacteria isolated from chicken gastrointestinal digestive tract. *World J. Microbiol. Biotechnol.*, 25(8): 1337-1345.
 38. Patterson, J.A. and Burkholder, K.M. (2003) Application of prebiotics and probiotics in poultry production. *Poult. Sci.*, 82(4): 627-631.
 39. Rammelsberg, M. and Radler, F. (1990) Antibacterial polypeptides of *Lactobacillus* species. *J. Appl. Bacteriol.*, 69(2): 77-184.
 40. Klaenhammer, T.R. (1993) Genetics of bacteriocins produced by lactic acid bacteria. *FEMS Microbiol. Rev.*, 12(1-3): 39-85.
 41. Charlier, C., Even, S., Gautier, M. and Le Loir, Y. (2008) Acidification is not involved in the early inhibition of *Staphylococcus aureus* growth by *Lactococcus lactis* in milk. *Int. Dairy J.*, 18(2): 197-203.
 42. Lin, W.H., Hwang, C.F., Chen, L.W. and Tsen, H.Y. (2006) Viable counts, characteristic evaluation for commercial lactic acid bacteria products. *Food Microbiol.*, 23(1): 74-81.
