





ISSN 0215-8930

MEDIA

Kedokteran Hewan

Veterinary Medicine Journal



MKH (Vet.Med.J.)	Vol. 26	No. 2	Pp. 76-138	Surabaya, May 2010	ISSN 0215-8930
------------------	---------	-------	------------	--------------------	----------------

Accreditation by Director General of Higher Education, No. 108/Dikti/Kep/2007, August 23, 2007

Veterinary Medicine Journal

Vol. 26 No. 2, May 2010

Veterinary Medicine Journal contains scientific literature in the field of
Veterinary Science and Animal Husbandry
it was first published in 1985 with a frequency published three times a year in the month:
January, May and September

Composition of the Board Editor

Chief Editor:

Kusnoto

Secretary:

Erma Safitri

Treasurer:

Lilik Maslachah

Ad and subscription:

Boedi Setiawan

Executive Editors:

Sri Subekti Bendryman Soedjoko

Sri Agus Sudjarwo

Suwarno

Epy Muhammad Luqman

Translation Teams:

Mas'ud Hariadi

Suzanita Utama

Muchammad Yunus

Mustofa Helmi Efendi

Editor supervisor:

Susilowati

Address: Faculty of Veterinary Medicine, Airlangga University
Kampus "C" Unair, Jl. Mulyorejo Surabaya 60115
Telp. (031) 5992785; 5993016; Fax. (031) 5993015
e-mail: mkh_ua@yahoo.com

Account: Tabungan Bisnis Mandiri No. 14100-07144132, (a.n. Media Kedokteran Hewan)

Veterinary Medicine Journal published by the Association of Indonesia Veterinarians (PDHI)
and the Faculty of Veterinary Medicine, Airlangga University, Surabaya

Accreditation by Director General of Higher Education, No.108/Dikti/Kep/2007.

Veterinary Medicine Journal

Vol. 26 No. 2, May 2010

AUTHOR GUIDELINES

1. General Regulation
 - a. Veterinary Medicine Media/MKH publishes research articles, reviews and case reports in the scopes of veterinary medicine and animal husbandry.
 - b. Manuscript/articles must be original and may not have been published. Manuscripts accepted for publication in Vet. Med. J. should not be submitted for any other publication.
2. Writing Standard
 - a. Manuscripts are typed in double-line spacing except for title, abstract, figure/table legends, table content, references and appendices.
 - b. New paragraphs use first line indentation by 0.3".
 - c. Standard Font for manuscript writing is 11 point Book Antiqua.
 - d. Manuscript should be printed on A4 HVS paper.
 - e. Manuscript could be submitted in Indonesian or English.
 - f. Tables / Figures must be very contrast and scanning files (photograph) must be enclosed separately in **JPG format**. Table/Figure Legends or other notes in Appendices should be typed in single-line spacing.
3. Manuscript writing guidelines
 - a. The whole manuscript should be composed of 12–14 pages.
 - b. Topics (Title, Author Names, Abstract, Introduction, Methods, etc.) should be written in Title Case, not Uppercase (all capitals) and are to be left-justified.
 - c. Manuscript should be written following this order: Title, Author names and Identity, Corresponding Address, Abstract with Keywords, Introduction, Methods, Results and Discussion, Conclusion, Acknowledgement (if any), References and Appendices (if any).
 - d. Title must be concise, specific, written in full (not abbreviated), informative, written in Indonesian and English.
 - e. Author names are placed beneath the title, author identity and corresponding address should be clear, written in full and placed after the author names.
 - f. Abstract is written in Indonesian and English, single-spaced and not exceed 200 (two hundred) words.
 - g. Keywords is placed after the abstract, list up to 5 (five) words.
 - h. Methods include equipments/materials which were used (particularly the specific ones), research procedure and statistical analysis (if used).
 - i. References should be listed in alphabetical order without numbering. Journal names should be abbreviated according to the system used by each journal. References should be single spaced with hanging indentation by 0.3" and before spacing by 3.6 pt. References should be composed of 60 % scientific articles and 40 % scientific text books. Below are the examples of references style to journals and books.

Roitt I, Brostoff J, and Male D. 1996. Immunology. 4th Ed. Blackwell Scientific Pub. Oxford.

Staropoli I, Clement JM, Frenkiel MP, Hofnung M, and Deuble V. 1996. Dengue-1 virus envelope glycoprotein gene expressed in recombinant baculovirus elicits virus neutralization antibody in mice and protects them from virus challenge. *Am J Trop Med Hyg*; 45: 159-167.
4. Manuscripts submission could be performed at any time by sending 3 print copies and one soft copy in a CD to Media Kedokteran Hewan, Faculty of Veterinary Medicine, Airlangga University, Kampus C Unair, Jalan Mulyorejo, Surabaya 60115, Telephone 031-5992785; 5993016; Fax. 031-5993015; e-mail: **mkh_ua@yahoo.com**
5. The editors reserve the right to
 - a. Publish manuscript/article without revision
 - b. Publish manuscript/article with revision
 - c. Reject manuscript/article
6. The editors are not responsible for all statements in published manuscripts.
7. Accepted manuscript for publication requires author to cover publishing and delivery cost.
8. Authors / subscribers may send publication or subscription cost by bank-transfer to **Media Kedokteran Hewan FKH UNAIR, on Tabungan Bisnis Mandiri, account no. 1410007144132.**
9. All editorial decisions are final; no further correspondence will be responded.

Veterinary Medicine Journal

Vol. 26 No. 2 May 2010

Published every 4 months, in January, May and September

Acknowledgements

Editors, writers and readers of Veterinary Medicine Journal award and thanks the highest to the experts below, as contribution partners who have examined all the good writing that was published or rejected according to the recommendations submitted to the editor of Volume 26, No.1 January 2010 edition.

Prof . Hj. Romziah Sidik, DVM., Ph.D.
(Faculty of Veterinary Medicine, Airlangga University)

Prof. Dr. Aulani'am, DVM., DEA.
(Faculty of Mathematic and Natural Science, Brawijaya University)

Prof. Dr. Imam Mustofa, DVM., M.Kes.
(Faculty of Veterinary Medicine, Airlangga University)

Prof. Dr. Bambang Sektiari, DVM., DEA.
(Faculty of Veterinary Medicine, Airlangga University)

Dr. Med.Vet. Denny Widaya Lukman, DVM, MSi.
(Faculty of Veterinary Medicine, Bogor Institute of Agriculture)

Dr. Ir. Osfar Sjojfan, M.Sc.
(Faculty of Husbandry, Brawijaya University)

Prof . Dr. Fedik Abdul Rantam, DVM.
(Faculty of Veterinary Medicine, Airlangga University)

Aris Junaidi, DVM., MP., Ph.D.
(Faculty of Veterinary Medicine, Gajah Mada University)

Ir. Marten Luther Mullik, Ph.D.
(Faculty of Husbandry, Nusa Cendana University)

Veterinary Medicine Journal

Vol. 26, No. 2, May 2010

Published every 4 months, in January, May and September

CONTENS

		Pages
12	<i>In Vitro</i> Nuclear Maturation of Domestic Cat (<i>Felis catus</i>) Ovarian Oocytes after being Stored at 4°C (Karja et al.)	76 – 79
13	Concentrations of Liver and Plasma Malondealdehyde (MDA) of Sprague Dawley Rats Injected by Ketamine-Xylazine HCl Anesthetics (Adji)	80 – 83
14	Protein Profile of Sporozoite of <i>Leucocytozoon</i> sp. from <i>Culicoides</i> sp. (Suwanti et al.)	84 – 87
15	Expression of CD ₄ ⁺ T cell, Interleukin-2 and Interleukin-4 on Splenectomy Balb/c Mice Post-Exposure <i>Salmonella typhi</i> (Aulanni'am et al.)	88 – 91
16	Detection of Cross Reaction between <i>Toxocara cati</i> Antigens against Anti- <i>Toxocara canis</i> Serum Used Western Blot Technique (Kusnoto)	92 – 99
17	The Profile of Antioxidant Superoxide Dismutase (SOD) in Liver of Isoflavone, Zn, and Vitamine E-treated Rats (Wresdiyati et al.)	98 – 105
18	Species Identification of coagulase positive staphylococci (CPS) by multiplex Polymerase Chain Reaction (PCR) (Effendi)	108 – 111
19	Mechanisms Underlying Abortion in Dexamethasone Treatment at Ewe in Midgestation (Srianto et al.)	112 – 117
20	Antibody Against Prolactin: as a Therapy for Molting Duck (Safitri)	118 – 123
21	Optimization of Temperature and Cellulase Enzyme pH from Rumen Bacteria Isolation of Beef Cattle (Lamid)	124 – 127
22	Antiproliferation Efect of the n-Heksana Extract of Kesum (<i>Polygonum minus</i>) at the Cells of the Rat Lung Cancer: Effort Exploiting of Kesum as Drug of Lung Cancer (Wibowo et al.)	128 – 133
23	Effects of Supplementation to Cows or Directly to Calves on Calf Performance in Bali Cattle Grazing Communal Pastures (Belli and Jelantik)	134 – 138

Optimization of Temperature and Cellulase pH from Rumen Bacteria Isolation of Beef Cattle

Mirni Lamid¹, Ni Nyoman Tri Puspaningsih², and Widya Paramita L¹

¹Department of Husbandry, Faculty of Veterinary Medicine, Airlangga University; ²Department of Chemistry, Faculty of Science and Technology, Airlangga University, C Campus, Jl Mulyorejo, Surabaya, Indonesia; e-mail: mirnylamid@yahoo.com

Abstract

Biotechnology in animal feed using biological treatment such as cellulase is mainly aimed to increase degrade the cellulose; the second most abundant polysaccharide in nature. Information of using cellulase as biocatalyst for increasing animal feed quality as ruminants feed is limited. Cellulose is the major component of cell wall agroindustry waste after hemicellulose. This cellulase comes from *Actinobacillus* sp. that were isolated from rumen fluid of beef cattle. The aim of this research was to determine activity of cellulase and characterization of optimum temperature and pH. In this research identification with carboxyl methylcellulose (CMC) has been done. Methods for cellulase activity was 3,5-dinitrosalicylic acid (DNS). The results showed that *Actinobacillus* which have positive activity of cellulase. Characterization of *Actinobacillus* sp. had optimum temperature 45 °C with activity 1.42 U/ml and optimum pH 6 with activity 0.57 U/ml.

Key words: cellulase, carboxyl methyl cellulose, temperature, beef cattle

Introduction

The potential of agro industry waste product is great, but the problem occurred was the low value of nutrition signed the high content of crude fiber (cellulose, hemicelluloses, and lignin) and the low value of crude protein. Cellulose and hemicelluloses provided used a little part only and the most become waste product. It is caused cell wall of agro industry waste product realized lignifications, make a complex bound with lignin. Cellulose as an organic material compound abundant in this nature has a sharp filament like; it is content of cell wall of plant. Cellulose is the highest component of the plant. Cellulose is the glucose polymer connected with glycoside no branch. Most of animal can not digest cellulose because they don't have cellulase enzyme in their digestive tract. Ruminant can use cellulose as their nutrient source because in their rumen contain cellulolytic bacteria produced cellulase enzyme to hydrolyze cellulose (Campbell, 1994).

Cellulose molecule is the polysaccharide with bound β -1-4 glycoside that is difficult to digested by rumen bacteria therefore digestibility of cellulose was very low (Mc. Donald *et al.*, 1995). Rumen microbes are main source of cellulase enzyme producer but digestibility of agro industry waste still low so need a

process before. Utilizing of cellulase enzyme group will increase nutrition value of agro industry waste product. Cellulase enzyme contains 3 enzyme components, that is: component C1 (β -1, 4-glucan cellobiohydrolase or exo- β -1, 4-glucanase), component Cc (endo- β -1, 4-glucanase) and component cellobyose (β -glucosidase). The centre point of cellulose degradation provide in the break of bound 1.4 β -glycoside. The breaking of bound 1.4 β -glycoside caused cellulase hydrolyzed become simpler component that is oligosaccharide (mainly cellobyose). Then it will hydrolyze polysaccharide (mainly glucose). The breaking of 1.4 β -glycoside bound done by cellulase enzyme group (Bondy, 1987).

Improvement of hydrolyze enzymatically was a new prospect to handle ligno cellulose become the feed for ruminant. Biodegradation of cellulase enzymatically accomplished by cellulase enzyme produced by much kind of microbes, such as: fungi, bacteria or *actinomycetes*, *alga*, *oomycetes*, ruminant (Campbell, 1995).

Cellulase enzyme has very big industry application that is textile industry, pharmacy, paper and pulp, food and animal feed, and chemistry industry. Application of cellulase enzyme becomes one of component in chicken rations (Sunna and

Antranikian, 1997). Cellulose enzyme gather with xylanase can break glucose reduce the fermentation result of structural polysaccharide of the plant cell wall at the grass cellulase process, thus utilizing of this enzyme can increase consumption and digestibility of ruminant (Spoelstra *et al.*, 1990). The giving of feed additive enzyme manly cellulase and xylanase can increase feed efficiency for beef cattle and dairy cattle (Beauchemin *et al.*, 2003).

The protein structure of enzyme can influence catalytic activities. Catalytic activities of enzyme will be loss if protein denaturizing occurred. The occurring cause of protein denaturizing is extreme pH, high temperature, and influencing of other component such as detergent, ionic, organic solution, concentrate urea, big anion of strong acid and chaotropic ion (I-,SCN-)(Ottoway and Apps, 1984). Considered from the biochemistry structural, there are some difference between enzyme termophilic and mesophilic. Beside from disulfide content, hydrophobic interaction level at the molecule has enzyme stability. The optimum temperature of termophilic enzyme activities is about 50°-80° C, and for termophilic enzyme is at the room temperature, about 25°-50° C. if the mesophilic enzyme placed at higher temperature will realize denaturizing.

This research used bacteria isolation from beef cattle rumen, *Actinobacillus* sp. The research was to investigate cellulase activities then accomplished characterization to determine optimal temperature and pH of cellulose enzyme from the bacteria isolation.

Materials and Methods

Sample

The sample used was bacteria isolation, *Actinobacillus* sp. produced from beef cattle rumen.

Production of fluid media

Fluid media contains of 0.2 g trypton; 0.1 yeast extract; 0.2 g NaCl. All materials dissolved at the Erlenmeyer tube 100 ml with 20 ml aquadest, and then closed by cotton and aluminum foil. After that it sterilized by autoclave for 45 minutes at 121°C, 1 atm.

Production of solid media

Fluid media contains of 0.2 g trypton; 0.1 yeast extract; 0.2 g NaCl; o, 3 g bacto agar. All materials dissolved at the Erlenmeyer tube 100 ml with 20 ml aquadest, and then closed by cotton and aluminum foil. After that it sterilized by autoclave for 45 minutes at 121°C, 1 tam. Then waited until lukewarm and poured into Petri disk then let it until cold and hard or solid.

Halo test

Bacteria isolation, *Actinobacillus* sp. planted in the solid media contained substrate Carboxyl Methyl Cellulose (CMC) by streak ose to the bacteria colony then streak again to the surface of solid media. Incubated at 37°C for 16 hours. Observation done to find Halo (transparent area) around the colony then choose the isolation had the biggest Halo and also shown the highest cellulase activities.

Then Halo gotten was examined by Congo-red 0.1% poured in to the isolation. Positive result signed with Halo area transparent still and the color became red (Puspaningsih, 2004).

Inoculums production

Isolation renewed was moved to the growth media 20 ml in the Erlenmeyer tube 100 ml, and then incubated with shaking 150 rpm at 39°C during optimal cultivation time received from growth curve. Cultivation result was cell suspension or inoculums

Enzyme production

Enzyme production accomplished by inoculated 1% inoculums in to 50 ml fluid media contained carboxyl methyl cellulose (CMC) in the Erlenmeyer tube 250 ml and incubated at 39°C for 16 hours with shaking 170 rpm. Incubation result taken the supernatant for tested the cellulase enzyme activities.

The cellulase activities tested by method 3,5-dinitro-salicylic (DNS) acid

A number 100µl enzyme and xilan substrate entered to the each Eppendorf tube then incubated at 39°C in the water heater for an hour. For control, the formula was same with sample without incubation. Then sample and control added 600µl DNS, after that heated in the boiled water for 15 minutes, then incubated in the ice bath for 20 minutes. Absorbance read in the wave length 550nm. 1 unit enzyme activities defined as a total enzyme produced 1µmol xylosa per minute for every ml enzyme, and the specific activities was enzyme unit divided protein content (Miller, 19960).

Characterization of Cellulose Enzyme

Determination of optimum temperature for cellulase done by determined activities of cellulase at many kind temperatures variant, that was at temperature: 35, 40, 45, 50, 55 and 60°C.

Determination of optimum temperature of cellulase determined by activity assay to cellulase at pH 5-9 used buffer phosphate citrate at optimum temperature of cellulase.

Results and Discussion

The research result of enzymes cellulase activities from bacteria *Actinobacillus* sp. planted in the medium contained MCM at the many kind of temperature presented at Table 1 and Fig. 1. The influence of temperature on the cellulase activities at this result investigated at 35-60°C. Optimization temperature result of cellulase received at 45°C with activities 1.42 U/ml, but at 50°C and 55°C cellulase still have good activities that reached 92.96% (1.32 U/ml) and 78.17% (1.11 U/ml). At the temperature above 55°C decreasing of cellulase activities occurred, because atoms in the enzyme molecule had big enough energy to move, it caused by the presence form changing of enzyme structure because increasing thermal tremble of atoms component, therefore protein as enzyme material denaturized.

The research of optimum temperature was lower then the research result reported by Lowe *et al.*, (1997) that rumen fungi *Neocallimastix frontalis* plant at the wheat straw substrate produced cellulase had optimum activities at 50°C.

Cellulase activities realized increasing from pH 5 reached the highest activities about 0.57 U/ml at pH 6, at pH 7 cellulase activities decreased reached 73.68% (0.42 U/ml) from activities at the optimum pH. It's probably caused by conformation changing between enzyme and substrate because pH decreasing (Fig. 2).

Generally, in the cattle rumen the optimum pH was 6-7. According to the research result, gotten optimum pH from cellulose enzyme was suitable with condition of rumen pH. It was one of cellulase characteristic that was total ions at that pH made confirmation of cellulase formed a pair with the

substrate. It caused also by enzyme received from rumen bacteria, thus pH gotten was suitable with bacteria's habitat. Optimum condition at pH 6-7 was suitable to growth cellulitic and hemicellulitic bacteria in the rumen to degraded fibrous feed produced VFA. This result synchronizes with the opinion of Owens and Goetsch (1988) that pH condition very influenced by microbes environmental in the feed degradation. That cellulolytic and hemicellulolytic bacteria need pH about 6.2-7.0 to develop fast (Orskov and Ryle, 1990).

Table 1. The Data of Optimum Temperature of Cellulase of Bacteria *Actinobacillus* sp.

Temperature (°C)	Cellulase activities (U/ml)
35	0.86
40	0.71
45	1.42
50	1.32
55	1.11
60	0.96

Table 2. The Data of Cellulase Optimum pH of Bacteria *Actinobacillus* sp.

pH	Cellulase Activities (U/ml)
5	0.53
6	0.57
7	0.42
8	0.16
9	0.15




Figure 1. The pattern of relationship between temperature and the cellulase enzyme activities.




Figure 2. The pattern of relationship between pH and the cellulase enzyme activities.

Conclusion

The characterization of cellulase enzyme has optimum temperature 45°C about 1.42 U/ml and optimum pH about 0.56 U/ml

References

- Bhat MK and Hazlewood GP. 2001. Enzymology and Other Characteristics of Cellulases and Xylanases, In: *Enzymes in Farm Animal Nutrition* M.R. Bedford and G.G. Partridge (Eds). CABI Publ, UK.
- Beauchemin KA, Colombatto D, Morgavi DP and Yang WZ. 2003. Use of exogenous fibrolytic enzymes to improve feed utilization by ruminant. *J Anim Sci*; 81: E37 – E47
- Bondi AA. 1987, *Animal Nutrition*, Wiley-Interscience Publication, London.
- Campbell Mary K. 1995. *Biochemistry*. 2nd Ed. Mount Holyoke College, Saunders College Publ, United States of America. Pp. 330-332
- Lowe SE, Michael K, Theodorou and Anthony PJT. 1987. Cellulases and Xylanases of an Anaerobic Rumen Fungus Grown on Wheat Straw, Wheat Straw Holocellulose, Cellulose, and Xylan. *Appl Environ Microbiol*.
- Mc. Donald P, Edwards RA and Greenhalgh JFD. 1995. *Animal Nutrition*. 3rd Ed. Logman, London and New York.
- Miller Gail Lorenz. 1960. Use of Dinitrosalicylic acid reagent for determination of reducing sugar. *Anal Chem*; 31: 426-428
- Orskov ER and Ryle. 1990. *Energy Nutrition in Ruminant*. Elsevier Application. Jhon Willey and Sons, New York.
- Owens FN and Goestch AL. 1988. *Ruminal Fermentation* In: D.C. Church (Ed), *The Ruminant Animal Digestive Physiology and Nutrition*. A Reston Book Prentice Hall. Englewood Cliffs, New Jersey.
- Ottaway JH and Apps DK. 1984. *Biochemistry*. 4th Ed. Cambridge, ELBS.
- Puspaningsih NNT. 2004. Isolation and characterization of β -Xylosidase and α -L-Arabinofurosidase from *Bacillus thermoleovorans* IT-08. biotechnology of lignocellulose degradation and biomass utilization. Mie University. Publ Co, Ltd.
- Spolestra SF, Steg A and Beuvink JMW. 1990. Application of cell wall degrading enzymes to grass silage. In: *Agricultural biotechnology in focus in The Netherlands*. J.J. Dekkers, H.C. van der Plas and D.H. Vuijk (Eds). Pudoc Wageningen.
- Sunna A and Antranikian G. 1997. Xylanolytic enzymes from fungi and bacteria. *Graham G. Stewart and Inge Russel (Eds)*. *Crit Rev Biotech*; 17(1): 39-67